



City Research Online

## City, University of London Institutional Repository

---

**Citation:** Mutsatsa, S., Mushore, M., Ncube, K. & Currid, T. J. (2013). Vitamin D: the role of the sunshine vitamin. *British Journal of Mental Health Nursing*, 2(4), pp. 182-187. doi: 10.12968/bjmh.2013.2.4.182

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/12391/>

**Link to published version:** <https://doi.org/10.12968/bjmh.2013.2.4.182>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

---

---

City Research Online:

<http://openaccess.city.ac.uk/>

[publications@city.ac.uk](mailto:publications@city.ac.uk)

---

## **Vitamin D and mental health: The role of the sunshine vitamin**

S. Mutsatsa<sup>1</sup>, M Mushore<sup>1</sup>, K Ncube<sup>1</sup> & T Currid<sup>1</sup>

<sup>1</sup>London south Bank University, Gubbins Lane, Romford, RM3 0BE

Correspondence: [mutsatss@lsbu.ac.uk](mailto:mutsatss@lsbu.ac.uk)

# Vitamin D and Mental health: The role of the “sunshine vitamin”

## *Abstract*

The importance of vitamin D in physical disorders has been well documented. Deficiency in this vitamin is associated with a wide range of physical and mental disorders that include, heart problems, hypertension, stroke, diabetes, various cancers, and asthma. In mental health vitamin D deficiency is associated with schizophrenia, depression, anxiety, Alzheimer’s disease among others. Risk factors for vitamin D deficiency include darker skin, lack of adequate sun exposure, autoimmune diseases, influenza, old age and the use of certain medicines like anticonvulsants. Vitamin D is likely to play an important role in the management and prevention of various mental health problems. In particular, adequate vitamin D during the perinatal stage is likely to impact positively on the long term mental health of a person.

## *Introduction*

Until recently, the importance of Vitamin D, nicknamed “the sunshine vitamin,” has not been fully appreciated with its relative importance only focusing on adult bone health (Holick 2005). However, current estimates suggest that over a billion people have either vitamin D insufficiency or deficiency (Holick 2007). Scientific evidence is mounting implicating its importance in a vast array of physical and mental health problems. In physical health, vitamin D deficiency (VDD) is associated with type 1 diabetes (Mathieu *et al.* 2005), cardiovascular diseases (Baz-Hecht and Goldfine 2010), multiple sclerosis (VanAmerongen *et al.* 2004), sleep disturbance, various cancers including colon, breast, prostate and lung (Pearce and Cheetham 2010). It is also associated with asthma, muscular degeneration, auto immune disturbance, hyperparathyroidism, osteoporosis, cell growth modulation, neuromuscular and immune function. Conversely, adequate levels of vitamin D are associated with a reduction of inflammation (Tariq *et al.* 2011), good bone formation, increasing the activities (upregulates) of some gene enzymes like osteocalcin, osteopontin, calbindin 24-hydroxylase among others. In addition, the hormonally active form of vitamin D called calcitriol, reduces the activities of inflammatory markers like

interleukin 1(IL-1) and IL-2. Calcitriol also stimulates cell differentiation, influences muscular function, and stimulates insulin secretion (Tariq *et al.* 2011). Vitamin D also plays a role in mortality and this is clearly underscored by a recent meta- analysis of 14 eligible studies with 62548 participants showing that, VDD is associated with higher incidence of mortality (Zittermann *et al.* 2012). VDD also causes muscle weakness in children and the elderly with affected children having difficulty in standing and walking (Holick 2006) whereas the elderly have increasing sway and frequent falls (Bischoff-Ferrari *et al.* 2009). Clearly, the importance of vitamin D in physical health is self evident and this is true for mental health.

In mental health, VDD is associated with a number of neurodevelopmental and endocrine disorders (Humble, 2010). Ailments include autism, depression, anxiety, schizophrenia, Alzheimer's disease, premenstrual problems, thyroid regulation and sleep disorders among others. Because of the public health importance of Vitamin D, we carried out a selective review of evidence supporting the role of Vitamin D in some mental health disorders.

In particular, the article reviews evidence supporting the role of VDD in serious mental illness as schizophrenia and depression. The article starts by reviewing the epidemiology of VDD before discussing the mechanism underlying Vitamin D manufacture in the skin and its role in neurodevelopment, schizophrenia and depression. Lastly, the implications for nursing and health in general are discussed.

### *What is Vitamin D?*

Vitamin D is a steroid hormone (neurosteroid), meaning it comes from cholesterol precursors and its traditional role is enhancing the absorption and metabolism of calcium and phosphorous in the body. It affects the key biological functions of over 2000 genes in the body responsible for hormone balance, cell growth and immune function. There are at least five known different forms of vitamin D but only Vitamin D<sub>2</sub> (ergocalciferol) and Vitamin D<sub>3</sub> (cholecalciferol) are clinically relevant. As a steroid hormone, it is formed from a substance in the skin called 7 dehydrocholesterol (7DHC) after skin is exposure to ultra violet beta (UVB) rays.

Intuitively, it is appealing that people living in sunny climates have sufficient Vitamin D levels but evidence suggest otherwise. Studies carried out in sunny climates like Hawaii, Turkey, India, Iran and Saudi Arabia show that there is a high prevalence of VDD in people living in these countries. The deficiency is almost similar to those living in high latitudes countries (Binkley *et al.* 2007; Lips 2007; Elsammak *et al.* 2011). There is growing tendency for indoor activities by people living in these countries and this partly explains these findings (Fields *et al.* 2011). Moreover, since the 1980's the public has been warned against sunlight exposure to reduce the risk of developing skin cancer and this has played a contributory role in high VDD(Ness *et al.* 1999). In this regard, mental health patients in inpatient settings are at risk of VDD, therefore replenishing this important vitamin is necessary.

### *Risk factors for vitamin D deficiency*

In high latitudes of the Northern hemisphere countries, seasons strongly affect levels of vitamin D in the body. Several studies have found that in these countries, levels of vitamin D decrease during the winter months and increase during summer months (Rapuri *et al.* 2002). Wearing a sunscreen with a sun protection factor of 30 reduces vitamin D synthesis in the skin by more than 95%. (Matsuoka *et al.* 1987). Patients on a wide variety of medications, including anticonvulsants and medications to treat AIDS/HIV, are at risk because these drugs enhance the breakdown of vitamin D in the body (Zhou *et al.* 2006). In addition to the factors discussed, age is also a risk factor for VDD.

Several studies have found an association between old age and VDD and this is due to atrophic changes of their skin which decreases their capacity to produce adequate amounts of vitamin D (Hirani and Primatesta 2005). Moreover, the capacity for humans to absorb dietary vitamin D through the gut diminish with age (Huotari and Herzig 2008). A study that examined the levels of vitamin D in older people (average age 69 years) in Guatemala found that despite living in optimal climate for vitamin D, older Guatemalans had suboptimal levels of vitamin D (Sud *et al.* 2010). This finding is in line with a UK study found worryingly low levels of vitamin D in old people of 65 years or over (Hirani and Primatesta 2005). A further recent study found decreased Vitamin D level in older people in nursing homes (Verhoeven *et al.* 2012). By extrapolation, elderly people with mental health problems are equally at risk of VDD. VDD is also common in people with darker skin.

Consistent evidence exist clearly linking VDD and skin colour (Rockell *et al.* 2005; Rockell *et al.* 2008). Some studies in the USA have found that African-Americans are eight times more likely to be vitamin D deficient compared to age-matched Caucasians (Harris 2006; Bodnar *et al.* 2007) and this is supported by a more recent study a multiethnic sample of 503 adults (European, Maori, Pacific and Asian) in New Zealand. The study found that amongst these ethnic groups, Asians had the lowest mean Vitamin D levels and Europeans with lighter coloured skin had the highest (Nessvi *et al.* 2011). The main reason why darker skinned people are prone to VDD is darker skin (highly melanised) is less efficient at manufacturing vitamin D

than lighter skin. In addition to skin colour, obesity is also known to be a risk factor for VDD.

There is an association between inadequate Vitamin D levels and obesity (Alemzadeh *et al.* 2008; Sanchez-Hernandez *et al.* 2005; Renzaho *et al.* 2011; Brock *et al.* 2010). A relatively recent systematic review of 14 studies found an association between VDD and obesity related disorders (Renzaho *et al.* 2011) and another study found a relationship between VDD and BMI index of above thirty (Brock *et al.* 2010). It is also known that obesity is a common problem in people with mental health ailments, therefore, VDD is likely to be prevalent in this population (Phelan *et al.* 2001). More importantly, emerging evidence suggest a link between VDD and the aetiology of various mental health problems such as schizophrenia, depression, Alzheimer's disease and anxiety. Specifically, there is a strong link between vitamin D and neurodevelopmental ailments. This is particularly provocative as we now consider a number of psychiatric illnesses neurodevelopmental in origin. For this reason, it is important to review evidence that support the link between VDD and neurodevelopment.

#### *Vitamin D and neurodevelopment*

The discovery of vitamin D metabolites in the cerebrospinal fluid offered the first direct clue of its functional role in the nervous system (Balabanova *et al.* 1984). Evidence from animal models suggests that *in utero* VDD leads to dysregulation of cell differentiation in the developing neonate rat brain. Maternal VDD effect on neonate brain development persist into adulthood even if sufficient vitamin D levels are restored after birth (Feron *et al.* 2005). Further evidence supporting VDD in brain function is the discovery of Vitamin D receptors in the brain of both rodents (Stumpf and O'Brien 1987) and human brain (Eyles *et al.* 2005).

Vitamin D receptors (VDR) start to function as early as the 12<sup>th</sup> day of gestation and this coincides with the formation of dopamine neurons, a neurotransmitter implicated in psychiatric disorders such as schizophrenia and depression. VDR are involved in the differentiation of many brain areas throughout gestation (Cui *et al.* 2007). Furthermore, there is evidence suggesting VDR is involved in natural cell elimination (Ko *et al.*, 2004). In addition, Vitamin D modulates nerve growth factors (NGF)

essential for the growth and survival of many neurons in the brain. Specifically, the cholinergic basal forebrain neurons are affected (Brown *et al.* 2003). This finding is in line with an earlier animal study that found that injecting vitamin D in ventricles of the brain induces nerve growth factor expression in the hippocampus of adult rats (Saporito *et al.* 1993). These findings have implications for pregnant women in human where an estimated 40% to 80% of pregnant women are vitamin D deficient. This VDD which has ramifications for subsequent brain function is more pronounced in dark skinned ethnic populations living at high latitudes countries. In combination with other factors, VDD during peri-natal life may establish a poor foundation that may produce long-term threats to human health that include risk for developing schizophrenia.

### *Schizophrenia and vitamin D*

We now accept that schizophrenia is a heterogeneous disorder in which genetic and environmental factors interact to contribute to the person's risk for developing the disorder (Eyles *et al.* 2009). To date, evidence linking VDD and schizophrenia is the tendency for people with schizophrenia to be born in the winter months or in higher latitudes countries (Torrey *et al.* 1997; Davies *et al.* 2003). In a review of 250 studies covering 29 Northern and five Southern Hemisphere countries, Torrey *et al.* (1997) found an excess of people with schizophrenia and depression to be born in winter/spring months. This excess was not seen in other mental health disorders. The investigators concluded that, statistical artefact and parental procreational habits are insufficient explanations for this excess, which was as high 8%. In darker skinned individuals living in high latitudes countries, VDD assumes a greater importance as there is a high and disproportionate number of people with schizophrenia in this population (Fearon *et al.* 2006).

More specifically, the relative risk of developing schizophrenia is higher in second than first generation darker skinned migrants who move to high latitude climates (Cantor-Graae and Selten 2005; Dealberto 2010). Because their vulnerability for VDD, darker skinned individuals increase the likelihood of VDD in their offsprings. This hypothesis is supported by several studies including one study that examined maternal VDD as a risk factor for schizophrenia in offsprings using banked maternal sera (McGrath *et al.* 2003). The study found that the offspring of

darker skinned mothers were more likely to develop schizophrenia if the mother had significantly low vitamin D levels especially in the third trimester of pregnancy. By contrast, the chances of the offspring developing schizophrenia were significantly reduced if their mothers had normal vitamin D level during the same period of pregnancy (McGrath et al 2003).

A study that provided direct evidence of the relationship between VDD and the risk of developing schizophrenia is the Northern Finnish cohort study (Ko *et al.* 2004). The study found a reduced risk of developing schizophrenia in male offspring who received adequate vitamin D supplementation than males who did not in their first year of life. This finding is supported by a relatively recent study that examined the link between neonatal vitamin D status and risk of schizophrenia (McGrath *et al.* 2010). The study assessed the concentration of Vitamin D from neonatal dried blood samples of individuals with schizophrenia and controls. Low or very high levels of vitamin D were associated with a risk for developing schizophrenia, suggesting a non linear relationship. Overall, VDD appear to have a link in mental health disorders like schizophrenia and depression.

### *Depression*

Evidence supporting the role of vitamin D in depression comes from three main sources namely, animal models, epidemiological and, randomised controlled studies. In animal models, mice lacking vitamin D receptor functioning show behavioural impairment that include, memory, perseverative responses and hypolocomotion and these behaviours are seen in humans with depression. Rodents with VDD show dysfunctional noradrenalin gene expression. In humans, noradrenaline is implicated in depression therefore lending indirect evidence for the role of vitamin D in depression. Animal studies also suggest a role for vitamin D in neuroprotection against the effects of dopamine toxins such as methamphetamine. Like noradrenaline, dopamine is implicated in depression (Cass *et al.* 2006). Cass et al (2006) exposed rats to the dopamine toxin, methamphetamine and the rats experienced a significant decrease in serotonin and dopamine concentration in the brain but rats treated with vitamin D did not demonstrate these effects. Cass et al(2006) study is compatible with an earlier finding that showed an increase in dopamine in the cortical region of the brain in rats that were fed a vitamin D replete diet compared to rats on VDD diet (Baksi and Hughes 1982). In people with

depression, both dopamine and serotonin levels are depleted. Taken together, animal studies support the vitamin D hypothesis of depression. In this regard, several epidemiological studies also lend credence to this hypothesis (Hoogendijk *et al.* 2008; Jorde *et al.* 2006; Ganji *et al.* 2010).

Ganji *et al.* (2010) found that those with low Vitamin D blood serum levels were nearly twice as likely to develop depression than those who had higher levels. By far, the strongest evidence supporting a causal link between depression and VDD comes from randomised controlled trials (Jorde *et al.* 2008). In a double blind randomised controlled trial of 441 participants, Jorde *et al.* (2008) compared the outcome of depression in three groups (two groups taking vitamin D of different strengths and one taking placebo). At one year, they found that those with low levels were significantly more depressed than those with adequate levels of vitamin D. Overall, the groups receiving vitamin D supplementation had improved depressive symptoms than those who received placebo. This finding among others has considerable implications for future public health and mental health nursing in particular.

#### *Implication for clinical practice*

Studies have demonstrated a clear link between VDD and mental health and its role as a therapeutic agent has attracted considerable interest. From a public health perspective, the use of vitamin D has the potential to alter health care in general and mental health nursing in particular. Vitamin D levels can be increased using inexpensive and well tolerated dietary supplements. In physical health, studies have demonstrated modest improvement in people who take vitamin D supplements towards health. Evidence suggest that taking vitamin D by older people has positive multiple outcomes including reduced falls, higher bone mineral density, and reduced fractures. There is potential therefore for a pivotal role in vitamin D supplementation as part of routine nursing care.

In physical health disorders such as diabetes, the use of vitamin D supplements again shows promising results (von Hurst *et al.* 2010). In a randomised controlled trial that sought to improve the vitamin D status of women who were insulin resistant, Von Hurst *et al.* (2010) found significant improvement in insulin sensitivity and a reduction in fasting insulin in the vitamin D supplementation group compared to

placebo. This finding is likely to find therapeutic application in mental health where a significant proportion of people also have type 2 diabetes.

In depression, this vitamin has the potential to play a useful therapeutic role as demonstrated by a several studies that recorded an improvement in depressive symptoms after vitamin D supplementation. (Hogberg *et al.* 2012; Bertone-Johnson *et al.* 2011). A Norwegian trial of overweight subjects showed that those receiving a high dose of vitamin D (20,000 or 40,000 IU weekly) had a significant improvement in depressive symptoms after 1 year versus those receiving placebo (Jorde *et al.* 2008). From a practical point of view, it is likely that in future, nurses will encourage patients to be exposed to light as part of a treatment plan. It is purported that a light exposure of between 20 to 30 minutes a day is sufficient for the skin to synthesise adequate vitamin. Moreover light therapy for non seasonal affective disorder shows promising results(Tuunainen *et al.* 2004). With regard to schizophrenia, there are currently several randomised controlled trials in process, testing the effect of vitamin D on numerous outcome measures including symptoms and cognition. Clearly, these prospects have the potential to modify nursing practice in future and this may include care for the pregnant.

During pregnancy, the foetus is entirely dependent on the mother for its supply of vitamin D. therefore maternal vitamin D supplementation during pregnancy is necessary and nurses have an important role to play in this regard. Relatively recent Cochrane review has provided evidence on the positive impact of vitamin D supplementation (De-Regil *et al.* 2012). Overall, it is likely that in future vitamin D may play a significant role in public mental health through primary preventative care of neurodevelopmental disorders like schizophrenia.

### *Conclusion*

There is ample biological evidence suggesting the importance of vitamin D in the neurodevelopmental process, physical health and mental health in particular. Current mental health policy emphasises an intergrated approach of physical and mental health care and, this has wider implications for future mental health nursing. Based on the evidence presented in this paper, Vitamin D is likely to have an important role in routine nursing care. From a preventative vantage point, the peri-natal stage of development and care is critical. Good peri-natal care is known to have a positive

and lasting impact on good physical and mental health (Huang 2011). It is likely that nutritional care as a means to promoting good mental health care is likely to be integral in future nursing interventions. In this regard mental health nurses ought to familiarise themselves more with multifunctional role of this important hormone.

#### Reference List

Alemzadeh R, Kichler J, Babar G, Calhoun M (2008) Hypovitaminosis D in obese children and adolescents: relationship with adiposity, insulin sensitivity, ethnicity, and season. *Metabolism* **57**, 183-191.

Baksi SN, Hughes MJ (1982) Chronic vitamin D deficiency in the weanling rat alters catecholamine metabolism in the cortex. *Brain Res* **242**, 387-390.

Balabanova S, Richter HP, Antoniadis G, Homoki J, Kremmer N, Hanle J, Teller WM (1984) 25-Hydroxyvitamin D, 24, 25-dihydroxyvitamin D and 1,25-dihydroxyvitamin D in human cerebrospinal fluid. *Klin.Wochenschr.* **62**, 1086-1090.

Baz-Hecht M, Goldfine AB (2010) The impact of vitamin D deficiency on diabetes and cardiovascular risk. *Curr.Opin.Endocrinol.Diabetes Obes.* **17**, 113-119.

Bertone-Johnson ER, Powers SI, Spangler L, Brunner RL, Michael YL, Larson JC, Millen AE, Bueche MN, Salmoirago-Blotcher E, Liu S, Wassertheil-Smoller S, Ockene JK, Ockene I, Manson JE (2011) Vitamin D intake from foods and supplements and depressive symptoms in a diverse population of older women. *Am.J Clin.Nutr.* **94**, 1104-1112.

Binkley N, Novotny R, Krueger D, Kawahara T, Daida YG, Lensmeyer G, Hollis BW, Drezner MK (2007) Low vitamin D status despite abundant sun exposure. *J Clin.Endocrinol.Metab* **92**, 2130-2135.

Bischoff-Ferrari HA, Dawson-Hughes B, Staehelin HB, Orav JE, Stuck AE, Theiler R, Wong JB, Egli A, Kiel DP, Henschkowski J (2009) Fall prevention with supplemental and active forms of vitamin D: a meta-analysis of randomised controlled trials. *BMJ* **339**, b3692.

Bodnar LM, Simhan HN, Powers RW, Frank MP, Cooperstein E, Roberts JM (2007) High prevalence of vitamin D insufficiency in black and white pregnant women residing in the northern United States and their neonates. *J Nutr.* **137**, 447-452.

Brock K, Huang WY, Fraser DR, Ke L, Tseng M, Stolzenberg-Solomon R, Peters U, Ahn J, Purdue M, Mason RS, McCarty C, Ziegler RG, Graubard B (2010) Low vitamin D status is associated with physical inactivity, obesity and low vitamin D intake in a large US sample of healthy middle-aged men and women. *J Steroid Biochem.Mol.Biol.* **121**, 462-466.

Brown J, Bianco JI, McGrath JJ, Eyles DW (2003) 1,25-dihydroxyvitamin D<sub>3</sub> induces nerve growth factor, promotes neurite outgrowth and inhibits mitosis in embryonic rat hippocampal neurons. *Neurosci.Lett.* **343**, 139-143.

- Cantor-Graae E, Selten JP (2005) Schizophrenia and migration: a meta-analysis and review. *Am.J Psychiatry* **162**, 12-24.
- Cass WA, Smith MP, Peters LE (2006) Calcitriol protects against the dopamine- and serotonin-depleting effects of neurotoxic doses of methamphetamine. *Ann.N.Y.Acad.Sci.* **1074**, 261-271.
- Cui X, McGrath JJ, Burne TH, Mackay-Sim A, Eyles DW (2007) Maternal vitamin D depletion alters neurogenesis in the developing rat brain. *Int.J Dev.Neurosci.* **25**, 227-232.
- Davies G, Welham J, Chant D, Torrey EF, McGrath J (2003) A systematic review and meta-analysis of Northern Hemisphere season of birth studies in schizophrenia. *Schizophr.Bull.* **29**, 587-593.
- De-Regil LM, Palacios C, Ansary A, Kulier R, Pena-Rosas JP (2012) Vitamin D supplementation for women during pregnancy. *Cochrane.Database.Syst.Rev.* **2**, CD008873.
- Dealberto MJ (2010) Ethnic origin and increased risk for schizophrenia in immigrants to countries of recent and longstanding immigration. *Acta Psychiatr.Scand.* **121**, 325-339.
- Elsammak MY, Al-Wossaibi AA, Al-Howeish A, Alsaeed J (2011) High prevalence of vitamin D deficiency in the sunny Eastern region of Saudi Arabia: a hospital-based study. *East Mediterr.Health J* **17**, 317-322.
- Eyles DW, Feron F, Cui X, Kesby JP, Harms LH, Ko P, McGrath JJ, Burne TH (2009) Developmental vitamin D deficiency causes abnormal brain development. *Psychoneuroendocrinology* **34 Suppl 1**, S247-S257.
- Eyles DW, Smith S, Kinobe R, Hewison M, McGrath JJ (2005) Distribution of the vitamin D receptor and 1 alpha-hydroxylase in human brain. *J Chem.Neuroanat.* **29**, 21-30.
- Fearon P, Kirkbride JB, Morgan C, Dazzan P, Morgan K, Lloyd T, Hutchinson G, Tarrant J, Fung WL, Holloway J, Mallett R, Harrison G, Leff J, Jones PB, Murray RM (2006) Incidence of schizophrenia and other psychoses in ethnic minority groups: results from the MRC AESOP Study 5. *Psychol.Med* **36**, 1541-1550.
- Feron F, Burne TH, Brown J, Smith E, McGrath JJ, Mackay-Sim A, Eyles DW (2005) Developmental Vitamin D3 deficiency alters the adult rat brain. *Brain Res Bull.* **65**, 141-148.
- Fields J, Trivedi NJ, Horton E, Mechanick JI (2011) Vitamin D in the Persian Gulf: integrative physiology and socioeconomic factors. *Curr.Osteoporos.Rep.* **9**, 243-250.
- Ganji V, Milone C, Cody MM, McCarty F, Wang YT (2010) Serum vitamin D concentrations are related to depression in young adult US population: the Third National Health and Nutrition Examination Survey. *Int.Arch.Med* **3**, 29.
- Harris SS (2006) Vitamin D and African Americans. *J Nutr.* **136**, 1126-1129.
- Hirani V, Primatesta P (2005) Vitamin D concentrations among people aged 65 years and over living in private households and institutions in England: population survey. *Age Ageing* **34**, 485-491.
- Hogberg G, Gustafsson SA, Hallstrom T, Gustafsson T, Klawitter B, Petersson M (2012) Depressed adolescents in a case-series were low in vitamin D and depression was ameliorated by vitamin D supplementation. *Acta Paediatr.*

- Holick MF (2005) The vitamin D epidemic and its health consequences. *J Nutr.* **135**, 2739S-2748S.
- Holick MF (2006) High prevalence of vitamin D inadequacy and implications for health. *Mayo Clin.Proc.* **81**, 353-373.
- Holick MF (2007) Vitamin D deficiency. *N.Engl.J Med* **357**, 266-281.
- Hoogendijk WJ, Lips P, Dik MG, Deeg DJ, Beekman AT, Penninx BW (2008) Depression is associated with decreased 25-hydroxyvitamin D and increased parathyroid hormone levels in older adults. *Arch.Gen.Psychiatry* **65**, 508-512.
- Huang LT (2011) The link between perinatal glucocorticoids exposure and psychiatric disorders. *Pediatr.Res* **69**, 19R-25R.
- Huotari A, Herzig KH (2008) Vitamin D and living in northern latitudes--an endemic risk area for vitamin D deficiency. *Int.J Circumpolar.Health* **67**, 164-178.
- Jorde R, Sneve M, Figenschau Y, Svartberg J, Waterloo K (2008) Effects of vitamin D supplementation on symptoms of depression in overweight and obese subjects: randomized double blind trial. *J Intern.Med* **264**, 599-609.
- Jorde R, Waterloo K, Saleh F, Haug E, Svartberg J (2006) Neuropsychological function in relation to serum parathyroid hormone and serum 25-hydroxyvitamin D levels. The Tromso study. *J Neurol.* **253**, 464-470.
- Ko P, Burkert R, McGrath J, Eyles D (2004) Maternal vitamin D3 deprivation and the regulation of apoptosis and cell cycle during rat brain development. *Brain Res Dev.Brain Res* **153**, 61-68.
- Lips P (2007) Vitamin D status and nutrition in Europe and Asia. *J Steroid Biochem.Mol.Biol.* **103**, 620-625.
- Mathieu C, Gysemans C, Giulietti A, Bouillon R (2005) Vitamin D and diabetes. *Diabetologia* **48**, 1247-1257.
- Matsuoka LY, Ide L, Wortsman J, MacLaughlin JA, Holick MF (1987) Sunscreens suppress cutaneous vitamin D3 synthesis. *J Clin.Endocrinol.Metab* **64**, 1165-1168.
- McGrath J, Eyles D, Mowry B, Yolken R, Buka S (2003) Low maternal vitamin D as a risk factor for schizophrenia: a pilot study using banked sera. *Schizophr.Res* **63**, 73-78.
- McGrath JJ, Eyles DW, Pedersen CB, Anderson C, Ko P, Burne TH, Norgaard-Pedersen B, Hougaard DM, Mortensen PB (2010) Neonatal vitamin D status and risk of schizophrenia: a population-based case-control study. *Arch.Gen.Psychiatry* **67**, 889-894.
- Ness AR, Frankel SJ, Gunnell DJ, Smith GD (1999) Are we really dying for a tan? *BMJ* **319**, 114-116.
- Nessvi S, Johansson L, Jopson J, Stewart A, Reeder A, McKenzie R, Scragg RK (2011) Association of 25-hydroxyvitamin D3]levels in adult New Zealanders with ethnicity, skin color and self-reported skin sensitivity to sun exposure. *Photochem.Photobiol.* **87**, 1173-1178.
- Pearce SH, Cheetham TD (2010) Diagnosis and management of vitamin D deficiency. *BMJ* **340**, b5664.

Phelan M, Stradins L, Morrison S (2001) Physical health of people with severe mental illness. *BMJ* **322**, 443-444.

Rapuri PB, Kinyamu HK, Gallagher JC, Haynatzka V (2002) Seasonal changes in calciotropic hormones, bone markers, and bone mineral density in elderly women. *J Clin.Endocrinol.Metab* **87**, 2024-2032.

Renzaho AM, Halliday JA, Nowson C (2011) Vitamin D, obesity, and obesity-related chronic disease among ethnic minorities: a systematic review  
3. *Nutrition* **27**, 868-879.

Rockell JE, Green TJ, Skeaff CM, Whiting SJ, Taylor RW, Williams SM, Parnell WR, Scragg R, Wilson N, Schaaf D, Fitzgerald ED, Wohlers MW (2005) Season and ethnicity are determinants of serum 25-hydroxyvitamin D concentrations in New Zealand children aged 5-14 y. *J Nutr.* **135**, 2602-2608.

Rockell JE, Skeaff CM, Williams SM, Green TJ (2008) Association between quantitative measures of skin color and plasma 25-hydroxyvitamin D. *Osteoporos.Int.* **19**, 1639-1642.

Sanchez-Hernandez J, Ybarra J, Gich I, De LA, Rius X, Rodriguez-Espinosa J, Perez A (2005) Effects of bariatric surgery on vitamin D status and secondary hyperparathyroidism: a prospective study. *Obes.Surg.* **15**, 1389-1395.

Saporito MS, Wilcox HM, Hartpence KC, Lewis ME, Vaught JL, Carswell S (1993) Pharmacological induction of nerve growth factor mRNA in adult rat brain. *Exp.Neurol.* **123**, 295-302.

Stumpf WE, O'Brien LP (1987) 1,25 (OH)<sub>2</sub> vitamin D<sub>3</sub> sites of action in the brain. An autoradiographic study. *Histochemistry* **87**, 393-406.

Sud SR, Montenegro-Bethancourt G, Bermudez OI, Heaney RP, Armas L, Solomons NW (2010) Older Mayan residents of the western highlands of Guatemala lack sufficient levels of vitamin D. *Nutr.Res* **30**, 739-746.

Tariq MM, Streeten EA, Smith HA, Sleemi A, Khabazghazvini B, Vaswani D, Postolache TT (2011) Vitamin D: a potential role in reducing suicide risk? *Int.J Adolesc.Med Health* **23**, 157-165.

Torrey EF, Miller J, Rawlings R, Yolken RH (1997) Seasonality of births in schizophrenia and bipolar disorder: a review of the literature. *Schizophr.Res* **28**, 1-38.

Tuunainen A, Kripke DF, Endo T (2004) Light therapy for non-seasonal depression  
2. *Cochrane.Database.Syst.Rev.* CD004050.

VanAmerongen BM, Dijkstra CD, Lips P, Polman CH (2004) Multiple sclerosis and vitamin D: an update. *Eur.J Clin.Nutr.* **58**, 1095-1109.

Verhoeven V, Vanpuyenbroeck K, Lopez-Hartmann M, Wens J, Remmen R (2012) Walk on the sunny side of life--epidemiology of hypovitaminosis D and mental health in elderly nursing home residents. *J Nutr.Health Aging* **16**, 417-420.

von Hurst PR, Stonehouse W, Coad J (2010) Vitamin D supplementation reduces insulin resistance in South Asian women living in New Zealand who are insulin resistant and vitamin D deficient - a randomised, placebo-controlled trial. *Br.J Nutr.* **103**, 549-555.

Zhou C, Assem M, Tay JC, Watkins PB, Blumberg B, Schuetz EG, Thummel KE (2006) Steroid and xenobiotic receptor and vitamin D receptor crosstalk mediates CYP24 expression and drug-induced osteomalacia. *J Clin. Invest* **116**, 1703-1712.

Zittermann A, Iodice S, Pilz S, Grant WB, Bagnardi V, Gandini S (2012) Vitamin D deficiency and mortality risk in the general population: a meta-analysis of prospective cohort studies. *Am.J Clin.Nutr.* **95**, 91-100.