



City Research Online

City St George's, University of London

Citation: Mulligan, K., McBain, H. B., Lamontagne-Godwin, F., Chapman, J., Flood, C., Haddad, M., Jones, J. & Simpson, A. (2018). A survey of factors that influence diabetes self-management in people with severe mental illness. *Diabetic Medicine*, 35(S1), p. 134. doi: 10.1111/dme.33_13571

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/19430/>

Link to published version: https://doi.org/10.1111/dme.33_13571

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

A survey of factors that influence diabetes self-management in people with severe mental illness.

Dr Kathleen Mulligan^{1,5}, Dr Hayley McBain¹, Frederique Lamontagne-Godwin², Jacqui Chapman³, Dr Chris Flood¹, Dr Mark Haddad¹, Dr Julia Jones⁴, Professor Alan Simpson^{1,5}.

¹ School of Health Sciences, City University of London

² College of Nursing, Midwifery and Healthcare, University of West London

³ Diabetes Centre, Homerton University Hospital NHS Foundation Trust

⁴ Centre for Primary & Community Health, University of Hertfordshire

⁵ East London NHS Foundation Trust

Abstract

Aim

To identify barriers and enablers to diabetes self-management experienced by people with SMI.

Methods

Adults with type 2 diabetes and SMI completed a survey that asked about their: diabetes self-management behaviours; diabetes healthcare received; psychological distress; barriers to and enablers of diabetes self-management.

Results

Seventy-seven people completed the survey; most had either bipolar disorder (42%) or schizophrenia (36%). Most participants (67.5%) took tablets to treat their diabetes. The number of days in the previous week that participants engaged in diabetes self-management behaviours was highest for medication (mean 6.1 (S.D. 2.0)) but lower for general diet (4.1 (2.3)), specific diet (3.6 (1.8)), blood glucose testing (3.8 (2.7)), exercise (2.4 (2.1)) and foot care (1.7 (1.8)). 44% of participants were smokers. Participants received a mean (S.D.) of 7.6 (3.0) of 14 diabetes healthcare essentials. At least moderate psychological distress was reported by 56% of participants.

In multivariate analyses, the significant ($p < 0.05$) explanatory variables for each diabetes self-management behaviour were:

Blood glucose testing – level of psychological distress; perceived capability to establish a routine to manage diabetes (behavioural regulation)

General Diet – goal priority; beliefs about ability to manage diabetes, beliefs about consequences of diabetes management

Specific diet – goal priority; behavioural regulation

Exercise – goal priority; availability of services

Foot care – emotion, diabetes treatment

Conclusions

Several aspects of diabetes healthcare and self-management were found to be suboptimal. This research identified factors that could be targeted to help support diabetes self-management in people with SMI.