



City Research Online

City St George's, University of London

Citation: Suttle, C. M., Barbur, J. L. & Conway, M. L. (2018). Author's reply. *Ophthalmic And Physiological Optics*, 38(4), 469. doi: 10.1111/opo.12569

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/20043/>

Link to published version: <https://doi.org/10.1111/opo.12569>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Authors' reply

We thank Dr Aleci for consideration of issues discussed in our paper¹ in which we reported that more than half of our sample of 21 adults and children with symptoms of visual stress chose different coloured overlays on two different occasions. We concluded that repeatability of colour for the relief of visual stress symptoms is poor.

We categorised consistency of coloured overlay choice as the Same (identical overlay colour chosen on two occasions), Similar (one of two overlay colours is identical on both occasions) or Different (different overlay colour chosen). Dr Aleci notes that almost all of the subjects in the Same or Similar categories chose overlays with maximum transmission on the left side of the wavelength spectrum (blue, aqua, purple or green), while most of those in the Different category chose overlays with transmission on the right side of the spectrum (yellow, orange, pink or rose). We will refer to these colour ranges here as short and long wavelength colours respectively.

Based on this observation and on the theories that visual stress symptoms are related to cortical hyperexcitation or a magnocellular deficit, Dr Aleci proposes that high repeatability may be found in people with visual stress who find short wavelength filters beneficial, because these may help to balance cortical excitation by modifying magnocellular and parvocellular responses.

It is not yet well established that coloured filters reduce cortical hyperexcitation in people with symptoms of visual stress. For example, reduced visual cortical activation has been found using functional magnetic resonance imaging (fMRI) in a small group of patients with migraine,² a condition in which visual stress symptoms often occur. However, in another small fMRI study³ cortical activation was *increased* in people with visual stress viewing through their chosen colour. Similarly, clear evidence for a magnocellular deficit in people with visual stress is lacking.⁴ Larger studies demonstrating clearly that coloured filters correct a neurophysiological imbalance in uncomplicated cases of visual stress (without dyslexia or migraine) would be needed to suggest that short wavelength colours alleviate symptoms in those people by addressing that imbalance.

A further consideration is that of colour preference. Previous work has shown a general preference for short wavelength colours among young Caucasian adults⁵ so the choice of blue, aqua and green coloured overlay choice by most of the participants in our study may have been influenced to some extent by preference.

References

1. Suttle CM, Barbur J, Conway ML. Coloured overlays and precision-tinted lenses: poor repeatability in a sample of adults and children diagnosed with visual stress. *Ophthalmic Physiol Opt* 2017; 37:542-548.
2. Huang J, Zong X, Wilkins A, et al fMRI evidence that precision ophthalmic tints reduce cortical hyperactivation in migraine. *Cephalgia*. 2011; 31(8): 925-936.
3. Kim JH, Seo H, Ha S et al Functional magnetic resonance imaging findings in Meares-Irlen syndrome: a pilot study. *K J Ophthalmol* 2015; 29: 121-125.
4. Conlon EG, Sanders MA, Wright CM Relationships between global motion and global form processing, practice, cognitive and visual processing in adults with dyslexia or visual discomfort. *Neuropsychologia* 2009; 47:907-915.
5. Hurlbert AC, Ling Y. Biological components of sex differences in color preference. *Curr Biol* 2007; 17:R623-5.