



City Research Online

City St George's, University of London

Citation: Anagnostopoulou, N., Alba, A., Gaete, J., Dima, D. & Kyriakopoulos, M. (2018). Prevalence of psychotic symptoms in children needing inpatient care: Relationship to diagnosis. *European Psychiatry*, 48, S521-S521. doi: 10.1016/j.eurpsy.2017.12.023

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/22499/>

Link to published version: <https://doi.org/10.1016/j.eurpsy.2017.12.023>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Prevalence of psychotic symptoms in children needing inpatient care: relationship to diagnosis

Anagnostopoulou N, Alba A, Gaete J, Dima D, Kyriakopoulos M

Background: Psychotic symptoms, marking impaired reality testing, are considered a hallmark of psychotic disorders. However, several lines of research point towards these symptoms also being highly prevalent both in non-clinical populations and in other psychiatric disorders. There is also evidence to suggest that their prevalence is even higher in children and young people, reaching up to 17% in epidemiological samples. There is limited research on the prevalence of psychotic symptoms in young clinical populations. The present study aimed to assess the prevalence of psychotic symptoms in a 6 to 12-year-old clinical population of an inpatient setting. **Method:** Clinical records of all patients of a national UK children's unit from 2009 to 2017 were examined for the presence of psychotic symptoms and their relationship to diagnosis at discharge. **Results:** One hundred and eighty-five (185) children were included in the study. Psychotic symptoms were highly prevalent in this young clinical population, reaching 63.8%, regardless of diagnosis at discharge. Hallucinations (58.4%) were more prevalent than delusions (35.1%). As expected, all children with schizophrenia spectrum disorders experienced psychotic symptoms. **Conclusions:** The present study is the first to examine the prevalence of delusions and hallucinations across diagnoses in children needing inpatient care. Psychotic symptoms were very common in children without a diagnosis of schizophrenia spectrum disorder. Given the high frequency of psychotic symptoms in this severely affected clinical population, it is possible that they represent a marker of clinical severity.

Table 1. Psychotic Symptoms Percentages among Diagnoses.

DIAGNOSIS	TOTAL
EARLY-ONSET SCHIZOPHRENIA	100%
INTELLECTUAL DISABILITY	86.7%
ANXIETY	67.2%
AUTISM SPECTRUM DISORDER	65.7%

<i>DEPRESSION</i>	65.5%
<i>HYPERKINETIC DISORDER</i>	65.4%
<i>EATING DISORDER</i>	60%
<i>OBSESSIVE COMPULSIVE DISORDER</i>	53.6%