



City Research Online

City, University of London Institutional Repository

Citation: Whittington, R., Bowers, L., Nolan, P., Simpson, A. & Neil, L. (2009). Approval Ratings of Inpatient Coercive Interventions in a National Sample of Mental Health Service Users and Staff in England. *Psychiatric Services*, 60(6), pp. 792-798. doi: 10.1176/ps.2009.60.6.792

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/7240/>

Link to published version: <https://doi.org/10.1176/ps.2009.60.6.792>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

1
2
3 **APPROVAL FOR DIFFERENT TYPES OF COERCIVE INTERVENTIONS**
4
5
6 **AMONGST A NATIONAL SAMPLE OF**
7
8 **MENTAL HEALTH SERVICE USERS AND STAFF IN ENGLAND**
9

10
11
12
13
14 Running head: Attitudes Toward Coercive Interventions
15
16

17 **Disclosures of interest:** None for any author.
18
19

20 **Acknowledgements:** this study was funded by the Service Delivery and Organisation
21 Research and Development Programme, Department of Health, England.
22
23

24 Revised version (in response to comments dated 20th. August 2008)
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Objective: To ascertain the degree of approval amongst service users and staff for various coercive measures commonly used in acute mental health care.

Methods: A cross-sectional design was adopted. The Attitudes to Containment Measures Questionnaire (ACMQ) was completed by 1,361 service users and 1,226 staff in acute care mental health services from three regions of England. This provided evaluation of eleven coercive measures (e.g. seclusion) on six dimensions of approval (e.g. indignity, safety) in a large national sample. Comparisons between groups were tested using independent samples t-tests, χ^2 or Spearman correlations.

Results: Both service users and staff disapproved strongly of mechanical restraint and expressed a relative preference for compulsory intramuscular medication and seclusion. Male staff, older service users and staff who had been involved in implementing coercion expressed greater approval of coercive measures.

Conclusion: Mechanical restraint remains highly objectionable to staff and service users in English mental health services despite its widespread acceptance elsewhere in the world.

Keywords: violence, coercion, attitude of health personnel, physical restraint, intramuscular injection.

Introduction

The aim of this study was to systematically ascertain the views of English service users and staff about the use of coercive measures in acute mental health care. It is widely accepted that effective mental health care sometimes includes the deployment of coercive measures such as seclusion and restraint by staff to contain dangerous or severely disruptive behaviour by a service user. Extensive efforts have been made in Europe and America over the past decade to assess rates of coercive measures systematically [1-3] and to reduce them [4]. D’Orio et al. [5], for example, report a 39% reduction in coercion following introduction of a package which included enhanced access to expertise during emergency situations. Involvement in the use of these measures or witnessing their use can be highly distressing for both service users and staff [6-8] and few are likely to remain neutral about them. Attitudes toward different types of coercive measure are likely to vary between and within service user and staff groups but little is known about the preferences of staff and service users when comparing different measures. Such preferences are likely to have some influence on the decision to deploy them and thus are worthy of the systematic investigation reported in this paper. Much research on subjective perceptions in this area has been qualitative and focused on emotional responses to the experience of coercion [9]. Qualitative studies conducted beyond the UK and American setting indicate that many of the responses of service users and staff are recognisable across cultures e.g. fear and anxiety [10, 11]. More structured approaches are increasingly being developed to survey staff and service user attitudes [12, 13] and have revealed intriguing patterns which may vary across cultures, e.g. some positive evaluations of

1
2
3 coercive measures amongst Chinese service users [14] However these instruments do
4
5 not enable the direct comparison between different types of coercive measure which
6
7 was the aim of this study. Two American studies do allow such comparisons and
8
9 found that staff in high secure care [15] and neuropsychiatric care [16] ranked the
10
11 following measures in the same order of preference: medication, seclusion and
12
13 mechanical restraint. No distinction was drawn in these studies however between
14
15 consensual PRN medication and coerced IM medication when these are clearly two
16
17 very different scenarios. A Canadian study [17] compared staff and service user
18
19 preferences in a small, purposive sample and found a number of patterns e.g. that
20
21 service users approved of PRN medication. much more than staff. However, the
22
23 questionnaire used presented the methods hierarchically, making interpretation of
24
25 relative preferences difficult. Bowers et al. [18] report the development of a new tool,
26
27 the Attitudes to Containment Measures Questionnaire (ACMQ) which enables direct
28
29 comparison between coercive measures. Coerced PRN medication is explicitly
30
31 identified to distinguish it from consensual medication in this instrument. Preliminary
32
33 data from student nurses in four countries suggest that English students had greater
34
35 reservations about coercive measures than their counterparts in the Netherlands and
36
37 Finland [19].
38
39
40
41
42
43
44
45
46
47

48 The aim of the study reported below was to move beyond previous research by
49
50 eliciting service user and staff preferences and approval for various coercive measures
51
52 when they were directly compared with each other using the ACMQ in a large
53
54 national sample. Whilst this country, England, markedly differs from North America
55
56 and the rest of continental Europe in at least one respect (intolerance of mechanical
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

restraint [3]) the aim was ultimately to provide a benchmark against which other national samples could be compared. A cross-sectional survey design was adopted.

For Review Only

Methods

Setting and sample

Staff and service user respondents were drawn from the 136 acute wards participating in the City 128 study [20]. The intention was to recruit 10 service users and all staff from each ward. Potential service user participants on each ward were identified by random sampling but, once identified, only those judged by staff as able to grant informed consent and participate were approached. After complete description of the study to the participants, written informed consent was obtained. Most service users were interviewed by a research assistant to aid completion of the ACMQ instrument (see below). All staff on each ward were sent a copy of the ACMQ instrument and those who completed it, returned it anonymously through an internal mailbox. The final sample consisted of 1,226 staff and 1,361 service users (see Table 1). 95% of staff respondents were nurses (68%) or health care assistants (27%) with the rest being from other occupations (occupational therapist, psychiatrist, psychologist, social worker). Data were collected in 2004 and 2005.

Insert Table 1 about here

Measure

The Attitudes to Containment Measures Questionnaire (ACMQ) [18] lists 11 coercive measures used widely either in the UK (PRN medication, compulsory IM medication; physical restraint; intermittent observation; constant observation; time out; PICU

1
2
3 transfer; locked door seclusion; open area seclusion) or elsewhere in Europe
4
5 (mechanical restraint or net bed, defined as a lockable metal frame with side netting
6
7 bolted to a bed [3, 21]). Each listed coercive measure is accompanied by a short
8
9 description and a visual illustration and then six dimensions of approval are assessed:
10
11 *effectiveness, acceptability, respectfulness, safety for service users, safety for staff,*
12
13 *willingness to undergo (service users) or use (staff).* The respondent is asked to
14
15 indicate their degree of approval on a 5-point Likert scale (strongly agree =5, to
16
17 strongly disagree=1) and then to indicate (yes/no) whether they have been involved in
18
19 implementing the measure (staff) or subjected to it (service users). Responses were
20
21 summed across approval ratings for each coercive measure and a high score indicates
22
23 approval as opposed to disapproval. Comparisons between groups were tested using
24
25 independent samples t-tests, χ^2 or Spearman correlations.
26
27
28
29
30
31
32
33

34 Ethical review

35
36 The study was approved by the National Health Service North West Multicentre
37
38 Research Ethics Committee.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

For those coercive methods in use in the UK, service users were asked whether they had been subjected to them and staff were asked whether they had used them. A summary of responses to these items is presented in Table 2. As might be expected, staff have greater experience of coercive methods, as they have a constant presence in the acute ward, whereas individual service users pass through for relatively short time spans.

Insert Table 2 about here

Figure 1 plots the sum total approval score for each group for comparison purposes. It can be seen that the service user group disapproved most strongly of net beds, mechanical restraint and IM medication and the staff group disapproved most strongly of net beds, mechanical restraint and open area seclusion. The service user group approved most strongly of intermittent observation, time out and PRN medication; and the staff group approved most strongly of PICU transfer, PRN medication and observation. A score of 18 in Figure 1 was adopted as a cut off to distinguish between ‘absolute’ approval and disapproval as this value lay at the midpoint of the modified likert scale. Using this cut off, both service users and staff disapproved of net beds and mechanical restraint and service users in addition disapproved of IM medication.

Insert Figure 1 about here

1
2
3 Tables 3 and 4 present the mean (sd) scores for service users and staff on each of the
4 six dimensions of approval with regard to the eleven coercive measures. Staff
5 approved of coercive measures overall more highly on each dimension of approval
6 though the dimension 'safe for staff' was endorsed at roughly equal levels. Most
7 aspects of net beds were strongly disapproved of by service users and staff and it is
8 noteworthy that there was a relatively strong endorsement of the item "I would not be
9 prepared to undergo mechanical restraint" by the service user group.
10
11
12
13
14
15
16
17
18

19
20
21
22 Insert Tables 3 and 4 here
23
24
25
26

27 Variations by gender

28
29
30
31 *Service users:* Approval ratings by male service users were consistently significantly
32 higher for manual restraint ($t = 2.26$, $df = 1339$, $p = 0.024$), seclusion ($t = 2.42$, $df =$
33 1330 , $p = 0.016$), mechanical restraint ($t = 3.16$, $df = 1318$, $p = 0.002$) and net beds (t
34 $= 3.79$, $df = 1308$, $p < 0.001$) compared to female service users. Female service users
35 were more likely to have had experience of being subject to intermittent observation
36 ($\chi^2 = 10.81$, $df = 1$, $p = 0.001$), and constant observation ($\chi^2 = 4.81$, $df = 1$, $p = 0.028$),
37 whereas male service users were more likely to have had experience of being subject
38 to seclusion ($\chi^2 = 5.48$, $df = 1$, $p = 0.019$), and psychiatric intensive care ($\chi^2 = 21.21$, df
39 $= 1$, $p < 0.001$).
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 *Staff:* There were also multiple differences within the staff group with male staff
55 consistently approving more highly of every coercive method (PRN medication, $t =$
56 2.14 , $df = 1153$, $p = 0.03$; manual restraint, $t = 3.01$, $df = 1159$, $p = 0.003$; intermittent
57 observation, $t = 2.5$, $df = 1157$, $p = 0.013$; seclusion, $t = 3.97$, $df = 10791$, $p < 0.001$;
58
59
60

1
2
3 mechanical restraint, $t = 4.14$, $df = 1080$, $p < 0.001$; constant observation, $t = 2.08$, df
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
mechanical restraint, $t = 4.14$, $df = 1080$, $p < 0.001$; constant observation, $t = 2.08$, df
= 1150, $p = 0.038$; net beds, $t = 2.6$, $df = 1033$, $p = 0.009$; open area seclusion, $t =$
3.05, $df = 1077$, $p = 0.002$) apart from time out, psychiatric intensive care and IM
medication. Male staff were more likely to have had experience of using seclusion (χ^2
= 9.17, $df = 1$, $p = 0.002$).

Variations by age

Service users: Most items showed a positive relationship to age, with older service
users expressing greater approval of many coercive methods (manual restraint $r =$
0.123, $n = 1349$, $p < 0.001$; seclusion $r = 0.083$, $n = 1340$, $p = 0.002$; IM medication,
 $r = 0.127$, $n = 1338$, $p = 0.077$; PICU, $r = 0.072$, $n = 1333$, $p = 0.008$; constant
observation, $r = 0.105$, $n = 1329$, $p < 0.001$). Younger service users were more likely
to have been subject to physical restraint ($\chi^2 = 11.67$, $df = 5$, $p = 0.04$), time out ($\chi^2 =$
20.44, $df = 5$, $p = 0.001$), and constant observation ($\chi^2 = 11.67$, $df = 5$, $p = 0.04$).

Staff: Younger staff were significantly more approving of mechanical restraint ($r = -$
0.175, $n = 1102$, $p < 0.001$) and net beds ($r = -0.117$, $n = 1057$, $p < 0.001$). There were
relationships between staff age and their experience of having used some coercive
measures (PRN medication, $\chi^2 = 18.8$, $df = 5$, $p = 0.002$; seclusion, $\chi^2 = 33.11$, $df = 5$,
 $p < 0.001$; IM medication, $\chi^2 = 16.09$, $df = 5$, $p = 0.007$), but these relationships were
not straightforward or in each case the same. For some measures younger and older
staff had greater experience but for other measures middle aged staff (30-49 years of
age) had greater experience. For staff, therefore, there is probably an interaction
between age, duration of time working in psychiatry, and cohort affecting approval of
coercive methods.

Variations according to personal experience

Service users: With regard to overall approval score, service users who had been subjected to PRN medication ($t = 6.29$, $df = 1342$, $p < 0.001$) and constant observation ($t = 2.78$, $df = 1327$, $p = 0.005$) approved of these measures more strongly than other non-subjected service users, and those who had been subjected to manual restraint ($t = 7.44$, $df = 1344$, $p < 0.001$) and compulsory IM medication ($t = 7.08$, $df = 1332$, $p < 0.001$) disapproved of these measures more strongly than non-subjected service users.

Staff: There was a universal tendency for staff who had been engaged in using a specific coercion measure approving of it more strongly than those staff who had not (e.g. PRN medication $t = 6.63$, $df = 1139$, $p < 0.001$; manual restraint $t = 6.13$, $df = 1161$, $p < 0.001$). Sample sizes for physical restraint and intermittent observation were highly unbalanced as only 10-15% of staff had never been involved in implementing these procedures.

Discussion

Overall, this survey has established a robust set of benchmark values with regard to how mental health service users and staff in England view various more-or-less controversial coercive measures. These norms can be used as a platform for further work in which the views of other groups of staff in the UK and internationally can be compared. They could also be used pre-post in evaluation studies in which attitudinal change is considered desirable.

This sample covering three regions of England is larger and more representative than that previously used for the ACMQ [18] based as that was on approximately one hundred student nurses at one university. Nevertheless it is worth noting that the relative order of approval of coercive methods from the students was slightly different when compared to the staff group in this study. PRN medication was the most approved method by the students and open area seclusion had a higher ranking. The students' ranking of methods was closer to that of the service user sample in this study, as were their overall approval scores.

The Canadian high secure staff studied by Harris et al. [17] showed the greatest approval rating for seclusion, in stark contrast to the English acute staff in this study, whose order of preferences broadly matched those of the English service users. This difference may be in part due to the different settings of the two studies. Canadian staff and service users disapproved equally of mechanical restraint as in this study. In contrast to the two American studies [16, 17], English staff (but not service users)

1
2
3 rated seclusion as preferable to coerced IM medication. However again there was
4
5 agreement between the American and English staff in that mechanical restraint was
6
7 the most disapproved coercive method in both groups.
8
9

10
11
12 Some significant findings within this study are worth highlighting. There is evidence
13
14 of strong disapproval amongst both staff and service users with regard to the
15
16 introduction of mechanical restraint. Attitudes toward other existing measures did not
17
18 differ hugely between the two groups although service users tended to be more
19
20 disapproving overall than staff. The staff responses varied according to age with older
21
22 staff tending to disapprove more strongly of coercive measures. The age effect, that in
23
24 which younger staff were less disapproving of mechanical restraint, may reflect a
25
26 generation change in which 'old-fashioned' prejudices against an apparently
27
28 legitimate technique are being shed or may result from a lack of exposure to concrete
29
30 examples of use in the real world.
31
32
33
34
35
36
37
38

39 There was a greater approval of coercive methods by men (whether they be staff or
40
41 service users), a finding that confirms a previous international study using the same
42
43 instrument [19]. This indicates the importance of gender roles, perceptions and
44
45 identity in this area. The UK is fairly unique in having a mixed gender qualified
46
47 nursing workforce in psychiatry. The gender effect could reflect any of a large
48
49 number of more general hypothesised gender-related differences (e.g. empathy,
50
51 emotional intelligence). It could be that a more female dominated nursing workforce
52
53 would result in less coercive practice. However in other countries where female
54
55 nursing staff predominate, this has led to the harsher coercive measures being
56
57
58
59
60

1
2
3 implemented by male security guards or by male unqualified nursing aides [22],
4
5 ultimately resulting in much higher levels of use.
6
7
8
9

10 Personal experience was associated with some heterogeneity in the service user group.
11
12 There seems to be a tendency for exposure to ‘gentler’ measures (e.g. observation) to
13
14 enhance approval and conversely, for exposure to ‘harsher’ measures (e.g. IM
15
16 medication) to lead to stronger rejection of the measure. Staff reported quite a
17
18 consistent tendency to approve of techniques once they had employed them in their
19
20 practice which may reflect a process of attitudinal adjustment in which the person
21
22 justifies the measure to themselves afterwards to avoid unpleasant feelings of
23
24 cognitive dissonance (Harmon Jones, 1999).
25
26
27
28
29
30
31

32 Various study limitations must be acknowledged. The sample was large and
33
34 representative of the three regions (although not necessarily of elsewhere in England)
35
36 but staff involvement in the selection of service users could have introduced bias. Due
37
38 to time constraints when assembling the large sample, it was not possible to assess
39
40 potentially important covariates such as service user diagnosis, type of unit or rates of
41
42 assault. In addition, the analysis presupposes the notion of a stable attitude toward a
43
44 coercive measure which is consistent across situations. However it is possible that
45
46 such attitudes are more fluid and inevitably specific incidents will require different
47
48 types of intervention at different times. Further testing of the instrument especially in
49
50 relation to test-retest reliability would be beneficial.
51
52
53
54
55

56 **Conclusions**


57
58
59
60

1
2
3 Service users and staff were both strongly disapproving of the use of any form of
4 mechanical restraint, although that disapproval was slightly stronger amongst the
5 staff. This result suggests that any endeavour to introduce the use of mechanical
6 restraint into adult acute psychiatry in the UK is likely to meet with significant
7 opposition. From the pattern of results we can also predict that if mechanical restraint
8 was introduced to the UK, staff who used it would approve of it more whilst service
9 users subjected to it would dislike it even more leading to a harmful schism between
10 service users and staff.
11
12
13
14
15
16
17
18
19
20
21
22
23

24 The greater approval of coercive methods by male staff, and in the case of the harsher
25 methods their greater involvement in the use of them, raise questions about gender
26 roles within psychiatric nursing. More attention to this aspect of psychiatric care and
27 the issues around it during nurses' training might be necessary.
28
29
30
31
32
33
34
35

36 For service users the most acceptable coercive measure was intermittent observation,
37 followed by time out and PRN medication. Ward regimes based on these methods
38 rather than others are likely to be better received by service users. The least
39 acceptable methods to service users (excluding those not in use in the UK) were
40 restraint, seclusion and coerced IM medication. For these methods disapproval
41 increased with experience, and their use should therefore be avoided as much as
42 possible.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1 Overall approval of coercive methods by service users and staff



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 1 Participant characteristics

		Patients		Staff	
		N	%	N	%
Female		648	48	782	67
Age	Under 20	57	4	13	1
	20-29	268	20	298	25
	30-39	346	26	361	30
	40-49	368	27	332	28
	50-59	198	15	167	14
	over 60	113	8	25	2
Region	North	470	35	411	34
	Central	438	32	469	38
	South	453	33	346	28
Total		1361	100	1226	100

Table 2: exposure to coercive measures (base 1,361 patients and 1,226 staff)

	Patients		Staff	
	N	%	N	%
PICU	367	27	858	70
PRN medication	871	64	883	72
Intermittent observation	966	71	1177	96
Constant observation	612	45	1189	97
Time out	504	37	834	68
IM medication	531	39	834	68
Physical restraint	585	43	1091	89
Seclusion	381	28	564	46

Table 3 Patient attitudes to containment scores: means and standard deviations

PATIENTS	Efficacy		Acceptability		Dignified		Safe for staff		Safe for patients		Prepared to undergo		Sum total approval	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
PRN	3.67	0.98	3.73	0.94	3.59	1.01	3.73	0.90	3.62	0.94	3.63	1.04	21.97	4.67
Physical restraint	3.42	1.13	3.34	1.13	2.91	1.17	3.19	1.08	3.12	1.12	2.89	1.26	18.87	5.60
Intermittent observation	3.78	0.96	3.86	0.90	3.60	1.05	3.89	0.82	3.88	0.88	3.69	1.03	22.69	4.68
Seclusion	3.07	1.19	2.99	1.21	2.78	1.20	3.57	1.01	3.13	1.18	2.62	1.27	18.16	5.78
Time out	3.63	0.99	3.76	0.89	3.67	0.97	3.77	0.83	3.69	0.90	3.61	1.02	22.13	4.67
IM medication	3.25	1.23	2.91	1.25	2.59	1.21	3.26	1.13	2.99	1.18	2.59	1.29	17.59	6.07
PICU	3.55	1.03	3.53	1.04	3.31	1.08	3.64	0.93	3.50	1.01	3.00	1.24	20.53	5.33
Mechanical restraint	2.59	1.27	2.28	1.19	2.11	1.11	3.03	1.23	2.57	1.21	1.99	1.12	14.59	5.90
Constant observation	3.71	1.02	3.66	1.03	3.32	1.15	3.66	0.95	3.73	0.95	3.36	1.19	21.44	5.33
Net bed	2.27	1.24	1.97	1.10	1.91	1.08	2.86	1.32	2.37	1.24	1.73	1.01	13.12	5.77
Open area seclusion	3.50	1.03	3.48	1.05	3.34	1.09	3.42	1.00	3.48	1.01	3.21	1.19	20.44	5.53
Summed total score	36.43	7.28	35.52	7.34	33.13	7.85	37.98	6.74	36.08	7.35	32.32	8.42		

Table 4 Staff attitudes to containment scores: means and standard deviations

STAFF	Efficacy		Acceptability		Dignified		Safe for staff		Safe for patients		Prepared to use		Sum total approval	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
PRN	4.27	0.65	4.24	0.59	4.14	0.69	4.01	0.75	4.03	0.66	4.20	0.77	24.95	3.27
Physical restraint	4.04	0.75	3.95	0.75	3.45	1.03	3.45	1.00	3.61	0.89	4.07	0.77	22.64	4.13
Intermittent observation	3.93	0.90	4.08	0.73	3.80	0.88	3.82	0.88	4.00	0.79	4.16	0.71	23.84	4.10
Seclusion	3.65	1.03	3.54	1.02	3.28	1.08	3.69	0.97	3.58	0.99	3.62	1.07	21.45	5.48
Time out	3.96	0.79	4.05	0.70	3.98	0.76	3.85	0.81	3.94	0.75	4.05	0.73	23.86	4.02
IM medication	4.13	0.77	3.91	0.82	3.32	1.01	3.72	0.84	3.69	0.81	3.96	0.85	22.78	4.19
PICU	4.35	0.66	4.32	0.63	4.09	0.80	4.10	0.80	4.17	0.74	4.29	0.69	25.39	3.75
Mechanical restraint	2.42	1.17	2.10	1.02	2.02	1.02	2.47	1.10	2.25	1.04	2.05	1.09	13.26	5.78
Constant observation	4.22	0.69	4.20	0.66	3.51	1.03	3.61	0.97	4.08	0.71	4.22	0.64	23.91	3.71
Net bed	2.27	1.06	1.98	0.96	1.93	0.97	2.36	1.07	2.20	1.01	1.90	0.98	12.58	5.42
Open area seclusion	3.48	0.95	3.48	0.93	3.42	0.95	3.12	1.02	3.43	0.94	3.36	1.03	20.28	5.39
Summed total score	41.11	5.21	40.22	5.16	37.36	6.25	38.44	5.90	39.38	5.23	40.34	5.43		

NB: range from 1 to 5 for each individual item, from 6 to 30 for Sum total approval and from 11 to 55 for Summed total score in both tables. Higher scores indicate a positive evaluation in all cases

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

For Review Only

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
1. Steinert, T., V. Martin, M. Baur et al., Diagnosis-related frequencies of compulsory measures in ten German psychiatric hospitals and correlates with hospital characteristics. *Social Psychiatry and Psychiatric Epidemiology*, 2007. 42: p. 140-145.
2. Donat, D.C., Special section on seclusion and restraint: Encouraging alternatives to seclusion, restraint, and reliance on PRN drugs in a public psychiatric hospital. *Psychiatric Services*, 2005. 56: p. 1105-1108.
3. Whittington, R., E. Baskind, and B. Paterson, Coercive measures in the management of imminent violence: restraint, seclusion and enhanced observation, in *Violence in Mental Health Settings: Causes, Consequences, Management*, D. Richter and R. Whittington, Editors. 2006, Springer Verlag: New York.
4. Huckshorn, K., Reducing seclusion restraint in mental health use settings: core strategies for prevention. *Journal of Psychosocial Nursing and Mental Health Services*, 2004. 42: p. 22-33.
5. D'Orio, B.M., D. Purselle, D. Stevens et al., Reduction of episodes of seclusion and restraint in a psychiatric emergency-service. *Psychiatric Services*, 2004. 55: p. 581-583.
6. Hopton, J., Control and restraint in contemporary psychiatric nursing: some ethical considerations. *Journal of Advanced Nursing*, 1995. 22: p. 110-115.
7. Lind, M., R. Kaltiala-Heino, T. Suominen, et al., Nurses' ethical perceptions about coercion. *Journal of Psychiatric and Mental Health Nursing*, 2004. 11: p. 379-385.
8. Bonner, G., T. Lowe, D. Rawcliffe et al., Trauma for all: a pilot study of the subjective experience of physical restraint for mental health inpatients and staff in the UK. *Journal of Psychiatric and Mental Health Nursing*, 2002. 9: p. 465-473.
9. Johnson, M., Being restrained: a study of power and powerlessness. *Issues in Mental Health Nursing*, 1998. 19: p. 191-206.
10. Wynn, R., Psychiatric inpatients' experiences with restraint. *Journal of Forensic Psychiatry and Psychology*, 2004. 15: p. 124-144.
11. Hoekstra, T., H. Lendemeijer, and M. Jansen, Seclusion: the inside story. *Journal of Psychiatric and Mental Health Nursing*, 2004. 11: p. 276-283.
12. Duxbury, J., An evaluation of staff and patient views of strategies employed to manage inpatient aggression and violence on one mental health unit: a pluralistic design. *Journal of Psychiatric and Mental Health Nursing*, 2002. 9: p. 325-337.

13. Meehan, T., H. Bergen, and K. Fjeldsoe, Staff and patient perceptions of seclusion: has anything changed? *Journal of Advanced Nursing*, 2004. 47: p. 33-38.
14. Chien, W.T., C. Chan, L. Lam et al., Psychiatric inpatients' perceptions of positive and negative aspects of physical restraint. *Patient Education And Counseling*, 2005. 59: p. 80-86.
15. Klinge, V., Staff Opinions About Seclusion and Restraint at a State Forensic Hospital. *Hospital and Community Psychiatry*, 1994. 45: p. 138-141.
16. Terpstra, T., T. Terpstra, E. Pettee et al., Nursing Staff's Attitudes Toward Seclusion and Restraint. *Journal of Psychosocial Nursing*, 2001. 29: p. 21-49.
17. Harris, G., M. Rice, and D. Preston, Staff and patient perceptions of the least restrictive alternatives for the short-term control of disturbed behaviour. *Journal of Psychiatry and Law*, 1989. 17: p. 239-263.
18. Bowers, L., J. Alexander, A. Simpson et al., Cultures of psychiatry and the professional socialization process: the case of containment methods for disturbed patients. *Nurse Education Today*, 2004. 24: p. 435-442.
19. Bowers, L., B. van der Werf, A. Vokkolainen, et al., International variation in attitudes to containment measures for disturbed psychiatric inpatients. *International Journal of Nursing Studies* 2007. 44: p. 357-364.
20. Bowers, L., R. Whittington, P. Nolan et al., The City 128 Study of Observation and Outcomes on Acute Psychiatric Wards. Report to the NHS SDO Programme. 2006, City University: London.
21. Tavcar, R., M.Z. Dernovsek, and V.N. Grubic, Use of coercive measures in a psychiatric intensive care unit in Slovenia. *Psychiatric Services*, 2005. 56: p. 491-492.
22. Morrison, E., G. Morman, G. Bonner et al., Reducing Staff Injuries and Violence in a Forensic Psychiatric Setting. *Archives of Psychiatric Nursing*, 2002. 16: p. 108-117.
23. Harmon-Jones, E., *Cognitive Dissonance: Progress on a Pivotal Theory in Social Psychology* 1999, Washington DC: American Psychological Association.