



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** McCann, E. (2010). The sexual and relationship needs of people who experience psychosis: quantitative findings of a UK study. *Journal of Psychiatric and Mental Health Nursing*, 17(4), pp. 295-303. doi: 10.1111/j.1365-2850.2009.01522.x

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/27927/>

**Link to published version:** <https://doi.org/10.1111/j.1365-2850.2009.01522.x>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

---

---

City Research Online:

<http://openaccess.city.ac.uk/>

[publications@city.ac.uk](mailto:publications@city.ac.uk)

---

# **The sexual and relationship needs of people who experience psychosis: quantitative findings of a UK study**

E. McCann PhD RMN FEHA

*Lecturer, School of Nursing and Midwifery, University of Dublin, Trinity College, Dublin, Ireland*

## **Accessible summary**

- Distinct lack of studies exist that explore sexual and relationship issues.
- Captures important experiences of people who use mental health services.
- Reveals potential obstacles to the expression of sexuality.
- Identifies a diversity of needs.
- Presents issues that may guide mental health practice, education and research.

## **Correspondence:**

E. McCann

Trinity College Dublin

School of Nursing and Midwifery 24 D'Olier Street

Dublin 2 Ireland

E-mail: [mccanned@tcd.ie](mailto:mccanned@tcd.ie)

## **Abstract**

Few studies have investigated the experiences of people regarding sexual and relationship issues in the area of mental health. This study presents the quantitative findings of a larger study that was conducted in London, UK. The aims of the study were to establish client's sexual and relationship experiences and perceived needs. A total of 30 people with a medical diagnosis of schizophrenia, living in the community, were interviewed using three questionnaires. The first related to demographics, the second used relevant parts of the Camberwell Assessment of Need (CAN) and the third looked at possible determinants of sexual behaviour. The CAN also captured keyworker responses to issues related to their clients sexual and relationship requirements. The results showed that 83% of the clients were currently experiencing sexual feelings. Some 90% of clients felt some need in relation to sexual expression and 83% for needs related to intimate relationships. Only 10% of staff recognized sexual expression as a need in clients in their care and 43% perceived a need for intimate relationships. Furthermore, most clients interviewed thought that their psychotropic medication caused sexual problems. Contrasts are made with other studies to help highlight the important issues that emerged for service users.

*Keywords:* psychosis, relationships, schizophrenia, sexuality

## Background

In the UK there have been government drives to make health and social care more responsive and inclusive. Earlier documents include the *NHS and Community Care Act* (Department of Health 1990a) and the Care Programme Approach (CPA) (Department of Health 1990b). More recent publications have outlined government strategies which aim to tackle issues pertinent to users, carers and significant others regarding mental health provision focussing on the process of recovery. Such documents include the *National Service Frameworks* (Department of Health 1999), *Vision for Change* (Department of Health and Children 2006), *Rights, Relationships and Recovery* (Government of Scotland 2006) and *A Recovery Approach within the Irish Mental Health Service* (Mental Health Commission 2008). Additionally, the Government has published a national strategy for sexual health and HIV (Department of Health 2001), a document addressing domestic and sexual violence (Department of Health 2006) and a proposal for potential therapeutic interventions (Department of Health 2007). As a result, more attention has been paid to detailed assessment of the needs of all individuals suffering from mental illnesses such as schizophrenia and psychosis. Topics of importance include housing, occupation, education, medical, social and psychiatric care, socialization, risk assessment and recovery (Davidson *et al.* 2006, Buchanan-Barker & Barker 2008). However, over time, the intimate sexual and interpersonal needs of affected individuals have emerged as a justifiable concern for mental health practitioners (Lewis & Scott 1997, McCann 2003, Higgins *et al.* 2009). The Health of the Nation (Department of Health 1992) targets compartmentalized sexual health and mental health as separate and distinct entities. The former concentrated on teenage pregnancies and sexually transmitted diseases and the latter examined ways of reducing suicides (Adler 1997). While these targets seem praiseworthy for the whole UK population, the sexual needs of individuals with psychosis appear to remain marginalized and neglected. In the UK, the *National Survey of Sexual Attitudes and Lifestyles* (Wellings *et al.* 1994), one of the largest sexual studies since the *Kinsey Report* (Kinsey *et al.* 1948), only made reference to physical health. (McCann 2003) carried out a systematic review of the available literature relating to sex and relationship issues and people with a diagnosis of schizophrenia or psychosis.

Studies mainly concerned mental health populations in the USA with very few studies, and no empirical research having been carried out in the UK. The subject of HIV/AIDS and perceived 'risky' behaviours received most coverage (McKinnon *et al.* 1993, Herman *et al.* 1994, McDermott *et al.* 1994, Gottesman & Groome 1997). There was some exploration into the effects of stigma and discrimination and the apparent damaging effects of such treatment of mentally ill people by society (Bacharach 1992, Rowlands 1995, Crisp *et al.* 2000). Several studies examined the effects of medication on a person's sexual functioning and included cognitive as well as physical responses (Bhui *et al.* 1997, Smith *et al.* 2002). Some support systems were studied along with notions about marriage (Lane *et al.* 1995), family planning (Miller & Finnerty 1996) and vulnerability (Read & Argyle 1999). Further, literature concerning possible professional responses, particularly the identification, assessment and planning of service and therapeutic provision was extrapolated (Park Dorsay & Forchuk 1994, Dilloway & Hildyard 1998). The final picture revealed distinct gaps in psychosexual knowledge in relation to patient needs and this provided the rationale for the present study.

### **Study aims**

1. To discover the clients' sexual and relationship experiences in the past and present and to elicit hopes and aspirations for the future.
2. To uncover some of the obstacles to the expression of sexuality in people with serious mental illness living in the community.
3. To present recommendations for mental health practice, education and research.

### **Methods in practice**

#### ***Sampling***

A recognized sampling procedure was followed (Kuzel 1992). The population under study consisted of people with a case-note diagnosis of schizophrenia, living in the community, and regularly attending a depot clinic in North London to receive anti-psychotic medication. The following selection criteria were

used:

- case note diagnosis of schizophrenia;
- aged between 16 and 64 years;
- taking neuroleptic medication;
- regularly attending the depot clinic.

The participants were recruited from one depot clinic. A list of attendees was obtained from the Locality Team administrator and totalled 47. Participants were conveniently recruited to take part in the study. An office was provided in the clinic that afforded privacy where participants were informed about the study and consent sought, The Community Psychiatric Nurses administering the injection would make a clinical judgement on the suitability or wellness of the person to take part in the study.

### ***Ethical considerations***

Approval for the study was gained from the local hospital ethics committee (Reference number: P98210). At all times the investigator was respectful of the participants while administering the questionnaires. Time was invested in explaining the purposes and aims of the research. Participants were assured that should they wish to stop the interview for any reason they would be free to do so. Furthermore, participants were guaranteed anonymity and were informed that names would not appear on any documents. They would not be identifiable. The data from the study, which included the completed questionnaires, were kept in secure locked cupboards within the University. Written consent was gained. Participants were invited to ask questions and seek clarification before beginning the interviews and at the end.

### **Data collection**

Three questionnaires were used to obtain data.

#### *A questionnaire relating to demographic data*

This included racial identity, gender, age, education, marital status, whether in a relationship or not, number of children, medical diagnosis, number of hospital admissions and last hospital admission (Table 1).

**\*\*Table 1 here\*\***

*Relevant parts of The Camberwell Assessment of Need (CAN) (Phelan et al. 1995)*

The CAN is a well-validated and reliable tool devised to assess the complex clinical and social needs of people with severe mental illness (Slade *et al.* 1999). The questionnaire incorporates both staff and user assessment of issues over the past month. There are 22 items in the full assessment schedule. The chosen parts used in the present study specifically address the issue of intimate relationships and sexual expression. Both user views (Table 2) and staff opinions (Table 3) were rated. The user and staff responses relating to need identification were collated (Table 4).

*A structured interview schedule incorporating the determinant factors of sexual behaviour through life (Pfeiffer & Davis 1972) modified version (McCann 2000)* The original determinants of sexual behaviour (DSB) questionnaire relates to interest, frequency and satisfaction of sexual activity in younger years and at the present time. It also records reasons for stopping sexual relations. Pfeiffer and Davis used the questionnaire on a population of 502 subjects aged between 46 and 69 years. The modified version uses additional questions related to views about medications (Table 5).

\*\*\*Tables 2,3,4,5, here \*\*\*

## **Results and data analysis**

### ***Data analysis***

The data were analysed using the Statistical Package for the Social Sciences (spss v9) (Brace *et al.* 2000).

The analysis is divided into two parts:

1. descriptive statistics which provide information relating to the sample characteristics;
2. inferential statistics investigating relationships between selected variables.

The data were manipulated after entry into spss, including some recoding to enable the execution of further inferential statistical analyses. It was necessary to split cases, e.g. male and female, to determine certain responses. Additionally, 2×2 contingency tables were constructed

to allow for the computation of specific statistical tests.

### ***Description of the client sample***

From a total of 47 patients attending a Health Centre in London to have depot medication administered, 30 people were interviewed. The participant characteristics are shown in Table 1. The age group of the sample varied between 22 and 57 years (mean = 40.93 years). Most people identified as White UK (46.7%), although within the sample, there was a rich mix of ethnic backgrounds, reflecting the diversity of the local population, including: Kurdish, Black African and Black Caribbean. A majority of the sample were heterosexual (73.3%), not in a relationship currently (60%) and had no children (66.7%). Most people had engaged in further education (70%). The entire sample had a clinical diagnosis of schizophrenia, schizotypal and delusional disorders, as defined by the International Classification of Diseases (World Health Organization 1992). A significant percentage of people had less than 10 years contact with mental health services and the average number of previous admissions was five and a half.

To allow for a further analysis of the data, chi-squared tests were used to compare relevant responses (Greene & d'Oliveira 1999). A cross-tabulation of the variables 'sex' and 'children' showed significant results, where more females than males were more likely to parent children ( $\chi^2 9.600$ , d.f. = 1,  $P = 0.002$ ). There were no other statistically significant relationships between demographic variables.

### ***Results of the CAN***

The CAN staff responses were completed by the 'named' CPA community keyworkers who worked in the locality team. With regard to intimate relationships, most users (70%) gave this maximum scoring in each of the three sections, seeing this as a serious unmet need. Staff (13%) and users (17%) perceived this as an area of no need. A large number of staff (43%) were unable to say if a need existed for their clients, with some (23%) perceiving this as a serious unmet need.

In response to the items on sexual expression, a very high proportion of staff (60%) could not say if there

was a need for clients in this area. Conversely, all users were able to articulate responses to every question. More than half of users (53%) reported serious unmet need compared with staff responses (7%). A larger proportion (37%) of users reported sexual expression as being a met or partially met need as opposed to (3%) of staff.

The user and staff responses relating to need identification were collated (Table 4). A high number of user respondents (90%) recognized that there was some need in the area of sexual expression, whereas very few staff (10%) thought that this was the case. With regard to intimate relationships, 83% of users identified some need, whereas 57% of staff respondents reported no need/not known. The merging of the responses 'no need' and 'not known' may be problematic as staff could not say whether there was an identified need or not. Presumably, they had never asked their patients about these specific areas of need.

### ***Results of the DSB interview***

More than half of the respondents (57%) had no enjoyment of sexual relations before receiving treatment with 43% reporting some degree of enjoyment. At the time of interview, 37% of the respondents were claiming to have had a satisfactory sex life. Only 13% reported having no enjoyment whatsoever. A significant number of people (83%) said they were experiencing sexual feelings at the present time although 49% reported that they were not having 'sex relations' at the moment. More than half (67%) noted a decline in sexual interest or activity with a majority of people reporting a diminution in the last 5 years. Moreover, sex ceased for 50% of the sample between 1 and 5 years ago. The main reasons given for stopping sexual relations were: illness of self (17%); separation/divorce (17%) and lack of opportunity (17%). A large number of respondents (77%) knew the names of the medications they were taking. Significantly, 60% of people interviewed, thought that medication caused sex problems.

Chi-squared tests were used to determine levels of significance between selected variables on the DSB questionnaire. The responses 'enjoyment of sex relations before receiving treatment' and 'frequency of sex relations now' were cross-tabulated and revealed significant results ( $\chi^2$  28.249, d.f. = 9,  $P = 0.001$ )

suggesting that people who never enjoyed sex pre-treatment were still not having sex. There were no other statistically significant results between the other variables.

### ***Relationships between the demographics and CAN***

Using the chi-squared test, relevant participant characteristics were compared with the results gained in the CAN questionnaire, to determine whether there were statistically significant differences in staff and user responses to sexual problems and relationship difficulties. No significant differences were noted on the variables *sex, ethnicity, age, children, diagnosis* and *service contact*. However, there was a strong significant difference between *heterosexual and non-heterosexual* perceptions of *intimate relationship problems* ( $\chi^2$  8.727, d.f. = 1,  $P$  = 0.003) which may suggest that gay and lesbian respondents were more satisfied with their relationships.

### ***Relationships between the demographics and DSB***

Chi-squared tests were applied to the demographic characteristics and the DSB results of the present sample ( $n$  = 30). The only significant differences between responses were found in the categories *in relationship now* and *frequency of sex relations at the present time* ( $\chi^2$  15.502, d.f. = 2,  $P$  = 0.000), suggesting that people in a current relationship were more likely to be having sex.

### ***Relationships between the CAN and DSB***

Chi-squared statistical tests were carried out between selected variables from both questionnaires. No statistically significant results were found between the responses.

### ***Comparison between the ward and community groups***

The data from the present study was compared with data obtained from an earlier hospital ward-based study (Table 5). The ward was situated in a large Victorian psychiatric hospital in the East End of London. There were fifteen patients being cared for in this locked acute rehabilitation environment. It

was a mixed-sex ward and culturally diverse. A total of 11 (73%) patients agreed to be interviewed from a possible 15. The entire group had a primary medical diagnosis of schizophrenia. Over half of the sample (64%) had spent between 1 and 5 years in psychiatric hospitals (McCann 2000).

In the community sample, a significant number (87%) said they enjoyed sexual relations at the present time compared with ward respondents (27%). More than 40% of the inpatients enjoyed sexual relations before hospitalization. Over 60% felt aware of a decline in sexual interest or activity, which almost equates with the community respondents. In both of the studies, over 50% reported a decline occurring within the last 10 years. Most could not remember exactly when, but over half stated ‘. . . since coming into hospital’. Sex relations stopped for over 70% of the sample since being hospitalized.

When asked why sex relations had stopped, the ward respondents mainly identified illness of self (36%) and lack of opportunity (36%). The community group identified illness of self (17%), separation/divorce (17%) and lack of opportunity (17%). A proportion of the community sample reported active sex relations (27%). Each of the client groups (over 70%) knew the medication they were taking. Of the 11 participants, two people said that their medication contributed to sex problems in contrast to a much higher proportion (60%) of community respondents. Chi-squared tests were run on the data to test whether there were significant differences between ward and community responses for each of the categories on the DSB questionnaire (Table 5). Questions 1, 5, 6 and 9 all had similar scores and showed no significant relationship between each of the variables. There were significant differences in response to the question *enjoyment of sex now*, with the ward sample scoring high on no sex (73%) compared with (13%) in the community. This may be attributed to the strict rules and regulations operating within ward environments. The question related to *sexual feelings now* showed significant differences. Both groups had strong sexual feelings with a high lack of opportunity expressed in the ward sample. Results from the question *frequency of sex now* revealed significance with 82% of ward respondents never having sex compared with 49% of the community group. A fair number of community respondents (22%) reported having sex once a week or more. Significant differences were shown in response to the question relating to *when sex*

*stopped*, with 27% of the community respondents saying sex had not stopped compared with sex ceasing to happen for the entire ward sample. Again, this is not surprising when consideration is given to the oppressive nature of some inpatient environments. Moreover, when asked about *reasons for stopping sex*, 36% of the ward sample stated lack of opportunity. Finally, significant differences were apparent in response to the last question about *medication and sex problems* with 60% of the community respondents stating that they felt that medication did contribute to sexual difficulties as opposed to 18% of the ward sample. This may be due to the fact that a relatively high number of people living in the community were perhaps involved in sexual activity and were experiencing problems.

### ***Comparison between CAN results and other studies***

A search was conducted to locate studies that had used the CAN (Table 6). In a Scottish study (Simons & Petch 2002), that examined needs assessment and discharge from hospital, the authors attempted a response to the National Service Framework for Mental Health (Department of Health 1999) Standard Five which states:

*Each service user who is assessed as requiring a period of time away from their home should have . . . a copy of a written care plan agreed on discharge, which sets out the care and rehabilitation to be provided, identifies the care-co-ordinator, and specifies the action to be taken in a crisis.*

In this study, the 22-item CAN was used to measure staff and patient perceptions of need. The patient responses ( $n = 173$ ) for sexual expression were rated zero. The reason given was that the interviewers were uneasy discussing the issue of sexuality with the respondents. Only 3% of staff ( $n = 98$ ) identified this as an unmet need. In the present study, 53% of patients felt their needs regarding sexual expression were not being met whereas only 7% of staff perceived this as an area of serious unmet need. Where intimate relationships were concerned, 10% of patients and 6% of staff respondents identified unmet needs in the Scottish study. This compares with patient responses (73%) and staff responses (23%) in the present study.

An earlier study (Slade *et al.* 1996) looked at the association between the assessment of need by staff

and by severely mentally ill patients. The results of the staff responses are not that dissimilar to those of the present study. However, patient identification of need varies greatly on both domains of sexual expression (83%, 20%) and intimate relationships (90%, 27%).

The high ratings in the present study may be due to the overall nature of the study and the focus upon sex and relationship issues, which was made explicit to the participants before the interviews began.

## **Discussion and conclusion**

In order to address the aims of the study, 30 people with a diagnosis of schizophrenia currently living in the community completed the questionnaires. Some had spent many years in hospital while others had less familiarity with life in institutional settings. The composition of the sample was ethnically diverse and appeared to be fairly representative of the service users within the local population. Just under half were currently in a relationship. The CAN was used to gauge levels of need from a staff and user perspective and is considered to be one of the first rating instruments to do so (Phelan *et al.* 1995). The CAN survey revealed dramatic differences between staff and user perceptions of need around both intimate relationships and sexual expression. Nearly all of the service users (90%) identified need in terms of sexual expression. Staff gave this very low priority. This apparent disparity perhaps highlights the concerns around attitudes of staff in relation to client sexuality needs and the potential to talk about the subject (McCann 2000, Higgins *et al.* 2009). A comparison between the present study and other studies that used the CAN revealed problems on behalf of the researchers (Simons & Petch 2002). The research interviewers appeared to have an inability to gain information concerning sexuality concerns. This, it could be argued, may be due to their own discomfort in addressing sexual issues that could result in this important area remaining neglected or ignored completely. The opposite was true in the present study. Participants were very forthcoming in their responses and were keen to share additional information that flags the opportunity for future therapeutic activity. The DSB survey uncovered some notions about individual's past and present sexual and relationship behaviours. Valuable information emerged in relation

to the earlier ward study and the present study. For instance, there was huge disparity between sexual activities on the ward compared with people living in the community. This is perhaps not so surprising as strict rules apply on many wards with sexual behaviour being problematized or even ignored (Alexander 2003). The issues around neuroleptic medication were elicited and a majority of respondents (60%) who lived in the community attributed sex problems to medication. The unwanted effects of medication and the related sexual problems, that often remain undetected, have been highlighted as one of the main reasons for non-concordance (Smith *et al.* 2002).

In conclusion, although some interesting and valuable findings emerged from the data, several limitations were acknowledged. The sample size was low in number and this reduced the power of statistical testing. Nevertheless, the data gathered and presented in this study informed the larger study that included semi-structured questionnaires in order that thoughts, attitudes and feelings of participants were determined. These findings are being presented elsewhere. However, it has become clear from the information gained in this study that nurses do not generally ask patients about sexual and relationship matters. All of the respondents, contrary to clinical opinion, were able to articulate their experiences around the issues of sexuality. Perhaps nurses' own anxieties create obstacles to the exploration of what may be deemed 'sensitive' topics. Additionally, societal pressures particularly surrounding the taboos of sex and the discussion of all matters sexual may obstruct the potential for dialogue between nurses and the people we care for and may impact upon a person's recovery trajectory.

## References

- Adler M.W. (1997) Sexual health – a Health of the Nation failure. *British Medical Journal* **314**, 1743–1751.
- Alexander J. (2003) *The relationship between flexibility/ inflexibility of ward nursing regimes and patient outcomes*. Unpublished PhD Thesis, City University, London.
- Bacharach L. (1992) Psychosocial rehabilitation and psychiatry in the care of long-term patients. *American Journal of Psychiatry* **149**, 11.
- Bhui K., Puffet A. & Strathdee G. (1997) Sexual and relationship problems amongst patients with severe

- chronic psychosis. *Social Psychiatry and Psychiatric Epidemiology* **32**, 459–467.
- Brace N., Kemp R. & Snelgar R. (2000) *SPSS for Psychologists: A Guide to Data Analysis Using SPSS for Windows*. Palgrave, New York.
- Buchanan-Barker P. & Barker P.J. (2008) The tidal commitments: extending the value base of mental health recovery. *Journal of Psychiatric and Mental Health Nursing* **15**, 93–100.
- Crisp A.H., Gelder M.G., Rix S., *et al.* (2000) Stigmatisation of people with mental illnesses. *British Journal of Psychiatry* **177**, 4–7.
- Davidson L., O’Connell M., Tondora J., *et al.* (2006) The top ten concerns about recovery encountered in mental health system transformation. *Psychiatric Services* **57**, 640–645. Department of Health (1990a) *NHS and Community Care Act*. HMSO, London.
- Department of Health (1990b) *The Care Programme Approach for People with Mental Illness*. HC (90) 23. HMSO, London.
- Department of Health (1992) *The Health of the Nation: A Strategy for Health in England*. HMSO, London.
- Department of Health (1999) *Modern Standards and Service Models – National Service Frameworks*. HMSO, London.
- Department of Health (2001) *The National Strategy for Sexual Health and HIV*. HMSO, London.
- Department of Health (2006) *Tackling the Health and Mental Health Effects of Domestic and Sexual Violence and Abuse*. HMSO, London.
- Department of Health (2007) *Commissioning A Brighter Future: Improving Access to Psychological Therapies – Positive Practice Guide*. HMSO, London.
- Department of Health and Children (2006) *A Vision for Change: Report of the Expert Group on Mental Health Policy*. Stationery Office, Dublin.
- Dilloway M. & Hildyard S. (1998) Female patients’ views on discussing sexual health. *British Journal of Community Nursing* **3**, 172–177.
- Gottesman I.I. & Groome C.S. (1997) HIV/AIDS risks as a consequence of schizophrenia. *Schizophrenia Bulletin* **23**, 675–684.
- Government of Scotland (2006) *Rights, Relationships and Recovery: The Report of the National Review of Mental Health Nursing in Scotland*. Scottish Executive, Edinburgh.
- Greene J. & d’Oliveira M. (1999) *Learning to Use Statistical Tests in Psychology*, 2nd edn. Open University Press, Buckingham.
- Herman R., Kaplan M., Satriano J., *et al.* (1994) HIV prevention with people with serious mental illness: staff training and institutional attitudes. *Psychosocial Rehabilitation Journal* **17**, 97–103.
- Higgins A., Barker P. & Begley C. (2009) Clients with mental health problems who sexualize the client-nurse encounter. *Journal of Advanced Nursing* **65**, 616–624.
- Kinsey A., Pomeroy W. & Martin C. (1948) *Sexual Behaviour in the Human Male*. W.B. Saunders, London.
- Kuzel A.J. (1992) Sampling in qualitative enquiry. In: *Doing Qualitative Research* (eds Crabtree, B.F. & Miller, W.L.), pp. 32–46. Sage Publications, London.
- Lane A., Byrne M., Mulvaney F., *et al.* (1995) Reproductive behaviour in schizophrenia relative to other mental disorders: evidence for increased fertility in men despite reduced marital rate. *Acta Psychiatrica Scandinavica* **91**, 222–228.
- Lewis J. & Scott E. (1997) The sexual education needs of those disabled by mental illness. *Psychiatric Rehabilitation Journal* **21**, 164–167.
- McCann E. (2000) The sexual and relationship needs of people with psychosis: breaking the taboos.

- Journal of Advanced Nursing* **32**, 132–138.
- McCann E. (2003) Exploring sexual and relationship possibilities for people with psychosis – a review of the literature. *Journal of Psychiatric and Mental Health Nursing* **10**, 640–649.
- McDermott B.E., Sautter F.J., Winstead D.K., *et al.* (1994) Diagnosis, health beliefs, and risk of HIV infection in psychiatric patients. *Hospital and Community Psychiatry* **45**, 580–585.
- McKinnon K., Cournos F., Meyer-Bahlburg H., *et al.* (1993) Reliability of sexual risk behaviour interviews with psychiatric patients. *American Journal of Psychiatry* **150**, 972–974.
- Mental Health Commission (2008) *A Recovery Approach within the Irish Mental Health Service: A Framework for Development*. Mental Health Commission, Dublin.
- Miller L.J. & Finnerty M. (1996) Sexuality, pregnancy and childrearing among women with schizophrenia-spectrum disorders. *Psychiatric Services* **47**, 502–506.
- Park Dorsay J. & Forchuk C. (1994) Assessment of the sexuality needs of individuals with psychiatric disability. *Journal of Psychiatric and Mental Health Nursing* **1**, 93–97.
- Pfeiffer R. & Davis G. (1972) Determinants of sexual behaviour in middle and old age. *Journal of the American Geriatric Society* **20**, 151–158.
- Phelan M., Slade M., Thornicroft G., *et al.* (1995) The Camberwell Assessment of Need: the validity and reliability of an instrument to assess the needs of people with severe mental illness. *British Journal of Psychiatry* **167**, 589–595.
- Read J. & Argyle N. (1999) Hallucinations, delusions, and thought disorder among adult psychiatric inpatients with a history of child abuse. *Psychiatric Services* **50**, 1467–1472.
- Rowlands P. (1995) Schizophrenia and sexuality. *Sexual and Marital Therapy* **10**, 47–61.
- Simons L. & Petch A. (2002) Needs assessment and discharge: a Scottish perspective. *Journal of Psychiatric and Mental Health Nursing* **9**, 435–445.
- Slade M., Phelan M., Thornicroft G., *et al.* (1996) The Camberwell Assessment of Need (CAN): comparison of assessments by staff and patients of the need of the severely mentally ill. *Social Psychiatry and Psychiatric Epidemiology* **31**, 109–113.
- Slade M., Thornicroft G., Loftus L., *et al.* (1999) *CAN: Camberwell Assessment of Need*. Gaskell, London.
- Smith S.M., O’Keane V. & Murray R. (2002) Sexual dysfunction in patients taking conventional antipsychotics medication. *British Journal of Psychiatry* **181**, 49–55.
- Wellings K., Field J., Johnson A., *et al.* (1994) *Sexual Behaviour in Britain*. Penguin, London.
- World Health Organization (1992) *ICD-10: The ICD-10 Classification of Mental and Behavioural Disorders: Clinical Descriptions and Diagnostic Guidelines*. WHO, Geneva.

**Table 1**  
Demographic characteristics

Variables		<i>n</i> (%)
Sex	Male	15 (50.0)
	Female	15 (50.0)
	Age: mean (SD)	40.93 (10.01)
Ethnicity	White UK	14 (46.7)
	Black Caribbean	8 (26.7)
	White European	5 (16.7)
	Black African	2 (6.7)
	Indian	1 (3.3)
Sexual orientation	Heterosexual	22 (73.3)
	Gay	2 (6.7)
	Lesbian	5 (16.7)
	Bisexual	1 (3.3)
Further education	College	14 (46.7)
	University	7 (23.3)
	None	9 (30.0)
Marital status	Single	16 (53.3)
	Ever married	14 (46.7)
Relationship now	Yes	12 (40.0)
	No	18 (60.0)
Children	Yes	10 (33.3)
	No	20 (66.7)
Clinical diagnosis	Schizophrenia	20 (66.7)
	Schizo-affective disorder	10 (33.3)
Service contact (years)		
	0-5	6 (20.0)
	6-10	10 (33.3)
	11-15	2 (6.7)
	16-20	4 (13.3)
	>20	8 (26.7)
Previous admissions: mean (SD)		5.53 (5.06)

**Table 2**User (*n* = 30) assessment of level of need (%)

CAN domain	No need, <i>n</i> (%)	Met or partially met need, <i>n</i> (%)	Serious unmet need, <i>n</i> (%)	Not known, <i>n</i> (%)
Intimate relationships	5 (17)	3 (10)	22 (73)	0
Sexual expression	3 (10)	11 (37)	16 (53)	0

CAN, Camberwell Assessment of Need.

**Table 3**Staff (*n* = 30) assessment of level of need (%)

CAN domain	No need, <i>n</i> (%)	Met or partially met need, <i>n</i> (%)	Serious unmet need, <i>n</i> (%)	Not known, <i>n</i> (%)
Intimate relationships	4 (13)	6 (21)	7 (23)	13 (43)
Sexual expression	9 (30)	1 (3)	2 (7)	18 (60)

CAN, Camberwell Assessment of Need.

**Table 4**Identified need by staff (*n* = 30) and users (*n* = 30)

CAN domain	Staff identifying need, <i>n</i> (%)	Not known/no need (staff), <i>n</i> (%)	Users identifying need, <i>n</i> (%)	Not known/no need (users), <i>n</i> (%)
Intimate relationships	13 (43)	17 (57)	25 (83)	5 (17)
Sexual expression	3 (10)	27 (90)	27 (90)	3 (10)
Mean	8 (26.5)	22 (73)	26 (86.5)	4 (13.5)

CAN, Camberwell Assessment of Need.

**Table 5**  
**Determinant factors of sexual behavior (%)**

	2000 study Ward (n = 11), n (%)	Present study Community (n = 30), n (%)	Chi-squared test	P value
1. Enjoyment of sex relations pre hospital/treatment			2.82	0.42
None	5 (46)	17 (57)		
Mild	2 (18)	5 (17)		
Moderate	2 (18)	4 (13)		
Very much	2 (18)	4 (13)		
2. Enjoyment of sex relations at the present time			86.78	0.01**
None	8 (73)	4 (13)		
Mild	1 (9)	7 (23)		
Moderate	2 (18)	8 (27)		
Very much	0 (0)	11 (37)		
3. Sexual feelings at present time			16.46	0.01**
Absent	1 (9)	5 (17)		
Weak	4 (36)	7 (23)		
Moderate	1 (9)	8 (27)		
Strong	5 (46)	10 (33)		
4. Frequency of sex relations at the present time			34.56	0.01**
Never	9 (82)	15 (49)		
Once a month	1 (9)	8 (27)		
Once a week	1 (9)	2 (7)		
More than once a week	0 (0)	5 (17)		
5. Awareness of any decline in sexual interest or activity			0.19	0.65
Yes	7 (64)	20 (67)		
No	4 (36)	10 (33)		
6. At what age first noted (5-year age bracket)			2.25	0.52
No problem	4 (36)	10 (33)		
1-5 years ago	4 (36)	9 (30)		
6-10 years ago	2 (18)	7 (24)		
More than 10 years ago	1 (9)	4 (13)		
7. If sex relations stopped, when stopped?			41.86	0.01**
Not stopped	0 (0)	8 (27)		
1-3 years ago	2 (18)	8 (27)		
2-5 years ago	3 (27)	7 (23)		
6-10 years ago	3 (27)	4 (13)		
11+ years ago	3 (27)	3 (10)		
8. Reasons for stopping sex relations			51.36	0.01**
Not stopped	0 (0)	8 (27)		
Death of partner	0 (0)	2 (7)		
Illness of self	4 (36)	5 (17)		
Illness of partner	0 (0)	1 (3)		
Self no longer able to perform sexually	1 (9)	4 (13)		
Separation or divorce from spouse	2 (18)	5 (17)		
Lack of opportunity	4 (36)	5 (17)		
9. What medications do you take?			0.43	0.51
Accurate response	8 (73)	23 (77)		
Don't know	3 (27)	7 (23)		
10. Do your medications cause any sex problems?			37.78	0.01**
Yes	2 (18)	18 (60)		
No	6 (55)	9 (30)		
Don't know	3 (27)	3 (10)		

Table 6

Needs identified by staff and patients in separate studies

	Present study ( <i>n</i> = 30)	Slade <i>et al.</i> 1996 ( <i>n</i> = 47)	Simons & Petch 2002 ( <i>n</i> = 173)
Staff responses, <i>n</i> (%)			
Intimate relationships	13 (43)	14 (29)	26 (27)
Sexual expression	3 (10)	8 (16)	0
Patient responses, <i>n</i> (%)			
Intimate relationships	25 (83)	10 (20)	26 (26)
Sexual expression	27 (90)	13 (27)	0







4. To uncover some of the obstacles to the expression of sexuality in people with serious mental illness living in the community.
5. To present recommendations for mental health practice, education and research.

## Methods in practice

### Sampling

A recognized sampling procedure was followed (Kuzel 1992). The population under study consisted of people with a case-note diagnosis of schizophrenia, living in the community, and regularly attending a depot clinic in North London to receive anti-psychotic medication.

The following selection criteria were used:

- case note diagnosis of schizophrenia;
- aged between 16 and 64 years;
- taking neuroleptic medication;
- regularly attending the depot clinic.

The participants were recruited from one depot clinic. A list of attendees was obtained from the Locality Team administrator and totalled 47. Participants were conveniently recruited to take part in the study. An office was provided in the clinic that afforded privacy where participants were informed about the study and consent sought, The Community Psychiatric Nurses administering the injection would make a clinical judgement on the suitability or wellness of the person to take part in the study.

### Ethical considerations

Approval for the study was gained from the local hospital ethics committee (Reference

number: P98210). At all times the investigator was respectful of the participants while administering the questionnaires. Time was invested in explaining the purposes and aims of the research. Participants were assured that should they wish to stop the interview for any reason they would be free to do so. Furthermore, participants were guaranteed anonymity and were informed that names would not appear on any documents. They would not be identifiable. The data from the study, which included the completed questionnaires, were kept in secure locked cupboards within the University. Written consent was gained. Participants were invited to ask questions and seek clarification before beginning the interviews and at the end.

\*\*\*Table 1 here \*\*\*

#### Data collection

Three questionnaires were used to obtain data.

**Table 1**  
Demographic characteristics

Variables	<i>n</i> (%)
Sex	
Male	15 (50.0)
Female	15 (50.0)
Age: mean (SD)	40.93 (10.01)
Ethnicity	
White UK	14 (46.7)
Black Caribbean	8 (26.7)
White European	5 (16.7)
Black African	2 (6.7)
Indian	1 (3.3)
Sexual orientation	
Heterosexual	22 (73.3)
Gay	2 (6.7)
Lesbian	5 (16.7)
Bisexual	1 (3.3)
Further education	
College	14 (46.7)
University	7 (23.3)
None	9 (30.0)
Marital status	
Single	16 (53.3)
Ever married	14 (46.7)
Relationship now	
Yes	12 (40.0)
No	18 (60.0)
Children	
Yes	10 (33.3)
No	20 (66.7)
Clinical diagnosis	
Schizophrenia	20 (66.7)
Schizo-affective disorder	10 (33.3)
Service contact (years)	
0-5	6 (20.0)
6-10	10 (33.3)
11-15	2 (6.7)
16-20	4 (13.3)
>20	8 (26.7)
Previous admissions: mean (SD)	5.53 (5.06)