



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** Dhakal, R., Huntjens, B., Shah, R., Lawrenson, J. & Verkicharla, P. K. (2022). Does the spectral composition of an ambient light vary between indoors and outdoors "in myopia perspective"? *Investigative Ophthalmology & Visual Science*, 63(7), ISSN 0146-0404

This is the published version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/29248/>

**Link to published version:**

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

City Research Online:

<http://openaccess.city.ac.uk/>

[publications@city.ac.uk](mailto:publications@city.ac.uk)

---

# Does the spectral composition of an ambient light vary between indoors and outdoors “in myopia perspective”? | IOVS

 [iovs.arvojournals.org/article.aspx](https://iovs.arvojournals.org/article.aspx)

ARVO Annual Meeting Abstract | June 2022

Does the spectral composition of an ambient light vary between indoors and outdoors “in myopia perspective”?

Rohit Dhakal; Byki Huntjens; Rakhee Shah; John Lawrenson; Pavan Kumar Verkicharla

Author Affiliations & Notes

Investigative Ophthalmology & Visual Science June 2022, Vol.63, 1890 – A0019. doi:

Abstract

**Purpose** : Considering that differences in the characteristics of ambient light between indoor and outdoor environment may impact myopia development in children, this study investigated the spectral composition of ambient light in different a) outdoor/indoor locations, b) times of a day, and c) seasons.

**Methods** : The spectral power distribution (SPD) was recorded using a hand-held spectrometer in three outdoor (‘open playground’, ‘under shade of tree’, and ‘canopy’) and three indoor locations (‘room with multiple windows’, ‘closed room’, and ‘closed corridor’) at five different time points (3-hour intervals between 6:30 and 18:00 clock-hours) on two days, each during summer and monsoon seasons. The pattern and the distribution of SPD across the spectrum, namely the short (380-500 nm), middle (505-565 nm) and long wavelengths (625-780 nm) were further assessed to understand the percentage composition in different locations.

**Results** : The overall median SPD (IQR [25th-75th percentile] W/nm/m<sup>2</sup>) across three outdoor locations (0.11 [0.09,0.12]) was 157 times higher than indoor locations (0.0007 [0.0001,0.001]). A considerable locational, diurnal and seasonal variation was observed in the distribution of median SPD value, with the highest value recorded in the ‘open playground’ (0.27 [0.21,0.28]) followed by ‘under shade of tree’ (0.083[0.074,0.09]), ‘canopy’ (0.014[0.012,0.015]), and ‘room with multiple windows’ (0.023[0.015,0.028]). While the spectral power of short, middle, long, and blue wavelengths was significantly lower in indoor compared to outdoor locations, the pattern and the relative percentage composition of short, middle and long wavelengths were similar in both the outdoor and indoor locations.

**Conclusions** : Irrespective of variation in SPD values with location, time, day and season, outdoor locations exhibited significantly higher SPD than indoor locations. The relative percentage composition of short, middle and long wavelengths of light are similar across all the locations. From a myopia perspective, our findings indicate a possibly

greater importance of the spectral power of ambient light rather than its spectral composition, as the latter was similar across outdoor and indoor locations. This finding warrants further research to understand its causal association with myopia.

This abstract was presented at the 2022 ARVO Annual Meeting, held in Denver, CO, May 1-4, 2022, and virtually.

This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/).



Copyright © 2015 Association for Research in Vision and Ophthalmology.

Copyright © 2015 Association for Research in Vision and Ophthalmology.