



City Research Online

City, University of London Institutional Repository

Citation: Cormican, A., Hirani, S. P. & McKeown, E. (2022). Healthcare professionals' perceived barriers and facilitators of implementing clinical practice guidelines for stroke rehabilitation: A systematic review. *Clinical Rehabilitation*, 37(5), pp. 701-712. doi: 10.1177/02692155221141036

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/29473/>

Link to published version: <https://doi.org/10.1177/02692155221141036>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Healthcare professionals' perceived barriers and facilitators of implementing clinical practice guidelines for stroke rehabilitation: A systematic review

Clinical Rehabilitation

1–12

© The Author(s) 2022



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/02692155221141036

journals.sagepub.com/home/cre

Adrienne Cormican¹ , Shashivadan P Hirani²,
and Eamonn McKeown³

Abstract

Objective: To identify healthcare professionals' perceived barriers and facilitators to clinical practice guideline implementation within stroke rehabilitation.

Data sources: CINAHL, MEDLINE, EMBASE, AMED, Cochrane library, Academic Search Complete and Scopus. Additional papers were identified through hand searching.

Review methods: The review followed the Preferred Reporting Item for Systematic Reviews and Meta-Analysis Protocols systematic review approach. Any empirical research that provided qualitative data on healthcare professionals' perceived factors influencing clinical guideline implementation in stroke rehabilitation was included. One reviewer screened all titles and abstract reviews ($n = 669$). Another two reviewers independently screened 30% of title and abstract reviews, followed by full-text reviews ($n = 61$). Study quality was assessed using the mixed-method appraisal tool.

Results: Data from 10 qualitative, six quantitative and six mixed-method studies published between 2000 and 2022, involving 1576 participants in total, were analysed and synthesised using modified thematic synthesis approach. The majority of participants were therapists $n = 1297$ (occupational therapists, physiotherapists, speech and language therapists). Organisational factors (time constraints, resources) alongside healthcare professionals' lack of knowledge and skills were the most cited barriers to guideline implementation. Contradictory attitudes and beliefs towards stroke guidelines applicability to real-life clinical practice and their evidence base were reported. Organisational support in the form

¹Occupational Therapy Department, Kings College Hospital, London, UK

²Centre for Health Services Research City, University of London, London, UK

³Health Services Research & Management Division, School of Health Sciences City, University of London, London, UK

Corresponding authors:

Adrienne Cormican, Occupational Therapy Department, Level 1 Neurosciences Gym, Ruskin Wing, Kings College Hospital, Denmark Hill, London, SE5 9RS, UK.

Email: acormican1@nhs.net

Shashivadan P Hirani, Centre for Health Services Research City, University of London, Northampton Square, London, EC1V 0HB, UK.

Email: shashi.hirani@city.ac.uk

of training, local protocols, performance monitoring and leadership were reported as perceived facilitators.

Conclusion: Barriers and facilitators are multifactorial and were identified at guideline, individual, team and organisational levels. There is a need to translate perceived barriers and facilitators into implementation interventions especially addressing organisational-level barriers.

Keywords

healthcare professional, clinical practice guideline, stroke rehabilitation, barriers, facilitators

Received December 6, 2020; accepted October 26, 2022

Introduction

Stroke clinical practice guidelines are supported by worldwide evidence of their positive impact on patient outcomes, healthcare costs, quality and effectiveness of stroke services.^{1–6} Such national and international guidelines are crucial components of the healthcare system's quality improvements.^{7–17} They are designed to support decision-making regarding the provision of evidence-based practice and its application^{7,9} thus improving rehabilitation practices. Despite their positive value, they are being underutilised,^{18–22} which can lead to suboptimal patient care with poorer functional recovery.^{2,4} This continues to be a pressing priority in research, management and policy.

An understanding of influential factors has long been advocated as an important component of guideline implementation.^{23,24} Healthcare professionals' perspectives and experiences represent as such influential factors influencing the effectiveness of clinical guideline implementation.²⁵ Comprehensive reviews^{26,27} and meta-reviews²⁸ found actual guideline, professional, patient and environmental characteristics all influence generic guideline implementation over a range of healthcare settings.

To date, research consists of studies evaluating stroke guideline adherence,⁶ intervention studies to improve adherence^{29–33} and cross-sectional surveys investigating implementation factors of stroke interventions.^{34–37} Recent systematic reviews on stroke clinical practice guidelines^{38,39} are limited in focus in acute stroke care and neither investigated healthcare professionals' views about stroke guideline implementation. Halls et al.⁴⁰ systematic review investigates

guideline use in stroke rehabilitation. However, it captures only the views of allied health professionals. No study has attempted to systematically analyse published primary studies of all healthcare professionals working in stroke rehabilitation and their perceived barriers and facilitators of guideline implementation to influence the adoption of guidelines in stroke rehabilitation.

Guideline factors were the most weighted domain influencing evidence-based acute stroke therapies³⁹ endorsing the need for further focused investigation. Effective stroke rehabilitation is multidisciplinary requiring input from several healthcare professionals. For successful guideline implementation, there is a need to better understand the complexity of changing clinical practice and guideline perceptions of all healthcare professionals working in stroke rehabilitation. Consequently, we conducted a systematic review to investigate and synthesise the current best evidence regarding the barriers and facilitators perceived by healthcare professionals in guideline implementation within stroke rehabilitation.

Methods

This systematic review was conducted according to the preferred reporting item for systematic reviews and meta-analysis protocols (PRISMA-P) systematic review approach⁴¹ as outlined in Supplemental Appendix 1 and complies with the Centre of Research and Dissemination Guidelines.⁴² A modified thematic synthesis⁴³ was adopted following the guidelines and reporting standards identified by PRISMA.⁴⁴ This approach⁴³ has been successfully

used within several systematic reviews involving both qualitative and quantitative study designs to address questions about people's perspectives and experiences.^{45,46}

To identify relevant studies, we searched CINAHL, MEDLINE, PsycINFO, EMBASE, AMED, Cochrane Library, Academic Search Complete and Scopus in June 2022. No date limits were applied and databases were searched from the conception date. Additional papers were identified through hand searching reference lists of papers selected for the full-text review. Our search strategy was generated following consultation with an experienced librarian and using search terms developed in previous studies.³⁹ A full search strategy is available in Supplemental Appendix 2. This search strategy was adapted for the other literature databases cited in Figure 1. MeSH terms (or equivalent) and free text keywords were used in combination (using Boolean operators) to systematically search the databases mentioned above. Keywords and phrases used included: 'Clinical practice guideline*' AND 'stroke' OR 'cerebrovascular accident' OR 'stroke rehabilitation' AND 'Barrier*' OR 'Factor*' OR 'Facilitat*' AND 'Perception*' OR 'Perspective*'.

A study was eligible for inclusion if it met the following selection criteria: (i) a fully published article in a peer-reviewed journal, (ii) qualitative, quantitative (cross-sectional studies) and mixed-methods empirical research, (iii) healthcare professionals working in any stroke rehabilitation setting, and (iv) focus on investigating perceived factors (barriers and/or facilitators) of clinical guideline implementation in stroke rehabilitation. Studies about stroke-specific rehabilitation interventions included in clinical practice guidelines were included if the study described the clinical guideline and its implementation. Studies were excluded if they were: (i) unavailable in English, (ii) a study population primarily of non-healthcare professionals (patients, families, carers), (iii) healthcare professionals' qualitative perceived factors were not extractable, and (iv) clinical practice guidelines that focused on the medical stroke management or prevention (drug therapy, surgery).

Results were downloaded and imported into Pro-Quest RefWorks citation management system

first to remove duplicates and then imported into Rayyan.⁴⁷ Study selection followed a two-step process. Initially, the lead author (AC) screened all titles and abstracts and identified studies for inclusion based on the eligibility criteria checklist. When titles and abstracts had insufficient information to support the screening, a full-text reading was conducted. Two independent reviewers (SH and EM) screened 30% of all titles and abstracts separately, followed by a consensus exercise involving all three authors to resolve discrepancies and arrive at a list of articles for full-text review.⁴⁸ In the second step, the lead author (AC) independently reviewed these full-text selected studies. Studies that met the inclusion criteria were selected for inclusion in the final analysis. The primary reason for the exclusion of full-text selected studies was recorded. See Figure 1.

The following relevant information was extracted from included studies and summated into one document before analysis: author(s), year of publication, country of study, clinical setting and clinical practice guidelines, intervention, study aim, design, data collection methods, participants and key findings on the barriers and facilitators identified to guideline implementation in stroke rehabilitation (see Supplemental Appendix 3). This approach was chosen to provide readers with a broad understanding of each article, how barriers and facilitators were examined, and the type of stroke guideline or implementation intervention being discussed.

As included studies were heterogeneous, the mixed-method appraisal tool⁴⁹ was used to assess the methodological quality of each study retained for this complex systematic review using different evaluation criteria for quantitative, qualitative and mixed-method studies. The specific criteria expose the factors that influence the risk of bias and transparency enabling the lead author (AC) to assess the relevance and rigour of all included studies (see Supplemental Appendix 4). While quality rating was not set as an inclusion criterion, this tool was used to help to inform the confidence with which study results could be interpreted in the synthesis of this review.

The guidelines for thematic synthesis⁴³ were followed to identify, extract and synthesise key factors from the selected papers according to our review

questions regarding the perceived barriers and facilitators of clinical practice guideline implementation in stroke rehabilitation. All perceived barriers and facilitators found in primary studies were labelled as such in the 'results' or 'findings' section of study reports. Barriers and facilitators were analysed and discussed collectively. The individual factors that represented barriers and facilitators were represented as initial codes. These initial codes were organised based on recurrency, similarities and differences to construct higher-level 'descriptive' themes, which remained close to the primary studies. Finally, overarching analytic themes were developed against the background of the review questions regarding how to promote effective guideline implementation amongst healthcare professionals in stroke rehabilitation. The lead author (AC) was responsible for all data extraction, quality appraisal, and data synthesis. Another author (SH) independently extracted and validated data from five randomly selected studies to ensure consistency with primary studies. SH also blindly assessed for matching themes. The agreement was 100%.

Results

Twenty-two studies published between 2000 and 2020 met the inclusion criteria.^{50–71} Their key characteristics are presented in Supplemental Appendix 3. Figure 1 depicts a PRISMA flow diagram documenting study selection and exclusion at all stages. The 22 studies provided data from healthcare professionals ($n = 1576$) from seven different countries (Australia, Canada, Ireland, Netherlands, New Zealand, UK, and Iceland). The majority of participants were therapists $n = 1297$ (occupational therapists, physiotherapists, speech and language therapists). The majority of therapists were physiotherapists with over 600 participants, followed closely by speech and language therapists ($n = 523$). Other participants included nurses ($n = 162$), doctors ($n = 31$), managers ($n = 35$), healthcare assistants ($n = 23$), therapy rehab assistants ($n = 7$), social workers ($n = 2$), orthoptics ($n = 2$) and dieticians ($n = 2$). Nine studies targeted single professional disciplines only; physiotherapists,^{60,62,70} speech and language therapists,^{51–54,71} nursing staff^{58,61,65} and other studies involved a combination of therapists.^{55,63,66,67,68} Six studies included a mix

of healthcare professionals working as part of multidisciplinary teams.^{50,56,57,59,64,69} Participants recruited in included studies worked in a range of clinical settings along the post-stroke continuum of care – see Figure 2.

Four studies investigated influential factors around national stroke guideline implementation in Ireland^{59,60} and Australia.^{51,53} The remaining studies investigated specific clinical practice guideline recommendation implementation,^{50,52,54–58,61–71} focusing on the upper limb, lower limb, aphasia, balance, communication, swallow, neglect, urinary continence, falls, depression, pain and education. All but one study⁵² examined both perceived barriers and facilitators of the guideline itself and the implementation process/interventions.^{56–58,61,64–66} Barriers and facilitators identified are consistent across quantitative, qualitative and mixed-method studies and were reported sometimes together as a continuum and not exclusively – see Figure 2. Thus, for this present review, healthcare professionals perceived barriers and facilitators to clinical practice guideline implementation were analysed and discussed jointly under themes. Figure 2 displays the six themes and their sub-themes that emerged.

Organisational factors were the most commonly reported factor influencing guideline implementation. Time and resources were the only factors to be reported in all 22 studies, predominantly as a barrier but also as a facilitator if the provision of time was allocated for guideline implementation and adequate availability of resources such as staffing, equipment, assessments, space, funding and educational material. Time constraints link directly with organisational processes and healthcare professionals' prioritisation of guideline recommendations. Unanimously, healthcare professionals' prioritisation of competing clinical and non-clinical tasks negatively interfered with guideline implementation.^{50–52,54,55,58,63,64,68,69,71} This was not just in acute settings where time and the acute medical model is perceived to be a significant limitation but across the continuum of stroke rehabilitation.

A lack of organisational processes acts as a negative influence on guideline implementation^{50,51,55,58–61,63,64,68–71} by way of insufficient training/education, lack of protocols to influence




Figure 1. Preferred reporting item for systematic reviews and meta-analysis (PRISMA) flowchart of the study retrieval and selection process.

Author	Setting	Guideline characteristics				Organisational support & capacity for change			Health care professionals' competence and beliefs			Professional interaction		Patient characteristics			Implementation intervention characteristics						
		CFG Specificity	CFG practicality	Applicability of CFG	CFG evidence base	Time & resources	Organisational Processes	Management & leadership	Organisational culture	Skills and knowledge	Prioritization	Attitudes on CFG/ Implementation	Teamwork	Professional collaboration	Professional roles	Impairments	Engagement	Access	Benefit	Facilitation	Tailored/practical intervention	Regular behaviour regulation	
Bayley [56]	RC			x	x	x		x		x	x	x	x	x	x					✓	✓		
Munsee [57]	RC			✓x		x	✓	x		✓x		x	✓x	✓x									
Donnellan [59]	Acute	x	✓x	✓x		✓x	✓x	✓x	✓			✓			x								
Mudge [63]	Acute	✓x	✓x	✓x	x	✓x	✓x	✓x		✓x	x	✓x	✓	✓	x			✓x					
Donohue [60]	Acute					x	x																
McChuskey [50]	Acute		x			✓x	✓x			✓x	x	✓x			x	✓							
Miao [51]	Acute	✓x	✓x	✓x	x	x	✓x	✓x		✓x	x	✓x	✓	✓	✓x	x							
Haalely [53]	Variety	✓	✓x	✓x	✓x	✓x	✓x	✓	✓x	✓x		✓x	✓	✓	✓x	x							
Bjartmarz [66]	Acute		✓	✓		✓x	✓	✓	✓x	✓x		✓	✓	✓x	x								
Hafsteinsdottir [61]	Variety		✓x	✓x		✓x	✓x	✓x						x		✓							
Shrubsole [54]	Acute, RC					✓x	✓	✓x	✓	✓x	x	✓x	✓x		✓x	x	✓x	✓					
Fisher [58]	Acute, RC			✓		✓x	✓x			✓x	x	✓x	✓	✓	✓x	✓					✓	✓	
Jolliffe [55]	Variety	x		x		✓x	✓x	✓	✓x	✓x	x	✓	✓x	✓	x	✓x	✓						
Clarke [64]	Acute				x	✓x	✓x			x	✓x	x	✓		x	✓							
Otterman [62]	Acute	✓	✓x	✓x		x				✓x													
Hickey [52]	Acute		x			x	✓		x		x	✓			x	x			✓			✓	✓
Jolliffe [66]	Variety					✓x	✓			✓x		✓								✓x	✓x	✓	✓
Connell [67]	Acute, RC			✓		✓x	✓	✓		✓x		✓x	✓							✓	✓	✓	✓
Cahill [68]	Acute, RC				✓	✓x	✓x	✓x		✓x	x	✓			x	✓x							
Inness [69]	Acute, RC				✓	✓x	✓x	✓	✓x	✓x	x	✓											
Auchstaetter [70]	Variety				✓	✓x	✓x	✓x	✓x	✓x	x	✓			✓	✓			✓x				
Chang [71]	Variety	x			✓	x	✓x	x	x	x	✓x	✓x			✓x	x	x	x					✓x

✓ indicates a particular facilitator reported by the authors. X indicates a particular barrier reported by the authors.
 RC indicates rehabilitation centre. Variety includes acute, rehabilitation centres and community stroke settings.

Figure 2. Domain of influential factors (perceived barriers and/or facilitators) under themes.

guideline use, lack of specific performance monitoring, poor service structure with daily processes that negatively influence staff prioritisations of recommended interventions and poor documentation systems. Organisational processes perceived as facilitators in 18 studies included organised dissemination of guideline information, consistent training of recommended interventions, alignment of services with guideline recommendations, local protocols, regular performance monitoring and action planning (audits, benchmarking, quality improvement projects), reminders and positive reinforcement for using recommended interventions.^{50,51,53–55,57–59,61,63–66,68–71}

Management and leadership were expressed as facilitators by actively supporting the organisational process mentioned above and protecting staff time for implementation creating a high level of organisational priority and commitment to guideline implementation resulting in a positive organisational culture.^{53–55,59,67,68,69}

Implementation interventions described in eight studies to increase the uptake of stroke guidelines included change management strategies,^{52,56,57} training and education for that specific stroke rehabilitation guideline,^{52,56–58,61,65,66} auditing,^{52,58,65} facilitators/opinion leaders/champions^{57,58,65–67} and

reminders.^{57,58,65} It is not the objective of this systematic review to determine which interventions are most effective, but due to the heterogeneity of the above-mentioned studies, it would be challenging to stipulate which interventions work better over others.

Healthcare professionals' competence (knowledge and skills) was reported as an influential factor in 17 studies. Knowledge and skills barriers identified ranged from a lack of familiarity with and theoretical knowledge behind guideline recommendations to a lack of knowledge and skills of how to modify and transfer guidelines to own local context and implement guideline recommendations.^{50,51,53,54,56,57,61–64} Greater knowledge of the guideline content, clinical skills and experience using recommendations facilitates guideline implementation in practice.^{50,51,53–55,62,63,65–67} Clinical experience and further education were found to be bi-directional. It acted as a facilitator^{51,53} with experienced clinicians being aware of and confident in implementing guidelines but also a barrier⁵¹ as changing their clinical practice habits can be difficult.

Healthcare professionals' beliefs and attitudes towards guideline implementation were heavily discussed in 19 studies. Eight studies recognised that healthcare professionals' attitude towards

guidelines was bi-directional and a further three studies found guidelines only viewed positively. Clinical practice guidelines were perceived as valuable; as a platform to set practice standards, as an audit tool,⁵⁹ as observed effective interventions,^{54,55,65} for improving patient outcomes,⁵⁰ in ratifying and advocating best practice^{51,53,67} and thus acted as a facilitator for their use. Observing the benefit of using guideline recommendations with patients^{52,55,61,64,68,70} provides healthcare professionals with positive reinforcement and thus acts as a facilitator for continued implementation. Healthcare professionals' negative attitudes related to their own lack of confidence^{54,55,67-71} in providing recommended interventions, patient safety concerns^{67,69,70} and effectiveness of recommended interventions.^{50,54,55} Therapists' mistrust and wariness regarding the evidence base supporting such guidelines was highlighted as a barrier to implementation in five studies.^{51,53,56,63,64}

Contradictory beliefs were identified in relation to guideline characteristics in 13 studies investigating both national and specific guideline recommendations. Guideline applicability was the most commonly described guideline characteristic often negatively perceived^{50,51,53,55,57,59,61-63} as 'too idealistic', not feasible, too time intensive for the 'real world', linking to a prominent issue of time. Reported facilitators for guideline application into clinical practice included clear, concise, easy to read and navigate recommendations, incorporating prioritised time frames for each recommendation, integrating recommendations into existing routines, and aligning guidelines with working routine practice and models of care. Guideline specificity, practicality and evidence base influenced clinical practice guideline applicability.

Patient characteristics as perceived barriers to guideline implementation were mainly due to the severity of clinical characteristics.^{51-57,64,68,71} The main patient-related facilitator identified was patient and family buy-in and engagement in recommended interventions.^{50,51,54,55,58,61,63,68,70,71} Patient access (their readiness and availability to participate in therapy) was briefly mentioned as a barrier in acute settings^{54,64,71} and as a facilitator across the continuum of stroke settings⁵⁵ with regular access and exposure to appropriate patients.

Professional interactions were the least described barriers and facilitations in included studies. In response to the lack of collaboration as a barrier, many studies described facilitators such as multidisciplinary working parties and guideline committees, and collaboration with patients and with healthcare professionals at different levels.^{51,53} The absence or presence of key authoritative individuals, for example, champions to coordinate implementation efforts were noted as perceived factors.^{55,57,58,67,68} Teamwork was recognised as facilitating guideline implementation through regular team discussions and communication, workload distribution, sharing responsibilities, understanding, engagement and commitment.

Discussion

The results of this systematic review substantiate the results of earlier stroke^{6,39} and generic literature^{23,27,28} highlighting translation of clinical guidelines into improved clinical practice remains challenging and perceived barriers and facilitators affecting stroke guideline implementation are multifactorial and multi-level. Organisational practice barriers (time, resources and a lack of supporting organisational processes) alongside professional barriers mainly healthcare professionals' competence (knowledge and skill) in guideline-recommended interventions were the most cited in this review. This is echoed in previous research relating to stroke guidelines and evidence-base practice adherence.^{6,39,40}

There remains an over-reliance on identifying and targeting professional level barriers only especially with educational approaches.^{63,68} Education in isolation is not an effective intervention for facilitating practice change.^{64,72} Indeed, this systematic review reflects an emphasis on education as a trialled implementation intervention^{52,56-58,61,65} with a lack of facilitators addressing organisational barriers. One study⁶⁶ builds on the evidence that active and multicomponent approaches are more effective in behaviour change and uptake of clinical guidelines over the educational approach alone.⁷² The findings also draw a parallel to Halls et al.'s⁴⁰ review, which noted that allied health professionals working on stroke are trying to use different active and multifaceted implementation strategies.

Interventions aimed at the organisational and managerial levels can facilitate guideline implementation.^{23,73–75} However, evidence on organisational-orientated interventions remains lacking in comparison with professional-oriented interventions.^{76,77} This review found organisational processes (training, dissemination, local protocols, performance monitoring) were heavily weighted influential factors in guideline implementation. Leadership and management were recognised as facilitators to support organisational processes and create a positive organisational culture. Leadership in the form of opinion leaders/champions/site facilitators are proven effective implementation tools in changing behaviour.^{66,73,77,78} Performance monitoring (auditing and feedback) has shown promise in changing healthcare professionals' behaviour^{31,66,72,79,80} and found effective in improving stroke clinical guideline application amongst allied healthcare professionals.⁴⁰ Such interventions not only minimise contextual barriers but also promote positive changes in attitudes, behaviours and organisational culture. A greater effort needs to be prioritised to research supporting organisational-level implementation interventions.

Stroke rehabilitation demands collaborative care and thus, team-based implementation interventions should be considered. Two studies in this review^{56,57} found decreased multidisciplinary team communication limited guideline implementation. Multidisciplinary team level interventions can develop a shared understanding, improve team functioning and effect change in guideline implementation.⁷³ Systemic challenges in organisational and team level barriers need to be addressed, then individual healthcare professionals will be more empowered to apply guideline recommendations.

The most frequently reported guideline characteristic in this review concerns 'applicability' to real-life stroke patients and clinical practice similar to previous literature.^{6,23,28,38,39} This review underlined the prominent issue of healthcare professionals' lack of time, mainly the prioritisation of competing clinical demands and the difficulties associated with impairment severity and co-morbidities. Halls et al.⁴⁰ endorse this finding of a perceived poor fit between stroke rehabilitation guidelines, patient characteristics

and healthcare systems' way of working. It is imperative for guideline developers to take into consideration the applicability of guidelines and present practical recommendations. Involving healthcare professionals in guideline development boosts the chances of successful implementation.⁸¹ An example of important collaborative work in the United Kingdom has been led by the James Lind Alliance.⁸² This national implementation strategy has helped to reveal a shared vision and facilitate collective action towards key priority areas in stroke rehabilitation as identified and shared by stroke clients, carers, researchers and healthcare professionals.

This review found contradictory attitudes and beliefs towards the evidence-base supporting stroke guidelines emphasising a continued relevant factor in guideline implementation research.^{6,23,28,38,39} Jolliffe et al.⁸³ and Halls et al.⁴⁰ also report concerns regarding the methodological quality of stroke guidelines evidence base and the implications it poses for their clinical use. Guideline compliance appears to be higher when a strong scientific evidence base underpins them.⁸⁴ Strengthening the evidence base may be influential.

Tailored multicomponent implementation strategies addressing perceived barriers in local contexts remain wanting despite their evidence in eliciting change in clinical practice.^{66,72,73,76,85,86} Only three studies in this review^{52,58,66} investigated potential barriers and facilitators pre-implementation to design their tailored implementation interventions. Implementation interventions informed by behaviour change theories⁸⁷ and stronger study designs are warranted to validate the effectiveness of specific implementation interventions as such evidence remains contradictory due to low-certainty evidence.^{72,73} Additional improvements to implementation design such as design based on known barriers,^{85,88–89} stakeholder engagement,⁹⁰ participatory action research approach⁹¹ and co-research⁹² should be considered to ensure designs are fit for purpose.

Despite this review following a systematic and transparent approach, it is possible that relevant studies were missed as not every available database was utilised. Databases were selected based on quality and relevancy. The substantial duplication in studies obtained gives a degree of confidence that the main papers that are indexed have been included. Guideline implementation efforts may have occurred

as local quality improvement projects and therefore, were not published in peer-reviewed journals. Boundaries set around inclusion criteria for this review were necessary and as a minimum quality standard. Peer-reviewed sources are likely to be larger and more impactful to the broader profession. The heterogeneity of the studies including the thin data and description of perceived barriers and facilitators in quantitative and mixed-method studies may have been a barrier. However, the included studies show a reasonable consistency of results, which was drawn out using an explicit approach to appraise, combine and synthesise complex data and thus adds to the confidence in the conclusions of this review.

The current review is the first to specifically detail healthcare professionals perceived barriers and facilitators of implementing clinical practice guidelines in stroke rehabilitation. The findings add to the knowledge base and can inform future clinical practice guideline implementation interventions in stroke rehabilitation. The results of this review stress the importance of healthcare professionals' experiences, views and their input in guideline policy and clinical management to address barriers and develop strategies to optimise the implementation of guidelines in stroke rehabilitation. Understanding perceived barriers and facilitators will enable managers, commissioners, and policymakers to deliver more focused and realistic recommendations, support and guidance to healthcare professionals.

Clinical messages

- Tailored multifaceted implementation approaches addressing local contextual barriers can improve guideline implementation.
- Perceived organisational facilitators for guideline implementation include consistent training, services and local protocols aligned with guideline recommendations, regular performance monitoring and action planning.
- Effective management and leadership (guideline champions) can facilitate organisational commitment and priority to guideline implementation.

Declaration of conflicting interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the National Institute for Health Research and the Stroke Association Medical Research Council through a Pre-Doctoral Academic Fellowship (grant number NIHR300389).

ORCID iD

Adrienne Cormican  <https://orcid.org/0000-0001-6508-8363>

Supplemental material

Supplemental material for this article is available online.

References

1. Brusamento S, Legido-Quigley H, Panteli D, et al. Assessing the effectiveness of strategies to implement clinical guidelines for the management of chronic diseases at primary care level in EU member states: A systematic review. *Health Policy* 2012; 107: 168–183.
2. Hubbard IJ, Harris D, Kilkenny MF, et al. Adherence to clinical guidelines improves patient outcomes in Australian audit of stroke rehabilitation practice. *Arch Phys Med Rehabil* 2012; 93: 965–971.
3. Cadilhac D, Ibrahim J, Pearce D, et al. Multicenter comparison of processes of care between stroke units and conventional care wards in Australia. *Stroke* 2004; 35: 1035–1040.
4. Duncan PW, Horner RD, Reker DN, et al. Adherence to post-acute rehabilitation guidelines is associated with functional recovery in stroke. *Stroke* 2002; 33: 167–178.
5. Stroke Unit Trialists' Collaboration. Organised inpatient (stroke unit) care for stroke. *Cochrane Database Syst Rev* 2013; 9: CD000197.
6. Donnellan C, Sweetman S and Shelley E. Health professionals' adherence to stroke clinical guidelines: A review of the literature. *Health Policy* 2013; 111: 245–263.
7. Royal College of Physicians, Intercollegiate Stroke Working Party. *National clinical guideline for stroke*. 5th ed. London, UK, 2016.
8. National Clinical Guideline Centre. *Stroke Rehabilitation. Long term rehabilitation after stroke clinical guideline 162 methods, evidence and recommendations*, May 2013.
9. Stroke Foundation. *Clinical guidelines for stroke management 2017*. Melbourne, Australia: Stroke Foundation, 2017.

10. Stroke Foundation of New Zealand and New Zealand Guidelines Group. *Clinical guidelines for stroke management 2010*. Wellington, New Zealand: Stroke Foundation of New Zealand, 2010.
11. Bryer A, Connor MD, Haug P, et al. The South African guideline for the management of ischemic stroke and transient ischemic attack: Recommendations for a resource constrained health care setting. *Int J Stroke* 2011; 6: 349–354.
12. Winstein CJ, Stein J, Arena R, et al. Guidelines for adult stroke rehabilitation and recovery: A guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2016; 47: e98–e169.
13. Jauch EC, Saver JL, Adams HP, et al. Guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American heart association/American stroke association. *Stroke* 2013; 44: 870–947.
14. Hebert D, Lindsay MP, McIntyre A, et al. Canadian Stroke best practice recommendations: Stroke rehabilitation practice guidelines, update 2015. *Int J Stroke* 2016; 11: 459–484.
15. Scottish Intercollegiate Guidelines Network (SIGN). *118, Management of patients with stroke: Rehabilitation, prevention and management of complications, and discharge planning*. Edinburgh, Scotland: A National Clinical Guideline, 2010.
16. Irish Heart Foundation: Council for Stroke. *National clinical guidelines and recommendations for the care of people with stroke and transient ischaemic attack*. Ireland 2010.
17. Royal Dutch Society for Physical Therapy. (Koninklijk Nederlands Genootschap voor Fysiotherapie, KNGF). KNGF Guideline Stroke, 2014.
18. Harris D, Cadilhac DA, Hankey GJ, et al. National stroke audit: The Australian experience. *Clin Audit* 2010; 2: 25–31.
19. Stroke Foundation. *National stroke audit – rehabilitation services report*. Melbourne, Australia, 2018.
20. National Stroke Foundation. *National action plan for stroke*. Melbourne, Australia, 2012.
21. Sentinel Stroke National Audit Programme (SSNAP). *National results clinical audit*. UK: Kings College London, 2019.
22. The Scottish Stroke Care Audit (SSCA). *2015 Annual national report stroke services in Scottish Hospitals*. Edinburgh, Scotland, 2015.
23. Cabana MD, Rand SC, Powe NR, et al. Why don't physicians follow clinical practice guidelines? A framework for improvement. *JAMA* 1999; 282: 1458–1465.
24. Mickan S, Burls A and Glasziou P. Patterns of 'leakage' in the utilisation of clinical guidelines: A systematic review. *Postgrad Med J* 2011; 87: 670–679.
25. Kortteisto T, Kaila M, Komulainen J, et al. Healthcare professionals' intentions to use clinical guidelines: A survey using the theory of planned behaviour. *Implement Sci* 2010; 5: 51.
26. Grimshaw J, Eccles M and Tetroe J. Implementing clinical guidelines: Current evidence and future implications. *J Contin Educ Health Prof* 2004; 24: S31–S37.
27. Prior M, Guerin M and Grimmer-Somers K. The effectiveness of clinical guideline implementation strategies – a synthesis of systematic review findings. *J Eval Clin Pract* 2008; 14: 888–897.
28. Francke A, Smit M, de Veer A, et al. Factors influencing the implementation of clinical guidelines for health care professionals: A systematic meta-review. *BMC Med Inform Decis Mak* 2008; 8: 38.
29. McCluskey A, Ada L, Kelly PJ, et al. A behaviour change program to increase outings delivered during therapy to stroke survivors by community rehabilitation teams: The out-and-about trial. *Int J Stroke* 2016; 11: 425–437.
30. Tistad M, Palmacrantz S, Wallin L, et al. Developing leadership in managers to facilitate the implementation of national guideline recommendations: A process evaluation of feasibility and usefulness. *Int J Health Policy Manag* 2016; 5: 477–486.
31. Singh M, Hynie M, Rivera T, et al. An evaluation study of the implementation of stroke best practice guidelines using a knowledge transfer team approach. *Can J Neurosci Nurs* 2015; 37: 24–33.
32. Pennington L, Roddam H, Burton C, et al. Promoting research use in speech and language therapy: A cluster randomized controlled trial to compare the clinical effectiveness and costs of two training strategies. *Clin Rehabil* 2005; 19: 387–397.
33. Salbach NM, Wood-Dauphinee S, Desrosiers J, et al. Facilitated interprofessional implementation of a physical rehabilitation guideline for stroke in inpatient settings: Process evaluation of a cluster randomized trial. *Implement Sci* 2017; 12: 100.
34. Van Peppen RPS, Maissan FJS, Van Genderen FR, et al. Outcome measures in physiotherapy management of patients with stroke: A survey into self-reported use, and barriers to and facilitators for use. *Physiother Res Int* 2008; 13: 255–270.
35. Van der Weijden T, Hooi JD, Groi R, et al. A multidisciplinary guideline for the acute phase of stroke: Barriers perceived by Dutch neurologists. *J Eval Clin Pract* 2004; 10: 241–246.
36. Arias M and Smith LN. Early mobilization of acute stroke patients. *J Clin Nurs* 2007; 16: 282–288.
37. Hart S and Morris R. Screening for depression after stroke: An exploration of professionals' compliance with guidelines. *Clin Rehabil* 2008; 22: 60–70.
38. Craig LE, McInnes E, Taylor N, et al. Identifying the barriers and enablers for a triage, treatment, and transfer clinical intervention to manage acute stroke patients in the emergency department: A systematic review using the theoretical domains framework (TDF). *Implement Sci* 2016; 11: 57.
39. Baatiema L, Otim ME, Mnatzaganian G, et al. Health professionals' views on the barriers and enablers to evidence-based practice for acute stroke care: A systematic review. *Implement Sci* 2017; 12: 74.
40. Halls D, Murray C and Sellar B. Why allied health professionals use evidence-based clinical guidelines in stroke

- rehabilitation: A systematic review and meta-synthesis of qualitative studies. *Clin Rehabil* 2021; 35: 1611–1626.
41. Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: Elaboration and explanation. *Br Med J* 2015; 349: g7647.
 42. Centre for Reviews and Dissemination. *Systematic reviews, CRD's guidance for undertaking systematic reviews in health care*. UK: CRD, University of York, 2009.
 43. Thomas J and Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Med Res Methodol* 2008; 8: 45.
 44. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med* 2009; 6: e1000097.
 45. Thomas J, Sutcliffe K, Harden A, et al. *Children and healthy eating: A systematic review of barriers and facilitators*. London, UK: EPPI-Centre, Social Science Research Unit, Institute of Education, University of London, 2003.
 46. Harden A, Garcia J, Oliver S, et al. Applying systematic review methods to studies of people's views: An example from public health. *J Epidemiol Commun Health* 2004; 58: 794–800.
 47. Ouzzani M, Hammady H, Fedorowicz Z, et al. Rayyan – a web and mobile app for systematic reviews. *Syst Rev* 2016; 5: 210.
 48. Furlan AD, Pennick V, Bombardier C, et al. Updated method guidelines for systematic reviews in the Cochrane back review group. *Spine* 2009; 34: 1929–1941.
 49. Pluye P. Critical appraisal tools for assessing the methodological quality of qualitative, quantitative and mixed methods studies included in systematic mixed studies reviews [letter]. *J Eval Clin Pract* 2013; 19: 122.
 50. McCluskey A, Vratsistas-Curto A and Schurr K. Barriers and enablers to implementing multiple stroke guideline recommendations: A qualitative study. *BMC Health Serv Res* 2013; 13: 323.
 51. Miao M, Power E and O'Halloran R. Factors affecting speech pathologists' implementation of stroke management guidelines: A thematic analysis. *Disabil Rehabil* 2015; 37: 674–685.
 52. Hickey J, Shrubsole K, Worrall L, et al. Implementing aphasia recommendations in the acute setting: Speech-language pathologists' perspectives of a behaviour change intervention. *Aphasiology* 2019; 33: 606–628.
 53. Hadely KA, Power E and O'Halloran R. Speech pathologists' experiences with stroke clinical practice guidelines and the barriers and facilitators influencing their use: A national descriptive study. *BMC Health Serv Res* 2014; 14: 110.
 54. Shrubsole K, Worrall L, Power E, et al. Barriers and facilitators to meeting aphasia guideline recommendations: What factors influence speech pathologists' practice? *Disabil Rehabil* 2019; 41: 1596–1607.
 55. Jolliffe L, Hoffmann T and Lannin NA. Increasing the uptake of stroke upper limb guideline recommendations with occupational therapists and physiotherapists. A qualitative study using the theoretical domains framework. *Aust Occup Ther J* 2019; 66: 603–616.
 56. Bayley MT, Hurdowar A, Richards CL, et al. Barriers to implementation of stroke rehabilitation evidence: Findings from a multi-site pilot project. *Disabil Rehabil* 2012; 34: 1633–1638.
 57. Munce SEP, Graham ID, Salbach NM, et al. Perspectives of health care professionals on the facilitators and barriers to the implementation of a stroke rehabilitation guidelines cluster randomized controlled trial. *BMC Health Serv Res* 2017; 17: 440.
 58. Fisher A. Development of clinical practice guidelines for urinary continence care of adult stroke survivors in acute and rehabilitation settings. *Can J Neurosci Nurs* 2014; 36: 16–31.
 59. Donnellan C, Sweetman S and Shelley E. Implementing clinical guidelines in stroke: A qualitative study of perceived facilitators and barriers. *Health Policy* 2013; 111: 234–244.
 60. Donohue A, McLaughlin C, Crowe M, et al. Clinical guideline adherence by physiotherapists working in acute stroke. *Ir Med J* 2014; 107: 287–289.
 61. Hafsteinsdóttir TB, Varekamp R, Rensink M, et al. Feasibility of a nursing rehabilitation guideline for patients with stroke: Evaluating the use by nurses. *Disabil Rehabil* 2013; 35: 939–949.
 62. Otterman NM, van der Wees PJ, Bernhardt J, et al. Physical therapists' guideline adherence on early mobilization and intensity of practice at Dutch acute stroke units a country-wide survey. *Stroke* 2012; 43: 2395–2401.
 63. Mudge S, Hart A, Murugan S, et al. What influences the implementation of the New Zealand stroke guidelines for physiotherapists and occupational therapists? *Disabil Rehabil* 2017; 39: 511–518.
 64. Clarke DJ, Burton LJ, Tyson SF, et al. Why do stroke survivors not receive recommended amounts of active therapy? Findings from the ReAcT study, a mixed-methods case-study evaluation in eight stroke units. *Clin Rehab* 2018; 32: 1119–1132.
 65. Bjartmarz I, Jónsdóttir H and Hafsteinsdóttir TB. Implementation and feasibility of the stroke nursing guideline in the care of patients with stroke: A mixed methods study. *BMC Nurs* 2017; 16: 72.
 66. Jolliffe L, Hoffmann T, Churilov L, et al. What is the feasibility and observed effect of two implementation packages for stroke rehabilitation therapists implementing upper limb guidelines? A cluster-controlled feasibility study. *BMJ Open Qual* 2020; 9: e000954.
 67. Connell LA, McMahon NE, Harris JE, et al. A formative evaluation of the implementation of an upper limb stroke rehabilitation intervention in clinical practice: A qualitative interview study. *Implement Sci* 2014; 9: 90.
 68. Cahill LS, Carey LM, Mak-Yuen Y, et al. Factors influencing allied health professionals' implementation of upper limb sensory rehabilitation for stroke survivors: A qualitative study to inform knowledge translation. *BMJ Open* 2021; 11: e042879.

69. Inness EL, Jagroop D, Andreoli A, et al. Factors that influence clinical implementation of aerobic exercise in stroke rehabilitation: A theory-informed qualitative study. *Phys Ther* 2022; 102: 1–12.
70. Auchstaetter N, Luc J, Lukye S, et al. Physical therapists' use of functional electrical stimulation for clients with stroke: Frequency, barriers and facilitators. *Phys Ther* 2016; 96: 995–1005.
71. Chang HF, Power E, O'Halloran R, et al. Stroke communication partner training: A national survey of 122 clinicians on current patterns and perceived implementation barriers and facilitators. *Int J Lang Commun Disord* 2018; 53: 1094–1109.
72. Cahill LS, Carey LM, Lannin NA, et al. Implementation interventions to promote the uptake of evidence-based practices in stroke rehabilitation. *Cochrane Database Syst Rev* 2020; 10: CD012575.
73. Bird ML, Miller T, Connell LA, et al. Moving stroke rehabilitation evidence into practice: A systematic review of randomized controlled trials. *Clin Rehab* 2019; 33: 1586–1595.
74. Grol R. Successes and failures in the implementation of evidence-based guidelines for clinical practice. *Med Care* 2001; 39: 46–53.
75. Ranta A, Dovey S, Weatherall M, et al. Cluster randomized controlled trial of TIA electronic decision support in primary care. *Neurology* 2015; 84: 1545–1551.
76. Field MJ and Lohr K. *Guidelines for clinical practice: From development to use*. Washington DC: National Academic Press, 1992.
77. Flodgren G, Hall AM, Goulding L, et al. Tools developed and disseminated by guideline producers to promote the uptake of their guidelines (Review). *Cochrane Database Syst Rev* 2016; 8: CD010669.
78. Hamilton S, McLaren S and Mulhall A. Multidisciplinary compliance with guidelines for stroke assessment: Results of a nurse-led evaluation study. *Clin Eff Nurs* 2006; 9S1: e57–e67.
79. Ivers N, Jamtvedt G, Flottorp S, et al. Audit and feedback: Effects on professional practice and healthcare outcomes. *Cochrane Database Syst Rev* 2012; 6: CD000259.
80. Kristensen H and Hounsgaard L. Evaluating the impact of audits and feedback as methods for implementation of evidence in stroke rehabilitation. *Br J Occup Ther* 2014; 77: 251–259.
81. Sachs M. Successful strategies and methods of nursing standards implementation. *Pflege* 2006; 19: 33–44.
82. James Lind Alliance Priority Setting Partnerships. The James Lind Alliance: National Institute for Health Research, 2022. Available at: <https://www.jla.nihr.ac.uk/priority-setting-partnerships/Stroke/>
83. Jolliffe L, Lannin NA, Cadilhac DA, et al. Systematic review of clinical practice guidelines to identify recommendations for rehabilitation after stroke and other acquired brain injuries. *BMJ Open* 2018; 8: e018791.
84. Saillour-Glenisson F and Michel P. Individual and collective facilitators of and barriers to the use of clinical practice guidelines by physicians: A literature review. *Rev Epidemiol Sante Publ* 2003; 51: 65–80.
85. Grimshaw JM, Eccles MP, Lavis JN, et al. Knowledge translation of research findings. *Implement Sci* 2012; 7: 50.
86. Baker R, Camosso-Stefinovic J, Gillies C, et al. Tailored interventions to address determinants of practice. *Cochrane Database Syst Rev* 2015; 4: CD005470.
87. Michie S, Johnston M, Abraham C, et al. On behalf of the “psychological theory” group. Making psychological theory useful for implementing evidence-based practice: A consensus approach. *Qual Saf Health Care* 2005; 14: 26–33.
88. Medves J, Godfrey C, Turner C, et al. Systematic review of practice guideline dissemination and implementation strategies for healthcare teams and team-based practice. *Int J Evid Based Healthc* 2010; 8: 79–89.
89. Cane J, O'Connor D and Michie S. Validation of the theoretical domain's framework for use in behaviour change and implementation research. *Implement Sci* 2012; 7: 37.
90. Graham ID and Tetroe J. Some theoretical underpinnings of knowledge translation. *Acad Emerg Med* 2007; 14: 936–941.
91. Donetto S, Tsianakas V and Robert G. *Using experience based co-design (EBCD) to improve the quality of health-care: Mapping where we are now and establishing future directions*. Report. UK: King's College London, 2014.
92. Casey M, O'Leary D and Coghlan D. Unpacking action research and implementation science: Implications for nursing. *J Adv Nurs* 2018; 74: 1051–1058.