



City Research Online

City, University of London Institutional Repository

Citation: Gallus, C., Blasiak, P. & Pothos, E. M. (2023). Winning a CHSH game without entangled particles in a finite number of biased rounds: how much luck is needed?. *Entropy*, 25(5), 824. doi: 10.3390/e25050824

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/30479/>

Link to published version: <https://doi.org/10.3390/e25050824>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Article

Winning a CHSH game without entangled particles in a finite number of biased rounds: how much luck is needed?

Christoph Gallus ^{1,*} , Pawel Blasiak ^{2,3}  and Emmanuel M. Pothos ⁴ ¹ THM Business School, Technische Hochschule Mittelhessen, D-35390 Gießen, Germany² Institute for Quantum Studies, Chapman University, Orange, CA, USA; blasiak@chapman.edu³ Institute of Nuclear Physics, Polish Academy of Sciences, PL-31342 Kraków, Poland; pawel.blasiak@ifj.edu.pl⁴ Psychology Department, City, University of London, London EC1V 0HB, UK; Emmanuel.Pothos.1@city.ac.uk

* Correspondence: christoph.gallus@w.thm.de

Abstract: Quantum games, like the CHSH game, are used to illustrate the puzzle and power of entanglement. These games are played over many rounds and in each round the participants, Alice and Bob, each receive a question bit to which they each have to give an answer bit, without being able to communicate during the game. When all possible classical answering strategies are analyzed, it is found that Alice and Bob cannot win more than 75% of the rounds. **A higher percentage of wins arguably requires^{CG}** an exploitable bias in the random generation of the questions bits or access to “non-local” resources, like entangled pairs of particles. However, in an actual game the number of rounds has to be finite and question regimes may come up with unequal likelihood, so there always is a possibility for Alice and Bob to win by pure luck. This statistical possibility has to be transparently analyzed for practical applications such as the detection of eavesdropping in quantum communication. Similarly, when Bell tests are used in macroscopic situations to investigate the connection strength between system components and the validity of proposed causal models, the available data is limited and the possible combinations of question bits (measurement settings) may not be controlled to occur with equal likelihood. In the present work we give a fully self-contained proof for a bound on the probability to win a CHSH game by pure luck without making the usual assumption of only small biases in the random number generators. We also show bounds for the case of unequal probabilities based on results by McDiarmid and Combes and illustrate certain exploitable biases numerically.

Keywords: CHSH Games; Bell Inequalities; Secure Quantum Communication; Quantum Information

Citation: Gallus, C.; Blasiak, P.; Pothos, E.M. Winning a CHSH game without entangled particles in a finite number of biased rounds: how much luck is needed?. *Entropy* **2023**, *1*, 0. <https://doi.org/>

Academic Editors: Andrei Khrennikov and Karl Svozil

Received: May 15, 2023

Accepted: ...

Published: ...

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

John Bell's seminal ideas [1–6] have allowed a precise and quantifiable illustration that quantum mechanical entanglement allows experimental outcomes which cannot be generated classically. Such presentations of the quantum edge can come in different forms, for example the Mermin-Peres magic square game, the Greenberger-Horne-Zeilinger (GHZ) game or the Clauser-Horne-Shimony-Holt (CHSH) game. They also come with different story lines, such as hypothetical TV quizzes, polls or trials [7–13]. The goals of these presentations are different as well. They may concern teaching quantum information theory or illustrating the power of entanglement, for example for uses in quantum cryptography or quantum computation, as well as in discussions of foundational questions of locality and free choice. These games provide a good and accessible illustration of the philosophical puzzle and practical power of entanglement while using only terms from macroscopic experience. **In describing these games, it is important to use precise formulations and to deal with all possible objections in an understandable way. Cheats, like classical communication between the players of the game or allowing for an undue influence on the generation of questions, can allow the generation of a quantum-like Bell statistic with classical means. The amount of cheating that is required**

is not very large. The Toner-Bacon protocol [14] shows, for example, that when Alice and Bob share two uniformly distributed random variables with values on the unit sphere of \mathbb{R}^3 , then sending a single bit can be enough: sending one bit of information from Alice to Bob is enough to simulate the statistic of a local projective measurement on an entangled Bell pair state. However, one should not underestimate what the exchange of one classical bit of information can do in terms of generating strong correlations [15]. In [16] the information to be shared between measurement settings is discussed further illustrating that small amounts of shared classical information suffice, in order to break the Bell bound. Such work is based on the implicit assumption that an arbitrarily large number of rounds is played and that the question regimes (measurement settings) are produced by unbiased random number generators with equal likelihood. In general, it is quickly pointed out, when describing such games to a scientifically trained audience, that additional randomness arises in practice, because the observed averages may deviate from the mathematical expectation values when only a finite number of rounds is used. Additionally, in the implementation of actual quantum devices and the practical use of quantum technologies imperfect probability distributions may arise, for example an undesired drifting of states, or crosstalk between different qubits [17]. Detection and the possible correction of such issues is important in applications like the detection of eavesdropping in quantum cryptography [18], the fidelity of quantum random number generators [19], or the general witnessing of entanglement [20].

When using games as illustrations, it is therefore important to also treat the effect of purely random fluctuations when only a finite number of rounds is played. Sweeping the issue of a finite number of rounds N under the rug takes away a lot from the usefulness of such presentations. Equally it is important to consider imperfect probability distributions. Our contribution lies in a rigorous and accessible treatment of CHSH games and the chances of winning in a finite number of rounds with the additional difficulty that measurement settings cannot be ensured to occur with equal probabilities. **This topic is not only relevant for illustrative games, but has practical implications as well: in the implementation of quantum technologies, imperfect joint probability distributions may arise, such as an undesired drifting of states, or crosstalk between different qubits [17]. Detection and the possible correction of such issues is important in various applications like the detection of eavesdropping in quantum cryptography [18,30], the fidelity of quantum random number generators [19], or the general witnessing of entanglement [20].** ^{CG}

In the present work we only discuss CHSH games, for which all questions are generated freely and independently. We assume a finite number of game rounds and we do not insist that all measurement settings occur with equal probability. We have made this choice of topic for three reasons. First, the CHSH form can be tested well with current experimental technology and is useful for cryptography and the detection of eavesdropping [30]. Second, the CHSH form of the Bell Inequalities can be formulated symmetrically with expectation values and is so open for applications to macroscopic systems outside physics [31,32]. In such applications not only the number of rounds is naturally finite, but also no a priori reason for measurement settings to occur with equal likelihood may exist. The number of rounds can be quite low so that random fluctuations play a greater role than in photon-based physical experiments. This is particularly problematic when historical data is considered where the causal regime at work is suspected to have changed over time. Last, CHSH games are rigid, so that strategies with maximum success probability are isomorphic [33–35] allowing a generic way of treatment.

Statistical effects from random fluctuations over a finite number of rounds have of course previously been discussed in the literature. The main focus has been the analysis of Bell experiments to disprove local hidden variable models while not allowing for loopholes. Usually statistical confidence in the conclusion from Bell experiments is derived by observing that violations of Bell inequalities happen with a certain number of standard deviations, but this line of argument typically requires the assumption of a Gaussian error distribution and the independence of the game rounds, which makes it problematic [23].

We therefore do not follow this line of argument here, but start from early work by Gill [24,25] on this topic, which was later extended in [26]. Such later work yielded sharper bounds, but references, usually without proof, very technical mathematical tools (based on concentration inequalities from Hoeffding-Azuma, McDiarmid and others) with the goal of aiding specialists in designing and evaluating Bell experiments. Putting such technical work to use even by specialists may still have its pitfalls, as pointed out in [27]. Here we strive for an accessible presentation that yields interesting and illustrative bounds for situations with unequal probabilities for the measurement settings, when games are played over a finite number of rounds. In contrast to [26], we are interested in random number generator biases that can be very large and we want to avoid linear programming or martingale methods. ~~Our extension of the result in [25] is shown by a fully self-contained proof in the present paper. Striking a balance between the strength of the results and the accessibility of the required tools to obtain them is delicate: multipartite games do not always allow a quantum advantage [28] and general problems around multiplayer games with an advantage, like computing the maximum winning probability given an arbitrary amount of entangled particles, can be very complex and even QMA-hard [29].~~

The present paper is organized as follows: In Section 2, we present two versions of the CHSH game, one in which the players have to achieve a high percentage of winning rounds and one in which the players have to achieve a high S -value, and comment on the equivalence of the versions. In Section 3.1 we prove an extension of Gill's result [25] for the case of unequal probabilities. The proof is self contained and requires only an elementary knowledge of probability. We continue in Section 3.2 by proving a bound for unequal probabilities based on McDiarmid Inequality. We also simulate certain game strategies in Section 4 and illustrate some exploitable biases. The last section concludes with a short summary and discussion.

2. The CHSH Game

Using the notion of the CHSH game, the additional resource that entangled particles may bring can be easily illustrated without the need to understand quantum mechanics. To this end, imagine a game, which is played over many rounds, in which Alice and Bob work as partners trying to win. **We assume that they can^{CG} jointly receive a prize, which is awarded to them^{CG} by a Verifier based on a clear statistical pre-agreed rule after all game rounds have been played.** Alice and Bob may discuss various pre-arranged strategies before the game commences, but they are isolated and not allowed to communicate once the game has started. The game shall be played over N rounds and in each round Alice and Bob will each receive one of two possible inputs (called *questions*) to which they have to give one of two possible outputs (called *answers*). Once the information is brought together, the complete list of questions and answers can be used by an independent Verifier to generate a statistic from which he will determine whether Alice and Bob deserve the prize. We will present two versions of the CHSH game, one in which Alice and Bob have the task of winning in more than 75% of the rounds and one in which they have the task of bringing a statistical quantity (the so-called S -value, computed from four expectation values that can be formed from the game statistic), above the value of 2. We will then show how these two versions relate to each other.

2.1. The Game with the goal of achieving a high percentage of winning rounds

In this version of the game the task of Alice and Bob is presented as receiving local input bits x and y (*questions*) from which they have to produce local output bits a and b (*answers*), such that

$$x \cdot y = a \oplus b \quad (1)$$

holds. Whenever that happens, Alice and Bob have won the round, otherwise they have lost the round. In this equation, the symbol \oplus denotes addition modulo 2 and \cdot denotes ordinary multiplication. All variables x, y, a, b have to be in the set $\{0, 1\}$.

This means that in each round there is a specific question regime, i.e. a pair (x, y) , which we write shorthand as $xy := (x, y)$. The key point is that the question regime in a given round $xy \in \{00, 01, 10, 11\}$ is not known fully to Alice and Bob during the game, and can only be determined when the data is brought together and analyzed jointly. During the game, Alice has to determine a based on x without knowing y , and Bob has to determine b based on y without knowing x .

Therefore, the set of all possible *elementary strategies*, i.e. input-output relations between questions and answers in a given round that Alice and Bob may utilize, can be encoded in four single bits (A_0, A_1, B_0, B_1) . In this notation, the number $A_0 \in \{0; 1\}$ is to be understood as the value that Alice will give as her answer a in a round in which she receives the input bit $x = 0$. $A_1 \in \{0; 1\}$ is the value she will give as her output bit if she receives the input bit $x = 1$. Similarly the number $B_0 \in \{0; 1\}$ is to be understood as the value that Bob will give as his answer b in a round in which he receives the input bit $y = 0$, whereas $B_1 \in \{0; 1\}$ denotes the output Bob will give in case of $y = 1$. As shown in Table 1 each question regime has specific losing strategies.

Table 1. The following table illustrates the goal implied by Equation (1) for the four possible question regimes and it lists the elementary strategies that result in a loss under each question regime. Instead of showing all four bits of a losing elementary strategy (A_0, A_1, B_0, B_1) , only the bits corresponding to the relevant question regime are shown, because the value of the other bits is irrelevant. Note that in each round Alice only knows x and has to answer by giving the value of a , while Bob only knows y and has to give the answer b .

x	y	Regime	Needed to win	Losing Strategies
0	0	$xy = 00$	$0 = a \oplus b$	$(A_0, B_0) = (0, 1)$ or $(1, 0)$
0	1	$xy = 01$	$0 = a \oplus b$	$(A_0, B_1) = (0, 1)$ or $(1, 0)$
1	0	$xy = 10$	$0 = a \oplus b$	$(A_1, B_0) = (0, 1)$ or $(1, 0)$
1	1	$xy = 11$	$1 = a \oplus b$	$(A_1, B_1) = (0, 0)$ or $(1, 1)$

It is easy to see which elementary strategies produce the best results when the question regime is unknown to both Alice and Bob. For example, always using the elementary strategy $(A_0, A_1, B_0, B_1) = (1, 1, 1, 1)$, Alice and Bob would win the round in case of question regimes $xy = 00, 01, 10$, but lose in case of $xy = 11$, so this strategy produces a win in 3 out of 4 regimes.

In general, the performance of an elementary strategy can be checked easily by observing that the number of question regimes L , in which Alice and Bob lose with the elementary strategy (A_0, A_1, B_0, B_1) , can be computed as

$$L = A_0 \oplus B_0 + A_0 \oplus B_1 + A_1 \oplus B_0 + (1 - A_1 \oplus B_1). \tag{2}$$

The validity of Equation (2) is easy to see, because in the question regimes 00, 01, 10 the goal of Alice and Bob is to generate $a \oplus b = 0$, while in the question regime 11 Alice and Bob need to produce $a \oplus b = 1$. Running through the 16 possible elementary strategies shows that L can only take the values 1 or 3, noting that there is no elementary strategy that would produce $L = 0$. Therefore, the best that Alice and Bob can do with an elementary strategy is to ensure that they only lose in 1 of 4 possible question regimes, and with $(A_0, A_1, B_0, B_1) = (1, 1, 1, 1)$ we have already found such a strategy. If all question regimes occur with equal likelihood during the game, this achieves a win in 75% of the rounds.

Therefore, the challenge given by the Verifier to Alice and Bob in this version of the game is to produce a percentage of rounds of at least $75\% + \nu$, for which Equation (1) holds. Here, $\nu > 0$ is a threshold that is set before the game commences.

2.2. The Game with the goal of achieving a high S-value

In a different presentation of the game, Alice and Bob need to produce a high S-value. The S-value is a statistical quantity derived from expectation values following the approach of Clauser-Horne-Shimony-Holt in [36]. In this version of the game, Alice and Bob receive questions x, y which are taken from the set $\{0, 1\}$, to which they have to give answers a, b from the set $\{\pm 1\}$.

Again, Alice and Bob work as a team that may pre-agree strategies but cannot communicate during the game. When all rounds have been played the questions and answers will be put together and a list of quadruplets (a, b, x, y) results, which can be partitioned depending on the question regime. This gives a statistic $P(ab | xy)$ for the product value $ab \in \{\pm 1\}$, conditional on each of the four different question regimes $xy \in \{00, 01, 10, 11\}$. We can now compute expectation values for the product ab under these four question regimes xy by defining

$$\langle ab \rangle_{xy} := \sum_{ab} ab P(ab | xy).$$

We will assume that each question regime occurs at least once in the game, so that all four expectation values are well defined. Following [36], the combination of these four expectation values yield four S-values

$$S_1 = \langle ab \rangle_{00} + \langle ab \rangle_{01} + \langle ab \rangle_{10} - \langle ab \rangle_{11}, \tag{3}$$

$$S_2 = \langle ab \rangle_{00} + \langle ab \rangle_{01} - \langle ab \rangle_{10} + \langle ab \rangle_{11}, \tag{4}$$

$$S_3 = \langle ab \rangle_{00} - \langle ab \rangle_{01} + \langle ab \rangle_{10} + \langle ab \rangle_{11}, \tag{5}$$

$$S_4 = -\langle ab \rangle_{00} + \langle ab \rangle_{01} + \langle ab \rangle_{10} + \langle ab \rangle_{11}. \tag{6}$$

If Alice and Bob agree to always give the answers $a = b = 1$, no matter what question they receive, they are able to generate $S_i = 2$ for all $i = 1, 2, 3, 4$. However, it is unclear how higher S-values could be achieved. What makes winning the game hard is the fact that in the expression for any S_i , one of the correlations is subtracted from the other three. It is this feature of the S_i -values which precludes success using a simple strategy of always giving the same answer to the same question.

Therefore, the challenge given to Alice and Bob for this version of the game is to produce an S_1 -value of at least $2 + \eta$. Here, $\eta > 0$ is a threshold that is set before the game commences. Instead of S_1 one could also use one of the other values S_2, S_3, S_4 to formulate the challenge. As shown in Proposition 2 of [32], at most one of the four S-values may exceed 2. The well-known claim is that this game is impossible to win if an infinite number of rounds is played and if there are no additional resources, like pairs of entangled particles, available to Alice and Bob.

2.3. On the equivalence of the two game descriptions

We want to show that the two versions of the game presented above are essentially the same. Conceptually we therefore have to connect the probability of winning a single round to expectation values. To do this, we first note that after a game over N rounds, the Verifier can break the complete list of quadruplets (a, b, x, y) into four lists by conditioning on specific question regimes xy . In general, these four lists will have a different length, which we denote by N_{xy} , and we only know $N = N_{00} + N_{01} + N_{10} + N_{11}$. In each of those four sub-lists the Verifier can count how often the product value ab was positive and how often it was negative. Let us assume that he uses the symbol N_{00}^+ for the number of rounds in which $ab = +1$ occurred in the sub-list with $xy = 00$, and the symbol N_{00}^- for the number of rounds in which $ab = -1$ occurred in the sub-list with $xy = 00$. For the first expectation value term in Equation (3) we can therefore write

$$\langle ab \rangle_{00} = \frac{N_{00}^+ - N_{00}^-}{N_{00}} = 1 - 2 \frac{N_{00}^-}{N_{00}} = 2 \frac{N_{00}^+}{N_{00}} - 1,$$

because $N_{00} = N_{00}^+ + N_{00}^-$. Analogous expressions hold for $\langle ab \rangle_{01}, \langle ab \rangle_{10}$ and $\langle ab \rangle_{11}$. In particular, we have

$$\langle ab \rangle_{11} = \frac{N_{11}^+ - N_{11}^-}{N_{11}} = 2 \frac{N_{11}^+}{N_{11}} - 1.$$

From Equation (3) we therefore obtain

$$\begin{aligned} S_1 &= \langle ab \rangle_{00} + \langle ab \rangle_{01} + \langle ab \rangle_{10} - \langle ab \rangle_{11} \\ &= 1 - 2 \frac{N_{00}^-}{N_{00}} + 1 - 2 \frac{N_{01}^-}{N_{01}} + 1 - 2 \frac{N_{10}^-}{N_{10}} - \left(2 \frac{N_{11}^+}{N_{11}} - 1 \right) \\ &= 4 - 2 \left(\frac{N_{00}^-}{N_{00}} + \frac{N_{01}^-}{N_{01}} + \frac{N_{10}^-}{N_{10}} + \frac{N_{11}^+}{N_{11}} \right). \end{aligned} \tag{7}$$

To connect the description of the game in Section 2.2, where the requirement is $a, b \in \{\pm 1\}$, with the description in Section 2.1, where the answer values have to be in the set $\{0, 1\}$, we start by denoting the answers in the description of Section 2.1 with an asterisk $*$, so that the requirement for winning a round by fulfilling Equation (1) reads $x \cdot y = a^* \oplus b^*$. The transformation $a = 2a^* - 1$ and $b = 2b^* - 1$ bijectively maps the set $\{0, 1\}$ to the set $\{\pm 1\}$. With this transformation the 16 elementary strategies that Alice and Bob can use in a given round can be described as tuples $(A_0, A_1, B_0, B_1) \in \{\pm 1\} \times \{\pm 1\} \times \{\pm 1\} \times \{\pm 1\}$. To relate this to the percentage of winning rounds, we define

$$\mathcal{L}_{xy} := \frac{\text{number of losing rounds played in question regime } xy}{\text{total number of round played in question regime } xy}.$$

In question regime $xy = 00$, losing a round is equivalent to $a^* \oplus b^* = 1$, which is equivalent to $ab = -1$. Therefore, $\mathcal{L}_{00} = N_{00}^-/N_{00}$ is the percentage of losing rounds in question regime $xy = 00$. Applying the same argument to the question regimes 01 and 10, we see that the losing percentages for these regimes are $\mathcal{L}_{01} = N_{01}^-/N_{01}$ and $\mathcal{L}_{10} = N_{10}^-/N_{10}$, respectively. For the question regime $xy = 11$ the value of the product $x \cdot y$ is equal one, so Alice and Bob lose a round if and only if $a^* \oplus b^* = 0$. In this question regime, losing is equivalent to $ab = +1$, which gives the losing percentage $\mathcal{L}_{11} = N_{11}^+/N_{11}$. From Equation (7) we therefore obtain

$$\begin{aligned} S_1 &= 4 - 2 \left(\frac{N_{00}^-}{N_{00}} + \frac{N_{01}^-}{N_{01}} + \frac{N_{10}^-}{N_{10}} + \frac{N_{11}^+}{N_{11}} \right) \\ &= 4 - 2(\mathcal{L}_{00} + \mathcal{L}_{01} + \mathcal{L}_{10} + \mathcal{L}_{11}). \end{aligned} \tag{8}$$

The last equation shows that maximizing the value of S_1 is tantamount to minimizing the losing percentages in the question regimes.

As discussed in Section 2.1, with every elementary strategy Alice and Bob face at least one losing question regime. Using the same elementary strategy in all game rounds and assuming that the Quiz Master uses all question regimes during the game, this implies $\mathcal{L}_{00} + \mathcal{L}_{01} + \mathcal{L}_{10} + \mathcal{L}_{11} \geq 1$ and hence $S_1 \leq 2$ by Equation (8).

The situation becomes more involved if Alice and Bob switch between different elementary strategies during the game. In some rounds they may be lucky to have chosen the right elementary strategy to produce a win. Intuitively, when the game is played over many rounds and all question regimes come up unpredictably with non-negligible frequency, then one would expect the chances of being lucky enough to produce $\mathcal{L}_{00} + \mathcal{L}_{01} + \mathcal{L}_{10} + \mathcal{L}_{11} < 1$ to diminish. If Alice and Bob randomly switch between elementary strategies and if the Quiz Master randomly switches between the four question regimes, we expect equal percentages of losing rounds in all question regimes, i.e. $\mathcal{L}_{00} = \mathcal{L}_{01} = \mathcal{L}_{10} = \mathcal{L}_{11} =: \mathcal{L}$, over the long run. Under this additional assumption Equation (8) reads

$S_1 = 4 - 8\mathcal{L}$, which shows that $S_1 = 2$ is equivalent to $\mathcal{L} = 25\%$, so $S_1 = 2$ translates to a winning percentage of 75%, which makes the challenges given at the end of Sections 2.1 and 2.2 equivalent (with $\eta = 8\nu$).

However, winning is subject to statistical fluctuations and Alice and Bob may try to benefit from unequal frequencies of question regimes if they are aware of them. In the next chapter we will give bounds for the probability of winning by pure luck if the question regimes occur with unequal probabilities.

3. Bounds for the likelihood of being lucky

The well-known claim is that the CHSH game is impossible to win if the questions are asked randomly and Alice and Bob have no additional (non-local) resources, like pairs of entangled particles, at their disposal. In fact, this is the reason which makes CHSH games useful tools for an explanation of the power of entanglement.

To make the claim true, we first have to insist on ruling out cheats, in which Alice and Bob may prompt some of the questions themselves or acquire knowledge about the question regime they will receive during the game. Such cheats could have consequences which are quantitatively equal to a violation of locality [31,37]. Going forward we insist on the *free* and *independent* generation of all questions, although we do not insist that all question regimes occur with equal likelihood.

The objections that Alice and Bob may win by luck or clever play still have to be dealt with, and we will deal with this now. In this, one needs to keep in mind that they can play in different ways and “being lucky” is a difficult term, as the effect of luck may in principle depend on how they play the game. In particular, Alice and Bob can

- (i) draw up a list, which specifies in advance which elementary strategy they will use in a game round based only on the question posed in this round,
- (ii) create a deterministic algorithm that determines how they will answer in a given round based on the information about all questions they have received so far locally,
- (iii) use the information about all answers they have given so far locally and all questions they have received so far locally as an input to an algorithm, which may use classical local independent randomization procedures while the game is ongoing, to generate their answers, or
- (iv) determine how they will answer in a given round using a *momentary hunch* (i.e. *gut feeling*) while the game is ongoing.

These four approaches will be discussed to determine the effect of luck. Randomness may lie in the generation of the questions x, y , but under Approach (iii) and (iv), also in the generation of the answers. We will now argue that Approaches (i) to (iii) result in the answers being drawn independently from two $N \times 2$ tables filled with numbers of plus or minus one (i.e. from two “spreadsheets”).

In Approach (i), by definition Alice will have a list of pairs $(A_0, A_1) \in \{\pm 1\} \times \{\pm 1\}$ to determine her answers in each round and Bob will have another list of pairs $(B_0, B_1) \in \{\pm 1\} \times \{\pm 1\}$ to determine his answers. The output bit a given by Alice in round n of the game is read out from row n of her list as

$$a := \begin{cases} A_0 & \text{if } x = 0, \\ A_1 & \text{if } x = 1, \end{cases} \quad (9)$$

and the output bit b given by Bob in round n is determined from row n of his list by

$$b := \begin{cases} B_0 & \text{if } y = 0, \\ B_1 & \text{if } y = 1. \end{cases} \quad (10)$$

Prior to the start of the game Alice and Bob may create and discuss their lists together and Alice may even take a copy of Bob’s list into her room, but that will not make a difference as she is only allowed to have local knowledge during the game. The important part is that

we have counter-factual definiteness for the answers of Alice and Bob as they are read off from a table that covers all situations.

Approach (ii) generalizes Approaches (i), but can still be formalized by pre-agreeing on a set of elementary strategies for each possible path the game may take up to a given round n . Written formally, in each round Alice receives a question and gives an answer so that the locally available information for her in round n consists of a question list $x_1 x_2 \dots x_n$ and a list of the answers $a_1 a_2 \dots a_{n-1}$, which she has already given. Based on this information she has to determine her answer a_n for the current round n , in which question x_n was posed. In principle, assuming sufficient memory, Alice and Bob could prepare a game over a total of N rounds by pre-agreeing on a large collection of tables to formalize their strategy under Approach (ii). For example, based on the information up to and including round $n - 1$, there are 4^{n-1} possible combinations of (global) question regimes that could have been used, out of which Alice can differentiate 2^{n-1} based on her local knowledge. To determine her next answer a_n she would pick her table corresponding to the history observed by her so far locally and read out row n . If that row reads (A_0, A_1) , she will use the current question x_n and give her answer using Equation (9) with $x := x_n$. Bob will work analogously on his side. Again the key point is that assuming the existence of a collection of tables corresponds to counter-factual definiteness as far as the answers of Alice and Bob are concerned. The questions they will receive may not be pre-determined, but Alice and Bob can state how they would have answered if they would have received different questions during the game. At the end of the day, the answers will have been generated by independent draws from two $N \times 2$ tables. Conceptually two $N \times 2$ tables could, of course, be seen as $N \times 4$ tables, as in [25], but if Alice and Bob do not reveal their strategies to each other, then none of the players has access to the full $N \times 4$ tables. If one wishes to discuss “realism” the collection of tables would correspond to elements of reality that could be read out by asking the right sequence of questions despite the fact that not all tables are read out in an actual game.

Approach (iii) generalizes Approach (ii) and now the algorithm may be thought of as a collection of tables corresponding to a locally observed history, but allowing several tables that correspond to the same history. Out of these table Alice will pick the table she will use in the current round by casting in her room a classical die, where the number of sides equals the number of her tables that is consistent with her locally observed history. This randomization shows Alice which table to use in the current round. If in that table she finds the entry (A_0, A_1) in row n , she will use the current question x_n and give her answer using Equation (9) with $x := x_n$. Bob would proceed in an analogous way and the important assumption is that Bob’s die is cast independently from Alice’s die.

For the Approaches (i) to (iii) the answers can be regarded as being generated from two $N \times 2$ tables by independent draws. This allows the derivation of the CHSH Inequality: We start by observing that when a combined row (A_0, A_1, B_0, B_1) is created from the two tables of numbers ± 1 , then we always get

$$A_0 B_0 + A_0 B_1 + A_1 B_0 - A_1 B_1 = A_0(B_0 + B_1) + A_1(B_0 - B_1) \in \{\pm 2\}.$$

Taking averages over all rounds, we conclude with the usual Bell-type argument

$$S_1 = \langle ab \rangle_{00} + \langle ab \rangle_{01} + \langle ab \rangle_{10} - \langle ab \rangle_{11} \leq 2. \quad (11)$$

We remark that moving from individual game rounds to averages and expectation values does require additional thoughts about convergence of relative frequencies, because the answers given during the game do not have to be independent over the game rounds. We refer to [38] for details and give numerical examples in Section 4.

Approach (iv) is outside the scope of mathematical procedures, because humans determining answers “on a momentary hunch” seems to be a concept that is impossible to formalize. In this context, some people would treat Approach (iv) just as a form of randomness and automatically subsume it under Approach (iii), but it is important to keep

in mind that such a treatment is an assumption based on certain philosophic positions. While the hunch of Alice experienced in round n can be influenced by the history of questions observed by her previously, some people would argue that this history alone does not allow a comprehensive description of her possibilities. **They might argue that^{CG}** the local future of Alice or non-local connections to Bob cannot be ruled out a priori in the creation of a hunch. To see if something like Approach (iv) can provide more than Approach (iii) in winning a CHSH game without having entanglement as a resource, actual experiments would have to be performed, which naturally can only occur over a finite number of rounds. The burden of proof certainly lies on those, who claim that Approach (iv) can yield more than Approach (iii) and bounds for the likelihood of winning by pure luck are given below.

Generally it is believed that Alice and Bob cannot win CHSH the game unless they have additional non-local resources, like pairs of entangled particles. However, if only a finite number of rounds N of the game is played, Alice and Bob may be declared winners, because the losing percentages that were actually realized may happen to be such that $S_1 \geq 2 + \eta$ occurs in Equation (8) by pure chance. If Alice and Bob switch between elementary strategies between the rounds e.g. under Approach (iii), then after N rounds they may have been lucky to have picked many winning ones even if the four question regimes xy were selected at random and with equal probability. We want to determine how much luck is needed for this to happen. Formulated more precisely, we want to prove statistical bounds for the probability that winning happens by pure luck. Mathematically, such proofs depend on so-called concentration inequalities, which are bounds for the probability that a sum of random variables deviates from its expected value, by a certain amount. In this area, mathematical contributions were made by Hoeffding, Bernstein, Chernoff, Azuma, Bentkus, McDiarmid, Combes and others and many technical formulations exist. We will use Hoeffding Inequalities [39] and an extension of McDiarmid Inequalities [40], giving a self-contained proof of the required results of the former for completeness of presentation.

In what follows it is important to note that the number of rounds N has to be fixed in advance and agreed between the players and the Verifier. If the number of rounds is not fixed beforehand, then one can be tempted to play until a desired level of violation has been reached. **Such an^{CG}** approach can invalidate the entire statistical analysis, see [27], because the probability distribution for the maximum of a stochastic process is different from the probability distribution of that process.

3.1. Bounds based on Hoeffding Inequalities

As shown by Gill as Theorem 1 in [25], when the observed answers are selected completely at random by fair, i.e. 50:50, coin tosses, the probability of generating an S_1 -values above 2 is limited by

$$\Pr\{S_1 \leq 2 + \eta\} \geq 1 - 8e^{-N(\eta/16)^2} \quad (12)$$

for any $\eta \geq 0$. Mathematically, the proof of the bound in Equation (12) in [25] is essentially a repeated application of Hoeffding's inequalities [39]. The right-hand side of the inequality requires a larger number of rounds to give meaningful bounds as the column (a) in Table 2 shows.

Table 2. The following table illustrates the bounds for the certainty that the CHSH game is not won by chance, assuming four equally probable question regimes. Column (a) illustrates the bound obtained with Equation (12), Column (b) illustrates the best available bounds from Theorem 1 and Column (c) the best available bounds from Theorem 2.

N	η	(a)	(b)	(c)
500	0.5	negative	negative	37.73%
1000	0.5	negative	13.51%	69.66%
2500	0.5	30.37%	93.33%	97.23%
5000	0.5	93.94%	99.90%	99.95%

The derivation of Equation (12) in [25] is based on assuming that starting from a spreadsheet containing a table with N rows and 4 columns, fully populated with entries of ± 1 , for each row "two fair coins are tossed independently of one another, independently over all the rows". Depending on the outcomes of the two coin tosses for a round, one value from first two columns and one value from the last two columns is observed in the round corresponding to the specific row. While this gives a clear picture of the assumed randomness, it creates an operational problem, because no one in the CHSH game may have this full spreadsheet: Alice and Bob can work with different lists which they did not share and the Verifier only has access to the list of answers that were actually given.

We will now give a self-contained proof for a bound that is stronger than Equation (12) and that does not require the assumption of a fair coin toss for the selection of the question regime. Relaxing this assumption is important, because the four question regimes $\{00, 01, 10, 11\}$ may occur with a somewhat unequal likelihood in physical Bell experiments given biases of random number generators. In a game between humans, in which a Quiz Master is free to choose the questions, equal likelihood of question regimes cannot be assumed a priori. In macroscopic situations, the available table of observations may not necessarily be assumed to contain the four regimes with equal probability. For example, if in the macroscopic application described in [31], large orders occur rarely and there are few rumours, regime $xy = 00$ will be prevalent. In this case, the first expectation term in the expression $S_1 = \langle ab \rangle_{00} + \langle ab \rangle_{01} + \langle ab \rangle_{10} - \langle ab \rangle_{11}$ will be computed from the observations of ab over many rounds while the other three expectation values will result from the observations over relatively few rounds. In such a case of highly asymmetric frequency for the different regimes, randomness may have a stronger influence on the S_1 -value. In an extreme case, where each of the three regimes $\{01, 10, 11\}$ occurs only once during the game rounds and chance happens to produce $\langle ab \rangle_{01} = 1, \langle ab \rangle_{10} = 1, \langle ab \rangle_{11} = -1$, we get $S_1 = \langle ab \rangle_{00} + 3$.

Therefore, we will with the subsequent Theorem 1 extend the work in [25] to situations where the four regimes may occur with different probabilities. We will, however, stick with the key idea of the proof made by Gill, i.e. a repeated application of Hoeffding's inequalities. Before stating and proving Theorem 1 we start by proving Hoeffding's inequality by elementary means, to provide the reader with a fully self-contained presentation.

Proposition 1. Let X_1, \dots, X_N be independent random variables with values X_i in the interval $[0; 1]$. Then for all $\delta > 0$ we have

$$Pr \left\{ \frac{X_1 + \dots + X_N}{N} > \mu + \delta \right\} \leq \exp(-2N\delta^2) \tag{13}$$

and

$$Pr \left\{ \frac{X_1 + \dots + X_N}{N} < \mu - \delta \right\} \leq \exp(-2N\delta^2) \tag{14}$$

with $\mu := \mathbb{E}[(X_1 + \dots + X_N)/N]$.

Proof. To give a self-contained proof, we start by making the following observation: If Y is a random variable with nonnegative values, then for each positive real number a we trivially may write

$$Y \geq a1_{\{Y>a\}}$$

using the indicator function $1_{\{\dots\}}$. Taking the expectation value is a monotonic and linear operation, so

$$\mathbb{E}[Y] \geq \mathbb{E}\left[a1_{\{Y>a\}}\right] = a\mathbb{E}\left[1_{\{Y>a\}}\right] = a \Pr\{Y > a\}$$

follows, which proves the so-called Markov Inequality

$$\Pr\{Y > a\} \leq \frac{1}{a}\mathbb{E}[Y]. \tag{15}$$

Second, we observe that for arbitrary real numbers r and each constant $c \in [0; 1]$ we have

$$1 + (e^r - 1)c \leq e^{rc+r^2/8}, \tag{16}$$

which can be shown as follows: $\color{red}{\in \text{CG}}$ consider the logarithm of the left-hand side and define the function $f(r) := \ln(1 + (e^r - 1)c)$. This function is twice continuously differentiable with

$$f'(r) = \frac{e^r c}{1 + (e^r - 1)c} = \frac{c}{c + (1 - c)e^{-r}} \text{ and } f''(r) \leq \frac{1}{4}.$$

Because of $f(0) = 0$ and $f'(0) = c$, the second order Taylor expansion of f yields

$$f(r) = f(0) + rf'(0) + \frac{1}{2}r^2 f''(r\theta) \leq rc + \frac{1}{8}r^2,$$

with some $\theta \in [0; 1]$, which proves Equation (16).

Third, we observe that for all real numbers r the exponential function $x \rightarrow e^{rx}$ is convex and hence for all $x \in [0; 1]$ we have

$$e^{rx} \leq xe^r + (1 - x)e^0 = 1 + (e^r - 1)x.$$

So, for each random variable X_i we get

$$e^{rX_i} \leq 1 + (e^r - 1)X_i,$$

which leads to

$$\mathbb{E}[e^{rX_i}] \leq 1 + (e^r - 1)\mathbb{E}[X_i] \stackrel{\text{Eq. (16)}}{\leq} e^{r\mathbb{E}X_i+r^2/8}.$$

This directly implies the so-called Hoeffding Lemma

$$\mathbb{E}[e^{r(X_i - \mathbb{E}X_i)}] \leq e^{r^2/8}. \tag{17}$$

After these three preparatory steps we can make the following estimate, which holds for all positive real numbers r

$$\begin{aligned}
 \Pr\left\{\frac{X_1 + \dots + X_N}{N} > \mu + \delta\right\} &= \Pr\{r(X_1 + \dots + X_N) > rN(\mu + \delta)\} \\
 &= \Pr\left\{e^{r(X_1 + \dots + X_N) - rN\mu} > e^{r\delta N}\right\} \\
 &\stackrel{\text{Eq.(15)}}{\leq} e^{-r\delta N} \mathbb{E}\left[e^{r(X_1 + \dots + X_N) - rN\mu}\right] \\
 &= e^{-r\delta N} \mathbb{E}\left[\prod_{i=1}^N e^{r(X_i - \mathbb{E}X_i)}\right] \\
 &\stackrel{\text{indep.}}{=} e^{-r\delta N} \prod_{i=1}^N \mathbb{E}\left[e^{r(X_i - \mathbb{E}X_i)}\right] \\
 &\stackrel{\text{Eq.(17)}}{\leq} e^{-r\delta N} \prod_{i=1}^N e^{r^2/8} = e^{-r\delta N} e^{Nr^2/8} = e^{(r^2 - 8r\delta)N/8} \\
 &= \exp\left(\left((r - 4\delta)^2 - 16\delta^2\right)N/8\right).
 \end{aligned}$$

The exponent in the last expression is minimized with the choice $r := 4\delta$. This proves Equation (13).

Knowing that Equation (13) holds true, it is easily seen that Equation (14) holds true as well: $w\mathbb{W}^{\text{CG}}$ only have to define $Y_i := 1 - X_i$ which gives a collection of independent random variables Y_1, \dots, Y_N with values in the interval $[0; 1]$ to which Equation (13) can be applied. \square

Now we can prove

Theorem 1. Assume that for each round of the N rounds of the CHSH game one of the four question regimes xy is selected at random with constant probability p_{xy} independently of the selection for the other rounds of the game. Assume further that there are $N \times 2$ tables of numbers ± 1 , from which the answers of Alice and Bob in each given round are determined as described earlier under the Approaches (i), (ii) or (iii) leading to the expectation of $S_1 \leq 2$.

Then, for any positive number η the probability of observing an S_1 -value above $2 + \eta$ at the end of the CHSH game is limited by

$$\Pr\{S_1^{\text{obs}} \leq 2 + \eta\} \geq 1 - q,$$

where the bound q is given by

$$q := 4 \exp\left(-2N\delta^2\right) + \sum_{xy} \exp\left(-2N(p_{xy} - \delta)(\eta/8)^2\right). \tag{18}$$

In this bound the number $\delta > 0$ can be chosen freely to minimize the value of q .

Proof. The proof uses arguments similar to those made by Gill in [25]. We use the notation introduced in Section 2.3 and add the superscript “(obs)” to those numbers that are calculated from the question and answer pattern that was actually observed in the CHSH game.

Let two positive numbers η and δ be given and let us write $\tilde{\delta} := \eta/8$. Applying Equation (14) for each question regime $xy = 00, 01, 10, 11$, we see that the probability of a

deviation of the sample mean from the theoretical probability p_{xy} of the question regime is bounded by

$$\Pr \left\{ \frac{N_{xy}^{\text{obs}}}{N} \leq p_{xy} - \delta \right\} \leq \exp(-2N\delta^2). \tag{19}$$

So ignoring the union of all sets for which (19) does not hold we have

$$N_{xy}^{\text{obs}} > N(p_{xy} - \delta) \text{ for all } xy = 00, 01, 10, 11, \tag{20}$$

except possibly on a set that has a probability of at most $4 \exp(-2N\delta^2)$.

Conditional on N_{xy}^{obs} observations being made in question regime xy , we have to compare the observed sample mean for a positive product $N_{xy}^{\text{obs},+} / N_{xy}^{\text{obs}}$ with its expected value N_{xy}^+ / N_{xy} . Again applying Hoeffding’s inequalities [39,41] we get the bound

$$\Pr \left\{ \frac{N_{xy}^{\text{obs},+}}{N_{xy}^{\text{obs}}} \geq \frac{N_{xy}^+}{N_{xy}} + \tilde{\delta} \right\} \leq \exp(-2N_{xy}^{\text{obs}}\tilde{\delta}^2), \text{ for all } xy = 00, 01, 10$$

and similarly

$$\Pr \left\{ \frac{N_{11}^{\text{obs},-}}{N_{11}^{\text{obs}}} \geq \frac{N_{11}^-}{N_{11}} + \tilde{\delta} \right\} \leq \exp(-2N_{11}^{\text{obs}}\tilde{\delta}^2).$$

Using (20) the upper bound for these probabilities can be dominated by

$$\exp(-2N_{xy}^{\text{obs}}\tilde{\delta}^2) \leq \exp(-2N(p_{xy} - \delta)\tilde{\delta}^2).$$

So, except on a small set, the observed frequency of each question regime and the observed frequency of a positive or negative product ab in each question regime may not deviate too far from their overall mean. As shown in Section 2.3, we have

$$\begin{aligned} \langle ab \rangle_{00} &= \frac{N_{00}^+ - N_{00}^-}{N_{00}} = 2 \frac{N_{00}^+}{N_{00}} - 1, \\ \langle ab \rangle_{01} &= \frac{N_{01}^+ - N_{01}^-}{N_{01}} = 2 \frac{N_{01}^+}{N_{01}} - 1, \\ \langle ab \rangle_{10} &= \frac{N_{10}^+ - N_{10}^-}{N_{10}} = 2 \frac{N_{10}^+}{N_{10}} - 1, \\ -\langle ab \rangle_{11} &= -\frac{N_{11}^+ - N_{11}^-}{N_{11}} = 2 \frac{N_{11}^-}{N_{11}} - 1. \end{aligned}$$

These equations remain true, when reference is made to observed quantities only, i.e. when $\langle ab \rangle_{xy}$ is replaced by $\langle ab \rangle_{xy}^{\text{obs}}$ and N_{xy} by N_{xy}^{obs} , etc. Therefore, we conclude by using (11) that the inequality

$$\begin{aligned} S_1^{\text{obs}} &= \langle ab \rangle_{00}^{\text{obs}} + \langle ab \rangle_{01}^{\text{obs}} + \langle ab \rangle_{10}^{\text{obs}} - \langle ab \rangle_{11}^{\text{obs}} \\ &\leq \langle ab \rangle_{00} + \langle ab \rangle_{01} + \langle ab \rangle_{10} - \langle ab \rangle_{11} + 8\tilde{\delta} = S_1 + 8\tilde{\delta} \\ &\leq 2 + 8\tilde{\delta} = 2 + \eta \end{aligned}$$

has to hold, except possibly on a set with a probability of not more than

$$4 \exp(-2N\delta^2) + \sum_{xy} \exp(-2N(p_{xy} - \delta)(\eta/8)^2).$$

Note that in the last expression we are free to choose any positive δ and the sum has to be taken over all four question regimes $xy = 00, 01, 10, 11$. \square

For given parameters N, p_{xy}, η the value for δ , which minimizes q in Equation (18) and hence gives the best bound, can be found numerically, for example by solving $0 = dq/d\delta$. Performing the minimization, for example for the case of equally likely question regimes $p_{xy} = 0.25$ with $N = 5,000$ and $\eta = 0.5$, an optimal value is found at $\delta \approx 0.03361$ is found. With this value Theorem 1 provides the bound $\Pr\{S_1^{\text{obs}} \leq 2.5\} \geq 99.90\%$, which is shown in Column (b) in the last row of Table 2.

3.2. Bounds based on an extended McDiarmid Inequality

To the extent that one is interested in some ideas behind the mathematics needed for more advanced bounds, an extension of McDiarmid Inequality can be of interest. In the proof of Theorem 1, Proposition 1 is applied separately to each of the four expectation values that occur in the definition of S_1 in Equation (3). The probabilities of the four exception sets, on which the observed averages deviate substantially from the theoretical expectation value, are then simply added. ~~However, even though~~^{CG} the exception sets overlap and cancellations may be expected to occur. In this section, we will work directly with the definition of the S_1 -value as a linear combination of random variables with values of ± 1 . ~~We will also~~^{CG} use a different concentration inequality, namely an extension of McDiarmid Inequality, to arrive at a powerful bound for the probability of winning the CHSH game by pure chance.

McDiarmid’s Inequalities build on the work of Hoeffding and Azuma and give concentration inequalities for situations in which differences for the evaluating function are bounded, i.e. satisfy a Lipschitz condition, see [42]. They were extended in [40] by Combes to cover situations in which the differences are not bounded everywhere, but only on a set that has a high probability. Of course, better bounds usually come at the price of higher complexity, so we will not attempt a self-contained presentation of results, but rather summarize McDiarmid and Combes’ result in the following

Proposition 2. *Let $\mathcal{X}_1, \dots, \mathcal{X}_N$ be sets. Define $\mathcal{X} := \mathcal{X}_1 \times \dots \times \mathcal{X}_N$ and let \mathcal{Y} be a subset of \mathcal{X} . Let $f : \mathcal{X} \rightarrow \mathbb{R}$ be a function that assigns a real number $f(x)$ to every $x = (x_1, \dots, x_N) \in \mathcal{X}$. Assume that on the set \mathcal{Y} the function f has bounded differences, i.e. there are constants c_1, \dots, c_N such that*

$$|f(x) - f(x')| \leq c_i \text{ for all } (x, x') \in \mathcal{Y} \times \mathcal{Y} \text{ with } x_j = x'_j \text{ for } j \neq i. \tag{21}$$

Let X_1, \dots, X_N be independent random variables with values X_i in the set \mathcal{X}_i and let us write $X := (X_1, \dots, X_N)$ as well as $m := \mathbb{E}[f(X)|X \in \mathcal{Y}]$. Then for all $\eta \geq 0$ we have

$$\Pr\{f(X) - m \geq \eta\} \leq p + \exp\left(-\frac{2 \max(\eta - p \sum_{i=1}^N c_i, 0)^2}{\sum_{i=1}^N c_i^2}\right) \tag{22}$$

with $p := \Pr\{X \notin \mathcal{Y}\}$.

Proof. See [40]. \square

While a proof cannot be given here, Proposition 2 is not so difficult to understand intuitively: If one has a set \mathcal{Y} on which the function is well-behaved, i.e. small random fluctuations in the inputs do not create a big difference in its value, then one can estimate the probability for a large deviation from the mean by an $\exp(-\dots)$ term, vaguely reminiscent of Hoeffding’s Inequality. Outside the set nothing can be said, but that is not a problem as long as the probability of being in \mathcal{Y} can be made very high by playing a sufficient number of rounds.

Proposition 2 can be applied to the CHSH game in the following way: ~~uU~~^{CG}nder Approach (iii) there are a $N \times 2$ spreadsheets that specify how Alice and Bob will answer.

Knowing all questions for all rounds, S_1 can be computed via Equation (3) and this procedure for obtaining S_1 plays the role of the function f in Proposition 2. The random variable X_i determines which question regime applies in round i , so it can take any of the four values in the set $\mathcal{X}_i := \{00, 01, 10, 11\}$. The assumption that in each round of the game one of the four question regimes xy is selected at random with probability p_{xy} independently from the selection for the other rounds of the CHSH game gives the independence property for the random variables X_1, \dots, X_N .

To show that computing S_1 has bounded differences on an appropriately defined set, we first note that the expectation values $\langle ab \rangle_{xy}$ in (3) are taken over N_{xy} observations. The numbers N_{xy} , which were introduced in Section 2.3, count how often question regime xy has occurred in the CHSH game. Formally, these numbers N_{xy} result via

$$N_{xy} = \sum_{i=1}^N 1_{\{X_i = xy\}}$$

from the random variables X_i . To show that S_1 has bounded differences on a subset of \mathcal{X} , we define such a subset \mathcal{Y} by making sure that inside it all four question regimes occur sufficiently often. This is achieved by

$$\mathcal{Y} := \mathcal{X} \setminus (\{N_{00} \leq K + 1\} \cup \{N_{01} \leq K + 1\} \cup \{N_{10} \leq K + 1\} \cup \{N_{11} \leq K + 1\}),$$

for some natural number K . We compute

$$Pr\{N_{xy} \leq K + 1\} = \sum_{k=0}^{K+1} \binom{N}{k} p_{xy}^k (1 - p_{xy})^{N-k}$$

for all $xy = 00, 01, 10, 11$, which allows the following crude upper bound for its probability

$$\begin{aligned} Pr\{X \notin \mathcal{Y}\} &= Pr\left(\bigcup_{xy} \{N_{xy} \leq K + 1\}\right) \leq \sum_{xy} Pr\{N_{xy} \leq K + 1\} \\ &= \sum_{xy} \sum_{k=0}^{K+1} \binom{N}{k} p_{xy}^k (1 - p_{xy})^{N-k}. \end{aligned} \tag{23}$$

An exact computation of $Pr\{X \notin \mathcal{Y}\}$ would be possible by using properties of the multinomial distribution, see [43,44], but we want to avoid such technical arguments.

As we will show in the subsequent proof, the function $S_1 : \mathcal{X} \rightarrow [-4; +4]$ has bounded differences, i.e. obeys Equation (21), on the set \mathcal{Y} . ~~We will show below that whenbecause~~ ^{if^{CG}} we change the outcome of just one random variable X_i , then the resulting change in the value of S_1 is bounded by $c_i = 4/K$. ~~as will be shown below.~~ ^{CG} On this basis, an application of Proposition 2 yields the following

Theorem 2. Assume that for each round of the N rounds of the CHSH game one of the four question regimes xy is selected at random with constant probability p_{xy} independently of the selection for the other rounds of the game. Assume further that there are $N \times 2$ tables of numbers ± 1 , from which the answers of Alice and Bob in each given round are determined as described earlier under the Approaches (i), (ii) or (iii) leading to the expectation of $S_1 \leq 2$.

Then, for any positive number η the probability of observing an S_1 -value above $2 + \eta$ after N rounds of the CHSH game is limited by

$$Pr\{S_1^{obs} \leq 2 + \eta\} \geq 1 - q$$

with

$$q := p + \exp\left(-\frac{2 \max(\eta - 4pN/K, 0)^2}{16N} K^2\right), \quad p := \sum_{xy} \sum_{k=0}^{K+1} \binom{N}{k} p_{xy}^k (1 - p_{xy})^{N-k} \quad (24)$$

for all $K < N/4$.

Proof. We apply Proposition 2 with $f(X) = S_1^{\text{obs}}$. The derivation of the expectation $S_1 \leq 2$ under Approach (i), (ii) or (iii) also works on the subset \mathcal{Y} , which implies

$$m = \mathbb{E}[f(X)|X \in \mathcal{Y}] \leq 2.$$

With the definition $\tilde{p} := \Pr\{X \notin \mathcal{Y}\}$ Proposition 2 now provides the estimate

$$\begin{aligned} \Pr\{S_1^{\text{obs}} > 2 + \eta\} &= \Pr\{f(X) - 2 > \eta\} \leq \Pr\{f(X) - m > \eta\} \\ &\leq \tilde{p} + \exp\left(-\frac{2 \max(\eta - \tilde{p} \sum_{i=1}^N c_i, 0)^2}{\sum_{i=1}^N c_i^2}\right). \end{aligned}$$

From this we obtain the claimed Equation (24) with $c_i := 4/K$ for all i , because we know $\tilde{p} \leq p$ from Equation (23).

To complete the proof of Theorem 2, it therefore only remains to be shown that the function $S_1 : \mathcal{X} \rightarrow [-4; +4]$ obeys Equation (21) with constants $c_i = 4/K$ on the set \mathcal{Y} that was defined above. To prove this, let us, to avoid notation that is too complex, for example, look at $i = 42$ and assume that x_{42} changes its value from $x_{42} = 00$ to $x'_{42} = 01$ while the other variable $x_1, \dots, x_{41}, x_{43}, \dots, x_N$ remain unchanged. This leads to a change from S_1 from S_1 and we need to estimate the magnitude of this change on the set \mathcal{Y} . As a result of this change from x_{42} to x'_{42} , the number N_{00} changes to $N'_{00} = N_{00} - 1$ and N_{01} changes to $N'_{01} = N_{01} + 1$, while N_{10} and N_{11} remain unaffected. As a consequence, the difference between S_1 and S'_1 can be written as

$$\begin{aligned} S_1 - S'_1 &= \frac{N_{00}^+ - N_{00}^-}{N_{00}} + \frac{N_{01}^+ - N_{01}^-}{N_{01}} - \left(\frac{(N_{00}^+)' - (N_{00}^-)'}{N'_{00}} + \frac{(N_{01}^+)' - (N_{01}^-)'}{N'_{01}} \right) \\ &= \left(\frac{N_{00}^+ - N_{00}^-}{N_{00}} - \frac{(N_{00}^+)' - (N_{00}^-)'}{N_{00} - 1} \right) + \left(\frac{N_{01}^+ - N_{01}^-}{N_{01}} - \frac{(N_{01}^+)' - (N_{01}^-)'}{N_{01} + 1} \right) \\ &= \frac{(N_{00} - 1)(N_{00}^+ - N_{00}^-) - N_{00}((N_{00}^+)' - (N_{00}^-)')}{N_{00}(N_{00} - 1)} \\ &\quad + \frac{(N_{01} + 1)(N_{01}^+ - N_{01}^-) - N_{01}((N_{01}^+)' - (N_{01}^-)')}{N_{01}(N_{01} + 1)} \\ &= \frac{N_{00}(N_{00}^+ - (N_{00}^+)') - N_{00}(N_{00}^- - (N_{00}^-)') - (N_{00}^+ - N_{00}^-)}{N_{00}(N_{00} - 1)} \\ &\quad + \frac{N_{01}(N_{01}^+ - (N_{01}^+)') - N_{01}(N_{01}^- - (N_{01}^-)') + (N_{01}^+ - N_{01}^-)}{N_{01}(N_{01} + 1)} \quad (25) \end{aligned}$$

We introduce the following definitions for the change in the number of winners and losers in each question regime xy when we move from x_{42} to x'_{42} , namely

$$\Delta_{xy}^+ := N_{xy}^+ - (N_{xy}^+)', \quad \Delta_{xy}^- := N_{xy}^- - (N_{xy}^-)'$$

Using this notation we can write (25) as

$$S_1 - S'_1 = \frac{1}{N_{00} - 1} \left(\Delta_{00}^+ - \Delta_{00}^- - \frac{N_{00}^+ - N_{00}^-}{N_{00}} \right) + \frac{1}{N_{01} + 1} \left(\Delta_{01}^+ - \Delta_{01}^- + \frac{N_{01}^+ - N_{01}^-}{N_{01}} \right)$$

and hence

$$|S_1 - S'_1| \leq \frac{1}{N_{00} - 1} \left(|\Delta_{00}^+ - \Delta_{00}^-| + \frac{|N_{00}^+ - N_{00}^-|}{N_{00}} \right) + \frac{1}{N_{01} + 1} \left(|\Delta_{01}^+ - \Delta_{01}^-| + \frac{|N_{01}^+ - N_{01}^-|}{N_{01}} \right)$$

The change from $x_{42} = 00$ to $x'_{42} = 01$ means that when we move from the computation of S_1 to the computation of S'_1 we have one outcome less in regime 00 implying $\Delta_{00}^+ \geq 0$ and $\Delta_{00}^- \geq 0$. Depending on the table that Alice and Bob use, the change in row 42 may have eliminated a loser or a winner, so there are only two possibilities

$$(\Delta_{00}^+ = 0 \text{ and } \Delta_{00}^- = 1) \text{ or } (\Delta_{00}^+ = 1 \text{ and } \Delta_{00}^- = 0),$$

which means $|\Delta_{00}^+ - \Delta_{00}^-| = 1$. Similarly, as a result of the change there is one additional outcome for regime 01, implying $\Delta_{01}^+ \leq 0$ and $\Delta_{01}^- \leq 0$. The additional outcome maybe a winner or a loser, so similarly there are only two possibilities

$$(\Delta_{01}^+ = 0 \text{ and } \Delta_{01}^- = -1) \text{ or } (\Delta_{01}^+ = -1 \text{ and } \Delta_{01}^- = 0),$$

which implies $|\Delta_{01}^+ - \Delta_{01}^-| = 1$. This simplifies the above estimate to

$$\begin{aligned} |S_1 - S'_1| &\leq \frac{1}{N_{00} - 1} \left(1 + \frac{|N_{00}^+ - N_{00}^-|}{N_{00}} \right) + \frac{1}{N_{01} + 1} \left(1 + \frac{|N_{01}^+ - N_{01}^-|}{N_{01}} \right) \\ &\leq \frac{2}{N_{00} - 1} + \frac{2}{N_{01}} \leq \frac{4}{K}, \end{aligned} \tag{26}$$

because on the set \mathcal{Y} we have $N_{xy} - 1 \geq K$ for all $xy = 00, 01, 10, 11$.

This line of reasoning is also valid for other changes than the specific change in $i = 42$ from 00 to 01. The fact that the definition of S_1 has a minus sign in front of the expectation value for regime $xy = 11$ does not pose difficulties. Therefore, we see that $c_i = 4/K$ provides a general bound in Equation (21) and the proof of Theorem 2 is complete. \square

In contrast to Equation (12), Theorems 1 and 2 allow situations where the question regimes do not occur with equal probability. The following Table 3 gives illustrative bounds for these cases. Both Theorems can be useful depending on how the probabilities for the question regimes are set. Theorem 2 can in principle still be improved by using the exact multinomial distribution instead of the estimate made in Equation (23).

Table 3. The following table illustrates the bounds for the certainty that the CHSH game is not won by chance over N rounds with threshold η assuming unequal probabilities for the four question regimes. Columns (a) and (c) illustrate the bounds obtained from Theorem 1 and Columns (b) and (d) the bounds from Theorem 2. The values in Columns (a) and (b) were computed for a weakly asymmetric situation with $p_{00} = 22\%$ and $p_{01} = p_{10} = p_{11} = 26\%$. The values in Columns (c) and (d) were computed for a strongly asymmetric situation with $p_{00} = 10\%$ and $p_{01} = p_{10} = p_{11} = 30\%$.

N	η	(a)	(b)	(c)	(d)
500	0.5	negative	30.91%	negative%	4.04%
1000	0.5	12.71%	60.45%	negative%	12.35%
2500	0.5	92.91%	93.82%	69.61%	37.20%
5000	0.5	99.87%	99.75%	93.97%	66.35%
500	0.75	14.51%	57.42%	negative%	9.36%
1000	0.75	85.39%	87.98%	52.34%	26.32%
2500	0.75	99.92%	99.79%	93.37%	65.46%
5000	0.75	99.99%	99.99%	99.73%	91.58%




Figure 1. Two probability distributions for S_1^{obs} generated by Monte Carlo simulation of 10,000 CHSH games of $N = 500$ rounds each. The threshold of $2 + \eta = 2.25$ is shown in red. (a) The graph on the **left-handleftthand^{CG}** side has been generated with Alice and Bob randomly picking elementary strategies with $L = 1$, while the regimes xy were generated by independent, unbiased coin tosses. The simulated probability is $\Pr\{S_1^{\text{obs}} \geq 2.25\} = 4.5\%$ with a maximum value of $S_{1,\text{max}}^{\text{obs}} = 2.52$ observed in a single CHSH game. (b) The graph on the **right-handrighthand^{CG}** side has been generated with Alice and Bob randomly picking elementary strategies that win in regime $xy = 00$ and also satisfy $L = 1$. Here, all regimes xy were generated by independent, but biased coin tosses with $P(xy = 00) = 0.7$ and $P(xy = 01) = P(xy = 10) = P(xy = 11) = 0.1$. The simulated probability is $\Pr\{S_1^{\text{obs}} \geq 2.25\} = 14\%$ with a maximum value of $S_{1,\text{max}}^{\text{obs}} = 2.86$ observed in a single CHSH game.

4. Monte Carlo Simulations and exploitable biases

The probability $\Pr\{S_1^{\text{obs}} \geq 2 + \eta\}$ can be estimated by performing Monte Carlo simulations on a classical computer. **For this we assume assuming^{CG}** an operationally well defined strategy that Alice and Bob use to produce their answers a, b and a well defined mechanism by which the questions x, y are generated. This allows the numerical estimation of bounds without the need for the analytical arguments given in the previous section, but it is of course dependent on specific assumptions regarding the operational procedures.

As discussed in Section 2.1 there are 16 elementary strategies available to Alice and Bob, half of which produce $L = 3$ while the other half produce $L = 1$. If the question regimes are generated fully at random by independent unbiased coin tosses, i.e. $P(xy) = 1/4$ for all $xy = 00, 01, 10, 11$ in all rounds, and if Alice and Bob just pick randomly from a set of elementary strategies, the probability density for S_1^{obs} will approach the normal distribution with a maximum in the range from -2 to $+2$. The position of the maximum depends on how often Alice and Bob pick strategies with $L = 3$ versus strategies with $L = 1$.

If Alice and Bob want to win and aim for a high number S_1^{obs} while assuming unbiased question regimes, they should use elementary strategies with $L = 1$ only. Assuming that Alice and Bob pick randomly only from those elementary strategies and assuming that the questions are generated fully at random by independent unbiased coin tosses, then a typical probability distribution for S_1^{obs} , as shown on the **left-handleftthand^{CG}** side of Figure 1, is the result. Here, it is extremely unlikely to win by chance if the threshold is set high enough.

If the question regimes are stochastically independent between game rounds but not distributed equally, which may happen in applications of Bell correlations outside physics [32], the chances of winning improve. If, for example, $P(xy = 00) =: c > 0.25$ and $P(xy = 01) = P(xy = 10) = P(xy = 11) = (1 - c)/3$ and Alice and Bob are aware of this bias, they should only play elementary strategies with $L = 1$ that are sure to win in regime $xy = 00$. Knowing about such a bias improves the chances of winning considerably, as shown on the righthand side of Figure 1.

The simulated probability for $S_1^{\text{obs}} \geq 2 + \eta$ remains substantially below the theoretical bounds. This is consistent with what [26,38] report, because the bounds based on Hoeffding-Azuma and McDiarmid Inequalities are generally not very tight. In addition,

some simplifications were made in the proof of Theorem 2, for example to avoid the use of multinomial distributions.

Clearly, a bias with $P(xy = 00) = c \gg 0.25$ is easily detectable in the data. However, in case the probability for question regimes varies over the game rounds, things become more tricky. First, note that for the results in the previous sections it is not required that the answers from Alice and Bob are stochastically independent over the game rounds. In fact, Alice could simply decide that she will never give the same answer in three consecutive rounds. In this case a is not stochastically independent over the different game rounds, but Theorems 1 and 2 can still be applied. The important assumption only concerns the question regimes.

Systematic biases in random numbers generated from quantum processors do occur in real implementations, so random numbers generated by a quantum computer may not be that random in practice. In [45] various tests were used on random numbers generated by a cloud quantum computer (IBM 20Q Poughkeepsie) and the authors conclude "As a result, we observed that some qubits were more biased than others". Errors and biases can change over time and are difficult to correct, see [17] stating "But errors typically fluctuate over time" and [46] stating "While some of the protocols extract quantum randomness and discard deterministic components arisen due to quantum processes implementation imperfections, the fidelity of such procedures is not ideal". This clearly is an important consideration when a Bell test together with a statistical confidence interval is used to exclude the possibility of eavesdropping in real life quantum communication.

We illustrate this problem by a simple scheme, in which the probabilities of the question regimes change predictably between the game rounds. Here we assume that there is a recurring pattern in the random number generator, such that stochastic patterns repeat every four rounds as illustrated in the following Table 4.

Table 4. An example for a systematic cyclical bias in the random question regimes that repeats over the round numbers n . The number k is understood to run through the natural numbers $k = 1, 2, 3, \dots$

Round n	$P(xy = 00)$	$P(xy = 01)$	$P(xy = 10)$	$P(xy = 11)$
...				
$4k$	c	$(1 - c)/3$	$(1 - c)/3$	$(1 - c)/3$
$4k + 1$	$(1 - c)/3$	c	$(1 - c)/3$	$(1 - c)/3$
$4k + 2$	$(1 - c)/3$	$(1 - c)/3$	c	$(1 - c)/3$
$4k + 3$	$(1 - c)/3$	$(1 - c)/3$	$(1 - c)/3$	c
...				

This creates an exploitable pattern as the bias rotates over time through the question regimes in a predictable manner. Note that even in this situation, knowing the question regime drawn in an earlier round does still not give predictive power about the question regimes in later rounds, the predictive power just comes from knowing the round number. As Alice and Bob for example know that in round number n , where n is divisible by 4, regime $xy = 00$ is more likely to occur, they can just agree to use elementary strategies that maximize the winning probability in such rounds. So, it is easy to see that Alice and Bob can exploit this cyclicity by playing appropriate elementary strategies in each game round n simply by looking at n modulo 4. With a high number of c , Alice and Bob thus get a good chance of producing values for S_1^{obs} that can be well above 3.

Summing over the columns of Table 4 it is clear that all four question regimes 00, 01, 10, 11 will still occur with equal probability of around 25% over the entire CHSH game, so the bias only becomes visible when testing subsets of rounds. Note that Theorems 1 and 2 and related work are not applicable here, because p_{xy} does not stay constant over the game rounds.

5. Discussion^{CG} and Summary

Cheats, like classical communication between the players of the CHSH game or allowing for an undue influence on the generation of questions, can allow the generation of a quantum-like Bell statistic with classical processes. The amount of cheating that is required is not very large. The Toner-Bacon protocol [14] shows, for example, that when Alice and Bob share two uniformly distributed random variables with values on the unit sphere of \mathbb{R}^3 , then sending a single bit can be enough: one bit of information from Alice to Bob suffices to simulate the statistic of a local projective measurement on an entangled Bell pair state. The exchange of one classical bit of information can do a lot in terms of generating strong correlations [15] and in [16] the information to be shared between measurement settings is discussed illustrating that small amounts of shared classical information suffice in order to break the Bell bound. However, this position [14–16] is based on the implicit assumption that an arbitrarily large number of rounds is played and that the question regimes (measurement settings) are produced by unbiased random number generators with equal likelihood. The practical requirement of having a finite number of game rounds and the possibility of biases were not considered in these contributions. Biases were considered elsewhere [21,22], but with a different focus. The present work addresses statistical questions in a situation where the number of rounds is finite and where biases may be present.

This has practical implications:^{CG} when CHSH inequalities are used to certify that no eavesdropping took place in quantum communication over a shorter time period, a natural question is how likely any observed violations of CHSH inequalities could have occurred by chance alone. In a pedagogical presentation of entanglement and CHSH games a similar question may arise: “How much luck is needed to win a CHSH game without entangled particles?”^{CG} The rehearsed opinion that the observed violation could only have occurred with a high standard deviation (“a n -sigma violation”) is not a good answer as the a priori assumption of a normal distribution can be dangerous. Good answers do exist in the literature, see e.g. the review in [26], but they tend to be based on highly technical tools and they focus on small biases in the random number generators that provides the question regimes.

In the present paper we focussed on an accessible development of bounds for the probability of being lucky in the form of

$$\Pr\{S_1 \leq 2 + \eta\} \geq 1 - q$$

for a given threshold $\eta > 0$. We discussed different forms of game play and provided three formulae for a straightforward computation of q , namely Equations (12), (18) and (24). In contrast to most of the literature we also covered situations in which the probabilities for different question regimes do not have to be more or less equal. Here, the chances for Alice and Bob to win can improve considerably.

Our extension of the result in [25] is presented with a fully self-contained proof. We did not consider multipartite games, which do not always allow a quantum advantage [28] and where general problems, like computing the maximum winning probability given an arbitrary amount of entangled particles, can be very complex and even QMA-hard [29].^{CG}

We also^{CG} analyzed different game strategies in Monte Carlo simulations. We would like to point to the possibility of winning the game by exploiting systematic biases in the generation of question regimes. In such a situation, Alice and Bob can have a good chance to produce S_1 values substantially above 2 without entangled particles. Therefore, when the security of quantum communication is based on observed violations of CHSH inequalities it is important to check the question regimes for systematic biases.

Author Contributions: Conceptualization, methodology, formal analysis, proofs, simulations and writing the original draft presentation, C.G.; writing, review and editing, C.G., P.B. and E.M.P. All authors have read and agreed to the published version of the manuscript.

Funding: E.M.P. and P.B. were supported by ONRG grant N62909-19-1-2000. E.M.P. was also supported by AFOSR grant FA8655-23-1-7220.

Acknowledgments: C.G. would like to thank Andrei Khrennikov and Jan-Åke Larsson for helpful conversations at the QIP22 Conference on Quantum Foundations and at other occasions.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

1. Bell, J.S. On the Einstein-Podolsky-Rosen paradox. *Physics* **1964**, *1*, 195.
2. Bell, J.S. On the Problem of Hidden Variables in Quantum Mechanics. *Rev. Mod. Phys.* **1966**, *38*, 447.
3. Bell, J.S. Introduction to the hidden variables question. Technical Report Ref.TH.1220-CERN, CERN, 1970.
4. Bell, J.; Shimony, A.; Horne, M.; Clauser, J. An Exchange on Local Beables. *Dialectica* **1985**, *39*, 85–110.
5. Bell, J.S. Free variables and local causality. In *Speakable and unspeakable in quantum mechanics*; Cambridge University Press, 1987; chapter 12.
6. Bell, J.S. *Speakable and unspeakable in quantum mechanics*; Cambridge University Press, 1987.
7. Brassard, G.; Cleve, R.; Tapp, A. Cost of Exactly Simulating Quantum Entanglement with Classical Communication. *Phys. Rev. Lett.* **1999**, *83*, 1874.
8. Gisin, N. *Quantum Chance: Nonlocality, Teleportation and Other Quantum Marvels*; Springer, 2014.
9. Brunner, N.; Cavalcanti, D.; Pironio, S.; Scarani, V.; Wehner, S. Bell nonlocality. *Rev. Mod. Phys.* **2014**, *86*, 419.
10. Scarani, V. *Bell Nonlocality*; Oxford University Press, 2019.
11. López-Incera, A.; Hartmann, A.; Dür, W. Encrypt me! A game-based approach to Bell inequalities and quantum cryptography. *European Journal of Physics* **2020**, *41*, 065702. doi:10.1088/1361-6404/ab9a67.
12. Yuen, H. The Complexity of Entanglement. Lecture notes for CSC2429/MAT1751 (Fall 2020) at the University of Toronto, 2020.
13. Welsch, B.; Thron, C. (November 4, . The 'Quantum Game Show': A Very Simple Explanation of Bell's Theorem in Quantum Mechanics. SSRN: <https://ssrn.com/abstract=3956512> or <http://dx.doi.org/10.2139/ssrn.3956512>, 2021.
14. Toner, B.F.; Bacon, D. Communication Cost of Simulating Bell Correlations. *Phys. Rev. Lett.* **2003**, *91*, 187904.
15. Svozil, K. Communication cost of breaking the Bell barrier. *Phys. Rev. A* **2005**, *72*, 050302. doi:10.1103/PhysRevA.72.050302.
16. Hall, M.J.W.; Branciard, C. Measurement-dependence cost for Bell nonlocality: Causal versus retrocausal models. *Phys. Rev. A* **2020**, *102*, 052228.
17. Proctor, T.; Revelle, M.; Nielsen, E.; Rudinger, K.; Lobser, D.; Maunz, P.; Blume-Kohout, R.; Young, K. Detecting and tracking drift in quantum information processors. *Nature Communications* **2020**, *11*, 5396. doi:10.1038/s41467-020-19074-4.
18. Broadbent, A.; Schaffner, C. Quantum cryptography beyond quantum key distribution. *Designs, Codes and Cryptography* **2016**, *78*, 351–382. doi:10.1007/s10623-015-0157-4.
19. Huang, L.; Zhou, H.; Feng, K.; Xie, C. Quantum random number cloud platform. *npj Quantum Information* **2021**, *7*, 107. doi:10.1038/s41534-021-00442-x.
20. Dirkse, B.; Pompili, M.; Hanson, R.; Walter, M.; Wehner, S. Witnessing entanglement in experiments with correlated noise. *Quantum Sci. Technol.* **2020**, *5*.
21. Lawson, T.; Linden, N.; Popescu, S. Biased nonlocal quantum games. arXiv:1011.6245, 2010.
22. Singh, P.; Faujdar, J.; Sarkar, M.; Kumar, A. Bell's inequality with biased experimental settings. *Quantum Information Processing* **2022**, *21*, 167. doi:10.1007/s11128-022-03504-x.
23. Zhang, Y.; Glancy, S.; Knill, E. Asymptotically optimal data analysis for rejecting local realism. *Physical Review A* **2011**, *84*, 062118.
24. Gill, R.D. Accardi contra Bell (cum mundi): The Impossible Coupling. In *Mathematical Statistics and Applications: Festschrift for Constance van Eeden, Hayward, CA: Institute of Mathematical Statistic*; Moore, M.; Froda, S., Eds.; Number 42 in Lecture notes-monograph series, Institute of Mathematical Statistic, 2003; pp. 133–154.
25. Gill, R.D. Statistics, Causality and Bell's Theorem. *Statist. Sci.* **2014**, *29*, 512–528.

26. Elkouss, D.; Wehner, S. (Nearly) optimal P values for all Bell inequalities. *npj Quantum Information* **2016**, *2*, 16026. doi:10.1038/npjqi.2016.26.
27. Adenier, G.; Khrennikov, A. Test of the no-signaling principle in the Hensen loophole-free CHSH experiment. *Fortschr. Phys.* **2017**, *65*.
28. Almeida, M.L.; Bancal, J.D.; Brunner, N.; Acín, A.; Gisin, N.; Pironio, S. Guess Your Neighbor's Input: A Multipartite Nonlocal Game with No Quantum Advantage. *Phys. Rev. Lett.* **2010**, *104*, 230404. doi:10.1103/PhysRevLett.104.230404.
29. Natarajan, A.; Vidick, T. Low-Degree Testing for Quantum States, and a Quantum Entangled Games PCP for QMA. 2018 IEEE 59th Annual Symposium on Foundations of Computer Science (FOCS), 2018, pp. 731–742. doi:10.1109/FOCS.2018.00075.
30. Ekert, A.K. Quantum Cryptography Based on Bell's Theorem. *Physical Review Letters* **1991**, *67*, 661–663.
31. Gallus, C.; Blasiak, P.; Pothos, E.M. Quantifying and Interpreting Connection Strength in Macro- and Microscopic Systems: Lessons from Bells Approach. *Entropy* **2022**, *24*. doi:10.3390/e24030364.
32. Gallus, C.; Pothos, E.M.; Blasiak, P.; Yearsley, J.M.; Wojciechowski, B.W. Bell correlations outside physics. *Scientific Reports* **2023**, *13*, 4394. doi:10.1038/s41598-023-31441-x.
33. Summers, S.J.; Werner, R. Maximal violation of Bell's inequalities is generic in quantum field theory. *Communications in Mathematical Physics* **1987**, *110*, 247 – 259. doi:cmp/1104159237.
34. McKague, M.; Yang, T.H.; Scarani, V. Robust self-testing of the singlet. *Journal of Physics A: Mathematical and Theoretical* **2012**, *45*, 455304. doi:10.1088/1751-8113/45/45/455304.
35. Grilo, A. A simple protocol for verifiable delegation of quantum computation in one round. arXiv:1711.09585v2, 2020.
36. Clauser, J.F.; Horne, M.A.; Shimony, A.; Holt, R.A. Proposed Experiment to Test Local Hidden-Variable Theories. *Phys. Rev. Lett.* **1969**, *23*, 880–884.
37. Blasiak, P.; Pothos, E.M.; Yearsley, J.M.; Gallus, C.; Borsuk, E. Violations of locality and free choice are equivalent resources in Bell experiments. *Proc. Natl. Acad. Sci. USA* **2021**, *118*, e2020569118.
38. Bierhorst, P. A Rigorous Analysis of the Clauser–Horne–Shimony–Holt Inequality Experiment When Trials Need Not be Independent. *Foundations of Physics* **2014**, *44*, 736–761. doi:10.1007/s10701-014-9811-3.
39. Hoeffding, W. Probability Inequalities for Sums of Bounded Random Variables. *Journal of the American Statistical Association* **1963**, *58*, 13–30.
40. Combes, R. An extension of McDiarmid's Inequality. arXiv:1511.05240v1, 2015.
41. Chvátal, V. The tail of the hypergeometric distribution. *Discrete Mathematics* **1979**, *25*, 285–287. doi:https://doi.org/10.1016/0012-365X(79)90084-0.
42. McDiarmid, C., On the method of bounded differences. In *Surveys in Combinatorics, 1989: Invited Papers at the Twelfth British Combinatorial Conference*; Siemons, J., Ed.; London Mathematical Society Lecture Note Series, Cambridge University Press: Cambridge, 1989; pp. 148–188. doi:DOI: 10.1017/CBO9781107359949.008.
43. Corrado, C.J. The exact distribution of the maximum, minimum and the range of Multinomial/Dirichlet and Multivariate Hypergeometric frequencies. *Statistics and Computing* **2011**, *21*, 349–359. doi:10.1007/s11222-010-9174-3.
44. Bonetti, M.; Cirillo, P.; Ogay, A. Computing the exact distributions of some functions of the ordered multinomial counts: maximum, minimum, range and sums of order statistics. *R. Soc. open sci.* **2019**, *6*.
45. Tamura, K.; Shikano, Y. Quantum Random Numbers Generated by a Cloud Superconducting Quantum Computer. International Symposium on Mathematics, Quantum Theory, and Cryptography; Takagi, T.; Wakayama, M.; Tanaka, K.; Kunihiro, N.; Kimoto, K.; Ikematsu, Y., Eds.; Springer Singapore: Singapore, 2021; pp. 17–37.
46. Jacak, J.E.; Jacak, W.A.; Donderowicz, W.A.; Jacak, L. Quantum random number generators with entanglement for public randomness testing. *Scientific Reports* **2020**, *10*, 164. doi:10.1038/s41598-019-56706-2.