

City Research Online

City, University of London Institutional Repository

Citation: Wilson, C. M., Thomson, W. & D'Ath, P. (2013). Seeing you through London 2012: eye care at the Olympics. British Journal of Sports Medicine, 47(7), pp. 463-466. doi: 10.1136/bjsports-2013-092152

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: https://openaccess.city.ac.uk/id/eprint/5549/

Link to published version: https://doi.org/10.1136/bjsports-2013-092152

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

 City Research Online:
 http://openaccess.city.ac.uk/
 publications@city.ac.uk

74

75

76

77

78

79

80

81

82



¹Department of Visual Neuroscience, UCL Institute of Ophthalmology, London, UK ²Division of Optometry & Visual Science, City University London, London, UK

Correspondence to

 $\begin{array}{c} Q1_{2}^{1} \\ \Lambda_{3}^{2} \\ Q2_{4}^{2} \\ \Lambda_{5}^{2} \\ Q3^{6} \\ \Lambda_{7}^{7} \\ Q6^{8} \\ \Lambda_{9}^{9} \end{array}$

Q40

11

12

13

14

15

16

17

18

19

20

21

22

23

24

2.5

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

Dr Clare M Wilson, Department of Visual Neuroscience, UCL Institute of Ophthalmology, 11-43 Bath Street, London EC1V 9EL, UK; clarewil25@yahoo.com

Accepted 13 June 2013

Seeing you through London 2012: eye care at the Paralympics

Clare M Wilson,¹ W David Thomson,² Penny J D'Ath²

Background The provision of eye care services for competitors and support teams is integral to the modern Olympic Games. The eye clinic for the London 2012 Paralympic Games employed a multidisciplinary team of eye care professionals using state-of-the-art instrumentation to provide the highest level of eye care. The detailed organisation of the eye care clinic at London 2012 is described in a companion paper which summarises the eye care clinic during the London 2012 Olympic Games. These two reports will aid in planning eve care clinics at future Games.

Aim This paper summarises the organisation of the eye clinic and provides outline audit data relating to eye conditions encountered during the Paralympic Games. **Results** A total of 870 patients representing 102 countries attended the eye clinic. 274 (31.5%) were competitors: the remainder were trainers and support staff. No serious ocular injuries resulted from competitor injury in the field of play during the Paralympic Games, although seven patients were referred urgently to hospital eye services for conditions including orbital cellulitis, retinal detachment, exudative macular degeneration, corneal ulcer, Stevens-Johnson syndrome and macular oedema. A total of 749 spectacles, 14 contact lenses and 7 low-vision aids were dispensed. **Conclusions** By combining excellent facilities and equipment with a multidisciplinary team of eye care professionals, we feel we provided the highest level of eye care, providing a legacy for future Games.


INTRODUCTION

Since the first modern Olympic Games held in Athens in 1896, the Olympic Charter has grown to include the provision of many allied services for athletes and their support teams; one of which is the eye clinic. The Paralympic Games benefits from the same healthcare provision for its athletes and entourage. At the London 2012 Paralympics, 164 countries and over 4000 competitors¹ competed in front of sell-out crowds.

In 2009, one ophthalmologist and two optometrists were appointed (the authors CMW, WDT and PJD) to lead the eye care service. Little information is published regarding eye care services at previous Paralympic Games, although a small but useful amount of information was obtained from personal communication from the Committee of the Paralympic Games.

83 Many systemic diseases have ocular complica-84 tions.² Paralympians tend to have more complex 8.5 ocular pathology than Olympians. Indeed, some 86 competitors are eligible to compete as Paralympians 87 due solely to visual impairment. Paralympians not 88 competing in visual impairment categories may also 89 have ocular conditions related to their underlying 90 systemic condition. For example, those with cerebral 91 palsy may have cerebral visual impairment, whereas 92 those with multiple sclerosis may have optic neur-93 opathy and competitors with polio may have ocular 94 motility disturbance. As many people of the support 95 team were former Paralympians, these patients 96 also had more complex ocular needs for the same 97 reasons. An unpublished report from Sydney stated 98 that many patients seen during the Paralympics had 99 a range of eye conditions varying from 'optic 100 neuritis secondary to malaria, sickle-cell retinopathy, 101 and there were a number of patients with corneal 102 conditions caused by birth trauma or infantile infec-103 tions' (Personal communication from LOCOG. 104 Unpublished report "Eye Service Sydney") resulting 105 in a 'higher level of ophthalmic complexity' than 106 found during the Olympic Games. (Personal 107 communication from LOCOG. Unpublished report 108 $\dot{\mathbf{Q}}_{09}$ "UPDATED—Service Specification—Optometry).

On the basis of figures from the London 2012 Paralympics, around 19% of athletes are competing with visual impairments³ (Personal communication



To cite: Wilson CM, Thomson WD, D'Ath PJ. Br J Sports Med Published Online First: [please include Day Month Year] doi:10.1136/ bjsports-2013-092614

110

111

112

113 114

115

116

117

118

119

120

121

122 123

124

125

126

127

128

Short report

Characteristic	Competitors	Non-competitors
Total number seen (%)	274 (31.5)	596 (68.5)
Male (M):female (F)	170:104	445:151
Age (mean: range: n)	M: 33.9: (18–56): n=170; F: 32.7: (17–51): n=104	M: 49.7: (19–75): n=44 F: 45.6: (19–79): n=151

from IPC, 27 May 2013). Visual impairment categories exist for the following sports: athletics, cycling, equestrian, football 5-a-side, goalball, judo, rowing, sailing and swimming.

This is the first paper to analyse eye care data from any Paralympic Games.

AIM

This paper aims to provide outline audit data relating to the patients attending the eye clinic during the 22 days of Paralympic Games.

METHODS

We have assimilated data on the usage of the eye care clinic at London 2012 with reference to demographics, reason for attendance. injuries among competitors and spectacles dispensed.

Layout, equipment and staffing

Figure 2 Number of patients

attending eye clinic per day.

As described in detail in our companion paper,⁵ the eve clinic formed part of a purpose-built polyclinic situated in the Athletes' Village and was designed to accommodate both com-petitors and their support teams. Details of room sizes, equip-ment, diagnostic drugs and volunteers are listed in this paper. The main findings from the Olympic Games were that 1406 patients from 154 countries were seen. No serious eye injuries or referrals occurred, but a number of eye diseases including glaucoma, diabetic retinopathy and macular degeneration were detected. Patients predominantly attended the clinic for a full refractive status check and 973 pairs of spectacles and 50 pairs of contact lenses were dispensed.⁵

Of the 309 optometrists and 103 dispensing opticians who applied to become Games Makers, 104 optometrists and 53 dispensing opticians were shortlisted of whom 8 optometrists and 6 dispensing opticians were selected for the Paralympic Games. Six ophthalmologists were appointed as 'Specialists' for 8 days

Table 2 Reason for visit			
Characteristic	Competitors n=196	Non-competitors n=419	
Reduced vision	75 (38.3%)	167 (39.9%)	
Distance vision	31	32	
Near vision	30	98	
Distance and near vision	13	35	
Not specified	1	2	
Routine eye examination	27 (13.8%)	43 (10.3%)	
Replacement spectacles (lost/ broken/left at home)	24 (12.2%)	145 (34.6%)	

of the Paralympic Games period and were not subject to the normal Games Maker recruitment process.

The eye clinic was open for 22 days from 07:00 to 23:15 throughout the Paralympics Games period. The predicted staff numbers required throughout the Games period are shown in figure 1.

RESULTS

Audit of patients seen


A total of 870 patients representing 102 countries attended the eve clinic over the period of the Paralympic Games. Of these, almost one-third were competitors (n=274; 31.5%), and 596 (68.5%) comprised members of the support team.

Table 1 shows the demographic characteristics of patients who presented to the clinic.

Figure 2 shows the number of patients attending the clinic on each day throughout the period of the Paralympic Games. The maximum number of patients examined in 1 day occurred on day 5 of the competition when 76 patients were seen.

The peak times that competitors attended the clinic were 11:00 and 15:00 and, for non-competitors, were 11:00, 15:00 and 21:00.

Almost 40% of patients reported 'reduced vision' (competitor, 38%: non-competitor, 40%) (table 2). Among the noncompetitors, 59% of cases of reduced vision related to problems with reading/near vision. A total of 14% of the competitors and 10% of the non-competitors were asymptomatic and attended for a routine eye examination. Non-competitors (35%) were three times more likely to present requiring replacement spectacles compared with competitors (12%). There were four minor ocular injuries that required specialist eye care, one of which



Type of optical appliance	n=756 (%)
Distance vision	361 (47.8)
Near vision	312 (41.3)
Varifocals	60 (7.9)
Bifocals	16 (2.1)
Low vision aids	7 (0.9)

2.62

was a mild corneal thermal injury caused by debris from the fireworks at the opening ceremony.

Of the 870 patients who attended the clinic, 14 (2%) had contact lens related issues or required new lenses (one required a cosmetic glass design) and approximately 72 (8%) were referred for an ophthalmological opinion. Ophthalmologists saw between 6 and 12 patients per day over a period of 8 days. Only 8 days had on-site ophthalmic cover as organised by

linking to predicted demand. The majority of patients required a single visit to the clinic (excluding the collection of spectacles). Exceptions were patients with contact lens issues or those with conditions requiring ophthalmological management who had up to four follow-up visits.

There were 749 pairs of spectacles and 7 low-vision aids dis-pensed to seven patients (1%) reporting non-tolerance to their new spectacles. Spectacle type was determined for all 749 pairs (see table 3).

Table 4 shows the number of ocular conditions by visual impairment classifications/ sport. Of the 38 cases of visual impairment, 8 (21%) were caused by high myopia and 5 (13%) were caused by congenital nystagmus.

There were no serious ocular injuries during the Paralympic Games, although there were seven referrals to hospital eye services. Conditions requiring this extralevel of care included orbital cellulitis (patient required admission), retinal detachment, exudative macular degeneration, corneal ulcer, Stevens-Johnson syndrome and macular oedema. There was also one private referral for chronic bilateral epiphora.

Table 4 Ocular condition by visual impairment classification
--

Sport	Classification	М	Condition	F	Condition
Goalball (all players blindfolded to ensure fairness)		n=3		n=3	
				1	Unknown
Blind	B1	1	Prosthetic eye		
Visually impaired/partially sighted	B2	2	Bilateral optic atrophy	2	Rod cone dystrophyBilateral optic atrophy
			 End stage glaucoma and nystagmus 		
5-a-side football		n=1			
Visual impairment		1	Unknown		
Cycling		n=1			
Visual impairment	В	1	Retinitis pigmentosa		
Judo		n=3		n=2	
Blind	B1	1	Retinitis pigmentosa with macular dystrophy		
Visually impaired/partially sighted	B2	1	Amblyopia (—6D)	1	Unknown
Visually impaired / partially sighted	B3	1	Myopia and astigmatism (–6D)	1	High myopia (>—21D)
(higher number=better vision)					
Rowing		n=1		n=2	
	LTA-VIB2	1	High hyperopia (+13D)	1	Congenital nystagmus and myopia (-12
	LTA-VIB3			1	Congenital glaucoma
Swimming		n=1			
Visual impairment	S12	1	Congenital nystagmus		
Field sports		n=9		n=2	
Visual impairment	F12	3	 Congenital nystagmus 	2	 Congenital cataracts
			► Retinal detachment LE and anisometropic high myopia		▶ High myopia (-12D)
			(RE −16D, LE −9D) ► Unknown		
Visual impairment	F13	6	 RTA, one blind eye 		
	115	0	 RTA, one blind eye Bad fall as child, high myopia (-21D) 		
			 Previous retinoblastoma 		
			 Bilateral congenital optic disc atrophy and nystagmus 		
			 Congenital blindness 		
Track		n=5		N=5	
Visual impairment	T11	1	AION aged 21	1	Poor vision since measles aged 3
Visual impairment	T11/T12			1	End stage POAG
Visual impairment	T12	1	Unknown	2	 Congenital cataracts
					 Microphthal-mos and nystagmus
Minual immediate	T10	2	2. List muses (PE, 1CD LE, 10D, PE, 12D	1	2 Claurante
Visual impairment	T13	3	3×high myope (RE: –16D, LE –10D; RE –12D, LE –11D, RE –28D, LE –12)	1	? Glaucoma
Total		24	LL = 110, IL = 200, LL = 12	14	

Short report

385 Of the non-competitors, the most common reasons for referral 386 to the ophthalmologist were glaucoma (n=5), ocular complications of diabetes (n=3) and conjunctivitis (viral and bacterial: 387 n=2). We also encountered some more unusual pathologies such 388 389 as Leber's congenital amaurosis, Stevens-Johnson syndrome, 390 orbital cellulitis, nystagmus, rod cone dystrophies, previous retino-391 blastoma and congenital cataracts. In cases that required long-term 392 care in the patient's own country, a letter with the findings and **O8**3 appropriate images from the OCT or visual field analyser were 394 given to the patient in CD ROM format.

No adaptations were made to the clinic from the Olympic
Games, perhaps as the set-up had been designed with provisions
for Paralympic athletes in mind. All wheelchair patients transferred themselves to the main consulting room chair.

We recommend that a full service should be run at future Paralympics with four optometrists, two dispensing opticians and one ophthalmologist available throughout the Games period.

404 SUMMARY

402

403

417

418

419

420

421

422

423

424

42.5

426

427

42.8

429

430

431

432

433

434

435

436 437

438

439

440

441

442 443

444

445

A total of 870 patients from 102 countries attended the eye clinic over a period of 22 days with a peak of 76 attendees on day 5 of the competition. Of these, 274 attendees were competitors and the remainder were trainers and support staff.

A total of 749 pairs of spectacles were dispensed. Just over
50% were prescribed for near vision/reading. This was significantly more than at Sydney 2000 where 261 pairs of spectacles
were dispensed. Fourteen contact/therapeutic lenses were fitted
and seven low-vision devices were issued.

No major ocular injuries occurred from sports, although
seven patients required referral to the hospital eye service, one
requiring hospital admission.

We believe we have provided a legacy of eye care for future Paralympic Games to build on.

What are the new findings

- A total of 870 competitors and support staff from 102 countries attended the Paralympic eye clinic at London 2012 over a period of 22 days.
- There were no serious ocular injuries during the Paralympic Games. Seven patients required urgent referral to the hospital eye service for conditions including orbital cellulitis and retinal detachment.
- The majority of patients attended the clinic to have their refractive status checked, and a total of 749 pairs of spectacles, 14 pairs of contact lenses and 7 low-vision aids were dispensed.

How might it impact on clinical practice in the near future

- Approximately one in five of all Paralympians are classified as having a visual impairment.
- Patients seen at the eye care clinic had more complex optometric and ophthalmological needs (eg, Stevens-Johnson syndrome, orbital cellulitis, nystagmus, rod cone dystrophies, previous retinoblastoma, congenital cataracts, etc) than those found during the Olympic Games.
- ► For this reason, we suggest that a full service should be run with four optometrists, two dispensing opticians and one ophthalmologist available throughout the Paralympic Games period.
- It would have been beneficial to have had ophthalmologists on-site for longer periods of the day and for the entirety of the Paralympic Games period because of the complexity of ophthalmic problems.

Acknowledgements The authors would like to express their thanks to the volunteers of the London 2012 eye clinic: Caroline Christie (Contact Lens Coordinator), Katherine Anguige, Nathanael Anguige, Robin Baker, Liz Baron, David Bennett, Susan Blakeney, Chris Boyde, Gillian Bruce, Niral Charadva, Oliver Comyn, Suzy Connolly, Shaun Crome, Ruth Cuthbert, Sue Daniel, Ruth Davies, Lee Davis, Helen Denton, Michelle Derbyshire, Julian de Silva, Tessa Fayers, Jessica Fielding, Susan Gibbons, Lucy Hall, Laura Hing, Fiona Hiscox, Byki Huntjens, Hari Jayaram, Margaret Lawrence, Jennifer Lee, Jake Low-Beer, Hamish MacDonald, Andrew Mace, Anthony Martin, Sara McCullough, Michael Offord, David Parkins, Sheetal Patel, Catherine Porter, Yasmin Riaz, Scott Robbie, Chris Roberts, Sally Rosedale, Claire Ruddock, Bharat Rughani, Anaeka Sodha, Karen Sparrow, Elaine Styles, Vanessa Uden, Ellen Watkins, James Wolffsohn, Karen Wong.

Contributors CMW, WDT and PJD made a significant contribution to the conception and design of the eye clinic at London 2012, the collection and interpretation of data and the drafting and subsequent refinement of the paper. We confirm that all authors have approved the final version submitted.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Willick SE, Webborn N, Emery C, et al. The epidemiology of injuries at the London 2012 Paralympic Games. Br J Sports Med 2013;47:426–32.
- 2 Mohsenin A, Huang JJ. Ocular manifestations of systemic inflammatory diseases. Conn Med 2012;76:533–44.
- 3 Figure calculated from 'London 2012 Paralympic athletes: the full list of competitors and disciplines'. The Guardian. http://www.guardian.co.uk/news/datablog/2012/aug/ 30/every-paralympic-athlete-and-their-sport-data (accessed 7 May 2013).
- 4 http://www.paralympic.org/sites/default/files/document/120716152047682_ ClassificationGuide_2.pdf (accessed 7 May 2013).
- 5 D'Ath PJ, Thomson WD, Wilson CM. Seeing you through London 2012: eye care at the Olympics. *Br J Sports Med* 2013;47:463–6.

499 500

501

502

503

504

505 506

507

508

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

512

446 447 448