



City Research Online

City, University of London Institutional Repository

Citation: Anderson, L. B., Gray, J., Grayson, D., He, Y. & Lukas, A. (2010). Yukawa Couplings in Heterotic Compactification. *Communications in Mathematical Physics*, 297(1), pp. 95-127. doi: 10.1007/s00220-010-1033-8

This is the unspecified version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/883/>

Link to published version: <https://doi.org/10.1007/s00220-010-1033-8>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Yukawa Couplings in Heterotic Compactification

Lara B. Anderson^{1,5}, James Gray², Dan Grayson³, Yang-Hui He^{2,4}, André Lukas²

¹ Department of Physics and Astronomy, University of Pennsylvania,
209 South 33rd Street, Philadelphia, PA 19104-6396, USA.

²Rudolf Peierls Centre for Theoretical Physics, University of Oxford,
1 Keble Road, Oxford OX1 3NP, UK.

³Department of Mathematics, University of Illinois at Urbana-Champaign,
1409 W. Green Street, Urbana IL 61801, USA.

⁴Merton College, Oxford, OX1 4JD, UK.

⁵ Institute for Advanced Study, School of Natural Sciences,
Einstein Drive, Princeton, NJ, 08540, USA.

Abstract

We present a practical, algebraic method for efficiently calculating the Yukawa couplings of a large class of heterotic compactifications on Calabi-Yau three-folds with non-standard embeddings. Our methodology covers all of, though is not restricted to, the recently classified positive monads over favourable complete intersection Calabi-Yau three-folds. Since the algorithm is based on manipulating polynomials it can be easily implemented on a computer. This makes the automated investigation of Yukawa couplings for large classes of smooth heterotic compactifications a viable possibility.

email: ¹andlara@hep.physics.upenn.edu, ²j.gray1@physics.ox.ac.uk, hey@maths.ox.ac.uk,
lukas@physics.ox.ac.uk, ³dan@math.uiuc.edu

Contents

1	Introduction	2
2	Yukawa couplings in heterotic compactification	4
2.1	Rephrasing in terms of cohomologies	5
2.2	The arena: positive monad bundles over CICYs	7
3	Calculating Yukawa couplings: general procedure	9
3.1	Polynomial representation of line bundle cohomology	10
3.2	$SU(3)$ vector bundles and E_6 GUTS	11
3.2.1	Polynomial representatives for families in $H^1(X, V)$	11
3.2.2	Polynomial representatives for $H^3(X, \wedge^3 V)$	12
3.2.3	Computing Yukawa couplings	13
3.2.4	A simple E_6 example	14
3.3	$SU(4)$ vector bundles and $SO(10)$ GUTs	15
3.3.1	Polynomial representatives for Higgs multiplets in $H^1(X, \wedge^2 V)$	16
3.3.2	Computing Yukawa couplings	16
3.4	$SU(5)$ vector bundles and $SU(5)$ GUTs	17
3.4.1	Polynomial representatives for $H^1(X, \wedge^2 V)$	17
3.4.2	Polynomial representatives for $H^2(X, \wedge^4 V)$	18
3.4.3	Polynomial representatives for $H^2(X, \wedge^2 V)$	19
3.4.4	Computing Yukawa couplings	20
4	An example: One Higgs multiplet and one heavy family	20
4.1	The model	20
4.2	Engineering Higgs multiplets	21
4.3	The mass matrix	22
5	Conclusions	23
A	Koszul Complex and Polynomial Representatives	25
B	Proof of Equivalence of Formulations	27
B.1	Chain complexes and bicomplexes	27
B.2	Cup products in hypercohomology of sheaves	30
B.3	Symmetric and exterior powers of complexes	31

1 Introduction

Some of the most important pieces of data defining a phenomenological theory of particle physics are the Yukawa couplings. Since these parameters determine particle masses and interactions, no theory’s phenomenology can be understood in even a rudimentary manner without knowledge of their values. In string phenomenology, Yukawa couplings are usually some of the first things one attempts to calculate once the low energy particle spectrum of a model is known [1, 2, 3]. However, despite their importance, in many cases it is not known how to carry out the calculations in practice.

For compactifications of heterotic string theory and M-theory on Calabi-Yau three-folds, Yukawa couplings have been calculated in only a relatively small number of cases. Examples include orbifold compactifications and heterotic models with “standard embedding”, that is, models where the gauge bundle is chosen to be the tangent bundle of the Calabi-Yau manifold. Aside from these, only a few other, isolated, examples appear [4, 5, 6, 7], with some of these being closely related to the standard embedding.

In this paper, we considerably improve this situation by providing a simple, easy to implement, algorithm for calculating the Yukawa couplings in a large class of heterotic compactifications on smooth Calabi-Yau spaces with “non-standard embedding”. In such compactifications the gauge fields are defined in terms of a general, poly-stable holomorphic vector bundle [8]. Our approach is in the spirit of the recent papers [9, 10], where a systematic analysis of such general heterotic compactifications by means of computational algebraic geometry has been pursued.

The manifolds we will consider are the favourable complete intersection Calabi-Yau (CICY) manifolds [11], which comprise a set of 4515 three-folds. In addition, we consider vector bundles defined over these manifolds that are built using the monad construction [12]. For simplicity, we will present our method for the case where certain bundle cohomology groups vanish, as summarised in Table 2, but it is likely that the basic ideas can be extended to all stable bundles on CICY manifolds. These conditions are automatically satisfied for positive monad bundles and a large number of “not too negative” monad bundles. The class of positive monad bundles has been recently studied in references [10, 13, 14, 15], and the methods described in the present paper represent a further step towards a systematic analysis of their phenomenological properties.

Our method calculates the Yukawa couplings that appear in the superpotential of the four-dimensional theory. They are related to the physical Yukawa couplings by a field rotation that brings the matter field kinetic terms into canonical form. Unfortunately, the matter field kinetic terms and, hence, the required field redefinitions, are not explicitly known, so the physical Yukawa couplings cannot be computed directly. Numerical calculations [16] may be the only way to overcome this common limitation, and we have nothing more to say about it in the present paper. In practice, this means that only certain invariants of the Yukawa couplings, which are unchanged under redefinition of the matter fields, can be regarded as physical. For example, in cases where one can talk about a Yukawa matrix, such as in a model

with $SO(10)$ gauge group and a single Higgs representation in $\mathbf{10}$, the rank of this matrix is physically meaningful.

Of the many different methods of constructing vector bundles, the monad construction is one that has been of consistent interest in the physics literature over the years (see, for example, references [10, 13, 17, 18, 19]). These constructions, which will be reviewed in section 2.2, lend themselves nicely to methods of computational algebraic geometry. This feature, which allows us to systematically study large classes of such bundles at a time, is one of main motivations for focusing on monad bundles in this paper. We shall consider the cases of $SU(n)$ bundles, where $n = 3, 4, 5$, corresponding to the GUT visible sector gauge groups E_6 , $SO(10)$ and $SU(5)$, respectively. We will give examples throughout our discussion, but in particular, in section 4, we give a detailed presentation of an $SO(10)$ model. We show how to engineer models with a single $\mathbf{10}$ multiplet of $SO(10)$. In addition, we show that the rank of the Yukawa matrix for the $\mathbf{16}$ multiplets of a model engineered in this way is one. As a result, these cases correspond to compactifications with one heavy family.

Before we delve into technicalities let us briefly outline the basic method for computing Yukawa couplings, which is quite simple in principle. For a heterotic compactification on a Calabi-Yau three-fold X , the families can be identified as elements of the cohomology group $H^1(X, V)$ of the gauge bundle V . Anti-families correspond to $H^2(X, V) \simeq H^1(X, V^*)$, but our focus will be on models without anti-families, a property which is automatic for positive monad bundles, as we will review. To be specific, let us discuss the case of an $SU(3)$ bundle which leads to the low-energy gauge group E_6 and families in $\mathbf{27}$ representations. In this case, we are interested in the $\mathbf{27}^3$ Yukawa couplings, that is, we need to understand the map $H^1(X, V) \times H^1(X, V) \times H^1(X, V) \rightarrow H^3(X, \wedge^3 V) \simeq \mathbb{C}$ (the last equivalence holds because $\wedge^3 V \simeq \mathcal{O}_X$ for an $SU(3)$ bundle V and $h^3(X, \mathcal{O}_X) = 1$). It turns out, for monad bundles, that the “family cohomology group” $H^1(X, V)$ can be represented by a quotient of polynomial spaces, containing polynomials of certain, well-defined, degrees. Likewise, we can represent the “Yukawa cohomology group” $H^3(X, \wedge^3 V)$ by a quotient of polynomial spaces which, of course, must be one-dimensional. Let Q be a representative of the single class in this quotient and P_I , where $I, J, K, \dots = 1, \dots, h^1(X, V)$, a polynomial basis for the families. Then the Yukawa couplings λ_{IJK} are obtained by multiplying three family representatives. The result represents an element in the one-dimensional Yukawa quotient space and must, hence, be proportional to Q . The constant of proportionality is precisely the desired Yukawa coupling, so $[P_I P_J P_K] = \lambda_{IJK} [Q]$, where $[\cdot]$ denotes the class in the quotient space. Hence, calculating Yukawa couplings is reduced to a simple procedure of multiplying polynomials and projecting the result onto the class representative Q . For the cases of bundles with structure groups $SU(4)$ and $SU(5)$ the procedure is analogous although slightly more complicated.

The plan of this paper is as follows. In the next section, we introduce the general methodology available for computing Yukawa couplings in heterotic compactifications and the arena in which we shall be working: positive monad bundles over the complete intersection Calabi-Yau manifolds. In section

3, we proceed to outline the procedure for calculating Yukawa couplings in such compactifications. We split the discussion into several subsections - one for each of the possible visible sector gauge groups of interest (E_6 , $SO(10)$, and $SU(5)$). Section 4 contains a detailed discussion of a one-Higgs $SO(10)$ model. In section 5 we end with conclusions and prospects. A technical result required in the bulk of the text, as well as the proof that the polynomial-based procedure, outlined in section 3, indeed reproduces the physical Yukawa couplings are presented in the Appendix.

2 Yukawa couplings in heterotic compactification

After a brief review of heterotic compactifications and Yukawa couplings, in §2.1 we will describe how the problem of calculating Yukawa couplings can be rephrased in terms of bundle cohomology groups. We will also hint at our method for calculating these interactions based thereon, leaving the technical details of the actual procedure to the following section. In addition, in §2.2, we shall describe the basic geometrical setup of our class of Calabi-Yau manifolds and bundles.

Let us start by considering how Yukawa couplings are usually described in heterotic compactifications. The matter fields in Calabi-Yau compactifications of heterotic string theory and M-theory descend from the internal parts of the gauge fields and their superpartners. In the case where we have a visible sector gauge bundle V over the Calabi-Yau threefold X taking values in a subgroup G of E_8 , the low energy observable gauge group, H , is given by the commutant of G in E_8 . The matter fields arise in the decomposition of the adjoint of E_8 under $G \times H$: $\mathbf{248} = \sum_I (R_G^I, R_H^I)$ with R_G^I and R_H^I being representations of the groups G and H respectively, indexed by I . See Table 1 for a complete list of the decompositions of the $\mathbf{248}$ of E_8 and associated cohomologies for standard heterotic theories.

The reduction ansatz for the holomorphic part of the gauge field A in 10-dimensions is, to lowest order [20],

$$A = \sum_I C_I^i u_I^a T_{ai} + A_{\text{BG}} . \quad (2.1)$$

Here A_{BG} is the background gauge field vacuum expectation value satisfying the hermitian Yang-Mills equations. The first term in (2.1) gives rise to the four-dimensional matter fields C_I^i , where I is an index running over the terms in the decomposition of $\mathbf{248}$ above, and i runs over the dimension of each representation R_H^I . The u_I^a are bundle-valued harmonic 1-forms on X , taking values in the associated representation R_G^I of the bundle structure group V . Finally, the T_{ia} are the relevant generators of the broken part of the original E_8 gauge group, that is, those broken generators that are not part of the bundle group G . The objects of interest in this paper are the trilinear couplings between the low energy matter fields C_I^i .

A simple expression for the superpotential Yukawa couplings has been well known for a some time

$G \times H$	Breaking Pattern: $\mathbf{248} \rightarrow$	Particle Spectrum
$SU(3) \times E_6$	$(\mathbf{1}, \mathbf{78}) \oplus (\mathbf{3}, \mathbf{27}) \oplus (\bar{\mathbf{3}}, \bar{\mathbf{27}}) \oplus (\mathbf{8}, \mathbf{1})$	$n_{27} = h^1(V)$ $n_{\bar{27}} = h^1(V^*) = h^2(V)$ $n_1 = h^1(V \otimes V^*)$
$SU(4) \times SO(10)$	$(\mathbf{1}, \mathbf{45}) \oplus (\mathbf{4}, \mathbf{16}) \oplus (\bar{\mathbf{4}}, \bar{\mathbf{16}}) \oplus (\mathbf{6}, \mathbf{10}) \oplus (\mathbf{15}, \mathbf{1})$	$n_{16} = h^1(V)$ $n_{\bar{16}} = h^1(V^*) = h^2(V)$ $n_{10} = h^1(\wedge^2 V)$ $n_1 = h^1(V \otimes V^*)$
$SU(5) \times SU(5)$	$(\mathbf{1}, \mathbf{24}) \oplus (\mathbf{5}, \mathbf{10}) \oplus (\bar{\mathbf{5}}, \bar{\mathbf{10}}) \oplus (\mathbf{10}, \bar{\mathbf{5}}) \oplus (\bar{\mathbf{10}}, \mathbf{5}) \oplus (\mathbf{24}, \mathbf{1})$	$n_{10} = h^1(V)$ $n_{\bar{10}} = h^1(V^*) = h^2(V)$ $n_5 = h^1(\wedge^2 V^*)$ $n_{\bar{5}} = h^1(\wedge^2 V)$ $n_1 = h^1(V \otimes V^*)$

Table 1: A vector bundle V with structure group G can break the E_8 gauge group of the heterotic string into a GUT group H . The low-energy representations are found from the branching of the $\mathbf{248}$ adjoint of E_8 under $G \times H$ and the low-energy spectrum is obtained by computing the indicated bundle cohomology groups.

[8]:

$$\lambda_{IJK} \propto \int_X u_I^a \wedge u_J^b \wedge u_K^c \wedge \bar{\Omega} f_{abc} . \quad (2.2)$$

We have used a “proportional to” sign here to emphasise the fact that, without knowledge of the Kähler potential, we can not meaningfully make statements about the overall normalization. In Eq. (2.2), the holomorphic $(3,0)$ form has been denoted by Ω and the f_{abc} are constants descending from the structure constants of E_8 , designed to make the above expression invariant under the bundle group G . This is the form for these couplings in the low energy theory as given to us by direct dimensional reduction. Naively, the evaluation of (2.2) is computationally awkward. On a given Calabi-Yau manifold, one would have to find explicit expressions for all of the forms involved and then integrate over the manifold. For $(2,1)$ matter fields in standard embedding models this has been explicitly carried out in references [2, 21]. To repeat such an explicit calculation for non-standard embedding models would be technically very challenging and we will instead pursue a different, more algebraic approach.

2.1 Rephrasing in terms of cohomologies

The formula (2.2) has many appealing properties [8]. In particular, it is quasi-topological¹. It depends only on the cohomology class of the 1-forms u_I^a and not upon the actual representative form within that

¹ Indeed, for standard embedding models, the Yukawa couplings for $(1,1)$ matter fields are topological and are given by the triple intersection numbers of the Calabi-Yau manifold.

chosen class. Indeed, taking $u_I^a \rightarrow u_I^a + D\epsilon_I^a$, for example, one sees that the change to (2.2),

$$\int_X D\epsilon_I^a \wedge u_J^b \wedge u_K^c \wedge \bar{\Omega} f_{abc} , \quad (2.3)$$

vanishes upon integration by parts since both the 1-forms u^b and the holomorphic 3-form $\bar{\Omega}$ are D closed. Given this observation, one can regard the matter fields as being represented in the formula (2.2) by cohomology classes, and not just their harmonic representatives. This suggests that a simple description of Yukawa couplings in terms of topological quantities exists.

To pursue this idea, we begin by rewriting the formula for the Yukawa couplings in the case where the bundle structure group G is $SU(3)$. We can then calculate four dimensional couplings between three **27** multiplets of E_6 . The relevant structure constants in this case are $f_{abc} = \epsilon_{abc}$ and, hence, the combination $u_I^a \wedge u_J^b \wedge u_K^c \epsilon_{abc}$ is an $SU(3)$ invariant harmonic 3-form. Up to an overall constant multiple there is, of course, only one such form on a Calabi-Yau 3-fold, namely the $(3, 0)$ form Ω . Thus we have that,

$$u_I^a \wedge u_J^b \wedge u_K^c \epsilon_{abc} = K_{IJK} \Omega , \quad (2.4)$$

where K_{IJK} are complex numbers. From

$$\lambda_{IJK} \propto \int_X u_I^a \wedge u_J^b \wedge u_K^c \wedge \bar{\Omega} \epsilon_{abc} = K_{IJK} \int_X \Omega \wedge \bar{\Omega} \quad (2.5)$$

we see these numbers are proportional to the desired Yukawa couplings.

Referring to Table 1 once more, we see that the families in the **27** representation of E_6 can be identified with the cohomology group $H^1(X, V)$. Therefore, equation (2.4) defines a map that takes three of our bundle-valued 1-forms to a harmonic 3-form valued in the trivial bundle. Now, for an $SU(n)$ bundle V we have that $\wedge^n V \cong \mathcal{O}_X$, where \mathcal{O}_X is the trivial line-bundle on X . Thus, (2.4) defines a map of the form

$$H^1(X, V) \times H^1(X, V) \times H^1(X, V) \rightarrow H^3(X, \wedge^3 V) \cong H^3(X, \mathcal{O}_X) \cong \mathbb{C} , \quad (2.6)$$

where the last equivalence follows from the fact that $h^3(X, \mathcal{O}_X) = 1$.

The main point of this paper is that, for a large class of compactifications, when the above cohomologies are represented by certain polynomial equivalence classes, there is a mathematically natural proposal for what the map implicit in Eq. (2.6) is. It is essentially the unique possibility and simply involves polynomial multiplication of cohomology representatives. In the next section, we present this proposal in detail and show that the results to which it gives rise have all of the properties one would expect. The rigorous proof that our method for calculating Yukawa couplings does indeed reproduce the physical formula (2.2) is somewhat technical and is thus presented in Appendix B.

A similar procedure can be applied to the case of structure group $G = SU(4)$ and a visible gauge group $SO(10)$. For such models, we are interested in Yukawa couplings of the type **10 16 16**, between two families in **16** representations and a Higgs multiplet in a **10** representation of $SO(10)$. Note that the

absence of anti-families in our models means there are no $\overline{\mathbf{16}}$ representations. From Table 1, it is clear that families are still identified with the cohomology group $H^1(X, V)$ while Higgs multiplets correspond to $H^1(X, \wedge^2 V)$. The analogue of Eq. (2.6) is then

$$H^1(X, V) \times H^1(X, V) \times H^1(X, \wedge^2 V) \rightarrow H^3(X, \wedge^4 V) \cong H^3(X, \mathcal{O}_X) \cong \mathbb{C} . \quad (2.7)$$

The appearance of the fourth wedge power, $\wedge^4 V$, means that one has to deal with polynomials of quite high degree in practical calculations. For this reason, it is useful to slightly reformulate the above mapping to

$$H^1(X, V) \times H^1(X, V) \rightarrow (H^1(X, \wedge^2 V))^* \cong H^2(X, \wedge^2 V) . \quad (2.8)$$

where the final equivalence follows from Serre duality [22], $H^p(X, W) \simeq H^{3-p}(X, W^*)^*$, and the fact that $\wedge^2 V \cong \wedge^2 V^*$ for $SU(4)$ bundles. Hence, instead of mapping two families and a Higgs multiplet into a one-dimensional space of high degree we combine two families to represent an element in the Higgs cohomology group. The relevant Yukawa couplings are then given by expressing the result in terms of a basis of Higgs multiplets. In this case, we only need to deal with second wedge powers of V which, as we will see, implies lower polynomial degrees.

Finally, the case where $G = SU(5)$ can be dealt with in either of the two ways we have discussed so far. It turns out to be computationally more efficient to follow the second approach. From Table 1 we have three relevant multiplets, namely $\mathbf{10}$ multiplets associated to $H^1(X, V)$, $\mathbf{5}$ multiplets associated to $H^1(X, \wedge^2 V^*)$ and $\overline{\mathbf{5}}$ multiplets associated to $H^1(X, \wedge^2 V)$ (and since we are considering models without anti-families there are no $\overline{\mathbf{10}}$ representations present). This gives rise to two types of Yukawa couplings that are schematically of the form $\mathbf{10} \mathbf{10} \mathbf{5}$ and $\overline{\mathbf{5}} \overline{\mathbf{5}} \mathbf{10}$. The corresponding maps in cohomology are

$$H^1(X, V) \times H^1(X, V) \rightarrow (H^1(X, \wedge^2 V^*))^* \cong H^2(X, \wedge^2 V) \quad (2.9)$$

$$H^1(X, \wedge^2 V) \times H^1(X, \wedge^2 V) \rightarrow (H^1(X, V))^* \cong H^2(X, \wedge^4 V) \quad (2.10)$$

We now need to discuss how the maps implied in (2.6), (2.8), (2.9) and (2.10) can actually be carried out explicitly. As we will see, within our class of models provided by CICY manifolds and monad bundles, the various cohomology groups can be represented by quotient spaces of polynomials and the maps amount to polynomial multiplication. To explore this in detail we now briefly describe the technical arena we will be working in - that of positive monad bundles over CICY manifolds - before we return to the problem of calculating Yukawa couplings in §3.

2.2 The arena: positive monad bundles over CICYs

In this paper, we will focus on heterotic compactifications involving vector bundles built via the monad construction [12]. In particular, we consider the class of positive monads² defined over favourable CICY

²For reviews of this construction and some of its applications, see references [10, 12, 17].

manifolds [11]. A systematic analysis of the stability and spectrum of this class has recently been completed in [10, 13, 14, 15].

To begin, we recall that complete intersection CICY manifolds are defined by the zero loci of K polynomials $\{p_j\}_{j=1,\dots,K}$ in an ambient space $\mathcal{A} = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_m}$ given by a product of m projective spaces with dimensions n_r . We denote the projective coordinates of each factor \mathbb{P}^{n_r} by $(x_0^{(r)}, x_1^{(r)}, \dots, x_{n_r}^{(r)})$, its Kähler form by J_r , and the k^{th} power of the hyperplane bundle by $\mathcal{O}_{\mathbb{P}^{n_r}}(k)$. The manifold X is called a *complete intersection* if the dimension of X equals the dimension of \mathcal{A} minus the number of polynomials. To obtain three-folds X in this way we then need $\sum_{r=1}^m n_r - K = 3$.

Each of the defining homogeneous polynomials p_j can be characterised by its multi-degree $\mathbf{q}_j = (q_j^1, \dots, q_j^m)$, where q_j^r specifies the degree of p_j in the coordinates $\mathbf{x}^{(r)}$ of the factor \mathbb{P}^{n_r} in \mathcal{A} . These polynomial degrees are conveniently encoded in a configuration matrix

$$\begin{bmatrix} \mathbb{P}^{n_1} & \left| \begin{array}{cccc} q_1^1 & q_2^1 & \dots & q_K^1 \\ q_1^2 & q_2^2 & \dots & q_K^2 \\ \vdots & \vdots & \ddots & \vdots \\ q_1^m & q_2^m & \dots & q_K^m \end{array} \right. \\ \mathbb{P}^{n_2} & \\ \vdots & \\ \mathbb{P}^{n_m} & \end{bmatrix}_{m \times K} . \quad (2.11)$$

Note that the j^{th} column of this matrix contains the multi-degree of the polynomial p_j . The Calabi-Yau condition, $c_1(TX) = 0$, is equivalent to the conditions $\sum_{j=1}^K q_j^r = n_r + 1$. In terms of this data, the normal bundle \mathcal{N} of the CICY manifold X in \mathcal{A} can be written as

$$\mathcal{N} = \bigoplus_{j=1}^K \mathcal{O}_{\mathcal{A}}(\mathbf{q}_j) . \quad (2.12)$$

Here and in the following we employ the short-hand notation $\mathcal{O}_{\mathcal{A}}(\mathbf{k}) = \mathcal{O}_{\mathbb{P}^{n_1}}(k^1) \otimes \dots \otimes \mathcal{O}_{\mathbb{P}^{n_r}}(k^r)$ for line bundles on the ambient space \mathcal{A} . In the notation given above, the famous quintic hypersurface in \mathbb{P}^4 is denoted as “[4|5]” and its normal bundle is $\mathcal{N} = \mathcal{O}_{\mathbb{P}^4}(5)$.

CICY threefolds have been completely classified [11] and of the 7890 manifolds, 4515 are favourable, that is, all of their Kähler forms, J , descend from those of the ambient projective space. This means that favourable CICY manifolds defined in an ambient space with m projective factors are characterized by $h^{1,1}(TX) = m$. We will focus on these favourable CICY manifolds in the following.

A monad bundle, V , is defined by the short exact sequence

$$0 \rightarrow V \rightarrow B \xrightarrow{f} C \rightarrow 0 , \quad \text{where} \\ B = \bigoplus_{i=1}^{r_B} \mathcal{O}_X(\mathbf{b}_i) , \quad C = \bigoplus_{j=1}^{r_C} \mathcal{O}_X(\mathbf{c}_j) \quad (2.13)$$

are sums of line bundles with ranks r_B and r_C , respectively³. From the exactness of (2.13), it follows

³More generally, a monad bundle is defined as the middle homology of a sequence of the form $0 \rightarrow A \xrightarrow{m_1} B \rightarrow C \rightarrow 0$. This sequence is exact at A and C , and $\text{Im}(m_1)$ is a subbundle of B [12]. In this paper we restrict ourselves, as is often done in the physics literature, to the case where $\text{Im}(m_1)$ vanishes. We thus recover the description (2.13).

that the bundle V is given by

$$V = \ker(f) . \quad (2.14)$$

From the above sequence, the rank, n , of V is

$$n = \text{rk}(V) = r_B - r_C . \quad (2.15)$$

For the structure group to be $SU(n)$ rather than $U(n)$ we need the first Chern class of V to vanish, hence

$$c_1^r(V) = \sum_{i=1}^{r_B} b_i^r - \sum_{a=1}^{r_C} c_a^r = 0 . \quad (2.16)$$

The existence of sufficiently general maps f is guaranteed by demanding that $c_j^r \geq b_i^s \forall i, j, r, s$. We can think of f as a matrix f_{ai} of polynomials with multi-degree $\mathbf{c}_a - \mathbf{b}_i$. Furthermore, from Eq. (2.14), the bundle moduli of V can be identified as the coefficients parameterizing the possible maps f (see [13] for a discussion). The term ‘‘positive’’ refers to monad bundles satisfying $b_i^r > 0$ and $c_j^r > 0 \forall r, i, j$.

For the technical details of monad bundles, including the spectrum, moduli and such properties as slope-stability, we refer the reader to [10, 13, 14, 15]. Here we will review one feature of positive monad bundles that will be of use to us in the following sections:

Positive monads do not give rise to anti-generations, that is, $H^2(X, V) = H^1(X, V^) = 0$.*

To see this, we consider dual of the monad sequence (2.13)

$$0 \rightarrow C^* \rightarrow B^* \rightarrow V^* \rightarrow 0 , \quad (2.17)$$

which gives rise to a long exact sequence

$$\dots \rightarrow H^1(X, B^*) \rightarrow H^1(X, V^*) \rightarrow H^2(X, C^*) \rightarrow \dots . \quad (2.18)$$

Now, since B and C are sums of positive line bundles both $H^1(X, B^*)$ and $H^2(X, C^*)$ are zero from Kodaira’s vanishing theorem (see, for example, references [21, 22]) so that $H^1(X, V^*) = 0$ follows immediately. Hence, there are no anti-families.

With these preliminary definitions in hand we turn now to the calculation of Yukawa couplings.

3 Calculating Yukawa couplings: general procedure

We shall consider in turn the three types of theories with E_6 , $SO(10)$ and $SU(5)$ low-energy groups, corresponding respectively to the choices of an $SU(n)$ bundle structure group with $n = 3, 4, 5$. A concrete $SU(3)$ example will be presented in this section but, in the interests of brevity, we postpone doing the same for the more complicated $SO(10)$ case until the next section. We do not give a detailed $SU(5)$ example in this paper because the techniques are lengthy, while qualitatively the same as in the $SU(3)$ and $SU(4)$ cases.

Case	Cohomologies required to vanish
E_6	$H^1(X, B), H^3(X, \wedge^3 B), H^2(X, \wedge^3 B)$ $H^1(X, \wedge^2 B \otimes C), H^2(X, \wedge^2 B \otimes C), H^1(X, B \otimes S^2 C)$
$SO(10)$	$H^1(X, B), H^1(X, \wedge^2 B), H^2(X, \wedge^2 B), H^1(B \otimes C)$
$SU(5)$	$H^1(X, B), H^1(X, \wedge^2 B), H^1(X, \wedge^4 B)$ $H^2(X, \wedge^4 B), H^1(X, \wedge^3 B \otimes C)$ $H^2(X, \wedge^2 B), H^1(X, B \otimes C)$

Table 2: List of vanishing conditions on the sums of line bundles B and C , defining the monad bundle, required for our calculation. All conditions are automatically satisfied for positive monads, due to the Kodaira vanishing theorem.

While the idea of computing Yukawa couplings using polynomial methods is based on the sheaf-module correspondence and should be quite general and widely applicable, the specific realisation discussed in this paper relies on a number of vanishing properties which we summarise in Table 2. These conditions are all automatically satisfied for positive monad bundles V , that is, when the sums of line bundles B and C that enter the monad sequence (2.13) consist of positive line bundles only.

Given these conditions, we would like to derive polynomial representations for certain bundle cohomology groups and maps between them. It is useful to first discuss this problem for the main building blocks of the monad construction, line bundles.

3.1 Polynomial representation of line bundle cohomology

We begin with the simple case of a single projective space \mathbb{P}^n with projective coordinates $\mathbf{x} = (x_0, \dots, x_n)$ and an associated graded ring $R = \mathbb{C}[\mathbf{x}]$. It is well-known that the sections, $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(k))$ of the line bundle $\mathcal{O}_{\mathbb{P}^n}(k)$ can be identified with the degree k polynomials in R . We denote the degree k part of R by R_k and write $H^0(\mathbb{P}^n, \mathcal{O}_{\mathbb{P}^n}(k)) \cong R_k$.

The generalization to products of projective spaces, $\mathcal{A} = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_m}$, is straightforward. We denote the projective coordinates of the r^{th} projective space by $\mathbf{x}^{(r)}$ and the associated multi-graded ring by

$$R = \mathbb{C}[\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(m)}]. \quad (3.1)$$

Then the sections of the line bundle $\mathcal{O}_{\mathcal{A}}(\mathbf{k})$ can be identified with the multi-degree \mathbf{k} polynomials in R , so

$$H^0(\mathcal{A}, \mathcal{O}_{\mathcal{A}}(\mathbf{k})) \cong R_{\mathbf{k}}. \quad (3.2)$$

In our actual applications, we are of course interested in line bundles $\mathcal{O}_X(\mathbf{k})$ on the CICY manifold $X \subset \mathcal{A}$. They can be related to their ambient space cousins via a Koszul resolution and this leads to a method of calculating their cohomology and, in particular, their sections. The details of this argument are given in Appendix A but the final result is rather simple. Consider the polynomial ring (3.1), associated

to our ambient space \mathcal{A} , and the ideal $\langle p_1, \dots, p_K \rangle \subset R$ generated by the defining polynomials p_j of the CICY manifold X . Then we can form the coordinate ring

$$A = \frac{R}{\langle p_1, \dots, p_K \rangle} \quad (3.3)$$

of the CICY manifold X , which one can think of as the space of polynomials on X . In terms of the coordinate ring, the sections of the line bundle $\mathcal{O}_X(\mathbf{k})$ are given by

$$H^0(X, \mathcal{O}_X(\mathbf{k})) \cong A_{\mathbf{k}}, \quad (3.4)$$

where the $A_{\mathbf{k}}$ denotes the multi-degree \mathbf{k} part of A . This relation requires certain vanishing conditions, as detailed in Appendix A, which are all automatically satisfied for positive line bundles. The result (3.4) is in close analogy to its ambient space counterpart (3.2), so all that is required when dealing with line bundles on the CICY manifold X is passing from the full polynomial ring to the coordinate ring of X .

3.2 $SU(3)$ vector bundles and E_6 GUTS

We start by considering the case of $SU(3)$ bundles. From Table 1, the symmetry breaking pattern and decomposition of the matter field representations is

$$E_8 \supset SU(3) \times E_6 \quad (3.5)$$

$$\mathbf{248} = (\mathbf{8}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{78}) \oplus (\mathbf{3}, \mathbf{27}) \oplus (\bar{\mathbf{3}}, \bar{\mathbf{27}}) \quad (3.6)$$

The $(\mathbf{8}, \mathbf{1})$ term in this decomposition is associated with the cohomology group $H^1(X, V \otimes V^*)$ that counts the dimension of the bundle moduli space. Furthermore, when we are dealing with positive monads, as discussed above, anti-families in $\bar{\mathbf{27}}$, corresponding to $H^1(X, V^*)$, are absent. Hence, we are left with families in $\mathbf{27}$ multiplets, associated with the cohomology group $H^1(X, V)$. The only type of Yukawa coupling is, therefore, of the form $\mathbf{27} \mathbf{27} \mathbf{27}$ and it can be calculated from the map (2.6). To do this we require polynomial representatives for the two cohomology groups involved, namely for $H^1(X, V)$ and $H^3(X, \wedge^3 V)$.

3.2.1 Polynomial representatives for families in $H^1(X, V)$

Looking at the long exact sequence in cohomology associated to the short exact monad sequence (2.13), we find that

$$\begin{aligned} 0 &\rightarrow H^0(X, V) \rightarrow H^0(X, B) \xrightarrow{f} H^0(X, C) \\ &\rightarrow H^1(X, V) \rightarrow H^1(X, B) \rightarrow \dots \end{aligned} \quad (3.7)$$

For stable $SU(n)$ bundles we know that $H^0(X, V) = 0$. In addition, if we assume that $H^1(X, B) = 0$, a condition which is always satisfied for positive monads as a consequence of Kodaira vanishing, it follows

that

$$H^1(X, V) \cong \frac{H^0(X, C)}{f(H^0(X, B))}. \quad (3.8)$$

From Eq. (3.4), both cohomology groups on the RHS can be represented in terms of the coordinate ring A of X , so we finally have

$$H^1(X, V) \cong \frac{\bigoplus_{a=1}^{r_C} A_{\mathbf{c}_a}}{f\left(\bigoplus_{i=1}^{r_B} A_{\mathbf{b}_i}\right)}. \quad (3.9)$$

The map f in this quotient is induced from the monad map in (2.13). If we represent the monad map by a matrix f_{ai} of polynomials with multi-degrees $\mathbf{c}_a - \mathbf{b}_i$ then its action on a vector of polynomials $(q_i) \in \bigoplus_{i=1}^{r_B} A_{\mathbf{b}_i}$ is given by

$$f((q_i)) = \left(\sum_{i=1}^{r_B} f_{ai} q_i \right) \in \bigoplus_{a=1}^{r_C} A_{\mathbf{c}_a}. \quad (3.10)$$

This is the action of a polynomial matrix and it allows us to explicitly compute the polynomial quotient (3.9) once the monad map $f \sim (f_{ai})$ is specified. We note that the degrees of the various polynomials involved are given by the integer vectors \mathbf{b}_i and \mathbf{c}_a that define the monad bundle (2.13).

We have obtained explicit polynomial representatives for the families and now turn to the ‘‘Yukawa cohomology group’’ $H^3(X, \wedge^3 V)$.

3.2.2 Polynomial representatives for $H^3(X, \wedge^3 V)$

Taking the exterior power sequence associated to our monad, as described in appendix B of reference [13], and splitting it into short exact sequences we obtain

$$\begin{aligned} 0 &\rightarrow \wedge^3 V \rightarrow \wedge^3 B \rightarrow K_1 \rightarrow 0 \\ 0 &\rightarrow K_1 \rightarrow \wedge^2 B \otimes C \rightarrow K_2 \rightarrow 0 \\ 0 &\rightarrow K_2 \rightarrow B \otimes S^2 C \rightarrow S^3 C \rightarrow 0. \end{aligned} \quad (3.11)$$

Here we have introduced the (co)-kernels K_1 and K_2 .

The following pieces may be extracted from the associated long-exact sequences in cohomology.

$$\dots \rightarrow H^2(X, \wedge^3 B) \rightarrow H^2(X, K_1) \rightarrow H^3(X, \wedge^3 V) \rightarrow H^3(X, \wedge^3 B) \rightarrow 0 \quad (3.12)$$

$$\dots \rightarrow H^1(X, \wedge^2 B \otimes C) \rightarrow H^1(X, K_2) \rightarrow H^2(X, K_1) \rightarrow H^2(X, \wedge^2 B \otimes C) \rightarrow \dots \quad (3.13)$$

$$\dots \rightarrow H^0(X, B \otimes S^2 C) \rightarrow H^0(X, S^3 C) \rightarrow H^1(X, K_2) \rightarrow H^1(X, B \otimes S^2 C) \rightarrow \dots \quad (3.14)$$

We now assume that the following vanishing conditions

$$\begin{aligned} H^3(X, \wedge^3 B) &= 0, \quad H^2(X, \wedge^3 B) = 0 \\ H^1(X, \wedge^2 B \otimes C) &= 0, \quad H^2(X, \wedge^2 B \otimes C) = 0 \\ H^1(X, B \otimes S^2 C) &= 0, \end{aligned} \quad (3.15)$$

are satisfied. This is automatically the case for positive monad bundles as a consequence of the Kodaira vanishing theorem. Then one can combine the sequences (3.12), (3.13) and (3.14) to obtain,

$$\dots \rightarrow H^0(X, B \otimes S^2C) \xrightarrow{F} H^0(X, S^3C) \rightarrow H^3(X, \wedge^3V) \rightarrow 0 . \quad (3.16)$$

We therefore conclude that

$$H^3(X, \wedge^3V) \cong \frac{H^0(X, S^3C)}{F(H^0(X, B \otimes S^2C))} . \quad (3.17)$$

Expressing this in terms of the coordinate ring via Eq. (3.4) as before, leads to

$$H^3(X, \wedge^3V) \cong \frac{\bigoplus_{a \geq b \geq c} A_{\mathbf{c}_a + \mathbf{c}_b + \mathbf{c}_c}}{F\left(\bigoplus_{i, a \geq b} A_{\mathbf{b}_i + \mathbf{c}_a + \mathbf{c}_b}\right)} . \quad (3.18)$$

The map F is induced by the monad map $f \sim (f_{ai})$ and, acting on a tensor of polynomials $(q_{iab}) \in \bigoplus_{i, a \geq b} A_{\mathbf{b}_i + \mathbf{c}_a + \mathbf{c}_b}$, it can be written as

$$F((q_{iab})) = \left(\sum_{i=1}^{r_B} q_{i(abf_c)i} \right) \in \bigoplus_{a \geq b \geq c} A_{\mathbf{c}_a + \mathbf{c}_b + \mathbf{c}_c} , \quad (3.19)$$

where the brackets around the indices denote symmetrization. Since $h^3(X, \wedge^3V) = h^3(X, \mathcal{O}_X) = 1$ we know that this polynomial quotient must be one-dimensional, although this is by no means obvious from the RHS of Eq. (3.18). For the example below we will explicitly verify that this is indeed the case.

3.2.3 Computing Yukawa couplings

From Eq. (3.9) we know that families are represented by a vector of polynomials $(P_a)_{a=1, \dots, r_C}$ with multi-degrees \mathbf{c}_a , subject, of course, to the identifications implied by having to work in the coordinate ring of X and the quotient in Eq. (3.9). Let us pick a basis (P_a^I) , in family space, where $I, J, K, \dots = 1, \dots, h^1(X, V)$ are the family indices. We can then form all possible symmetrized products, $P_{(a}^I P_b^J P_c^K)$, of these polynomials which are of degree $\mathbf{c}_a + \mathbf{c}_b + \mathbf{c}_c$. For each choice, (I, J, K) , of three families, these products form a three-index symmetric tensor $(P_{(a}^I P_b^J P_c^K)$ which defines an element of $\bigoplus_{a \geq b \geq c} A_{\mathbf{c}_a + \mathbf{c}_b + \mathbf{c}_c}$ and, hence, from Eq. (3.18), an element of the Yukawa cohomology group $H^3(X, \wedge^3V)$. That the polynomial degrees match in this way is non-trivial and, of course, necessary for our method to work. We can now pick a representative, (Q_{abc}) , consisting of polynomials with multi-degree $\mathbf{c}_a + \mathbf{c}_b + \mathbf{c}_c$, whose class $[(Q_{abc})]$ spans the quotient (3.18). Since we are dealing with a one-dimensional quotient, the class, $[(P_{(a}^I P_b^J P_c^K)]$, defined by the product of three families, must be proportional to $[(Q_{abc})]$, so that we can write

$$\left[(P_{(a}^I P_b^J P_c^K) \right] = \lambda_{IJK} [(Q_{abc})] . \quad (3.20)$$

The complex numbers λ_{IJK} are of course the desired Yukawa couplings. Since the ‘‘comparison class’’ $[(Q_{abc})]$ was chosen arbitrarily this only defines the Yukawa couplings up to an overall normalization and, of course, relative to the chosen basis in family space, as expected.

3.2.4 A simple E_6 example

Let us illustrate this procedure by a simple example on the quintic in \mathbb{P}^4 . The coordinate ring of the quintic is given by

$$A = \frac{\mathbb{C}[x_0, \dots, x_4]}{\langle p \rangle}, \quad (3.21)$$

where (x_0, \dots, x_4) are projective coordinates on \mathbb{P}^4 and p is the defining quintic polynomial. We would like to consider the $SU(3)$ monad bundle defined by

$$0 \rightarrow V \rightarrow \mathcal{O}_X(1)^{\oplus 4} \xrightarrow{f} \mathcal{O}_X(4) \rightarrow 0 \quad (3.22)$$

which is perhaps the simplest positive monad on the quintic. Note that, in this case, the monad map f can be represented by a vector $f = (f_1, \dots, f_4)$ of four cubics in A . To make contact with the previous general notation, this means that the vectors \mathbf{b}_i and \mathbf{c}_a are, in fact, one-dimensional and explicitly given by $\mathbf{b}_1 = \mathbf{b}_3 = \mathbf{b}_3 = \mathbf{b}_4 = (1)$ and $\mathbf{c}_1 = (4)$.

From Eq. (3.9) it follows that the families are represented by the quotient

$$H^1(X, V) \cong \frac{A_4}{f(A_1^{\oplus 4})} \quad (3.23)$$

of quartic polynomials by the image of four linear polynomials. On a vector $(q_1, \dots, q_4) \in A_1^{\oplus 4}$ consisting of four linear polynomials, the map f acts as

$$f((q_1, \dots, q_4)) = \sum_{i=1}^4 f_i q_i. \quad (3.24)$$

It is easy to count the dimension of this quotient. In general, the number of degree k polynomials in $n+1$ variables is,

$$\dim(\mathbb{C}[x_0, \dots, x_n]_k) = \binom{n+k}{n}. \quad (3.25)$$

Hence, $\dim A_4 = 70$ and $\dim A_1^{\oplus 4} = 20$. (In general, one has to correct for the fact that one is working with the coordinate ring, rather than the ring of all polynomials. In the present case we are dividing by an ideal generated by a quintic polynomial so that degrees A_k , where $k < 5$ are not affected.) For sufficiently generic choices of polynomials f_i , the map f is injective and we conclude that the quotient (3.23) has dimension $70 - 20 = 50$. So we are dealing with a model with 50 families.

For the Yukawa cohomology group (3.18) we have in the present case

$$H^3(X, \wedge^3 V) \cong \frac{A_{12}}{F(A_9^{\oplus 4})}, \quad (3.26)$$

where F acts on a vector $(r_1, \dots, r_4) \in A_9^{\oplus 4}$ as

$$F((r_1, \dots, r_4)) = \sum_{i=1}^4 f_i r_i. \quad (3.27)$$

	$I = 1$	$I = 2$	$I = 3$		$i = 1$	$i = 2$	$i = 3$		$i = 1$	$i = 2$	$i = 3$
Y_{11I}	0	0	0	Y_{21I}	0	0	1	Y_{31I}	0	1	0
Y_{12I}	0	0	1	Y_{22I}	0	0	0	Y_{32I}	1	0	0
Y_{13I}	0	1	0	Y_{23I}	1	0	0	Y_{33I}	0	0	0

Table 3: The array of the $\mathbf{27}^3$ Yukawa couplings for the E_6 GUT associated to the $SU(3)$ monad given in (3.22) on the quintic. There are 50 families of $\mathbf{27}$ multiplets, represented by $H^1(X, V)$; we select three of these for illustrative purposes here, as indexed by $I = 1, 2, 3$. These are represented by the monomials x_4^4 , $x_2^2 x_3^2$ and $x_0^2 x_1^2$ respectively (that is, these are the normal forms of the equivalence class of polynomials representing these families). The normal form of the comparison class in this calculation was $x_0^2 x_1^2 x_2^2 x_3^2 x_4^4$. The monad map is given by $f = (x_0^3, x_1^3, x_2^3, x_3^3)$.

As the degrees involved exceed 5, counting polynomials to determine the dimension of this quotient is not so simple any more. However, it is relatively straightforward to extract this information from the relevant Hilbert series which can be computed with computer algebra packages such as Macaulay and Singular [23, 24]. It turns out that this dimension is indeed 1, as it must be from our general arguments. It should be noted that the computer algebra package Singular [24] is fast enough on a standard desktop machine to perform the calculation of the Yukawa couplings between all 50 families in a matter of minutes. A useful interface for Singular, designed for use by physicists, may be found here [25]. A sample of the result, for a given choice of family representatives and monad map, is given in Table 3.

3.3 $SU(4)$ vector bundles and $SO(10)$ GUTs

Having introduced our general method of computing Yukawa couplings for the case of $SU(3)$ bundles, let us move on to consider the case of $SU(4)$ bundles. From Table 1 we have the following symmetry breaking pattern and decomposition of the matter field representations

$$E_8 \supset SU(4) \times SO(10) \tag{3.28}$$

$$\mathbf{248} = (\mathbf{15}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{45}) \oplus (\mathbf{4}, \mathbf{16}) \oplus (\overline{\mathbf{4}}, \overline{\mathbf{16}}) \oplus (\mathbf{6}, \mathbf{10}) . \tag{3.29}$$

The $(\mathbf{15}, \mathbf{1})$ term corresponds to bundle moduli that are counted by the cohomology group $H^1(X, V \otimes V^*)$. As in the E_6 case anti-families in $(\overline{\mathbf{4}}, \overline{\mathbf{16}})$ multiplets are absent for positive monads. Therefore, the relevant Yukawa couplings are of the form $\mathbf{10} \mathbf{16} \mathbf{16}$ and couple two families in $\mathbf{16}$ multiplets, associated to the cohomology group $H^1(X, V)$, to a Higgs multiplet in $\mathbf{10}$, associated to the cohomology group $H^1(X, \wedge^2 V)$. The associated Yukawa coupling can be computed by considering the map (2.8), so we need polynomial representations for $H^1(X, V)$ and $H^2(X, \wedge^2 V)$.

Polynomial representatives for the families in $H^1(X, V)$ can be worked out in exactly the same way as for the E_6 case and the result is given by Eqs. (3.9) and (3.10).

3.3.1 Polynomial representatives for Higgs multiplets in $H^1(X, \wedge^2 V)$

For the **10** multiplets, corresponding to $H^1(X, \wedge^2 V) \simeq H^2(X, \wedge^2 V)$, we introduce an exterior power sequence associated to the defining sequence of the monad, (2.13). Splitting the sequence up using the (co)-kernel K_3 we obtain the following

$$\begin{aligned} 0 &\rightarrow \wedge^2 V \rightarrow \wedge^2 B \rightarrow K_3 \rightarrow 0 \\ 0 &\rightarrow K_3 \rightarrow B \otimes C \rightarrow S^2 C \rightarrow 0. \end{aligned} \quad (3.30)$$

These induce the following long exact sequences in cohomology

$$\dots \rightarrow H^1(X, \wedge^2 B) \rightarrow H^1(X, K_3) \rightarrow H^2(X, \wedge^2 V) \rightarrow H^2(X, \wedge^2 B) \rightarrow \dots \quad (3.31)$$

$$\dots \rightarrow H^0(X, B \otimes C) \xrightarrow{F} H^0(X, S^2 C) \rightarrow H^1(X, K_3) \rightarrow H^1(X, B \otimes C) \rightarrow \dots \quad (3.32)$$

Thus, if $H^1(X, \wedge^2 B) = H^2(X, \wedge^2 B) = 0$, which are two of our vanishing conditions in Table 2 satisfied for all positive monads, we have that $H^1(X, K_3) \cong H^2(X, \wedge^2 V)$. Together with the vanishing condition $H^1(X, B \otimes C) \cong 0$, again satisfied for all positive monads, this can be used in (3.32) to obtain

$$H^1(X, \wedge^2 V) \cong \frac{H^0(X, S^2 C)}{F(H^0(X, B \otimes C))}. \quad (3.33)$$

From Eq. (3.4) this translates to

$$H^1(X, \wedge^2 V) \cong \frac{\bigoplus_{a \geq b} A_{\mathbf{c}_a + \mathbf{c}_b}}{F\left(\bigoplus_{i,a} A_{\mathbf{b}_i + \mathbf{c}_a}\right)}. \quad (3.34)$$

The map F is induced by the monad map f and, acting on a tensor of polynomials $(q_{ia}) \in \bigoplus_{i,a} A_{\mathbf{b}_i + \mathbf{c}_a}$, it can be written as

$$F((q_{ia})) = \left(\sum_{i=1}^{rc} q_{i(a)fb} \right) \in \bigoplus_{a \geq b} A_{\mathbf{c}_a + \mathbf{c}_b}. \quad (3.35)$$

3.3.2 Computing Yukawa couplings

We would now like to compute Yukawa couplings by mapping in the way indicated in (2.8). We note that, from Eq. (3.9), a basis in family space takes the form (P_a^I) , where $I, J, K, \dots = 1, \dots, h^1(X, V)$ are family indices, and the polynomials are of multi-degree \mathbf{c}_a . A basis for the Higgs space (3.34) can be expressed in terms of multi-degree $\mathbf{c}_a + \mathbf{c}_b$ polynomials (H_{ab}^A) , where $A = 1, \dots, h^1(X, \wedge^2 V)$ numbers the Higgs multiplets and (ab) is a symmetrized index pair. Hence, the product of two polynomials representing families is precisely of the right multi-degree to be interpreted as an element of the Higgs polynomial space. We can, therefore, write

$$\left[(P_{(a)}^I P_b^J) \right] = \sum_A \lambda_{AIJ} \left[(H_{ab}^A) \right] \quad (3.36)$$

with λ_{AIJ} being the desired Yukawa couplings. An explicit example with just one Higgs multiplet will be discussed in the next section.

3.4 $SU(5)$ vector bundles and $SU(5)$ GUTs

The final case we shall consider is that of $SU(5)$ bundles. For this case we have the following symmetry breaking pattern and decomposition of the matter field representations (to avoid confusion we have marked the GUT $SU(5)$ group with a subscript GUT):

$$E_8 \supset SU(5) \times SU(5)_{\text{GUT}} \quad (3.37)$$

$$\mathbf{248} = (\mathbf{24}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{24}) \oplus (\mathbf{5}, \mathbf{10}) \oplus (\overline{\mathbf{5}}, \overline{\mathbf{10}}) \oplus (\overline{\mathbf{10}}, \mathbf{5}) \oplus (\mathbf{10}, \overline{\mathbf{5}}). \quad (3.38)$$

The absence of anti-generations for positive monad bundles implies that the $(\overline{\mathbf{5}}, \overline{\mathbf{10}})$ states in the decomposition above are not present in the low energy spectrum as $H^1(X, V^*) = 0$. The relevant Yukawa couplings are then of the two types $\mathbf{5} \mathbf{10} \mathbf{10}$ and $\mathbf{10} \overline{\mathbf{5}} \overline{\mathbf{5}}$ and they have to be computed from the maps (2.9) and (2.10). This means we must have polynomial representations for $H^1(X, V)$, $H^1(X, \wedge^2 V)$, $H^2(X, \wedge^4 V)$ and $H^2(X, \wedge^2 V)$.

We start, as in the other cases, by obtaining representatives for the cohomologies associated to the families residing in $\mathbf{10}$ multiplets. They correspond to the cohomology group $H^1(X, V)$ and can be dealt with in exactly the same way as the $\mathbf{16}$ multiplets in the $SO(10)$ case and the $\mathbf{27}$ multiplets for E_6 . Hence, their polynomial representatives are given by Eqs. (3.9) and (3.10).

3.4.1 Polynomial representatives for $H^1(X, \wedge^2 V)$

Polynomial representatives for the $\overline{\mathbf{5}}$ multiplets in $H^1(X, \wedge^2 V)$ may be obtained as for the $\mathbf{10}$ multiplets in the $SO(10)$ case, see Section 3.3.1. However, in the present case these particular representatives are not suitable for a calculation of Yukawa couplings following Eq. (2.10) since they do not square to the polynomial representatives for $H^2(X, \wedge^4 V)$, as determined below. We, therefore, have to follow a slightly more complicated approach. As usual, we use an exterior power sequence associated to the defining sequence of the monad. Splitting this sequence up, using the (co)-kernel K_4 , we obtain the two short exact sequences

$$0 \rightarrow \wedge^2 V \rightarrow \wedge^2 B \rightarrow K_4 \rightarrow 0 \quad (3.39)$$

$$0 \rightarrow K_4 \rightarrow B \otimes C \rightarrow S^2 C \rightarrow 0 .$$

The corresponding long exact sequences in cohomology contain the parts

$$\begin{aligned} \dots \rightarrow H^0(X, \wedge^2 B) \xrightarrow{f_1} H^0(X, K_4) \rightarrow H^1(X, \wedge^2 V) \rightarrow H^1(X, \wedge^2 B) \rightarrow \dots \\ 0 \rightarrow H^0(X, K_4) \rightarrow H^0(X, B \otimes C) \xrightarrow{f_2} H^0(X, S^2 C) \rightarrow \dots \end{aligned} \quad (3.40)$$

Given that $H^0(X, K_4) \cong \text{Ker}(f_2)$ it follows that $f_2 \circ f_1 = 0$ and with the polynomial representatives

$$H^0(X, \wedge^2 B) \cong \bigoplus_{i>j} A_{\mathbf{b}_i+\mathbf{b}_j} \quad (3.41)$$

$$H^0(X, B \otimes C) \cong \bigoplus_{i,a} A_{\mathbf{b}_i+\mathbf{c}_a} \quad (3.42)$$

$$H^2(X, S^2 C) \cong \bigoplus_{a \geq b} A_{\mathbf{c}_a+\mathbf{c}_b} \quad (3.43)$$

we have the complex

$$\bigoplus_{i>j} A_{\mathbf{b}_i+\mathbf{b}_j} \xrightarrow{f_1} \bigoplus_{i,a} A_{\mathbf{b}_i+\mathbf{c}_a} \xrightarrow{f_2} \bigoplus_{a \geq b} A_{\mathbf{c}_a+\mathbf{c}_b} . \quad (3.44)$$

On polynomial tensors (Q_{ij}) and (q_{ia}) the two maps above act as

$$f_1((Q_{ij})) = \left(\sum_j f_{ai} Q_{ij} \right) , \quad f_2((q_{ia})) = \left(\sum_i q_{i(a} f_{b)i} \right) , \quad (3.45)$$

which confirms explicitly that $f_2 \circ f_1 = 0$. The desired bundle cohomology $H^1(X, \wedge^2 V)$ is now given, if $H^1(X, \wedge^2 B) = 0$, by the cohomology of the above complex, that is,

$$H^1(X, \wedge^2 V) \simeq \frac{\text{Ker}(f_2)}{\text{Im}(f_1)} . \quad (3.46)$$

3.4.2 Polynomial representatives for $H^2(X, \wedge^4 V)$

Let us now obtain an appropriate polynomial description for the **10** multiplets in $H^2(X, \wedge^4 V)$ as required for calculating the **10** $\overline{\mathbf{5}} \overline{\mathbf{5}}$ Yukawa couplings from Eq. (2.10). Consider the exterior power sequence of the monad exact sequence, split by introducing (co)-kernels K_5, K_6 and K_7 .

$$0 \rightarrow \wedge^4 V \rightarrow \wedge^4 B \rightarrow K_5 \rightarrow 0 \quad (3.47)$$

$$0 \rightarrow K_5 \rightarrow \wedge^3 B \otimes C \rightarrow K_6 \rightarrow 0 \quad (3.48)$$

$$0 \rightarrow K_6 \rightarrow \wedge^2 B \otimes S^2 C \rightarrow K_7 \rightarrow 0 \quad (3.49)$$

$$0 \rightarrow K_7 \rightarrow B \otimes S^3 C \rightarrow S^4 C \rightarrow 0 \quad (3.50)$$

For our argument we require the following parts of the associated long exact sequences.

$$\dots \rightarrow H^1(X, \wedge^4 B) \rightarrow H^1(X, K_5) \rightarrow H^2(X, \wedge^4 V) \rightarrow H^2(X, \wedge^4 B) \rightarrow \dots \quad (3.51)$$

$$\dots \rightarrow H^0(X, \wedge^3 B \otimes C) \xrightarrow{f_3} H^0(X, K_6) \rightarrow H^1(X, K_5) \rightarrow H^1(X, \wedge^3 B \otimes C) \rightarrow \dots \quad (3.52)$$

$$0 \rightarrow H^0(X, K_6) \rightarrow H^0(X, \wedge^2 B \otimes S^2 C) \xrightarrow{f_4} H^0(X, K_7) \rightarrow \dots \quad (3.53)$$

$$0 \rightarrow H^0(X, K_7) \rightarrow H^0(X, B \otimes S^3 C) \rightarrow H^0(X, S^4 C) \rightarrow \dots \quad (3.54)$$

From our vanishing assumptions, which we remind the reader are automatically satisfied by the positive monads, $H^1(X, \wedge^4 B) = H^2(X, \wedge^4 B) = 0$ and, hence, the first of these sequences implies that

$H^2(X, \wedge^4 V) \cong H^1(X, K_5)$. The last two sequences tell us that $H^0(X, K_6)$ injects into $H^0(X, \wedge^2 B \otimes S^2 C)$ and $H^0(X, K_7)$ injects into $H^0(X, B \otimes S^3 C)$. Introducing the polynomial representatives

$$H^0(X, \wedge^3 B \otimes C) \cong \bigoplus_{i>j>k,a} A_{\mathbf{b}_i+\mathbf{b}_j+\mathbf{b}_k+\mathbf{c}_a} \quad (3.55)$$

$$H^0(X, \wedge^2 B \otimes S^2 C) \cong \bigoplus_{i>j,a \geq b} A_{\mathbf{b}_i+\mathbf{b}_j+\mathbf{c}_a+\mathbf{c}_b} \quad (3.56)$$

$$H^0(X, B \otimes S^3 C) \cong \bigoplus_{i,a \geq b \geq c} A_{\mathbf{b}_i+\mathbf{c}_a+\mathbf{c}_b+\mathbf{c}_c} , \quad (3.57)$$

we can therefore combine (3.52)–(3.54) to form the complex

$$\bigoplus_{i>j>k,a} A_{\mathbf{b}_i+\mathbf{b}_j+\mathbf{b}_k+\mathbf{c}_a} \xrightarrow{f_3} \bigoplus_{i>j,a \geq b} A_{\mathbf{b}_i+\mathbf{b}_j+\mathbf{c}_a+\mathbf{c}_b} \xrightarrow{f_4} \bigoplus_{i,a \geq b \geq c} A_{\mathbf{b}_i+\mathbf{c}_a+\mathbf{c}_b+\mathbf{c}_c} . \quad (3.58)$$

On polynomial tensors $(Q_{ijk a})$ and (q_{ijab}) the above maps f_3 and f_4 acts as

$$f_3((Q_{ijk a})) = \sum_k Q_{ijk}(a f_b)_k , \quad f_4((q_{ijab})) = \sum_j q_{ij}(ab f_c)_j . \quad (3.59)$$

As before, the desired cohomology $H^2(X, \wedge^4 V)$ is given, if $H^1(X, \wedge^3 B \otimes C) = 0$, by the cohomology of this complex, that is,

$$H^2(X, \wedge^4 V) \cong \frac{\text{Ker}(f_4)}{\text{Im}(f_3)} . \quad (3.60)$$

3.4.3 Polynomial representatives for $H^2(X, \wedge^2 V)$

Finally, we require polynomials to represent the **5** multiplets in $H^2(X, \wedge^2 V)$, to calculate the **5 10 10** Yukawa couplings from Eq. (2.9). We once again consider the long exact sequence in cohomology induced by (3.39). This contains the following pieces:

$$\dots \rightarrow H^1(X, \wedge^2 B) \rightarrow H^1(X, K_4) \rightarrow H^2(X, \wedge^2 V) \rightarrow H^2(X, \wedge^2 B) \rightarrow \dots \quad (3.61)$$

$$\dots \rightarrow H^0(X, B \otimes C) \xrightarrow{f_5} H^0(X, S^2 C) \rightarrow H^1(X, K_4) \rightarrow H^1(X, B \otimes C) \rightarrow \dots \quad (3.62)$$

Given the vanishing assumptions $H^1(X, \wedge^2 B) = H^2(X, \wedge^2 B) = H^1(X, B \otimes C) = 0$, which are automatically satisfied for positive monads, the first of these sequences implies that $H^2(X, \wedge^2 V) \cong H^1(X, K_4)$.

Using this in the second sequence leads to

$$H^2(X, \wedge^2 V) \cong \frac{H^0(X, S^2 C)}{f_5(H^0(X, B \otimes C))} . \quad (3.63)$$

Written in terms of polynomial representatives this means

$$H^2(X, \wedge^2 V) \cong \frac{\bigoplus_{a \geq b} A_{\mathbf{c}_a+\mathbf{c}_b}}{f_5 \left(\bigoplus_{i,a} A_{\mathbf{b}_i+\mathbf{c}_a} \right)} . \quad (3.64)$$

On polynomial tensors (q_{ia}) the map f_5 acts as

$$f_5((q_{ia})) = \left(\sum_i q_i(a f_b)_i \right) . \quad (3.65)$$

3.4.4 Computing Yukawa couplings

We begin by summarising the polynomial representations for the various multiplets. For the families in **10** multiplets we have a basis of polynomials (P_a^I) with multi-degrees \mathbf{c}_a , where $I, J, \dots = 1, \dots, h^1(X, V)$, as before. From Eq. (3.64), **5** multiplets are represented by multi-degree $\mathbf{c}_a + \mathbf{c}_b$ polynomials (H_{ab}^A) , where $A, B, \dots = 1, \dots, h^1(X, \wedge^3 V^*)$ and (ab) is a symmetric index pair. Eq. (3.46) shows that $\overline{\mathbf{5}}$ multiplets can be represented by polynomials $(\bar{H}_{i\bar{a}}^{\bar{A}})$ of multi-degree $\mathbf{b}_i + \mathbf{c}_a$, where $\bar{A}, \bar{B}, \dots = 1, \dots, h^1(X, \wedge^2 V)$. Finally, from Eq. (3.60) we have an alternative polynomial representation for the families in **10** by multi-degree $\mathbf{b}_i + \mathbf{b}_j + \mathbf{c}_a + \mathbf{c}_b$ polynomials (\tilde{P}_{ijab}^I) , where (ij) is an anti-symmetric and (ab) a symmetric index pair.

Given these polynomial representatives, the **5 10 10** Yukawa couplings λ_{AIJ} and the **10 $\overline{\mathbf{5}}$ $\overline{\mathbf{5}}$** Yukawa couplings $\lambda_{I\bar{A}\bar{B}}$ can be computed from

$$\left[(P_a^I P_b^J) \right] = \sum_A \lambda_{AIJ} \left[(H_{ab}^A) \right] \quad (3.66)$$

$$\left[(\bar{H}_{i\bar{a}}^{\bar{A}} \bar{H}_{j\bar{b}}^{\bar{B}}) \right] = \sum_I \lambda_{I\bar{A}\bar{B}} \left[(\tilde{P}_{ijab}^I) \right]. \quad (3.67)$$

This concludes our general discussion. We now move on to give a comprehensively worked example of some physical interest in the $SO(10)$ case.

4 An example: One Higgs multiplet and one heavy family

4.1 The model

As in our previous example, in section 3.2.4, we consider the quintic in \mathbb{P}^4 . The coordinate ring is given by

$$A = \frac{\mathbb{C}[x_0, \dots, x_4]}{\langle p \rangle}, \quad (4.1)$$

where (x_0, \dots, x_4) are the projective coordinates on \mathbb{P}^4 and p is the defining quintic polynomial. In this section, we will consider the following monad on the quintic.

$$0 \rightarrow V \rightarrow \mathcal{O}_X(1)^{\oplus 7} \xrightarrow{f} \mathcal{O}_X(2)^{\oplus 2} \oplus \mathcal{O}_X(3) \rightarrow 0. \quad (4.2)$$

This short exact sequence defines an $SU(4)$ bundle and thus we are discussing an $SO(10)$ GUT theory as in §3.3. The Yukawa couplings we shall calculate for this model are thus of the form **10 16 16**. The monad map f can be written as $f = (f_{1i}, f_{2i}, f_{3i})$, where $i = 1, \dots, 7$ runs over the seven $\mathcal{O}_X(1)$ line bundles and f_{1i}, f_{2i} are degree one polynomials in A while f_{3i} are degree two polynomials. From the general discussion in §3.3 it follows that the families in $H^1(X, V)$ can be represented by polynomials as

$$H^1(X, V) \cong \frac{A_2^{\oplus 2} \oplus A_3}{f(A_1^{\oplus 7})}. \quad (4.3)$$

Given that $\dim A_2 = 15$, $\dim A_3 = 35$ and $\dim A_1 = 5$, and choosing the map f sufficiently general so it is injective, it follows that this model has 30 families. From Eq. (3.34), the Higgs multiplets in $H^1(X, \wedge^2 V)$ can be represented by

$$H^1(X, \wedge^2 V) \cong \frac{A_4^{\oplus 3} \oplus A_5^{\oplus 2} \oplus A_6}{F(A_3^{\oplus 14} \oplus A_4^{\oplus 7})}, \quad (4.4)$$

where the map F has been defined in Eq. (3.35). If we write polynomials in the denominator of this quotient as $(q_{(3)i}, \tilde{q}_{(3)i}, q_{(4)i})^T$, where $i = 1, \dots, 7$ and the first index indicates the polynomial degree, and polynomials in the numerator as $(Q_{(4)1}, Q_{(4)2}, Q_{(5)1}, Q_{(4)3}, Q_{(6)}, Q_{(5)2})^T$, with the first index again indicating the polynomial degree, then this map can be explicitly written as

$$F \begin{pmatrix} q_{(3)i} \\ \tilde{q}_{(3)i} \\ q_{(4)i} \end{pmatrix} = \begin{pmatrix} Q_{(4)1} \\ Q_{(4)2} \\ Q_{(5)1} \\ Q_{(4)3} \\ Q_{(6)} \\ Q_{(5)2} \end{pmatrix} = \begin{pmatrix} f_{1i} & 0 & 0 \\ \frac{1}{2}f_{2i} & \frac{1}{2}f_{1i} & 0 \\ \frac{1}{2}f_{3i} & 0 & \frac{1}{2}f_{1i} \\ 0 & f_{2i} & 0 \\ 0 & 0 & f_{3i} \\ 0 & \frac{1}{2}f_{3i} & \frac{1}{2}f_{2i} \end{pmatrix} \begin{pmatrix} q_{(3)i} \\ \tilde{q}_{(3)i} \\ q_{(4)i} \end{pmatrix}. \quad (4.5)$$

We note that this is a 6×21 matrix of polynomials. We can use this explicit map to compute the dimension of the quotient (4.4). For generic choices of the monad map f it turns out that this dimension is zero, so there are no Higgs multiplets. This confirms the general result, found in references [10, 13], that $h^1(X, \wedge^2 V) = 0$ at a generic point in bundle moduli space. This generic case is of course of no interest in our context since Yukawa couplings of the form **10 16 16** are not present.

4.2 Engineering Higgs multiplets

To arrive at physically more interesting cases we have to understand how to engineer models with one (or possibly more than one) Higgs multiplet. This is typically not easy from a technical point of view and it was a particular challenge in the effort to find the exact MSSM spectrum from heterotic compactifications based on elliptically-fibered Calabi-Yau manifolds [26, 27]. In the present framework, it is at least straightforward to state what needs to be done in principle. We need to make special choices for the polynomials defining the monad map f in such a way that the induced map F in Eq. (4.5) leads to dimension-one quotient (4.4). At the same time, f still has to be sufficiently general so that V , as defined by the monad short exact sequence, is indeed a bundle rather than merely a sheaf.

To examine this in detail we can consider F as a map between modules $F : A(-3)^{\oplus 14} \oplus A(-4)^{\oplus 7} \longrightarrow A(-4)^{\oplus 3} \oplus A(-5)^{\oplus 2} \oplus A(-6)$ and then examine the Hilbert function of $\text{Coker}(F)$ at degree zero. As stated above, for generic choices of f , that is, at generic points in bundle moduli space, the Hilbert function at degree zero vanishes. Another way of stating the same fact is that the Hilbert functions of the ideals

$$\langle f_{1i} \rangle, \langle f_{2i}, f_{1i} \rangle, \langle f_{3i}, f_{1i} \rangle, \langle f_{2i} \rangle, \langle f_{3i} \rangle, \langle f_{3i}, f_{2i} \rangle. \quad (4.6)$$

of the Calabi-Yau's coordinate ring, that correspond to the images of the matrix rows in (4.5), are each individually zero at the appropriate degrees, that is at degrees (4, 4, 5, 4, 6). This suggests a simple way of engineering one Higgs multiplet. Rather than dealing with the full complication of the map (4.5) and its associated Hilbert function, we can focus on one row and produce a dimension one entry at the appropriate degree of the associated ideal, while keeping the dimensions zero for all other ideals. In particular, we can specialise the polynomials f_{1i} so that the ideal $\langle f_{1i} \rangle$ has dimension one at degree 4. Since all of the other ideals depend on polynomials other than f_{1i} , one finds, upon doing this, that the Hilbert function for the remaining ideals can be kept zero at the appropriate degrees. As a result, the dimension of the quotient (4.4) is one and we have engineered an example with one Higgs multiplet.

We still have to check that there exists a choice for f along the above lines that defines a bundle rather than just a sheaf. To do this we consider the explicit example

$$f_{1i} = (40x_3 + 94x_4, 117x_3 + 119x_4, 449x_3 + 464x_4 + 266x_0 + 195x_1 + 173x_2, \\ 306x_2, 273x_3, 259x_3 + 291x_4, 76x_3 + 98x_2), \quad (4.7)$$

with the remaining polynomials in f being left generic. This choice has been engineered in the way described above and it can be verified that it indeed leads to precisely one Higgs multiplet. In addition, one can check that the locus in \mathbb{P}^4 where the polynomial matrix f degenerates (that is, where its rank is not maximal) does not intersect a sufficiently general quintic and, hence, leads to a bundle on the quintic (although not to a bundle on \mathbb{P}^4).

4.3 The mass matrix

We now wish to calculate the Yukawa couplings in this class of examples with one Higgs multiplet. To do so we first pick 30 family representatives $P^I = (P_1^I, P_2^I, P_3^I) \in A_2^{\oplus 2} \oplus A_3$ whose associated classes form a basis of (4.3). Further we choose a Higgs representative $H = (H_1, \dots, H_6) \in A_4^{\oplus 3} \oplus A_5^{\oplus 2} \oplus A_6$ whose class spans the one-dimensional space (4.4). The Yukawa couplings then follow from Eq. (3.36) and form a symmetric matrix λ_{IJ} . Given that we do not know the matter field kinetic terms, the only physically significant property of this matrix is its rank. It turns out, with the map (4.7) this rank is precisely one.

The monad in (4.2) gives rise to one massive family in four dimensions at the point specified by (4.7) in its bundle moduli space.

In fact, this structure is somewhat more generic. Let us consider, more generally, an $SO(10)$ model with a basis $\{P_A\}$ of $\mathcal{F} \equiv \bigoplus_a A_{c_a}$, such that $\{P_I\} \subset \{P_A\}$ is a set of family representatives and a single Higgs multiplet represented by $H \in \mathcal{H} \equiv \bigoplus_{a \geq b} A_{c_a + c_b}$. Note that $\mathcal{H} = S^2 \mathcal{F}$, so the symmetric tensor products $\{P_A \otimes_S P_B\}$ form a basis of \mathcal{H} and we can introduce a hermitian scalar product, $\langle \cdot, \cdot \rangle$, on \mathcal{H} such that this basis is orthonormal. From Eq. (3.36) it then follows that the Yukawa matrix is given by the scalar product $\lambda_{IJ} \sim \langle P_I \otimes_S P_J, H \rangle$. The Higgs representative H can, of course, always be written as a linear

combination $H = \sum_{A,B} H_{AB} P_A \otimes_S P_B$. Inserting this into the above scalar product expression for the Yukawa couplings one finds that

$$\lambda_{IJ} \sim H_{IJ} . \quad (4.8)$$

This means, in a case where the Higgs representative can be expressed in terms of the family representatives, so that $H = \sum_{I,J} H_{IJ} P_I \otimes_S P_J$, the Yukawa matrix and the matrix representing the Higgs are proportional. In particular, their rank has to be the same. The method of engineering models with one Higgs multiplet described above typically leads to a Higgs representative which can be written as the square of a vector v_I , that is $H = \sum_{I,J} v_I v_J P_I \otimes_S P_J$ ⁴. This can be seen as follows.

Let us choose a Higgs representative by taking a so called “normal form” of a sufficiently generic linear combination of terms, $\sum_{A,B} c_{A,B} P_A \otimes_S P_B$ where $c_{A,B}$ are some randomly generated coefficients. We take this normal form by performing the Buchberger Algorithm [28, 29] on the linear combination relative to the module generated by the map polynomials defining the one dimensional class (4.4). Given our method of engineering a single Higgs, as discussed in the previous sub-section, the resulting Higgs representative will be of the form $(Q_{(4)1}, 0, 0, 0, 0)^T$. An inspection of the Buchberger algorithm [28, 29] reveals that $Q_{(4)1}$ in this expression will be a single monomial. It is in fact the “lagging monomial” of degree four that is not in the ideal $\langle f_{1i} \rangle$. That is, it is the degree four monomial that is lowest according to the monomial ordering used in the Buchberger algorithm, which does not appear as an element of $\langle f_{1i} \rangle \subset A$. If this lagging monomial is a square, then clearly our Higgs representative is the square of a family representative. This is always the case for the type of example considered in Sections 4.1 and 4.2. The f_{1i} are all linear polynomials for the monad given in (4.2). Given this, the lagging monomial is some variable to the fourth power and $H = c (x_i^2, 0, 0)^T \otimes_S (x_i^2, 0, 0)^T$ for some i and some constant c .

As a result, the matrix associated to the Higgs representative and, hence, the Yukawa matrix has rank one. We see that there is a close relation between our method for engineering one-Higgs models and obtaining precisely one heavy family. In conclusion we can state the following.

A model in which one Higgs is engineered in the manner described in §4.2 will generically have one heavy family.

5 Conclusions

In this paper we have introduced a simple algorithm for calculating Yukawa couplings in a wide class of heterotic models. The compactifications we have considered are on smooth Calabi-Yau spaces and are not restricted to the standard embedding. The methods can be used to calculate Yukawa couplings for a large class of bundles on complete-intersection Calabi-Yau manifolds. Such a systematic procedure for non-standard embedding models has not been presented in the literature before and, we believe, constitutes substantial advance.

⁴We would like to thank Tony Pantev for very helpful comments on this point.

The key to our methodology is to obtain polynomial representatives for family and other relevant multiplets whose degrees are compatible with one another. In practice, this requires finding polynomial representatives for various cohomology groups whose degrees are such that our procedure of polynomial multiplication and reduction to normal form may be carried out. Because of the simple, algebraic, nature of the resulting algorithm, the calculations can be carried out on a computer and we have done this in the text for a number of concrete examples.

We should stress again that we have calculated the *superpotential* contributions to the Yukawa couplings. The Kähler potential for the matter fields remains an unknown quantity in these non-standard embedding models. Nevertheless we have shown how some physically relevant information can be extracted from our results by focusing on quantities, such as the rank in the case of a Yukawa matrix, which are unaffected by the choice of basis in family space.

The final example, presented in section 4, demonstrates the power of these methods. This is an example of a smooth Calabi-Yau compactification leading to an $SO(10)$ GUT. We have shown how one may isolate loci in bundle moduli space where the model has precisely one Higgs multiplet, residing in the $\mathbf{10}$ representation. Our approach based on polynomial representatives makes this conceptually rather simple and merely requires making specific choices for the polynomials defining the bundle. In practice, it is not always straightforward to find these but we have described a simple method to engineer viable cases.

We have then shown that the structure of this one-Higgs model leads to a Yukawa matrix of rank one, and so to precisely one massive family. Moreover, the relation between our method of engineering one Higgs multiplet and obtaining one massive family seems to be more general, an observation which might be important for building heterotic models with a phenomenologically viable pattern of fermion masses.

While our method of computing Yukawa couplings by multiplying polynomial representatives has been presented in the context of a particular class of models, the underlying mathematical structure – the sheaf-module correspondence – is quite general and we expect related methods to work for other Calabi-Yau and bundle constructions. This work should be of considerable utility in checking conclusions about the vanishing of Yukawa couplings resulting from the research presented in references [30, 31]. Eventually, one would like to calculate Yukawa couplings in the context of more realistic models, where the GUT symmetry is broken due to Wilson lines. We expect that the methods described in this paper can be readily applied to such models, basically by projecting onto the various equivariant sub-spaces of the cohomology groups involved.

Acknowledgements

We gratefully acknowledge enlightening discussions with Philip Candelas and Tony Pantev. D. G. is partially supported by NSF grants DMS 08-10948 and DMS 03-11378, L. A. is supported by the DOE under contract No. DE-AC02-76-ER-03071, J. G. , by STFC, UK, Y.-H. H. , by an Advanced Fellowship

from the STFC, UK, as well as the FitzJames Fellowship of Merton College, Oxford, and A. L. , by the EC 6th Framework Programme MRTN-CT-2004-503369. L. A. , J. G. and A. L. would like to thank the organisers of the 2008 Vienna ESI workshop “Mathematical Challenges in String Phenomenology” where part of this work was completed. D. G. would like to thank Y.-H. H. for hospitality at the University of Oxford and Merton College, Oxford.

Appendices

A Koszul Complex and Polynomial Representatives

In this appendix we justify the relation (3.4) between sections of line bundles on a CICY manifold and its coordinate ring. First let us recall the general set-up and the notation. We work in an ambient space $\mathcal{A} = \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_m}$ with projective coordinates $(\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(m)})$. Line bundles on \mathcal{A} are denoted by $\mathcal{O}_{\mathcal{A}}(\mathbf{k}) = \mathcal{O}_{\mathbb{P}^{n_1}}(k_1) \otimes \dots \otimes \mathcal{O}_{\mathbb{P}^{n_m}}(k_m)$, where $\mathbf{k} = (k_1, \dots, k_m)$. The associated ring

$$R = \mathbb{C}[\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(m)}] \tag{A.1}$$

is multi-graded by an m -dimensional grade vector $\mathbf{k} = (k_1, \dots, k_m)$ where k_r specifies the degree in the projective coordinates $\mathbf{x}^{(r)}$ of \mathbb{P}^{n_r} . The multi-degree \mathbf{k} part of R is denoted by $R_{\mathbf{k}}$. Sections $H^0(X, \mathcal{O}_{\mathcal{A}}(\mathbf{k}))$ of the line bundle $\mathcal{O}_{\mathcal{A}}(\mathbf{k})$ can be represented by polynomials of multi-degree \mathbf{k} in R , so we write

$$H^0(X, \mathcal{O}_{\mathcal{A}}(\mathbf{k})) \cong R_{\mathbf{k}}. \tag{A.2}$$

A co-dimension K CICY manifold $X \subset \mathcal{A}$ is defined as the zero locus of homogeneous polynomials p_1, \dots, p_K and we denote the normal bundle of X in \mathcal{A} by \mathcal{N} . We define line bundles on X by restricting ambient space line bundles, that is $\mathcal{O}_X(\mathbf{k}) \equiv \mathcal{O}_{\mathcal{A}}(\mathbf{k})|_X$. Moreover, we assume that the CICY manifold is “favourable”, that is, all line bundles on X are obtained in this way. The coordinate ring of X is given by

$$A = \frac{R}{\langle p_1, \dots, p_K \rangle}, \tag{A.3}$$

and it inherits the multi-grading from R . We denote by $A_{\mathbf{k}}$ the multi-degree \mathbf{k} part of A .

For the purpose of this appendix, we focus on line bundles $\mathcal{L} = \mathcal{O}_{\mathcal{A}}(\mathbf{k})$ and their counterparts $L = \mathcal{O}_X(\mathbf{k})$ on X which satisfy the vanishing conditions ⁵

$$H^q(\wedge^{\kappa} \mathcal{N}^* \otimes \mathcal{L}) = 0 \tag{A.4}$$

for $q > 0$ and $\kappa = 0, \dots, K$. We note that, as a consequence of Kodaira’s vanishing theorem applied to line bundles \mathcal{L} on the ambient space \mathcal{A} , all positive line bundles fall into this class. Provided the above

⁵This can be slightly weakened without changing the result of this appendix. In fact we require the vanishing conditions stated in Table 4.

introducing co-kernels C_1, \dots, C_{K-1} . Here, we have $\kappa = 1, \dots, K-2$ in the middle sequence. From our vanishing condition, the first of these sequences implies that $H^q(\mathcal{A}, C_{K-1}) = 0$ for all $q > 0$. Further, from the long exact sequence associated to the middle sequence above and the vanishing conditions it follows that $H^{q-1}(\mathcal{A}, C_\kappa) \cong H^q(\mathcal{A}, C_{\kappa+1})$ for $\kappa = 1, \dots, K-2$ and $q = 2, \dots, K+3$. Together, this means that $H^1(\mathcal{A}, C_\kappa) = 0$ for $\kappa = 1, \dots, K-1$ and, hence, the long exact sequences associated to (A.10) all break after three terms. This leads to the recursion relations

$$H^0(X, L) \cong \frac{H^0(\mathcal{A}, \mathcal{L})}{H^0(\mathcal{A}, C_1)} \quad (\text{A.11})$$

$$H^0(\mathcal{A}, C_\kappa) \cong \frac{H^0(\wedge^\kappa \mathcal{N}^* \otimes \mathcal{L})}{H^0(\mathcal{A}, C_{\kappa+1})} \quad (\text{A.12})$$

$$H^0(\mathcal{A}, C_{K-1}) \cong \frac{H^0(\mathcal{A}, \wedge^K \mathcal{N}^* \otimes \mathcal{L})}{H^0(\mathcal{A}, \wedge^{K-1} \mathcal{N}^* \otimes \mathcal{L})}, \quad (\text{A.13})$$

where $\kappa = 1, \dots, K-2$, which allow one to express $H^0(X, L)$ as a ‘‘chain of quotients’’. However, since

$$0 \rightarrow H^0(\mathcal{A}, \wedge^K \mathcal{N}^* \otimes \mathcal{L}) \rightarrow H^0(\mathcal{A}, \wedge^{K-1} \mathcal{N}^* \otimes \mathcal{L}) \rightarrow \dots \rightarrow H^0(\mathcal{A}, \mathcal{N}^* \otimes \mathcal{L}) \rightarrow H^0(\mathcal{A}, \mathcal{L}) \rightarrow H^0(X, L) \rightarrow 0 \quad (\text{A.14})$$

is a complex it is sufficient to keep the first quotient in this chain. Hence, we have

$$H^0(X, L) \cong \frac{H^0(\mathcal{A}, \mathcal{L})}{p(H^0(\mathcal{A}, \mathcal{N}^* \otimes \mathcal{L}))} \quad (\text{A.15})$$

where p is the map induced by the defining polynomials p_1, \dots, p_K of the CICY manifold. Using the polynomial representatives (A.2) for sections of line bundles in the ambient space this implies the desired Eq. (A.5).

B Proof of Equivalence of Formulations

In this Appendix, as promised in the text, we give a formal mathematical proof of why calculating the Yukawa couplings using (2.2) is equivalent to the maps in cohomology as given in Section 2.1.

B.1 Chain complexes and bicomplexes

The standard way to convert a bicomplex⁶ $C = C_{p,q}$ (with horizontal differential $d' : C_{p,q} \rightarrow C_{p-1,q}$ and vertical differential $d'' : C_{p,q} \rightarrow C_{p,q-1}$ that commute) to a chain complex is to define the *total complex* $\text{Tot } C$ by setting $(\text{Tot } C)_n = \bigoplus_{p+q=n} C_{p,q}$ and setting the differential $d : \text{Tot } C \rightarrow \text{Tot } C$ of degree -1 to be $x \mapsto d'(x) + (-1)^p d''(x)$ for $x \in C_{p,q}$. This is in accordance with the principle of signs [33] if C is the tensor product of two chain complexes and we identify the symbols d , d' , and d'' .

⁶Warning: the standard definition of *double complex* in [35, p. 60] (see also [34, p. 174, Exer. 11] and [39, p. 8]) uses a sign convention different from the one we use here, namely $d'd'' = -d''d'$.

Given $m \in \mathbb{Z}$ and a chain complex B define the shifted chain complex $B[m]$ by setting $B[m]_p = B_{m+p}$; use the same differential, with no change in sign⁷. Similarly, given $m, n \in \mathbb{Z}$ and a bicomplex C define the shifted bicomplex $C[m, n]$ by setting $C[m, n]_{p,q} = C_{m+p, n+q}$.

The formula for the differential in $\text{Tot } C$ involves p but not q , so there is a simple isomorphism $\text{Tot}(C[0, n]) \cong (\text{Tot } C)[n]$ involving just direct sums of identity maps, with no minus signs involved. Thus, if we think of a bicomplex C as being assembled from its rows $C_{\cdot, q}$ for $q \in \mathbb{Z}$, reindexing the rows results in shifting the total complex. If the bicomplex is zero outside the range $0 \leq q \leq N$ we will use the pictorial notation $C = [C_{\cdot, 0} \leftarrow C_{\cdot, 1} \leftarrow \dots \leftarrow C_{\cdot, N}]$ to indicate its assembly from its rows. No minus signs are to be used when assembling a bicomplex from chain complexes and maps between them in this way.

In general, an isomorphism of chain complexes $\gamma : \text{Tot}(C[m, n]) \xrightarrow{\cong} (\text{Tot}(C))[m + n]$ can be defined as $(-1)^{mq}$ times the identity map on the component $(C[m, n])_{p,q} = C_{p+m, q+n}$. We omit the computation that γ is a chain map. A careful eye can discern something of degree m moving past something of degree q , in accordance with the principle of signs [33].

Given a map $f : B \rightarrow C$ of chain complexes we define the mapping cone⁸ by setting $\text{Cone } f = \text{Cone}(C \leftarrow B) = \text{Tot}[C \leftarrow B]$. There are isomorphisms $\text{Cone}(C \leftarrow 0) \cong C$ and $\text{Cone}(0 \leftarrow B) \cong B[-1]$, and the exact sequence $0 \rightarrow [C \leftarrow 0] \rightarrow [C \leftarrow B] \rightarrow [0 \leftarrow B] \rightarrow 0$ of bicomplexes leads to an exact sequence $0 \rightarrow C \rightarrow \text{Cone } f \rightarrow B[-1] \rightarrow 0$ of chain complexes. Given $c \in C_p$ and $b \in B_{p-1}$ the element $(c, b) \in (\text{Cone } f)_p$ satisfies $d(c, b) = (dc + (-1)^{p-1}fb, db)$.

A fundamental lemma in homological algebra states that a map $C \rightarrow D$ of first quadrant bicomplexes that is a quasi-isomorphism in each row induces a quasi-isomorphism on total complexes. The same statement applies to a map of third quadrant bicomplexes, or when rows are replaced by columns. A slightly stronger version, for filtered complexes, is proved in [32, Lemma 3.2]. This result is presented as the *acyclic assembly lemma* in [39].

Given a short exact sequence $E : 0 \rightarrow A \xrightarrow{f} B \xrightarrow{g} C \rightarrow 0$ of chain complexes, the corresponding map $[0 \leftarrow A] \rightarrow [C \leftarrow B]$ of bicomplexes is a quasi-isomorphism in each column, hence, according to the lemma, $A[-1] \rightarrow \text{Cone}(C \leftarrow B)$ is a quasi-isomorphism. Its inverse in the derived category composed with the map $C \rightarrow \text{Cone}(C \leftarrow B)$ gives a map $\rho = \rho_E : C \rightarrow A[-1]$ in the derived category. We would like to compare the induced map $\rho : H_p C \rightarrow H_{p-1} A$ with the connecting homomorphism $\partial = \partial_E : H_p C \rightarrow H_{p-1} A$ that appears in the long exact homology sequence. Given cycles $c \in C_p$ and $a \in A_{p-1}$, $\partial[c] = [a]$ means that there is an element $b \in B_p$ such that $gb = c$ and $fa = db$. The element $(0, b) \in \text{Cone}(C \leftarrow B)_{p+1}$ satisfies $d(0, b) = ((-1)^p gb, db) = ((-1)^p c, fa)$, so $((-1)^{p-1}c, 0)$ and $(0, fa)$ are homologous elements of $\text{Cone}(C \leftarrow B)_p$, which tells us that $\rho[c] = (-1)^{p-1}[a]$, and thus $\rho = (-1)^{p-1}\partial$ on

⁷Warning: this sign convention differs from the standard one implied by [35, p. 72, Exercise 1] and explicitly presented in [39, p. 9]. There the differential on $B[m]$ is equal to $(-1)^m$ times the differential on B . Better notation for that concept, compatible with the principle of signs [33], would be $[m]B$. An isomorphism $[m]B \xrightarrow{\cong} B[m]$, also compatible with the principle of signs, can be defined by $x \mapsto (-1)^{mp}x$ for $x \in B_p$.

⁸Our definition of the mapping cone is not the usual one, see [39, p. 18, 20].

$H_p C$.

Now we consider longer extensions in the sense of Yoneda. Suppose we have an exact sequence $E : 0 \rightarrow A \rightarrow B_n \rightarrow \cdots \rightarrow B_2 \rightarrow B_1 \rightarrow C \rightarrow 0$ of chain complexes. The commutative diagram

$$\begin{array}{cccccccccccc} \cdots & \longleftarrow & 0 & \longleftarrow & 0 & \longleftarrow & 0 & \longleftarrow & \cdots & \longleftarrow & 0 & \longleftarrow & A & \longleftarrow & 0 & \longleftarrow & \cdots \\ & & \downarrow \\ \cdots & \longleftarrow & 0 & \longleftarrow & C & \longleftarrow & B_1 & \longleftarrow & \cdots & \longleftarrow & B_{n-1} & \longleftarrow & B_n & \longleftarrow & 0 & \longleftarrow & \cdots \end{array}$$

of chain complexes can be regarded as a map of chain complexes of chain complexes. The corresponding map $[0 \leftarrow \cdots \leftarrow A] \rightarrow [C \leftarrow B_1 \leftarrow \cdots \leftarrow B_n]$ of bicomplexes is a quasi-isomorphism in each column (by exactness of E), hence the map $A[-n] \rightarrow \text{Tot}[C \leftarrow B_1 \leftarrow \cdots \leftarrow B_n]$ is a quasi-isomorphism. Its inverse composed with the map $C \rightarrow \text{Tot}[C \leftarrow B_1 \leftarrow \cdots \leftarrow B_n]$ gives a map $\rho = \rho_E : C \rightarrow A[-n]$ in the derived category.

Suppose we have another exact sequence $F : 0 \rightarrow C \rightarrow P_m \rightarrow \cdots \rightarrow P_2 \rightarrow P_1 \rightarrow Q \rightarrow 0$ of chain complexes, and consider the associated map $\rho_F : Q \rightarrow C[-m]$. Let $E * F : 0 \rightarrow A \rightarrow B_n \rightarrow \cdots \rightarrow B_2 \rightarrow B_1 \rightarrow P_m \rightarrow \cdots \rightarrow P_2 \rightarrow P_1 \rightarrow Q \rightarrow 0$ be the exact sequence obtained by splicing E to F along C ; the differential in the middle is the composite map $B_1 \rightarrow C \rightarrow P_m$. The following commutative diagram of bicomplexes, in which quasi-isomorphisms are indicated by \sim , shows that $\rho_{E * F} = \rho_E[-m] \circ \rho_F$; the simplicity of this formula and the absence of signs in it is a consequence of our choices above.

$$\begin{array}{ccc} & & A[-m-n] \\ & & \downarrow \sim \\ C[-m] & \longrightarrow & [C \leftarrow B_1 \leftarrow \cdots \leftarrow B_n][-m] \\ \downarrow \sim & & \downarrow \sim \\ Q \longrightarrow [Q \leftarrow P_1 \leftarrow \cdots \leftarrow P_m] & \longrightarrow & [Q \leftarrow P_1 \leftarrow \cdots \leftarrow P_m \leftarrow B_1 \leftarrow \cdots \leftarrow B_n] \end{array}$$

Now let's decompose our original sequence E by writing it as $E = E_1 * \cdots * E_n$, where $E_i : 0 \rightarrow D_i \rightarrow B_i \rightarrow D_{i-1} \rightarrow 0$, and $D_n = A$, $D_0 = C$, and $D_i = \text{im}(B_{i+1} \rightarrow B_i)$ for $0 < i < n$. Then $\rho_E = \rho_{E_1}[-(n-1)] \circ \cdots \circ \rho_{E_{n-1}}[-1] \circ \rho_{E_n}$. The resulting map $H_p(\rho_E) : H_p C \rightarrow H_{p-n} A$ is thus a composite of connecting homomorphisms, up to sign. More precisely, $H_p(\rho_E) = H_p(\rho_{E_1}[-(n-1)]) \circ \cdots \circ \rho_{E_{n-1}}[-1] \circ \rho_{E_n} = H_p(\rho_{E_1}[-(n-1)]) \circ \cdots \circ H_p(\rho_{E_{n-1}}[-1]) \circ H_p(\rho_{E_n}) = H_{p-n+1}(\rho_{E_1}) \circ \cdots \circ H_{p-1}(\rho_{E_{n-1}}) \circ H_p(\rho_{E_n}) = ((-1)_{p-n} \partial_{E_1}) \circ \cdots \circ ((-1)_{p-2} \partial_{E_{n-1}}) \circ ((-1)^{p-1} \partial_{E_n}) = (-1)^{(p-n)+\cdots+(p-2)+(p-1)} \partial_{E_1} \circ \cdots \circ \partial_{E_{n-1}} \circ \partial_{E_n} = (-1)^{pn+n(n+1)/2} \partial_{E_1} \circ \cdots \circ \partial_{E_{n-1}} \circ \partial_{E_n}$. (This result was proved in [35, Chap. V, Prop. 7.1, p. 92]. See also the application in [35, Chap. V, Exer. 8, p. 105].) In particular, $H_0(\rho_E) = (-1)^{n(n+1)/2} \partial_{E_1} \circ \cdots \circ \partial_{E_{n-1}} \circ \partial_{E_n}$.

Now suppose our chain complexes are bounded above and have their components drawn from an abelian category \mathcal{C} with enough injectives, and suppose we are studying the right derived functors $R^p F$ of a left exact additive functor $F : \mathcal{C} \rightarrow \mathcal{V}$, where \mathcal{V} is an abelian category. A chain complex B with B_p injective for each p is called *injective*. If $E : \cdots \rightarrow C_2 \rightarrow C_1 \rightarrow C_0 \rightarrow \cdots$ is a chain complex of such complexes

(each bounded above), then it maps (injectively) to a chain complex $E' : \cdots \rightarrow C'_2 \rightarrow C'_1 \rightarrow C'_0 \rightarrow \cdots$ of injective chain complexes (each bound above), so that for each p the map $C_p \rightarrow C'_p$ is a quasi-isomorphism; moreover, if E is exact, then E' may be chosen to be exact; also, E' may be chosen so that $C'_p = 0$ for all p with $C_p = 0$.⁹

In particular, a chain complex C maps via an injective quasi-isomorphism to an injective chain complex C' (an injective resolution). We set $RF(C) = F(C')$ and $R^pF(C) = H^p(F(C'))$, thereby extending the usual definition of RF for objects (*cohomology*) of our category to chain complexes (*hypercohomology*). The usual arguments that show this definition is independent of the choice of injective resolution can be extended to cover this case. See [35, Chap. XVII] for a detailed discussion of hyperhomology and hypercohomology. See also [34, p. 183, Exer. 17] for a discussion of resolutions of complexes.

When working with chain complexes of sheaves of abelian groups on a space X , we will use the same notation for sheaf cohomology and for sheaf hypercohomology, writing $H^p(X, C)$ whether C is a sheaf or a complex of sheaves (bounded above). In this context, one may use the flasque resolution $C \rightarrow G(C)$ constructed by Godement in [36, Chap. II, Sect. 4.3, p. 167]. The construction gives an exact functor from sheaves to flasque resolutions of them, hence, by applying it to each sheaf in a chain complex and then taking the total complex, it gives an exact functor from chain complexes to injective resolutions of them.

Whether we use injective resolutions or flasque resolutions, the formula above for $H_0(\rho_E)$ leads to an analogous formula on sheaf cohomology for $H^0(X, \rho_E) : H^0(X, C) \rightarrow H^n(X, A)$ as a composite of connecting homomorphisms with a sign.

B.2 Cup products in hypercohomology of sheaves

In this section, the tensor product $B \otimes C$ of sheaves B and C on X may denote either: tensor product of sheaves of abelian groups; tensor product of sheaves of R -modules, where R is a commutative ring; or tensor product of sheaves of \mathcal{O} -Modules, where \mathcal{O} is a sheaf of rings on X . It is an additive functor in each variable.

Godement's canonical flasque resolution $G(C)$ of a sheaf C in [36, Chap. II, Sect. 4.3, p. 167] begins with the map $\eta : C \rightarrow G^0(C) = \prod_{x \in X} (i_x)_* C_x$, where $i_x : \{x\} \rightarrow X$ is the inclusion map. The stalk of η at any point $y \in X$ can be split by projecting onto the factor corresponding to y in the product, and that shows η is injective. Moreover, if B is another sheaf, then $B \otimes \eta$ is an injective map, for the same reason. The second step in Godement's construction is $G^1(C) = G^0(\text{coker } \eta)$, and the pattern continues. Because tensor product is always right exact, it follows that $B \otimes C \rightarrow B \otimes G(C)$ is a quasi-isomorphism, and that is

⁹We only sketch the proof. As in the construction of a Cartan-Eilenberg resolution of a chain complex, one writes the chain complex in terms of short exact sequences with maps from the tail end of one to the start of the next. Then one modifies the proof that the modules in a short exact sequence have injective resolutions that fit into a short exact sequence by replacing cokernels by pushouts at a certain point. In any case, for our intended application to sheaves on a topological space, we don't really need this abstract formulation, because of the canonical Godement flasque resolution.

also true when C is a chain complex of sheaves. This fact, which we may call (*universal exactness*), makes cup product operations possible, as we shall now see. (Our approach differs slightly from Godement's original approach to cup products in [36, Chap. II, Sect. 6.1, p. 238], in that he emphasized the role of external tensor product sheaves on $X \times X$. For a thorough and modern approach, see [38].)

If B and C are chain complexes, the bicomplex $B \otimes C$ is defined by setting $(B \otimes C)_{p,q} = B_p \otimes C_q$. The vertical and horizontal differentials come from those of B and C , with no signs introduced, contrary to the standard convention [35, Chap. 4, Sect. 5].

By universal exactness of the Godement resolution, the map $\text{Tot}(B \otimes C) \rightarrow \text{Tot}(G(B) \otimes G(C))$ is a quasi-isomorphism, hence the identity map on $\text{Tot}(B \otimes C)$ can be lifted to a map from $\text{Tot}(G(B) \otimes G(C))$ to an injective resolution of $\text{Tot}(B \otimes C)$, unique up to homotopy. The resulting pairing $H^p(X, B) \otimes H^q(X, C) \rightarrow H^{p+q}(X, \text{Tot}(B \otimes C))$ is the *cup product* in hypercohomology. If a map $\text{Tot}(B \otimes C) \rightarrow D$ is given, the resulting composite pairing $H^p(X, B) \otimes H^q(X, C) \rightarrow H^{p+q}(X, D)$ may also be called a cup product pairing. We may also assemble these maps into a single map $H^*(X, B) \otimes H^*(X, C) \rightarrow H^*(X, D)$ of graded groups.

Let's examine compatibility of cup products with shifting. Composition with the map γ introduced above gives a map $\text{Tot}(B[m] \otimes C[n]) \rightarrow D[m+n]$ that leads to a cup product pairing $H^p(X, B[m]) \otimes H^q(X, C[n]) \rightarrow H^{p+q}(X, D[m+n])$. Identity maps can be used to compare this with the original cup product pairing $H^{p-m}(X, B) \otimes H^{q-n}(X, C) \rightarrow H^{p+q-m-n}(X, D)$, and the resulting discrepancy is the factor $(-1)^{mq}$ appearing in the definition of γ .

B.3 Symmetric and exterior powers of complexes

Suppose $k \geq 0$ and C is a chain complex. Let $C^{\otimes k}$ denote the tensor product $C \otimes \cdots \otimes C$ of k copies of C . The symmetric group Σ_k acts on $C^{\otimes k}$ by permuting the factors, but a sign must be inserted to get an action on $\text{Tot } C^{\otimes k}$, in accordance with the principle of signs [33]. Transposing adjacent factors involves a minus sign exactly when the two factors are both of odd degree, and the total sign can be determined by writing an arbitrary permutation as a product of adjacent transpositions. Another way of saying it that one excises the factors of even degree, collapsing to a possibly shorter tensor product, and then takes the sign of the residual permutation on the factors of odd degree.

To see that that works, it suffices to consider the case $k = 2$. Let $\tau : C_p \otimes C_q \rightarrow C_q \otimes C_p$ denote the (signed) transposition map defined by $x \otimes y \mapsto (-1)^{pq} y \otimes x$. If $x \in C_p$ and $y \in C_q$, then $\tau(d(x \otimes y)) = \tau(dx \otimes y + (-1)^p x \otimes dy) = (-1)^{(p+1)q} y \otimes dx + (-1)^{p+p(q+1)} dy \otimes x = (-1)^{pq} dy \otimes x + (-1)^{pq+q} y \otimes dx = d((-1)^{pq} y \otimes x) = d(\tau(x \otimes y))$.

Assume for the rest of the section that we are working with coherent sheaves on a variety X over a field of characteristic 0, and that the tensor product $B \otimes C$ of sheaves denotes tensor product of sheaves of \mathcal{O}_X -Modules. Assume that B is a locally free finitely generated \mathcal{O} -Module (vector bundle). Then $B \otimes C$ is an exact functor of the sheaf C . Hence, if C is an acyclic chain complex, so is $B \otimes C$. If $C \rightarrow D$ is a quasi-

isomorphism, then so is $B \otimes C \rightarrow B \otimes D$ (because being a quasi-isomorphism is determined by whether the mapping cone is acyclic, and formation of the mapping cone commutes with tensor product by B). Alternatively, assume that B is a chain complex of locally free sheaves, and that all our chain complexes are bounded above. Then if $C \rightarrow D$ is a quasi-isomorphism, then so is $\text{Tot}(B \otimes C) \rightarrow \text{Tot}(B \otimes D)$. Finally, if $C \rightarrow D$ is a quasi-isomorphism of chain complexes of locally free sheaves, then $\text{Tot } C^{\otimes k} \rightarrow \text{Tot } D^{\otimes k}$ is a quasi-isomorphism.

Now let C be a chain complex, let $S^k C$ denote the part of $\text{Tot } C^{\otimes k}$ upon which Σ_k acts trivially, and let $\wedge^k C$ denote the part of $\text{Tot } C^{\otimes k}$ upon which Σ_k acts by the sign of permutations. The projection operators $(1/k!) \sum_{\sigma \in \Sigma_k} \sigma$ and $(1/k!) \sum_{\sigma \in \Sigma_k} (-1)^\sigma \sigma$ show that $S^k C$ and $\wedge^k C$ appear functorially as direct summands of $\text{Tot } C^{\otimes k}$. Moreover, if $C \rightarrow D$ is a quasi-isomorphism, then so are the induced maps $S^k C \rightarrow S^k D$ and $\wedge^k C \rightarrow \wedge^k D$.

Let's compute symmetric and exterior powers of complexes of length 0 and length 1 in terms of symmetric and exterior powers of sheaves. Suppose C is a sheaf. When suggested by the notation, we convert C to a chain complex of length 0 by putting it in degree 0 and putting zeroes in the other positions; thus $C[m]$ will denote the chain complex of length 0 with C in position $-m$ and zeroes in the other positions. With this notation, we see that $S^k(C[m]) = (S^k C)[km]$ if m is even and $S^k(C[m]) = (\wedge^k C)[km]$ if m is odd, and $\wedge^k(C[m]) = (\wedge^k C)[km]$ if m is even and $\wedge^k(C[m]) = (S^k C)[km]$ if m is odd. Suppose now that $C = [A \xleftarrow{d} B]$ is a complex of length 1; recall that this notation puts A in degree 0 and B in degree 1. We wish to compute $(S^k C)_q$ and $(\wedge^k C)_q$ for $q \in \mathbb{Z}$; for this purpose we may assume $d = 0$, so that $C \cong A \oplus B[-1]$. Then $S^k C \cong S^k(A \oplus B[-1]) \cong \bigoplus_{p+q=k} S^p A \otimes S^q(B[-1]) \cong \bigoplus_{p+q=k} S^p A \otimes (\wedge^q B)[-q] \cong \bigoplus_{p+q=k} (S^p A \otimes \wedge^q B)[-q]$. The general conclusion is that $(S^k C)_q = S^{k-q} A \otimes \wedge^q B$, and a similar argument shows that $(\wedge^k C)_q = \wedge^{k-q} A \otimes S^q B$.

Suppose $E : 0 \rightarrow V \rightarrow B \rightarrow C \rightarrow 0$ is a short exact sequence of vector bundles. Then the map $V \rightarrow [C \leftarrow B][1]$ is a quasi-isomorphism, and hence so are the maps $S^k V \rightarrow S^k([C \leftarrow B][1])$ and $\wedge^k V \rightarrow \wedge^k([C \leftarrow B][1])$. In other words, we have long exact sequences

$$S^k E : 0 \rightarrow S^k V \rightarrow S^k B \rightarrow S^{k-1} B \otimes C \rightarrow \dots \rightarrow S^{k-p} B \otimes \wedge^p C \rightarrow \dots \rightarrow \wedge^k C \rightarrow 0$$

and

$$\wedge^k E : 0 \rightarrow \wedge^k V \rightarrow \wedge^k B \rightarrow \wedge^{k-1} B \otimes C \rightarrow \dots \rightarrow \wedge^{k-p} B \otimes S^p C \rightarrow \dots \rightarrow S^k C \rightarrow 0.$$

(Suppose alternatively that $0 \rightarrow A \rightarrow B \rightarrow V \rightarrow 0$ is an exact sequence of vector bundles. Then we have long exact sequences

$$0 \rightarrow S^k A \rightarrow S^{k-1} A \otimes B \rightarrow \dots \rightarrow S^{k-p} A \otimes \wedge^p B \rightarrow \dots \rightarrow \wedge^k B \rightarrow \wedge^k V \rightarrow 0$$

and

$$0 \rightarrow \wedge^k A \rightarrow \wedge^{k-1} A \otimes B \rightarrow \dots \rightarrow \wedge^{k-p} A \otimes S^p B \rightarrow \dots \rightarrow S^k B \rightarrow S^k V \rightarrow 0.)$$

For an alternative presentation of the portion of these results that hold in any characteristic, see [37, Sect. 2]. For the case where $V = 0$ see the exposition in [34, p. 151, Ex. 1].

Our goal now is to relate ρ_E to $\rho_{S^k E}$. We have a commutative diagram

$$\begin{array}{ccc}
(H^1(X, V))^{\otimes k} & \longrightarrow & H^k(X, S^k V) \\
\downarrow \cong & & \downarrow \cong \\
(H^1(X, [C \leftarrow B][1]))^{\otimes k} & \longrightarrow & H^k(X, S^k([C \leftarrow B][1])) \\
\uparrow & & \uparrow \\
(H^1(X, C[1]))^{\otimes k} & \longrightarrow & H^k(X, S^k(C[1]))
\end{array}$$

where the horizontal maps are obtained by iterating the cup product pairings. Let $E^k = S^k$ if k is even, and $E^k = \wedge^k$ if k is odd. Shifting the bottom row of the diagram above yields the following diagram. The vertical maps arise from identity maps, and the diagram commutes up to sign of $(-1)^{((k-1)+(k-2)+\dots+1)} = (-1)^{k(k-1)/2}$, using our earlier computation.

$$\begin{array}{ccc}
(H^1(X, C[1]))^{\otimes k} & \longrightarrow & H^k(X, S^k(C[1])) \\
\cong \uparrow & & \cong \uparrow \\
(H^0(X, C))^{\otimes k} & \longrightarrow & H^0(X, E^k C)
\end{array}$$

Splicing the two diagrams together and retaining only the top and bottom rows yields the following diagram, which commutes up to a sign of $(-1)^{k(k-1)/2}$.

$$\begin{array}{ccc}
(H^1(X, V))^{\otimes k} & \longrightarrow & H^k(X, S^k V) \\
\uparrow (\rho_E)^{\otimes k} & & \uparrow \rho_{S^k E} \\
(H^0(X, C))^{\otimes k} & \longrightarrow & H^0(X, E^k C)
\end{array}$$

The vertical maps can also be expressed in terms of connecting homomorphisms, as we have seen before.

References

- [1] B. R. Greene, K. H. Kirklin, P. J. Miron and G. G. Ross, “ 27^3 Yukawa Couplings For A Three Generation Superstring Model,” *Phys. Lett. B* **192** (1987) 111.
- [2] P. Candelas, “Yukawa Couplings Between (2,1) Forms,” *Nucl. Phys. B* **298** (1988) 458.
- [3] P. Candelas and S. Kalara, “Yukawa couplings for a three generation superstring compactification,” *Nucl. Phys. B* **298** (1988) 357.
- [4] J. McOrist and I. V. Melnikov, “Summing the Instantons in Half-Twisted Linear Sigma Models,” arXiv:0810.0012 [hep-th].
- [5] R. Donagi, R. Reinbacher and S. T. Yau, “Yukawa couplings on quintic threefolds,” arXiv:hep-th/0605203.
- [6] R. Donagi, Y. H. He, B. A. Ovrut and R. Reinbacher, “The particle spectrum of heterotic compactifications,” *JHEP* **0412**, 054 (2004) [arXiv:hep-th/0405014].
- [7] P. Berglund, L. Parkes and T. Hubsch, “The Complete Matter Sector In A Three Generation Compactification,” *Commun. Math. Phys.* **148**, 57 (1992).
- [8] M. B. Green, J. H. Schwarz and E. Witten, “Superstring Theory. Vol. 2: Loop Amplitudes, Anomalies And Phenomenology,” *Cambridge, Uk: Univ. Pr. (1987) 596 P. (Cambridge Monographs On Mathematical Physics)*
- [9] M. Gabella, Y. H. He and A. Lukas, “An Abundance of Heterotic Vacua,” *JHEP* **0812**, 027 (2008) [arXiv:0808.2142 [hep-th]].
- [10] L. B. Anderson, Y. H. He and A. Lukas, “Heterotic compactification, an algorithmic approach,” *JHEP* **0707**, 049 (2007) [arXiv:hep-th/0702210].
- [11] P. Candelas, A. M. Dale, C. A. Lutken and R. Schimmrigk, “Complete Intersection Calabi-Yau Manifolds,” *Nucl. Phys. B* **298**, 493 (1988).
- [12] C. Okonek, M. Schneider, H. Spindler, ”Vector Bundles on Complex Projective Spaces”, Birkhauser Verlag, 1988.
- [13] L. B. Anderson, Y. H. He and A. Lukas, “Monad Bundles in Heterotic String Compactifications,” arXiv:0805.2875 [hep-th].
- [14] L. B. Anderson, “Heterotic and M-theory Compactifications for String Phenomenology,” Oxford University DPhil Thesis, 2008. arXiv:0808.3621 [hep-th].

- [15] L. B. Anderson, Y. H. He, and A. Lukas, “Vector bundle stability in heterotic monad models”, In preparation.
- [16] S. K. Donaldson, “Some numerical results in complex differential geometry,” `math.DG/0512625`.
M. R. Douglas, R. L. Karp, S. Lukic and R. Reinbacher, “Numerical solution to the hermitian Yang-Mills equation on the Fermat quintic,” *JHEP* **0712**, 083 (2007) [arXiv:hep-th/0606261].
M. R. Douglas, R. L. Karp, S. Lukic and R. Reinbacher, “Numerical Calabi-Yau metrics,” *J. Math. Phys.* **49**, 032302 (2008) [arXiv:hep-th/0612075].
V. Braun, T. Brelidze, M. R. Douglas and B. A. Ovrut, “Calabi-Yau Metrics for Quotients and Complete Intersections,” *JHEP* **0805**, 080 (2008) [arXiv:0712.3563 [hep-th]].
- [17] R. Blumenhagen, S. Moster and T. Weigand, “Heterotic GUT and standard model vacua from simply connected Calabi-Yau manifolds,” *Nucl. Phys. B* **751**, 186 (2006) [arXiv:hep-th/0603015].
- [18] R. Blumenhagen, G. Honecker and T. Weigand, “Loop-corrected compactifications of the heterotic string with line bundles,” *JHEP* **0506**, 020 (2005) [arXiv:hep-th/0504232].
- [19] J. Distler and B. R. Greene, “Aspects of (2,0) String Compactifications,” *Nucl. Phys. B* **304**, 1 (1988).
- [20] A. Lukas, B. A. Ovrut and D. Waldram, “On the four-dimensional effective action of strongly coupled heterotic string theory,” *Nucl. Phys. B* **532**, 43 (1998) [arXiv:hep-th/9710208].
- [21] T. Hubsch, “Calabi-Yau manifolds: A Bestiary for physicists,” *Singapore, Singapore: World Scientific (1992) 362 p*
- [22] R. Hartshorne, “Algebraic Geometry, Springer,” GTM 52, Springer-Verlag, 1977. P. Griffith, J. Harris, “Principles of algebraic geometry,” 1978.
- [23] D. Grayson and M. Stillman, “Macaulay 2, a software system for research in algebraic geometry.” Available at <http://www.math.uiuc.edu/Macaulay2/>
- [24] G.-M. Greuel, G. Pfister, and H. Schönemann, *Singular: a computer algebra system for polynomial computations*, Centre for Computer Algebra, University of Kaiserslautern (2001). Available at <http://www.singular.uni-kl.de/>
- [25] J. Gray, Y. H. He, A. Ilderton and A. Lukas, “STRINGVACUA: A Mathematica Package for Studying Vacuum Configurations in String Phenomenology,” arXiv:0801.1508 [hep-th].
J. Gray, Y. H. He, A. Ilderton and A. Lukas, “A new method for finding vacua in string phenomenology,” *JHEP* **0707** (2007) 023 [arXiv:hep-th/0703249].
J. Gray, Y. H. He and A. Lukas, “Algorithmic algebraic geometry and flux vacua,” *JHEP* **0609** (2006) 031 [arXiv:hep-th/0606122].

The Stringvacua Mathematica package is available at:

<http://www-thphys.physics.ox.ac.uk/projects/Stringvacua/>

- [26] V. Braun, Y. H. He, B. A. Ovrut and T. Pantev, “A heterotic standard model,” *Phys. Lett. B* **618**, 252 (2005) [arXiv:hep-th/0501070].
– “The exact MSSM spectrum from string theory,” *JHEP* **0605**, 043 (2006) [arXiv:hep-th/0512177].
- [27] R. Donagi, Y. H. He, B. A. Ovrut and R. Reinbacher, “Moduli dependent spectra of heterotic compactifications,” *Phys. Lett. B* **598**, 279 (2004) [arXiv:hep-th/0403291].
- [28] B. Buchberger, “An Algorithm for Finding the Bases Elements of the Residue Class Ring Modulo a Zero Dimensional Polynomial Ideal” (German), Phd thesis, Univ. of Innsbruck (Austria), 1965.
B. Buchberger, “An Algorithmical Criterion for the Solvability of Algebraic Systems of Equations” (German), *Aequationes Mathematicae*, 4(3):374-383,1970.
English translation can be found in: B. Buchberger and F. Winkler, editors, “Gröbner Bases and Applications,” volume 251 of the L.M.S. series, Cambridge University Press, 1998. Proc. of the International Conference “33 Years of Gröbner bases”.
See B. Buchberger, “Gröbner Bases: A Short Introduction for Systems Theorists,” p1-19 Lecture Notes in Computer Science, Computer Aided Systems Theory - EUROCAST 2001, (2001 Springer Berlin/Heidelberg) for an elementary introduction.
- [29] For a recent brief review of the application of this algorithm in string phenomenology see J. Gray, “A Simple Introduction to Grobner Basis Methods in String Phenomenology,” arXiv:0901.1662 [hep-th].
- [30] L. B. Anderson, J. Gray, A. Lukas and B. Ovrut, “The Edge Of Supersymmetry: Stability Walls in Heterotic Theory,” arXiv:0903.5088 [hep-th].
- [31] L. B. Anderson, J. Gray, A. Lukas and B. Ovrut, “Stability Walls in Heterotic Theories,” to appear.
- [32] Luchezar L. Avramov and Daniel R. Grayson, *Resolutions and cohomology over complete intersections*, Computations in algebraic geometry with Macaulay 2, Algorithms Comput. Math., vol. 8, Springer, Berlin, 2002, pp. 131–178.
- [33] J. M. Boardman, *The principle of signs*, Enseignement Math. (2) **12** (1966), 191–194.
- [34] N. Bourbaki, *Éléments de mathématique. Algèbre. Chapitre 10. Algèbre homologique*, Springer-Verlag, Berlin, 2007, Reprint of the 1980 original [Masson, Paris; MR0610795].
- [35] Henri Cartan and Samuel Eilenberg, *Homological algebra*, Princeton University Press, Princeton, N. J., 1956.
- [36] Roger Godement, *Topologie algébrique et théorie des faisceaux*, Actualit’es Sci. Ind. No. 1252. Publ. Math. Univ. Strasbourg. No. 13, Hermann, Paris, 1964.

- [37] Daniel R. Grayson, *Adams operations on higher K-theory*, *K-Theory* **6** (1992), no. 2, 97–111.
- [38] Richard G. Swan, *Cup products in sheaf cohomology, pure injectives, and a substitute for projective resolutions*, *J. Pure Appl. Algebra* **144** (1999), no. 2, 169–211.
- [39] Charles A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994.