



City Research Online

City, University of London Institutional Repository

Citation: Heesen, C., Bruce, J., Gearing, R., Moss-Morris, R., Weinmann, J., Hamalainen, P., Motl, R., Dalgas, U., Kos, D., Visioli, F., Feys, P., Solari, A., Finlayson, M., Eliasson, L., Matthews, V., Bogosian, A., Liethmann, K., Kopke, S. and Bissell, P. (2015). Adherence to behavioural interventions in multiple sclerosis: Follow-up meeting report (AD@MS-2). *Multiple Sclerosis Journal - Experimental, Translational and Clinical*, 1, p. 2055217315585333. doi: 10.1177/2055217315585333

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/11846/>

Link to published version: <http://dx.doi.org/10.1177/2055217315585333>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Adherence to behavioural interventions in multiple sclerosis: Follow-up meeting report (AD@MS-2)

Christoph Heesen, Jared Bruce, Robert Gearing, Rona Moss-Morris, John Weinmann, Paivi Hamalainen, Robert Motl, Ulrik Dalgas, Daphne Kos, Francesco Visioli, Peter Feys, Alessandra Solari, Marcia Finlayson, Lina Eliasson, Vicki Matthews, Angeliki Bogossian, Katrin Liethmann, Sascha Köpke, Paul Bissell for the Adherence in Multiple Sclerosis Study Group (AD@MS group)

Multiple Sclerosis Journal – Experimental, Translational and Clinical

1: 1–4

DOI: 10.1177/
2055217315585333

© The Author(s), 2015.
Reprints and permissions:
[http://www.sagepub.co.uk/
journalsPermissions.nav](http://www.sagepub.co.uk/journalsPermissions.nav)

Abstract

After an initial meeting in 2013 that reviewed adherence to disease modifying therapy, the AD@MS group conducted a follow-up meeting in 2014 that examined adherence to behavioural interventions in MS (e.g. physical activity, diet, psychosocial interventions). Very few studies have studied adherence to behavioural interventions in MS. Outcomes beyond six months are lacking, as well as implementation work in the community. Psychological interventions need to overcome stigma and other barriers to facilitate initiation and maintenance of behaviour change. A focus group concentrated on physical activity and exercise as one major behavioural intervention domain in MS. The discussion revealed that patients are confronted with multiple challenges when attempting to regularly engage in physical activity. Highlighted needs for future research included an improved understanding of patients' and health experts' knowledge and attitudes towards physical activity as well as a need for longitudinal research that investigates exercise persistence.

Keywords: Multiple sclerosis, rehabilitation, adherence, behavioural interventions, exercise, quality of life

Date received: 29 January 2015; accepted: 2 April 2015

Introduction

Adherence to medical interventions is a global problem, particularly in chronic diseases such as multiple sclerosis (MS). Participants of a first international meeting on Adherence in MS (AD@MS) in 2013 under the auspices of the European Rehabilitation in MS (RIMS) Network concluded that there is a lack of rigorous and robust research examining adherence to medication regimens in MS.¹ The group further highlighted the importance of research examining the preferences, attitudes and needs of patients regarding behaviour change decisions, and the importance of investigating the provision of high quality patient information as a prerequisite for optimal decision making.

The goal of the second AD@MS meeting (5 June 2014) was to summarize the current knowledge of adherence to behavioural interventions in MS and to identify core areas for future research. In the 2013

consensus we defined adherence as active agreement, consent and involvement of patients in their medical treatment. This definition of adherence can include drug treatments and medically oriented diagnostic tests, but also substantial and subtle lifestyle changes. We now operationally define behavioural interventions as educational and interactive treatments designed to induce healthy non-pharmacologic behaviour change. Behavioural interventions that successfully promote a healthy lifestyle can reduce the risk and/or severity of many chronic conditions, including cardiovascular disease, cancer, depression, dementia and sexually transmitted disease.²

Rehabilitation may be the most complex behavioural intervention. It is a problem-solving educational process designed to change behaviour and adherence to these changes via enhanced activity and health participation.³ Understanding rehabilitation adherence requires the examination of multiple distinct

Correspondence to:
Christoph Heesen
Institute of
Neuroimmunology and
Multiple Sclerosis,
University Medical Centre
Hamburg-Eppendorf,
Martinistrasse 52, D-20246
Hamburg, Germany.
Email:
heesen@uke.de

Jared Bruce
University of Missouri –
Kansas City, Department of
Psychology, USA

Robert Gearing
Columbia University, New
York, USA

**Rona Moss-Morris
John Weinmann**
Health Psychology Section,
Psychology Department,
Institute of Psychiatry,
Psychology and
Neuroscience, KCL, Guy's
Hospital Campus London,
UK



Paivi Hamalainen
Director Masku Neurological
Rehabilitation Centre,
Finland

Robert Motl
Department of Kinesiology
and Community Health,
University of Illinois at
Urbana-Champaign, USA

Ulrik Dalgas
Department of Public Health
– Sport Science, Aarhus,
Denmark

Daphne Kos
Department of Rehabilitation
Sciences, KU Leuven and
Department of Health and
Social Care, AP University
College Antwerp, Belgium

Francesco Visioli
IMDEA-Food, Madrid,
Spain; University of Padova,
Italy

Peter Feys
REVAL, Biomedical
Research Institute
(BIOMED), Hasselt
University, Belgium

Alessandra Solari
Neuroepidemiology Unit,
Foundation Neurological
Institute C. Besta, Milan,
Italy

Marcia Finlayson
School of Rehabilitation
Therapy, Faculty of Health
Sciences, Queen's
University, Kingston,
Canada

Lina Eliasson
Atlantis Healthcare, Summit,
USA; Centre for
Haematology, Imperial
College London,
Hammersmith Hospital, UK

Vicki Matthews
MS Trust, Letchworth, UK

Angeliki Bogossian
Health Psychology Section,
Psychology Department,
Institute of Psychiatry,
Psychology and
Neuroscience, KCL, Guy's
Hospital Campus London,
UK

Christoph Heesen
Katrin Liethmann
Institute of
Neuroimmunology and
Multiple Sclerosis and
Department of Neurology,
UMC Hamburg Eppendorf,
Germany

Sascha Köpke
Institute for Social Medicine
and Epidemiology,
University of Lübeck,
Germany

Paul Bissell
Section of Public Health,
School of Health and Related
Research (ScHARR), The
University of Sheffield, UK

treatment approaches and processes that are commonly oriented by a set of overlapping goals. Whereas adherence to single components of a rehabilitation process (e.g. exercise, diet or use of specific self-management skills) can be studied, examining adherence to the whole approach presents significant methodological challenges.

Although adherence is a multi-dimensional construct, it is often conceptualized as unitary in drug treatment and behavioural intervention studies. Multimodal measurement of adherence may be especially important in behavioural interventions to obtain a detailed estimate of patient engagement.⁴ Moreover, the effectiveness of behavioural interventions frequently depends upon the continued application of newly acquired skills. As such, adherence research is at the core of all behavioural interventions.

The first part of the second AD@MS meeting consisted of a series of short presentations by participants detailing behavioural adherence to psychosocial interventions, diet and exercise. Next, a structured focus group ($n = 20$) was used to compile future research prospects for physical activity adherence in MS (for detailed group information see the participant list below). A list of core questions was devised by the facilitator (PB) and the focus group lasted for 90 minutes.

Summary of major topics

Diet change and adherence

As evidenced by rising rates of obesity in the western world, adherence to healthy diets requires substantial behavioural control. While there is some scientific agreement surrounding what constitutes optimal nutrition in the general population,⁵ there is no specific consensus in MS. Obesity causes fatigue and mobility problems in the general population and worsens mobility symptoms in MS. Moreover, new data suggest that obesity is associated with disease onset and symptoms expression. As such, dietary counselling in MS is highly relevant, but rarely discussed in the scientific literature. MS patients commonly seek an array of dietary interventions and supplements. Although many different diets such as vitamin D supplementation, low salt, and ketogenic diets have been touted for their possible therapeutic effects, current evidence may only justify supplementation with polyunsaturated fatty acids.^{6,7} Given the emerging importance of obesity as a risk factor for developing MS and the value placed on nutritional supplementation by MS patients, the

paucity of valid studies in this area is possibly one the most unmet needs in MS lifestyle research. Studies promoting diets in MS beyond supplementary interventions are largely absent. With improved life expectancy, addressing co-morbidities like obesity becomes increasingly relevant to patients with MS.

Behavioural interventions to manage neuropsychiatric symptoms and distress

An increasing number of studies show beneficial effects of psychosocial interventions to control and manage common mental health difficulties in MS, including fatigue, depression and cognitive dysfunction.^{8–10} Despite these successes, psychosocial interventions generally show drop-out rates from 25% to 75% and lack long-term follow-up data.⁴ Measuring adherence to these interventions is a complex endeavour. In addition to missing appointments outright, patients fail to actively engage during, between or after sessions. In contrast to drug regimens for chronic disease that typically must be taken indefinitely, psychosocial interventions may be applied in a low frequency for a defined number of sessions, some with occasional follow-up or 'booster' sessions. While session attendance can easily be measured, intra- and inter-session involvement depends on surrogates such as questionnaires, interviews and audio/video ratings. These interventions commonly promote adherence by using reminder calls, motivational interviewing, SMS texting, email, motivational enhancement techniques, and concrete support (e.g. scheduling, transportation). Homework completion has been shown to contribute substantially to treatment effects in psychosocial interventions.¹¹ In addition, promoting a strong therapeutic alliance between patients and their providers based on trust may play a substantial role in fostering adherence.

Based on the cognitive problems in MS, specific neuropsychological interventions are increasingly studied.¹⁰ They need to handle not only the stigma of psychiatric disease but also the stigma of cognitive dysfunction.

Finally, although seldom studied, acceptance-oriented psychological interventions might improve openness to behaviour change.

Physical activity

Among other benefits, regular physical activity and exercise improve functional mobility and overall quality of life in MS.¹² On the other hand, physical activity rates are reduced in MS leading to concerns about exercise adherence. Bodily limitations,

Table 1. Research issues derived from the focus group regarding physical activity (PA), exercise and adherence.

<p>PA and exercise adherence mechanisms</p> <ul style="list-style-type: none"> • There is a need for greater understanding of the mechanisms by which PA and exercise impact on outcomes ranging from quality of life through MS pathophysiology processes (i.e. inflammation, degeneration, progression). Without a better understanding of neurobiologic mechanisms of exercise, fostering adherence will remain difficult. • There is a need to design longitudinal studies to understand adherence to PA and exercise over the long term. This includes selection and application of suitable psychological models that explain health behaviour change. • It is acknowledged as a limitation that MS patients with excessive fatigue and/or significant cognitive deficits have not been included in exercise trials. Although these patients might have specific problems adhering to the training, they might also experience specific benefits. <p>Patient focused questions</p> <ul style="list-style-type: none"> • What do MS patients know and understand about the importance of PA and exercise in maintaining good health? • What priority do MS patients give to physical activity and exercise, given all the other adherence related issues they need to deal with? What do they expect to result from engaging in a physically active lifestyle and exercise? • How do MS patients' socio-economic and familial/social networks influence perception of and adhere to exercise interventions? <p>Health professional focused questions</p> <ul style="list-style-type: none"> • What do health care professionals know about the benefits and wider determinants of PA and exercise in MS, what sources of evidence do they draw on and how are these issues communicated to patients (especially in more disabled or fatigued patients)?

difficulties with service access and limited transportation are substantial barriers. Exercise studies in MS have shown as high as 80% adherence rates in the core study phase¹³ but data six months or more post intervention are scarce. Moreover, how adherence is defined in exercise studies is highly variable and no gold standard exists for the requisite percentage of training sessions completed or the amount of in-session effort required. Implementation of exercise programmes within the community is one major key to increasing physical activity among patients with MS. This can be addressed by identifying theoretically-based determinants of physical activity (e.g. self-efficacy or outcome expectations as facilitators and bodily limitations as impediments) that can then be targeted by behavioural interventions or integrated into exercise training programmes.

There is limited evidence regarding optimal approaches for initiating and then maintaining changes in physical activity and very little research has been conducted among MS patients with a progressive course. Various approaches to improving exercise adherence have been discussed, including group training, socializing via internet exchanges,

and the application of gaming settings. Health education and financial incentives have also been proposed as possible means of improving exercise adherence. However, more research is needed to fully ascertain the efficacy of these approaches. Collectively, these voids in the literature served as a major impetus for the focus group to discuss exercise and physical activity.

Focus group on adherence to physical activity in MS

The focus group had one facilitator (PB); it lasted for 90 minutes, and was audio-recorded. The focus group data were analysed iteratively for emergent themes using qualitative interpretative analysis.¹⁴ The 20 participants were multidisciplinary (see listing below). Table 1 illustrates the main findings emerging from the focus group analysis, performed by PB, JB and CH.

Conclusions

Drug and behavioural therapies both play a major role in the optimal management of MS. While a plethora of data suggest short-term effects on patient-reported outcomes, the long-term impact of behavioural interventions on psychological

wellbeing, symptom management, illness-related impairment, activity and healthcare participation is not clear. The group agreed that any large behavioural intervention study should include maintenance data beyond month 6 post treatment. In addition, it was agreed that multiple levels of adherence data should be obtained and reported for all behavioural trials. Finally, there was consensus that there is a lack of information regarding how patients and health professionals perceive nonadherence to behavioural interventions.

Acknowledgements

Participants: neurologists: A Solari, C Heesen; psychologists: P Hamalainen, J Bruce, L Eliasson, P Weinmann, R Moss-Morris, A Bougoussian, K Liethmann, A Giordano, R Gearing; health scientist: S Köpke; physiotherapists: P Feys, P van Asch; occupational therapists: D Kos, M Finlayson; sport scientists: R Motl, U Dalgas; sociologist: P Bissell; MS nurse: V Mathews; nutrition researcher: F Visioli.

Funding

This work was supported by Novartis Pharma (unconditional grant).

Conflict of interest

The authors declare that there is no conflict of interest.

References

1. Heesen C, Bruce J, Feys P, et al. Adherence in multiple sclerosis (ADAMS): Classification, relevance, and research needs. A meeting report. *Mult Scler* 2014; 20: 1795–1798.
2. Bauer UE, Briss PA, Goodman RA, et al. Prevention of chronic disease in the 21st century: Elimination of the leading preventable causes of premature death and disability in the USA. *Lancet* 2014; 384: 45–52.
3. Wade DT. Stroke: Rehabilitation and long-term care. *Lancet* 1992; 339: 791–793.
4. Gearing RE, Townsend L, Elkins J, et al. Strategies to predict, measure, and improve psychosocial treatment adherence. *Harv Rev Psychiatry* 2014; 22: 31–45.
5. Fardet A and Rock E. Toward a new philosophy of preventive nutrition: From a reductionist to a holistic paradigm to improve nutritional recommendations. *Adv Nutr* 2014; 5: 430–446.
6. Von Geldern G and Mowry EM. The influence of nutritional factors on the prognosis of multiple sclerosis. *Nat Rev Neurol* 2012; 12: 678–689.
7. Farinotti M, Vacchi L, Simi S, et al. Dietary interventions for multiple sclerosis. *Cochrane Database Syst Rev* 2012; 12: CD004192.
8. Thomas PW, Thomas S, Hillier C, et al. Psychological interventions for multiple sclerosis. *Cochrane Database Syst Rev* 2006; 1: CD004431.
9. Asano M and Finlayson ML. Meta-analysis of three different types of fatigue management interventions for people with multiple sclerosis: Exercise, education, and medication. *Mult Scler Int* 2014; 2014: 798285.
10. Rosti-Otajärvi EM and Hämäläinen PI. Neuropsychological rehabilitation for multiple sclerosis. *Cochrane Database Syst Rev* 2014; 2: CD009131.
11. Kazantzis N, Ronan KR and Deane FP. Concluding causation from correlation: Comment on Burns and Spangler (2000). *J Consult Clin Psychol* 2001; 69: 1079–1083. (discussion 1084–1086).
12. Motl RW and Pilutti LA. The benefits of exercise training in multiple sclerosis. *Nat Rev Neurol* 2012; 8: 487–497.
13. Latimer-Cheung AE, Pilutti LA, Hicks AL, et al. Effects of exercise training on fitness, mobility, fatigue, and health-related quality of life among adults with multiple sclerosis: A systematic review to inform guideline development. *Arch Phys Med Rehabil* 2013; 94: 1800–1828.
14. Seale C. Quality in qualitative research. *Qual Inquiry* 1999; 5: 465–478.