



City Research Online

City, University of London Institutional Repository

Citation: Wickwar, S., McBain, H. B., Ezra, D.G., Hirani, S. P., Rose, G.E. & Newman, S. P. (2015). Which factors are associated with quality of life in patients with Graves' orbitopathy presenting for orbital decompression surgery?. *Eye*, 29, pp. 951-957. doi: 10.1038/eye.2015.76

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/11874/>

Link to published version: <https://doi.org/10.1038/eye.2015.76>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

1 **Title: Which factors are associated with quality of life in patients with Graves'**
2 **orbitopathy presenting for orbital decompression surgery?**

3 **Authors:** Sadie Wickwar^{1,2}, Hayley B McBain^{1,3}, Daniel G Ezra^{2,4}, Shashivadan P Hirani¹, Geoffrey
4 E Rose^{2,4}, Stanton P Newman¹

5 ¹ City University London, School of Health Sciences, London, UK

6 ² Moorfields Eye Hospital, London, UK

7 ³ East London Foundation Trust, London, UK

8 ⁴ UCL Institute of Ophthalmology, NIHR Biomedical Research Centre for Ophthalmology, London,
9 UK

10 **Corresponding author and person to whom reprint requests should be addressed:**

11 Professor Stanton Newman

12 City University

13 10 Northampton Square

14 London

15 EC1V 0HB

16 Tel: +44 (0)207 040 5755

17 Email: Stanton.newman.1@city.ac.uk

18 **Running Title:** Factors associated with quality of life in GO

19 **Acknowledgements:** The authors would like to acknowledge City University London and the Special
20 Trustees at Moorfields Eye Hospital for jointly funding this research project. DGE and GER receive
21 partial funding from the Department of Health's NIHR Biomedical Research Centre for
22 Ophthalmology at Moorfields Eye Hospital and UCL Institute of Ophthalmology. The views
23 expressed in this publication are those of the authors and not necessarily those of the Department of
24 Health.

25 **Word count:** 2,673

26

27 **ABSTRACT**

28 **Aims:** Graves' orbitopathy (GO) is associated with changes in the appearance of the eyes and visual
29 dysfunction. Patients report feeling socially isolated and unable to continue with day-to-day activities.
30 This study aimed to investigate the demographic, clinical and psychosocial factors associated with
31 quality of life in patients presenting for orbital decompression surgery.

32 **Methods:** One hundred and twenty-three adults with GO due for orbital decompression at Moorfields
33 Eye Hospital London were recruited prospectively. Clinical measures including treatment history,
34 exophthalmos, optic neuropathy and diplopia were taken by an ophthalmologist. Participants
35 completed psychosocial questionnaires, including the Graves' Ophthalmopathy Quality of Life Scale
36 (GO-QOL), the Hospital Anxiety and Depression Scale (HADS) and the Derriford Appearance Scale
37 (DAS24). Hierarchical multiple regression analyses were used to identify predictors of quality of life.

38 **Results:** Higher levels of potential cases of clinical anxiety (37%) and depression (26%) were found
39 in this study sample than in patients with other chronic diseases or facial disfigurements. A total of
40 55% of the variance in GO-QOL *visual function* scores was explained by the regression model and
41 age, asymmetrical GO and depressed mood were significant unique contributors. 75% of the variance
42 in GO-QOL *appearance* scores was explained by the regression model and gender, appearance-
43 related cognitions and depressed mood were significant unique contributors.

44 **Conclusions:** Appearance-related quality of life and mood were particularly affected in this sample.
45 Predominantly psychosocial characteristics were associated with quality of life. It is important when
46 planning surgery for patients that clinicians are aware of factors that could potentially influence
47 outcomes.

48 INTRODUCTION

49 Graves' orbitopathy (GO) is an autoimmune thyroid disorder that affects the eyes. An estimated 25%
50 to 50% of patients with Graves' disease (GD) develop GO¹. The commonest early symptom of GO is a
51 noticeable change in the appearance of the eyes, including redness of the eyelids, swelling, and
52 disfiguring proptosis².

53 Patients with GO report feeling stared at by others and socially isolated as a consequence of their
54 changed appearance³ this having a significant impact on mood⁴. There is also growing evidence that
55 GO has a detrimental impact on vision-related daily functioning including reading, watching TV, and
56 driving^{5,6,7}. People with GO have been found to have a poorer quality of life than patients with other
57 chronic conditions including diabetes, emphysema and heart failure^{4,8}. Although it is important to
58 establish the impact GO may have on a patient's well-being, it is equally important to understand
59 what factors explain how some people live within normal levels of mood and experience a better
60 quality of life than others.

61 There is mixed evidence to support an association between clinical factors and quality of life in GO.
62 For instance, whilst Park et al.⁷ found that poorer quality of life was associated with more severe
63 disease, including diplopia and dry eyes, Kulig et al.⁹ failed to replicate these findings either before or
64 after treatment for GO. Recent literature about appearance and disfigurement has suggested
65 significant variability amongst individuals with disfiguring conditions – with many adjusting
66 positively to living with a visible difference – and that psychological processes rather than objective
67 measurements can better explain this variability in adjustment^{10,11}. In GO, Kahaly et al.⁴ found that
68 depressive coping, trivialising the condition and higher levels of emotional distress were associated
69 with poorer physical and mental quality of life. However, other psychosocial variables, identified
70 within a framework for adjustment to disfigurement as proposed by The Appearance Research
71 Collaboration¹¹, have not been investigated within the GO population. The framework suggests a
72 number of intervening cognitive processes, individual to each patient, might help to explain quality of
73 life in people with a disfiguring condition.

74 It was hypothesised that there will be large variation in quality of life between individuals with GO,
75 and that intervening psychosocial processes would better explain this variation than demographic or
76 clinical factors.

77 **MATERIALS & METHODS**

78 **Participants**

79 Patients were recruited from Moorfields Eye Hospital, London. Eligible patients aged 18 years or over
80 with a consultant-led diagnosis of GO, and having been listed for orbital decompression surgery, were
81 invited to participate in the study by a researcher (SW). Patients were excluded if they were
82 considered by the consultant ophthalmologist to have inadequate comprehension of written and
83 spoken English, or were suffering from psychiatric or co-morbid health conditions that rendered them
84 too ill or distressed to take part.

85 **Study design**

86 A prospective cross-sectional design was used.

87 **Measures**

88 Demographics

89 Self-reported age, gender, marital status and ethnicity were collected.

90 Clinical measures

91 The clinical measures assessed when patients were listed for surgery included ophthalmic disease
92 duration, thyroid function, treatment history, laterality of GO and planned surgery, smoking status,
93 upper and lower margin-reflex distance (MRD1 and MRD2; mm), and the presence of corneal
94 superficial punctate keratopathy (SPK), diplopia and/or signs of hydraulic orbital disease. Disease
95 activity was measured using the Clinical Activity Scale (CAS;¹²), a 10-item measure covering four of
96 the five classic signs of inflammation (pain, redness, swelling, and impaired ocular function). Visual
97 acuity was measured for each eye using a Snellen Chart. This was converted to the log of the minimal
98 angle of resolution (LogMAR), ranging between -0.20 and 2.1, with a score of 2.2 assigned to patients

99 with vision of counting fingers or worse. Optic neuropathy was identified using Ishihara colour
100 testing and, the presence of a relative afferent pupillary defect (RAPD). Proptosis was measured
101 using an Oculus exophthalmometer (in mm) and the degree of asymmetry gauged from the difference
102 between each eye (in mm).

103 Psychosocial measures

104 Self-report questionnaires were completed by participants at the time they were listed for surgery. The
105 questionnaires chosen for this study were based on a proposed framework of adjustment to GO
106 developed by the research team (Figure 1) adapted from The Appearance Research Collaboration's
107 framework¹¹. Existing validated measures were used where possible, and brief versions of
108 questionnaires were adopted to reduce participant burden.

109 Primary outcome measure

110 Quality of life was measured using the Australian version of Graves' Ophthalmopathy Quality of Life
111 Questionnaire (GO-QOL)⁷. The GO-QOL is made up of two subscales: "vision-related" and
112 "appearance-related" quality of life⁵. The GO-QOL has been found to be a valid and reliable disease-
113 specific measure of quality of life with high internal consistency ($\alpha=0.86$ for the visual function scale
114 and $\alpha=0.82$ for the appearance scale)¹³. Subscale scores were calculated following the questionnaire
115 guidelines¹³ and higher scores on each subscale indicate better health-related quality of life.

116 *Socio-cognitive factors*

117 *The Brief Fear of Negative Evaluation scale (BFNE)*¹⁴ was used to measure anxiety about others'
118 opinions. This 12-item brief version correlates highly with the original scale ($\alpha = 0.96$) and higher
119 scores indicate a greater fear of negative evaluation from others.

120 *The Iowa-Netherlands Comparison Orientation Measure (INCOM)*¹⁵ measures how well respondents
121 feel they are doing in life when comparing him or herself to others. This 11-item scale has been
122 demonstrated to have good internal consistency ($\alpha = 0.83$) and higher scores indicate a greater
123 tendency to make social comparisons.

124 *The Multidimensional Scale of Perceived Social Support (MSPSS)*¹⁶ measures subjective levels of
125 social support from family, friends and significant others. The overall scale has demonstrated good

126 internal consistency ($\alpha=0.88$) and test-retest reliability ($r=0.85$). Higher scores on each 4-item
127 subscale indicate greater perceived social support.

128 ***Patient expectations of treatment***

129 *Expectations of GO Surgery:* In the absence of a GO-specific measure of a patients' expectations of
130 treatment, an existing questionnaire – the Expectations of Strabismus Surgery Questionnaire (ESSQ)¹⁷
131 – was adapted with the items being reviewed for relevance to GO. The 23-item questionnaire
132 assessed patients' expectations in relation to three domains: “appearance concerns”, “visual
133 functioning”, and “intimacy and social relationships”. Participants were instructed to rate how they
134 expected surgery to change these aspects of their lives, for instance “my vision” on a Likert scale
135 from 1 (‘Made considerably worse’) to 5 (‘Considerably improve’). Higher scores indicate a greater
136 expectation for these areas to improve after surgery.

137 ***Appearance-specific cognitions***

138 *The Derriford Appearance Scale (DAS24)*¹⁸ measures the impact of appearance-related distress
139 including social anxiety and avoidance. This measure has demonstrated high internal consistency (α
140 = .92) and good test-retest reliability ($r=0.82$) and higher scores represent greater levels of
141 appearance-related distress and social avoidance.

142 *The Valence and Salience of Appearance scales (CARVAL & CARSAL)*¹⁹ measure how an individual
143 evaluates his or her own physical appearance (CARVAL) and the extent to which physical appearance
144 is important to the individual (CARSAL). Higher scores on each brief measure indicate a more
145 negative self-evaluation of appearance and that greater value is placed on appearance, respectively.
146 Both questionnaires have demonstrated high internal consistency (Pearson's r correlations between
147 0.72 and 0.84).

148 *Perceived Visibility of GO:* Patients were asked to rate how visible they felt their proptosis was to
149 other people on a 7-point Likert scale from 1 (‘Not at all visible’) to 7 (‘Extremely visible’).

150 ***Mood***

151 *The Hospital Anxiety & Depression scale (HADS)*²⁰ has been designed to screen for depression and
152 anxiety in patients with health problems. Higher total subscale scores on this valid and reliable
153 measure indicate greater levels of anxious or depressed mood. Cut-off scores were also applied to
154 identify non-cases (0 to 7), doubtful cases (8 to 10), and cases of possible clinical anxiety or
155 depression (scores of 11 and over).

156 **Statistical analysis**

157 Using G*Power (version 3.1.7), it was estimated that between 64 and 97 patients would be needed to
158 achieve a power of 90% with effect sizes of 0.45 and 0.9 for the GO-QOL appearance and GO-QOL
159 visual function subscales, respectively¹³.

160 All other statistical analyses were performed using IBM SPSS 21. Levels of missing data, analysed
161 using Little's Missing Completely at Random (MCAR) test were shown to be MCAR ($\chi^2 = 7127$, $df =$
162 8177 , $p = 1.000$), with 11.9% of the data missing at item level. Multiple imputation was undertaken
163 and 10 imputed datasets were generated. Scale scores for the psychosocial variables were re-
164 calculated and the analysis was conducted on all 10 datasets and the results pooled.

165 Univariate linear regressions were performed to explore the relationship between each of the
166 independent variables and the GO-QOL subscale scores (dependent variables). Hierarchical multiple
167 regressions were conducted using only the variables found to be significantly associated with each
168 GO-QOL subscale. The hierarchy used to enter the predictors into the regression was based on the
169 framework outlined in Figure 1. Cohen's f^2 was used to calculate effect sizes for each of these
170 regressions²¹. The variables were also examined for multicollinearity, linearity and homoscedasticity.
171 Multicollinearity was identified using VIF scores provided in SPSS after each regression analysis,
172 with scores above 10 indicating multicollinearity²². Histograms and normal probability plots were
173 assessed for linearity and homoscedasticity.

174 **Statement of ethics**

175 Ethical approval was obtained from the North London Research Ethics Committee (Reference
176 11/H0724/6). We certify that all applicable institutional and governmental regulations concerning the
177 ethical use of human volunteers were followed during this research

178 **RESULTS**

179 Of 192 patients identified as eligible for the study, 135 (70%) agreed to take part, and 123 of the 135
180 enrolled (91%) returned their questionnaire. Two participants' data was removed from analysis
181 because of high proportions of missing data (>50%).

182 The descriptive characteristics of the study population are shown in Table 1.

183 Summary statistics for the psychosocial variables are shown in Table 2. Possible cases of clinical
184 depression were detected in 26% of patients and 37% had possible clinical levels of anxiety; 25 (21%)
185 participants experienced both. The large standard deviations (SDs) for both GO-QOL subscales
186 indicate great variability in adjustment from patient to patient.

187 Ten of the original 36 variables were significantly associated with the GO-QOL *visual function*
188 subscale using univariate analyses: age ($F_{1,119} = 16.6, p < 0.001, f^2 = 0.14$), optic neuropathy ($F_{1,119} =$
189 $15.8, p < 0.001, f^2 = 0.15$), LogMAR ($F_{1,119} = 15.6, p < 0.001, f^2 = 0.12$), previous immunosuppression
190 ($F_{1,119} = 11.1, p = 0.001, f^2 = 0.09$), asymmetrical GO ($F_{1,119} = 6.12, p = 0.015, f^2 = 0.05$), hydraulic
191 orbit ($F_{1,119} = 9.22, p = 0.003, f^2 = 0.06$), diplopia ($F_{1,119} = 7.77, p = 0.006, f^2 = 0.07$), CAS ($F_{1,119} =$
192 $6.22, p = 0.014, f^2 = 0.05$), appearance-related social anxiety and avoidance ($F_{1,119} = 3.95, p = 0.049,$
193 $f^2 = 0.06$), anxiety ($F_{1,119} = 12.9, p < 0.001, f^2 = 0.11$), and depression ($F_{1,119} = 41.6, p < 0.001, f^2 =$
194 0.36).

195 After entry of these variables into the model in the order shown in Figure 1, 55% of the observed
196 sample variation in GO-QOL visual function score was accounted for ($R^2=0.55, F_{1,119} = 9.89, p <$
197 $0.001, f^2 = 0.8$). Beta-coefficients indicated that age, asymmetrical GO and depression made
198 significant unique contributions to the model, above other factors (Table 3).

199 Univariate analyses indicated that 13/36 variables were significantly associated with GO-QOL
200 *appearance*: age ($F_{1,119} = 5.42, p = 0.022, f^2 = 0.05$), gender ($F_{1,119} = 8.13, p = 0.005, f^2 = 0.07$),
201 previous surgery ($F_{1,119} = 5.55, p = 0.020, f^2 = 0.06$), family support ($F_{1,119} = 5.12, p = 0.025, f^2 =$
202 0.04), friends support ($F_{1,119} = 7.39, p = 0.008, f^2 = 0.06$), fear of negative evaluation ($F_{1,119} = 58.8, p <$
203 $0.001, f^2 = 0.52$), social comparison ($F_{1,119} = 12.2, p = 0.001, f^2 = 0.11$), appearance-related social
204 anxiety and avoidance ($F_{1,119} = 60.0, p < 0.001, f^2 = 0.59$), salience of appearance ($F_{1,119} = 64.6, p <$
205 $0.001, f^2 = 0.51$), valence of appearance ($F_{1,119} = 98.9, p < 0.001, f^2 = 0.76$), perceived visibility ($F_{1,119}$
206 $= 27.5, p < 0.001, f^2 = 0.24$), anxiety ($F_{1,119} = 42.2, p < 0.001, f^2 = 0.39$), and depression ($F_{1,119} = 70.5,$
207 $p < 0.001, f^2 = 0.57$).

208 After entry of the variables using the same model as before, 75% of the observed sample variation in
209 GO-QOL appearance scores was accounted for ($R^2 = 0.75, F_{13,107} = 20.7, p < 0.001, f^2 = 2.3$). Beta-
210 coefficients indicated that gender, appearance-related social anxiety and social avoidance, salience of
211 appearance, valence of appearance, perceived visibility of GO, and depression all made significant
212 contributions to the model (Table 4).

213 **DISCUSSION**

214 This study investigated the factors that may be associated with quality of life in patients with GO
215 presenting for orbital decompression surgery. It was found that being older, having asymmetrical
216 proptosis and having higher levels of depressed mood were associated with poorer vision-related
217 quality of life. Likewise, a greater value being placed on appearance, a more negative evaluation of
218 appearance, greater perceived visibility of GO, and having higher levels of depressed mood were all
219 associated with poorer appearance-related quality of life.

220 Participants in this study experienced levels of anxiety and depression greater than the general
221 population²³ and those living with other visibly disfiguring conditions²⁴. GO-QOL visual-function
222 scores were comparable to normative values established in a previous GO sample¹³, but GO-QOL
223 appearance scores were considerably lower. These results suggest that, for many, the changed

224 appearance caused by GO has a greater impact and is more debilitating than previously reported in the
225 literature.

226 Appearance-related quality of life was significantly associated with gender. For women the eyes
227 might be regarded as central in perceived attractiveness, and changes in ocular appearance could have
228 a detrimental influence on self-confidence and willingness to appear in photographs. Recent studies
229 have suggested that women with visible differences, including strabismus, may experience higher
230 levels of appearance-related distress than men^{25,26}, which in turn could impact on their quality of life
231 in this domain. Furthermore age was found to be associated with vision-related quality of life, which
232 might reflect the greater disease severity often found in older age.

233 Appearance-related cognitions were particularly important in predicting appearance-related quality of
234 life. Increased social anxiety was associated with both poorer vision-related and appearance-related
235 quality of life in this study, analogous to strabismus²⁷. Increased importance of appearance-related
236 information, as well as having a poorer evaluation of one's own appearance, were also associated with
237 quality of life in this sample. Terwee et al.²⁸ found in a study investigating perceptions of the severity
238 of GO in different groups of observers and patients themselves that clinicians tended to under-rate,
239 and patients over-rate, the severity of GO: This emphasises the importance of eliciting a patient's
240 perspective during pre-surgical assessment to improve the chance of generating realistic patient
241 expectations about what surgery can achieve.

242 A limitation of the study is the cross sectional design, which does not enable causal direction to be
243 established and longitudinal studies that follow patients over time are needed. If patients in this study
244 were not euthyroid, present hyper- or hypothyroidism could have impacted on their quality of life.
245 However, recent research found no difference in the quality of life of people with thyroid dysfunction
246 compared to people with normal thyroid levels²⁹ and it is possible that this may not have biased the
247 results of this study. It is also possible that quality of life may predict mood in GO. However, mood
248 has been found to be a strong predictor of quality of life in strabismus²⁴, supporting the current
249 findings. Furthermore, by exploring other factors that might explain variance in quality of life in this

250 population, rather than examining quality of life and mood in isolation, this study has expanded on
251 previous studies and has provided a new insight into the experiences of patients with GO.

252 In conclusion, there was significant variation in quality of life in this sample suggesting that some
253 people adjust successfully to living with GO, but for others the impact is extreme. Contrary to
254 conventional medical perspectives, this variation was predominantly accounted for by intervening
255 cognitive processes, rather than objective measures. There was however evidence that older age and
256 asymmetrical disease were associated with poorer vision-related quality of life. The high proportion
257 of patients with potentially diagnosable clinical depression and anxiety should be of concern to
258 clinicians and highlights the need for additional psychosocial support.

259 **ACKNOWLEDGEMENTS**

260 The authors would like to acknowledge those who funded the research (City University London and
261 Moorfields Eye Hospital Special Trustees). The authors also wish to thank colleagues at Moorfields
262 Eye Hospital who assisted with recruitment and data collection for the study.

263 **COMPETING INTERESTS**

264 The authors have no competing interests.

265

REFERENCES

- 1 Bahn RS, Heufelder AE. Pathogenesis of Graves' ophthalmopathy. *N Engl J Med* 1993; 329:1468-1475
- 2 Dickinson AJ. Clinical Manifestations. In Wiersinga WM, Kahaly GJ (eds). *Graves' Orbitopathy: A Multidisciplinary Approach - Questions and Answers*. Basel: Karger, 2010, pp1-25.
- 3 Jensen AL, Harder I. The impact of bodily change on social behaviour in patients with Thyroid-Associated Ophthalmopathy. *Scand J Car Sci* 2011; 25 (2):341-349
- 4 Kahaly GJ, Hardt J, Petrak F, et al. Psychosocial factors in subjects with thyroid-associated ophthalmopathy. *Thyroid* 2002; 12 (3):237-239
- 5 Choi YJ, Lim HT, Lee SJ, et al. Assessing Graves' ophthalmopathy-specific quality of life in Korean patients. *Eye* 2012; 26 (4):544-551
- 6 Ponto KA, Hommel G, Pitz S, et al. Quality of life in a German Graves orbitopathy population. *Am J Ophthalmol* 2011; 152 (3):483-490
- 7 Park JJ, Sullivan TJ, Mortimer RH, et al. Assessing quality of life in Australian patients with Graves' ophthalmopathy. *Br J Ophthalmol* 2004; 88:75-78
- 8 Gerding MN, Terwee CB, Dekker FW, et al. Quality of life in patients with Graves' ophthalmopathy is markedly decreased: Measurement by the medical outcomes study instrument. *Thyroid* 1997; 7 (6):885-889

9 Kulig G, Andrysiak-Mamos E, Sowinska-Przepiera E, et al. Quality of life assessment in patients with Graves' disease and progressive infiltrative ophthalmopathy during combined treatment with methylprednisolone and orbital radiotherapy. *Endokrynol Pol* 2009; 60 (3):158-165

10 Rumsey N, Clarke A, White P, et al. Altered body image: appearance-related concerns of people with visible disfigurement. *J Adv Nurs* 2004; 48 (5):443-453

11 Clarke A, Thompson AR, Jenkinson E, et al. *CBT for Appearance Anxiety*. Chichester: John Wiley & Sons, 2014.

12 Mourits MP, Prummel MF, Wiersinga WM, et al. Clinical activity score as a guide in the management of patients with Graves' ophthalmopathy. *Clin Endocrinol* 1997; 47:9-14

13 Terwee CB, Dekker FW, Mourits MP, et al. Interpretation and validity of changes in scores on the Graves' ophthalmopathy quality of life questionnaire (GO-QOL) after different treatments. *Clin Endocrinol* 2001; 54 (3):391-398

14 Leary MR. A brief version of the Fear of Negative Evaluation Scale. *Pers Soc Psychol Bull* 1983; 9 (3):371-375

15 Gibbons FX, Buunk BP. Individual differences in social comparison: development of a scale of social comparison orientation. *J Pers Soc Psychol* 1999; 76 (1):129-142

16 Zimet GD, Dahlem NW, Zimet SG, et al. The Multidimensional Scale of Perceived Social Support. *J Pers Assess* 1988; 52 (1):30-41

17 McBain HB, MacKenzie KA, Hancox J, et al. Why do patients with strabismus choose to undergo realignment surgery and what are their expectations of post-surgical outcomes? Development of two questionnaires. *Invest Ophthalmol Vis Sci* 2014: In Press

18 Carr T, Moss TP, Harris DL. The DAS24: a short form of the Derriford Appearance Scale DAS59 to measure individual responses to living with problems of appearance. *Br J Health Psychol* 2005; 10:285-298

19 Moss TP, Rosser BA. The moderated relationship of appearance valence on appearance self-consciousness: development and testing of new measures of appearance schema components. *PLoS ONE* 2012; 7 (11):1-7

20 Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand* 1983; 67:361-370 1987.

21 Cohen J. A power primer. *Psychol Bull* 1992; 112 (1):155-159

22 Field A. *Discovering Statistics Using SPSS*, 3rd ed. London: Sage Publications Ltd, 2009.

23 Crawford J, Henry J, Crombie C, et al. Brief report normative data for the HADS from a large non-clinical sample. *Br J Clin Psychol* 2001; 40 (4):429- 434

24 Martin CR, Newell RJ. Factor structure of the Hospital Anxiety and Depression Scale in individuals with facial disfigurement. *Psychol, Health & Med* 2004; 9 (3):327-336

25 Rumsey N, Clarke A, White P. Exploring the psychosocial concerns of outpatients with disfiguring conditions. *Journal of Wound Care* 2003; 12, (7): 247-252

26 James H, Jenkinson E, Harrad R, Ezra DG, Newman S. Appearance concerns in ophthalmic patients. Eye 2011; 25: 1039-1044

27 McBain H, MacKenzie KA, Au C, et al. Factors associated with quality of life and mood in adults with strabismus. Br J Ophthalmol 2012; 98 (4):550-555

28 Terwee CB, Dekker FW, Bonsel GJ, et al. Facial disfigurement: Is it in the eye of the beholder? A study in patients with Graves' ophthalmopathy. Clin Endocrinol 2003; 58 (2):192-198

29 Klaver EI, van Loon HCM, Stienstra R, Links TP, Keers JC, Kema IP, Kobold ACM, van der Klauw MM, Wolffenbuttel BHR. Thyroid hormone status and health-related quality of life in the lifelines cohort study. Thyroid, 2013; 23, (9): 1066-1073

TABLES

Table Error! Main Document Only.. Demographic and clinical characteristics of the study

sample

Variable	n (%)	Range	Mean±SD
Age (years)		22 - 79	47.1 ± 12.3
Gender			
<i>Male</i>	33 (27)		
<i>Female</i>	88 (73)		
Ethnicity			
<i>Asian</i>	11 (9.1)		
<i>White British/Irish/Other</i>	95 (78)		
<i>Black African/Caribbean/Other</i>	15 (12)		
Relationship Status			
<i>Married/Living with partner</i>	73 (60)		
<i>Single/Other</i>	48 (40)		
Disease duration (months)		4 - 336	62.01 ± 42
Laterality of GO			

	<i>Bilateral</i>	101 (83)		
	<i>Unilateral</i>	20 (17)		
Laterality of planned surgery				
	<i>Bilateral</i>	79 (65)		
	<i>Unilateral</i>	42 (35)		
Treatment history				
	<i>Previous immunosuppressants</i>	58 (48)		
	<i>Previous radiotherapy</i>	18 (15)		
	<i>Previous eyelid or orbital surgery</i>	14 (12)		
Thyroid function				
	<i>Stable</i>	106 (88)		
	<i>Unstable</i>	15 (12)		
Visual acuity (LogMAR conversion)†			0 - 2	0.1 ± 0.4
Superficial punctate keratopathy		39 (32)		
Hydraulic orbital signs		25 (21)		
Optic neuropathy		15 (12)		
Diplopia		62 (51)		
Marginal reflex distance 1 (mm)†			1.5 - 13	5.9 ± 2.1
Marginal reflex distance 2 (mm)†			4 - 11	6.7 ± 1.4
Exophthalmometry (mm)†			15 - 33	23.7 ± 2.7
Asymmetry (mm)			0 - 8	1.8 ± 1.8
Clinical Activity Score			0 - 9	1.12 ± 1.9
Smokers		38 (31)		

† A worst eye analysis was conducted on these variables, based on amount of proptosis

Table Error! Main Document Only.. Scores for the psychosocial measures at baseline for the study sample

Variable	Min	Max	Max Possible	Mean	SD	
GO-QOL Appearance Score	0	93.8	100	36.3	24.1	
GOQOL Visual function Score	2.8	100	100	64.0	26.7	
BFNE	14	60	60	38	9.2	
INCOM	16	54	55	36	7.2	
MSPSS Family	4	20	20	15.3	4.5	
MSPSS Friends	4	20	20	15.2	4.2	
MSPSS Significant other	4	20	20	15.6	5.2	
CARSAL	5	30	30	25.2	4.7	
CARVAL	11	48	48	38.7	8.4	
DAS24	22	83	96	51.3	13	
	<i>Men (n = 33)</i>	22	83	96	50.7	15
	<i>Women (n = 88)</i>	22	83	96	51.6	12
Visibility	1	7	7	5.7	1.5	
HADS Depression	1	21	21	9.2	4.9	

HADS Anxiety	0	19	21	7.6	4.7
---------------------	---	----	----	-----	-----

Table Error! Main Document Only.. The final step of a hierarchical multiple regression model, with GO-QOL visual function score as the dependent variable

	B	SE B	t	p
(Constant)	119.49	13.78	8.67	0.000**
Age	-0.50	0.19	-2.63	0.009*
LogMAR	-11.28	6.45	-1.75	0.080
CAS	-1.92	1.31	-1.46	0.144
Asymmetry	-2.25	1.13	-2.00	0.046*
Optic neuropathy	-4.09	9.54	-0.43	0.669
Hydraulic orbit	2.55	6.47	0.39	0.694
Previous immunomodulation	-6.79	4.90	-1.38	0.168
Diplopia	-4.39	4.29	-1.02	0.307
DAS24	0.05	0.21	0.23	0.815
HADS Anxiety	-0.69	0.58	-1.20	0.231
HADS Depression	-2.41	0.59	-4.10	0.000**

* $p < 0.05$, ** $p < 0.001$

Table Error! Main Document Only.. The final step of a hierarchical multiple regression model, with GO-QOL appearance-related score as the dependent variable

	B	SE B	t	p
(Constant)	132.09	13.84	9.55	0.000**
Age	0.11	0.11	0.94	0.346
Gender	6.56	3.02	2.17	0.03*
Previous surgery	-5.04	5.08	-0.99	0.325
BFNE	-0.23	0.22	-1.03	0.302
INCOM	0.01	0.23	0.06	0.950
MSPSS Family	-0.29	0.34	-0.87	0.382
MSPSS Friends	0.56	0.37	1.53	0.127
DAS24	-0.39	0.13	-2.93	0.004*
CARSAL	-1.23	0.33	-3.69	0.000**
CARVAL	-0.58	0.21	-2.76	0.006*
Perceived visibility	-2.75	0.96	-2.86	0.004*
HADS Anxiety	-0.05	0.39	-0.14	0.891
HADS Depression	-1.12	0.43	-2.60	0.009*

* $p < 0.05$, ** $p < 0.001$

