



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** Zu, Y. (2015). A note on asymptotic normality of kernel deconvolution density estimator with logarithmic Chi-square noise: with application in volatility density estimation. *Econometrics*, 3(3), pp. 561-576. doi: 10.3390/econometrics3030561

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/12211/>

**Link to published version:** <https://doi.org/10.3390/econometrics3030561>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

---



A note on asymptotic normality of kernel  
deconvolution density estimator with logarithmic  
Chi-square noise: with application in volatility density  
estimation

Yang Zu\*

City University London

July 15, 2015

**Abstract**

This paper studies the asymptotic normality for kernel deconvolution estimator when the noise distribution is logarithmic Chi-square, both identical and independently distributed observations and strong mixing observations are considered. The dependent case of the result is applied to obtaining the pointwise asymptotic distribution of the deconvolution volatility density estimator in a discrete-time stochastic volatility models.

JEL Classification: C13, C22, C46, C58.

Keywords: kernel deconvolution estimator, asymptotic normality, volatility density estimation.

---

\*Email: yang.zu@city.ac.uk. Department of Economics, City University London, Northampton Square, EC1V 0HB London, United Kingdom. Telephone: +44 (0)20 7040 8619. Fax: +44 (0)20 7040 8580.

# 1 Introduction

Consider the measurement error model

$$Y = X + \varepsilon,$$

where  $X$  is the signal while  $\varepsilon$  is the noise. Assume  $X$  is independent with  $\varepsilon$ ,  $X$  has density  $f_X$ ,  $\varepsilon$  has density  $k$ , so the density of  $Y$ , denoted as  $f_Y$ , is the convolution of  $f_X$  and  $k$

$$f_Y = f_X * k,$$

where the  $*$  denotes convolution.

Assume we observe the realizations  $Y_1, \dots, Y_n$  of  $Y$ , and the function  $k$  is fully known, one possible estimator for  $f_X$  from the noisy observations  $Y_1, \dots, Y_n$  is the kernel deconvolution estimator

$$\hat{f}_X(x) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-itx} \frac{\phi_K(th) \widehat{\phi_{f_Y}}(t)}{\phi_k(t)} dt, \quad (1)$$

where

$$\widehat{\phi_{f_Y}}(t) = \frac{1}{n} \sum_{j=1}^n e^{itY_j},$$

is the empirical characteristic function of density  $f_Y$ ,  $K(x)$  is a kernel function,  $\phi_K$  and  $\phi_k$  are the Fourier transform of  $K$  and  $k$ , respectively<sup>1</sup>. The kernel deconvolution estimator was first proposed for the measurement error model by Carroll and Hall (1988) and Stefanski and Carroll (1990).

Define the kernel deconvolution function as follows:

$$\nu_h(x) := \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{\phi_K(t)}{\phi_k(t/h)} e^{-itx} dt,$$

---

<sup>1</sup>Characteristic function of a random variable with density  $f$  is defined as  $\phi_f = \int_{\mathbb{R}} e^{itx} f(x) dx$ .

the kernel deconvolution estimator can be written compactly as

$$\hat{f}_X(x) = \frac{1}{nh} \sum_{j=1}^n \nu_h \left( \frac{x - Y_j}{h} \right). \quad (2)$$

In this paper, I show the asymptotic normality for the estimator  $\hat{f}_X(x)$  when the distribution of  $\varepsilon$  is logarithmic Chi-square. The asymptotic distribution of the kernel deconvolution estimator has been considered in Fan (1991a), Fan and Liu (1997), Van Es and Uh (2004), and Van Es and Uh (2005) for identically independently distributed (*i.i.d.*) observations. Masry (1993) and Kulik (2008) consider various cases for the weakly dependent observations. However, none of the above research allows the error distribution to be the logarithmic Chi-square distribution. I consider both the identical and independently distributed (*i.i.d.*) observations and strong mixing observations in this paper, which complements the above mentioned literature.

The results obtained in this paper can be applied to obtaining the asymptotic distribution of deconvolution volatility density estimator. The problem of estimating volatility density has been gaining increasing interest in econometrics in recently years, see e.g. Van Es, Spreij, and Van Zanten (2003) and Van Es, Spreij, and Van Zanten (2005) for the kernel deconvolution estimator, Comte and Genon-Catalot (2006) for the penalized projection estimator, and Todorov and Tauchen (2012) for the study in the context of high-frequency data. Kernel deconvolution with logarithmic Chi-square noise arises naturally when estimating the volatility density in Stochastic Volatility (SV) models. Existing research (e.g. Van Es, Spreij, and Van Zanten (2003) and Van Es, Spreij, and Van Zanten (2005)) focuses on the convergence rates of the estimators, and the asymptotic distribution of the estimators is not available.

In Section 2, I review the probabilistic properties of logarithmic Chi-square distribution; Section 3 presents the asymptotic normality of the estimator, for both *i.i.d.* observations and dependent observations; Section 4 discusses the application of the results to volatility density estimation in SV models; Section 5 concludes the paper.

## 2 Logarithmic Chi-square distribution

The logarithmic Chi-square distribution is obtained by taking logarithm of a Chi-square distribution with degree of freedom 1. The density function of logarithmic Chi-square distribution is

$$k(x) = \frac{1}{\sqrt{2\pi}} e^{\frac{1}{2}x} e^{-\frac{1}{2}e^x}.$$

The density function of the logarithmic Chi-square distribution is asymmetric and is plotted in Figure 1.




Figure 1: Density function of the Logarithmic Chi-square distribution

The characteristic function of the logarithmic Chi-square distribution is

$$\phi_k(t) = \frac{1}{\sqrt{\pi}} 2^{it} \Gamma\left(\frac{1}{2} + it\right),$$

where  $\Gamma(\cdot)$  is the gamma function.

Fan (1991a) studies the quadratic mean convergence rate of the kernel deconvolution estimator, it turns out that the convergence rate of the estimator depends heavily on the type of the error distribution. In particular, it is determined by the tail behavior of the modulus of the characteristic function of the error distribution - the faster of the modulus function goes to zero in the tail, the slower the converge rate. The following Lemma, which is from Van Es, Spreij, and Van Zanten (2005), gives the tail behavior of  $|\phi_k(t)|$ .

**Lemma 1** (*Lemma 5.1 of Van Es, Spreij, and Van Zanten (2005)*) For  $|t| \rightarrow \infty$ , we have

$$|\phi_k| = \sqrt{2}e^{-\frac{1}{2}\pi|t|} \left( 1 + O\left(\frac{1}{|t|}\right) \right), \quad (3)$$

and

$$\operatorname{Re} \phi_k(t) = |\phi_k| \left( \cos \left[ t \log \left( \sqrt{1 + 4t^2} - t \right) \right] + O\left(\frac{1}{|t|}\right) \right), \quad (4)$$

$$\operatorname{Im} \phi_k(t) = |\phi_k| \left( \sin \left[ t \log \left( \sqrt{1 + 4t^2} - t \right) \right] + O\left(\frac{1}{|t|}\right) \right). \quad (5)$$

From (3), it is known that the modulus of  $\phi_k(t)$  decays exponentially fast as  $|t| \rightarrow \infty$ . It thus belongs to the supersmooth density according to the classification in Fan (1991b). According to Fan (1991b), the optimal convergence rate of the estimator is  $(\log n)^{-2}$ , when  $h = (\log n)^{-1}$ . Figure 2 plots the modulus function  $|\phi_k|$  and its approximation  $\sqrt{2}e^{-\frac{1}{2}\pi|t|}$ , we notice that the two functions almost coincide at both tails.




Figure 2: Modulus function of the characteristic function of logarithmic Chi-square distribution, and its approximation: the higher peak curve is the approximating function  $\sqrt{2}e^{-\frac{1}{2}\pi|t|}$

From (4) and (5), it is known that in both tails, neither the real part nor imaginary part of the characteristic function can dominate the other, this violates the assumptions in the previous works by, e.g. Fan (1991a) and Masry (1993), on studying the asymptotic normality - for supersmooth error distributions, these papers assume either the real part or the imaginary part to be dominant.

### 3 Asymptotic normality

In this paper I consider one particular kernel function, namely the sinc kernel function:

(C1) The sinc kernel function is defined as

$$K(x) = \frac{\sin(x)}{\pi x},$$

with Fourier transform<sup>2</sup>

$$\phi_K(t) = I\{|t| \leq 1\}.$$

The sinc kernel function is favored in theoretical literature because of the simplicity of its Fourier transform and is thus used here.<sup>3</sup>

#### 3.1 *i.i.d.* observations

In this section, I prove the asymptotic normality of the estimator when the observations are *i.i.d.*.

**Theorem 1** *When the observations are i.i.d., and  $\varepsilon$  is distributed as logarithmic Chi-square, if assumption (C1) holds, when  $\exp(1/h)/n \rightarrow 0$  as  $n \rightarrow \infty$  and  $h \rightarrow 0$ , it holds that,*

$$\frac{\hat{f}_X(x) - K_h * f_X(x)}{\sqrt{\frac{1}{2\pi^2 n} \exp(\pi/h) f_Y(x)}} \rightarrow^d N(0, 1),$$

where  $K_h(x) := (1/h)K(x/h)$ .

---

<sup>2</sup>In this paper, I follow the convention to define the Fourier transform of a function  $f$  to be  $\phi_f = \int_{-\infty}^{+\infty} e^{itx} f(x) dx$ .

<sup>3</sup>Usually for practical implementations, the following kernels

$$K_1(x) = \frac{48 \cos x}{\pi x^4} \left(1 - \frac{15}{x^2}\right) - \frac{144 \sin x}{\pi x^5} \left(2 - \frac{5}{x^2}\right),$$

with Fourier transform

$$\phi_{K_1}(t) = I\{|t| \leq 1\} (1 - t^2)^3,$$

is used because it has better numerical properties, see Delaigle and Gijbels (2004) for the discussions.



**Proof** Denote

$$Z_j = \frac{1}{h} \nu_h \left( \frac{x - Y_j}{h} \right),$$

then

$$\hat{f}(x) = \frac{1}{n} \sum_{j=1}^n Z_j.$$

First

$$\begin{aligned} E\hat{f}(x) &= EZ_1 \\ &= E \left[ \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-itx} \frac{\phi_K(th) \widehat{\phi_{f_Y}}(t)}{\phi_k(t)} dt \right] \\ &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-itx} \frac{\phi_K(th) E[\widehat{\phi_{f_Y}}(t)]}{\phi_k(t)} dt \\ &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-itx} \frac{\phi_K(th) \phi_{f_Y}(t)}{\phi_k(t)} dt \\ &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{-itx} \phi_K(th) \phi_{f_X}(t) dt \\ &= K_h * f_X(x), \end{aligned}$$

Second, I evaluate  $\text{Var } Z_1$ ,

$$\begin{aligned} \text{Var } Z_1 &= \text{Var} \left( \frac{1}{h} \nu_h \left( \frac{x - Y_1}{h} \right) \right) \\ &= \frac{1}{h^2} \left( E \nu_h \left( \frac{x - Y_1}{h} \right)^2 - \left( E \nu_h \left( \frac{x - Y_1}{h} \right) \right)^2 \right) \\ &= \frac{1}{h^2} \left( \int \nu_h \left( \frac{x - y}{h} \right)^2 f_Y(y) dy - (K_h * f_X(x))^2 \right) \\ &= \frac{1}{h^2} \left( h \int \nu_h(y)^2 dy f_Y(x) - (K_h * f_X(x))^2 \right) \\ &= \frac{1}{2\pi^2} \exp \left( \frac{\pi}{h} \right) f_Y(x) (1 + o(1)), \end{aligned} \tag{6}$$

where the last equality is obtained because  $K_h * f_X(x) \rightarrow f_X(x)$  as  $h \rightarrow 0$ , and  $\int |\nu_h(x)|^2 dx = \frac{h}{2\pi^2} \exp\left(\frac{\pi}{h}\right) (1 + o(1))$ . The latter result is showed as follows,

$$\begin{aligned}
\int |\nu_h(x)|^2 dx &= \frac{1}{2\pi} \int |\phi_{\nu_h}(u)|^2 du \\
&= \frac{1}{2\pi} \int \left| \frac{\phi_K(u)}{\phi_k(u/h)} \right|^2 du \\
&= \frac{h}{\pi} \int_0^{1/h} \left| \frac{1}{\phi_k(u)} \right|^2 du \\
&= \frac{h}{\pi} \left( \int_0^M \left| \frac{1}{\phi_k(u)} \right|^2 du + \int_M^{1/h} \left| \frac{1}{\phi_k(u)} \right|^2 du \right),
\end{aligned}$$

where  $M$  is a very big number. The first term in the bracket is a constant depending on  $M$ ; the order of the second term can be evaluated as follows,

$$\begin{aligned}
\int_M^{1/h} \left| \frac{1}{\phi_k(u)} \right|^2 du &= \frac{1}{2\pi} \left( \exp\left(\frac{\pi}{h}\right) - \exp(\pi M) \right) \\
&= \frac{1}{2\pi} \exp\left(\frac{\pi}{h}\right) (1 + o(1)),
\end{aligned}$$

where I use the fact that when  $M$  is big  $|\phi_k(u)|$  can be replaced by its asymptotic approximation. The second term clearly dominates the first term, which is a constant, such that

$$\int |\nu_h(x)|^2 dx = \frac{h}{2\pi^2} \exp\left(\frac{\pi}{h}\right) (1 + o(1)). \quad (7)$$

Here I use the argument of Butucea (2004) to split the integral and show the tail part of the integral dominates.

A sufficient condition for asymptotic normality is the Lyapounov condition, which reduces to

$$\frac{E|Z_1 - EZ_1|^{2+\delta}}{n^{\delta/2} [\text{Var}(Z_1)]^{1+\delta/2}} \rightarrow 0, \quad (8)$$

for *i.i.d.* data.

For an upper bound for the numerator,

$$\begin{aligned}
E|Z_1 - EZ_1|^{2+\delta} &\leq E|Z_1|^{2+\delta} + |EZ_1|^{2+\delta} \\
&\leq 2E|Z_1|^{2+\delta} \\
&= \frac{2}{h^{2+\delta}} \int_{-\infty}^{+\infty} |\nu_h\left(\frac{x-y}{h}\right)|^{2+\delta} f_Y(y) dy \\
&\leq \frac{C}{h^{2+\delta}} \int_{-\infty}^{+\infty} |\nu_h\left(\frac{x-y}{h}\right)|^{2+\delta} dy
\end{aligned} \tag{9}$$

Now notice the result from Van Es, Spreij, and Van Zanten (2005) and Masry (1991) that, for  $p > 2$ ,<sup>4</sup>

$$\|\nu_h\|_p \leq \|\nu_h\|_\infty^{1-2/p} \|\nu_h\|_2^{2/p}.$$

An upper bound for  $\|\nu_h\|_\infty$  is easy to get, as<sup>5</sup>

$$\begin{aligned}
\|\nu_h\|_\infty &= \sup_x \left| \frac{1}{2\pi} \int \frac{\phi_K(t)}{\phi_k(t/h)} e^{-itx} dt \right| \\
&\leq \frac{1}{2\pi} \int \left| \frac{\phi_K(t)}{\phi_k(t/h)} \right| dt \\
&\leq \frac{\sqrt{2}}{\pi^2} h \exp\left(\frac{\pi}{2h}\right),
\end{aligned}$$

while  $\|\nu_h\|_2^2$  is known from (7), such that

$$\begin{aligned}
\int |\nu_h(z)|^p dz &\leq \|\nu_h\|_\infty^{p-2} \|\nu_h\|_2^2 \\
&\leq C \times h^{p-2} \exp\left(\frac{\pi(p-2)}{2h}\right) \times h \exp\left(\frac{\pi}{h}\right) \\
&= C \times h^{p-1} \exp\left(\frac{\pi p}{2h}\right),
\end{aligned}$$

for  $p > 2$ .

---

<sup>4</sup>This is easy to see by noticing  $\int |\nu_h(x)|^p dx \leq \int |\nu_h(x)|^2 |\sup_x \nu_h(x)|^{p-2} dx$  for  $p > 2$ .

<sup>5</sup>Here again the splitting integral argument as in proving (7) is used, I omit the details for ease of exposition.

So take  $p = 2 + \delta$  in (3.1), it then holds that

$$E|Z_1 - EZ_1|^{2+\delta} \leq C \times \exp\left(\frac{\pi(2+\delta)}{2h}\right), \quad (10)$$

this together with (6) imply the Lyapounov's condition (8) holds, which completes the proof.  $\square$

### 3.2 Strong mixing observations

In this section, I consider the model

$$Y = X + \varepsilon, \quad (11)$$

where  $X$ 's realizations of  $X_1, \dots, X_n$  are strictly stationary and strong mixing, while the noise realizations  $\varepsilon_1, \dots, \varepsilon_n$  are *i.i.d.* logarithmic Chi-square variables, independent with  $X$ , such that the observations  $Y_1, \dots, Y_n$  are also strictly stationary and strong mixing.

There are various concepts of dependence, here I consider the case of  $\alpha$ -mixing, also called strong mixing, which is the weakest among all the dependence concepts.

**Definition 1** *Let  $\{X_t\}$ ,  $t = \dots, -1, 0, 1, \dots$  be an infinite sequence of strictly stationary random variables, and  $\mathcal{F}_i^j$  be the  $\sigma$ -algebra generated by  $\{X_t, i \leq t \leq j\}$ , then the  $\alpha$ -mixing coefficient is defined as*

$$\alpha(k) = \sup_{A \in \mathcal{F}_{-\infty}^0, B \in \mathcal{F}_k^{+\infty}} |P(A)P(B) - P(AB)|.$$

*The sequence  $\{X_t\}$ ,  $t = \dots, -1, 0, 1, \dots$  is called  $\alpha$ -mixing if  $\alpha(k) \rightarrow 0$  as  $k \rightarrow \infty$ .*

For dependent case, a bounded assumption on the joint density of observations is also needed.

**(C2)** The probability density function of any joint distribution  $(Y_i, Y_j)$ ,  $1 \leq i < j \leq n$ , exist and bounded by a constant.

Now I give the asymptotic normality theorem. Notice that the mixing assumption here is a little weaker than that in Masry (1993).

**Theorem 2** *In model (11), let  $X_1, X_2, \dots, X_n$  be strictly stationary,  $\alpha$ -mixing with*

$$\sum_{k=1}^{\infty} \alpha(k)^{1-2/\delta} < \infty, \quad (12)$$

*for some  $\delta > 2$ ; the noise  $\varepsilon_1, \dots, \varepsilon_n$  are i.i.d. logarithmic Chi-square variables, independent with  $X$ ; if (C1) and (C2) hold, when  $\exp(1/h)/n \rightarrow 0$  as  $n \rightarrow \infty$  and  $h \rightarrow 0$ , then*

$$\frac{\hat{f}_X(x) - K_h * f_X(x)}{\sqrt{\frac{1}{2\pi^2 n} \exp(\pi/h) f_Y(x)}} \rightarrow^d N(0, 1).$$

**Proof** First by strictly stationarity and using Ergodic theorem for strong mixing sequences, similarly as in the proof of Theorem 1,

$$E\hat{f}(x) = K_h * f_X(x).$$

Next the variance of the estimator is evaluated, first

$$\text{Var}(\hat{f}(x)) = \frac{1}{n} \text{Var}(Z_1) + \frac{2}{n^2} \sum_{j=1}^{n-1} (n-j) \text{Cov}(Z_1, Z_{j+1}).$$

Knowing from Theorem 1 that the first term is

$$\frac{1}{n} \text{Var}(Z_1) = \frac{1}{2\pi^2 n} \exp\left(\frac{\pi}{h}\right) f_Y(x)(1 + o(1)). \quad (13)$$

For the covariance term, first notice

$$\begin{aligned} |\text{Cov}(Z_1, Z_{j+1})| &\leq |E(Z_1 Z_{j+1})| + (K_h * f_X(x))^2 \\ &\leq |E(Z_1 Z_{j+1})| + O(1), \end{aligned} \quad (14)$$

as  $h \rightarrow 0$ . Now because

$$\begin{aligned}
|E(Z_1 Z_{j+1})| &= \frac{1}{h^2} \left| E \left( \nu_h \left( \frac{x - Y_1}{h} \right) \nu_h \left( \frac{x - Y_{j+1}}{h} \right) \right) \right| \\
&= \frac{1}{h^2} \left| E \int \int \frac{\phi_K(t) \phi_K(t')}{\phi_k(t/h) \phi_k(t'/h)} e^{-it \frac{x - Y_1}{h}} e^{-it' \frac{x - Y_{j+1}}{h}} dt dt' \right| \\
&= \frac{1}{h^2} \left| \int \int \frac{\phi_K(t) \phi_K(t')}{\phi_k(t/h) \phi_k(t'/h)} E \left( E \left( e^{-it \frac{x - X_1 - \varepsilon_1}{h}} e^{-it' \frac{x - X_{j+1} - \varepsilon_{j+1}}{h}} \middle| X \right) \right) dt dt' \right| \\
&= \frac{1}{h^2} \left| \int \int \frac{\phi_K(t) \phi_K(t')}{\phi_k(t/h) \phi_k(t'/h)} \phi_k(t/h) \phi_k(t'/h) E \left( e^{-it \frac{x - X_1}{h}} e^{-it' \frac{x - X_{j+1}}{h}} \right) dt dt' \right| \\
&= \frac{1}{h^2} \left| \int \int \phi_K(t) \phi_K(t') E \left( e^{-it \frac{x - X_1}{h}} e^{-it' \frac{x - X_{j+1}}{h}} \right) dt dt' \right| \\
&\leq \frac{1}{h^2} \left| \int \int |\phi_K(t) \phi_K(t')| E \left( \left| e^{-it \frac{x - X_1}{h}} e^{-it' \frac{x - X_{j+1}}{h}} \right| \right) dt dt' \right| \\
&\leq \frac{1}{h^2} \left| \int \int |\phi_K(t) \phi_K(t')| dt dt' \right| \\
&\leq \frac{C}{h^2},
\end{aligned}$$

where  $C$  is a constant, continue on (14) I get

$$|\text{Cov}(Z_1, Z_{j+1})| \leq C \frac{1}{h^2} (1 + o(1)). \quad (15)$$

On the other hand, using the assumption on the  $\alpha$ -mixing coefficients and the covariance inequality for strong mixing sequence in Proposition 2.5 in Fan and Yao (2002), for  $\delta > 2$ ,

$$\begin{aligned}
\text{Cov}(Z_1, Z_{j+1}) &\leq 8\alpha(j)^{1-2/\delta} \left( E |Z_1|^\delta \right)^{1/\delta} \left( E |Z_{j+1}|^\delta \right)^{1/\delta} \\
&= 8\alpha(j)^{1-2/\delta} \left( E |Z_1|^\delta \right)^{2/\delta} \\
&\leq C' \alpha(j)^{1-2/\delta} \exp \left( \frac{\pi}{h} \right). \quad (16)
\end{aligned}$$

So using (15) and (16),

$$\begin{aligned}
& \left| \sum_{j=1}^{n-1} \text{Cov}(Z_1, Z_{j+1}) \right| \\
& \leq \left| \sum_{j=1}^{m_n} \text{Cov}(Z_1, Z_{j+1}) \right| + \left| \sum_{j=m_n}^{n-1} \text{Cov}(Z_1, Z_{j+1}) \right| \\
& \leq C \frac{1}{h^2} m_n + C \exp\left(\frac{\pi}{h}\right) \sum_{j=m_n}^{n-1} \alpha(j)^{1-2/\delta},
\end{aligned}$$

if one chooses  $m_n = \frac{1}{h|\log h|}$ , then  $m_n \rightarrow \infty$  and  $m_n h \rightarrow 0$ , then obviously the first term is  $o\left(\exp\left(\frac{\pi}{h}\right)\right)$ ; the second term is also  $o\left(\exp\left(\frac{\pi}{h}\right)\right)$  by noticing the mixing assumption in (12). Then it is shown that

$$\left| \sum_{j=1}^{n-1} \text{Cov}(Z_1, Z_{j+1}) \right| = o\left(\exp\left(\frac{\pi}{h}\right)\right). \quad (17)$$

From (13) and (17) it then follows that

$$\text{Var}\left(\hat{f}(x)\right) = \frac{1}{2\pi^2 n} \exp\left(\frac{\pi}{h}\right) f_Y(x) (1 + o(1)).$$

Now I prove the central limit theorem, for which I use the classical large block-small block argument of proving central limit theorem for dependent sequence. First I make some normalizations, define  $\sigma_0 = \left(\frac{1}{2\pi^2} \exp\left(\frac{\pi}{h}\right) f_Y(x)\right)^{1/2}$ , and

$$Z'_j = \frac{Z_j - K_h * f_X(x)}{\sigma_0},$$

then  $Z'_j$  has mean 0 and unit variance, and

$$\frac{1}{n} \sum_{j=1}^n Z'_j = \frac{\hat{f}(x) - K_h * f_X(x)}{\sigma_0},$$

and it will be shown that

$$\sqrt{n} \left( \frac{1}{n} \sum_{j=1}^n Z'_j \right) \rightarrow^d N(0, 1),$$

which is the result need to show.

First the set  $\{1, \dots, n\}$  is partitioned into  $2k_n + 1$  subsets with large blocks of size  $l_n$  and small blocks of sized  $s_n$ , such that  $k_n = \lfloor n/(l_n + s_n) \rfloor$ , so the last remaining block is having size  $n - k_n(l_n + s_n)$ . The size are such that  $l_n \rightarrow \infty$ ,  $s_n \rightarrow \infty$ ,  $l_n/s_n \rightarrow \infty$ . Then we can write

$$\sum_{j=1}^n Z'_j = \sum_{j=1}^{k_n} \xi_j + \sum_{j=1}^{k_n} \eta_j + \zeta,$$

where

$$\begin{aligned} \xi_j &= \sum_{j'=(j-1)(l_n+s_n)+1}^{(j-1)(l_n+s_n)+l_n} Z'_{j'} \\ \eta_j &= \sum_{j'=(j-1)(l_n+s_n)+l_n+1}^{j(l_n+s_n)} Z'_{j'} \\ \zeta &= \sum_{k_n(l_n+s_n)+1}^n Z'_j. \end{aligned}$$

which are sum of large blocks, small blocks, and the last block respectively. Then as a standard procedure for small block-big block argument, I show the following

$$\frac{1}{n} E \left( \sum_{j=1}^{k_n} \eta_j \right)^2 = o(1), \quad (18)$$

$$\frac{1}{n} E \zeta^2 = o(1), \quad (19)$$

$$\left| E \exp \left( i t \sum_{j=1}^{k_n} \xi_j / \sqrt{n} \right) - \prod_{j=1}^{k_n} E \exp (i t \xi_j / \sqrt{n}) \right| \rightarrow 0, \quad (20)$$

$$\frac{1}{n} \sum_{j=1}^{k_n} E \xi_j^2 \rightarrow 1, \quad (21)$$

$$\frac{1}{n} \sum_{j=1}^{k_n} E [\xi_j^2 I(|\xi_j| \geq \varepsilon n^{1/2})] \rightarrow 0, \quad (22)$$



for  $\forall \varepsilon > 0$ . (18) and (19) say the small blocks and the last block are of smaller order. (20) says that the large blocks are as if independent in the sense of characteristic function. Then (21) and (22) are Lindeberg-Feller condition for the asymptotic normality for  $\sum_{j=1}^{k_n} \xi_j$  under independence.

For (18) and (19), using the moment inequality for  $\alpha$ -mixing sequence in Proposition 2.7 (i) in Fan and Yao (2002), it can be shown that

$$\begin{aligned} E \left( \sum_{j=1}^{k_n} \eta_j \right)^2 &= O(k_n s_n), \\ E \zeta^2 &= O(n - k_n(l_n + s_n)), \end{aligned}$$

notice that the conditions for Proposition 2.7 (i) are satisfied - because by (10),  $E |Z'_j|^\delta < \infty$  for  $\delta > 2$ ; and the mixing assumption (12) implies that  $\alpha(j)^{1-2/a} = j^{-b}$  for  $b > 1$ , which is  $\alpha(j) = j^{-ab/(a-2)} = j^{-\frac{1}{2} \times \frac{1}{1/(2b)-1/(ab)}}$ , take  $\delta = ab$  and  $q = 2b$  so the mixing condition is also satisfied.

For (20), using the covariance inequality in Proposition 2.6 in Fan and Yao (2002), we have

$$\begin{aligned} & \left| E \exp \left( it \sum_{j=1}^{k_n} \xi_j / \sqrt{n} \right) - \prod_{j=1}^{k_n} E \exp (it \xi_j / \sqrt{n}) \right| \\ & \leq 16(k_n - 1)\alpha(s_n), \end{aligned}$$

this is  $o(1)$  by choosing for example  $l_n = (nh^{\gamma_1})^{1/2}$ ,  $s_n = (nh^{\gamma_2})^{1/2}$  for  $1 < \gamma_1 < \gamma_2$ . Then  $k_n = O(n^{1/2}h^{-\gamma_1/2})$ , such that for some  $q > 1$ ,

$$\begin{aligned} k_n \alpha(s_n) &= n^{1/2} h^{-\gamma_1/2} \frac{1}{(nh^{\gamma_2})^{q/2}} \\ &= n^{\frac{(1-q)}{2}} h^{-\frac{(\gamma_2 q + \gamma_1)}{2}}, \end{aligned}$$

obviously the above expression is  $o(1)$  by the assumption that  $\exp(1/h)/n \rightarrow 0$ , so (20) is proved.

For the Feller's condition (21), first use the same strategy as calculating the variance of the estimator, it holds that

$$E\xi_j^2 = l_n (1 + o(1)),$$

for any  $j$ , because  $\xi_j$  is also an infinite sum of the observations. So

$$\frac{1}{n} \sum_{j=1}^{k_n} E\xi_j^2 = \frac{1}{n} k_n l_n (1 + o(1)) \rightarrow 1.$$

Finally for the Lindeberg's condition (22), first observe that

$$\begin{aligned} E [\xi_j^2 I (|\xi_j| \geq \varepsilon n^{1/2})] &\leq (E\xi_j^4)^{1/2} P (|\xi_j| \geq \varepsilon n^{1/2}) \\ &\leq (E\xi_j^4)^{1/2} \frac{E\xi_j^2}{(\varepsilon\sqrt{n})^2}, \end{aligned}$$

where I first use Holder's inequality and then Markov's inequality. Using again the moment inequality for strong mixing sequence in Proposition 2.7 in Fan and Yao (2002),

$$\begin{aligned} (E\xi_j^4)^{1/2} \frac{E\xi_j^2}{(\varepsilon\sqrt{n})^2} &\leq (l_n^2)^{1/2} \times \frac{l_n}{(\varepsilon\sqrt{n})^2} \\ &= \frac{l_n^2}{\varepsilon^2 n}, \end{aligned}$$

so

$$\begin{aligned} \frac{1}{n} \sum_{j=1}^{k_n} E [\xi_j^2 I (|\xi_j| \geq \varepsilon n^{1/2})] &= O \left( \frac{k_n}{n} \times \frac{l_n^2}{\varepsilon^2 n} \right) \\ &= \frac{1}{\varepsilon^2} O \left( \frac{l_n}{n} \right) = o(1). \end{aligned}$$

Using the Lindeberg-Feller condition, and employ the standard argument for the proof

of central limit theorems, it can be shown that

$$\prod_{j=1}^{k_n} E \exp (it\xi_j/\sqrt{n}) \rightarrow \exp \left( -\frac{t^2}{2} \right),$$

this together with (18), (19) and (20) entail the stated result.  $\square$

## 4 Application to density estimation in stochastic volatility model

In this section, I consider applying the results of Theorem 2 to obtain the asymptotic distribution of the kernel deconvolution volatility density estimator in SV models. A generic SV model has the following form,

$$y_{t_i} = \sigma_{t_i} \varepsilon_{t_i}, \quad i = 1, \dots, n, \quad (23)$$

where  $\varepsilon_{t_i}$ ,  $i = 1, \dots, n$  are *i.i.d.*  $N(0, 1)$ ;  $\{\sigma_{t_i}\}$  is a latent stochastic process called volatility process;  $\{y_{t_i}\}$  is the observed financial returns. SV model is a popular model used in financial econometrics to describe the evolution of financial returns. Model (23) incorporates popular discrete-time SV models (e.g. Taylor (1982)) and discretized continuous-time SV model which assume the volatility process to be stationary as special cases (see e.g. Shephard (2005) for a review). Van Es, Spreij, and Van Zanten (2003) and Van Es, Spreij, and Van Zanten (2005) considered estimating the volatility density using kernel deconvolution estimator in this model.

**Remark 1** *By using the term "stochastic volatility" here, I consider the so-called "genuine stochastic volatility" models, where the volatility process has a separate stochastic driving factor (see e.g. Shephard and Andersen (2009) and Andersen, Bollerslev, Diebold, and Labys (2009) for detailed discussions). It thus does not include the ARCH/GARCH class models, where one has explicitly specified one-step-ahead predictive densities. Van Es,*

*Spreij, and Van Zanten (2005) considered estimating volatility density in the context of ARCH/GARCH class of models and had given rate of convergence for their estimator.*

To apply the general theory derived in Section 3, it is further assumed that the volatility process  $\{\sigma_{t_i}\}$  is strictly stationary, and it is independent with  $\varepsilon_{t_i}$  for  $i = 1, \dots, n$ . The independence assumption rules out the leverage effect in stochastic volatility models and thus suitable to apply to say, exchange rate data, where the leverage effect is rarely observed. Extending the model to allow for the leverage effect is an important yet challenging task, which is thus left for future research.

The SV model can be written as a measurement error model (11) by taking squares and logarithms on both sides of equation (23),

$$\log y_{t_i}^2 = \log \sigma_{t_i}^2 + \log \varepsilon_{t_i}^2, i = 1, \dots, n, \quad (24)$$

such that the variable  $\log y_i^2$  is the convolution of  $\log \eta_i$  with a completely known distribution logarithmic Chi-square. Following the notations in the previous sections, write the density functions of  $\log y_{t_i}^2$ ,  $\log \sigma_{t_i}^2$  and  $\log \varepsilon_{t_i}^2$  to be  $f_y$ ,  $f_\sigma$  and  $k$  respectively.

If we want to recover the density  $f_\sigma$  of  $\log \sigma_{t_i}^2$  from the observations  $\{\log y_{t_i}^2\}$ , this is a problem of deconvolution with logarithmic Chi-square error, and the kernel deconvolution estimator can be used. Van Es, Spreij, and Van Zanten (2003) and Van Es, Spreij, and Van Zanten (2005) first noticed this connection. Define  $Z_j := \log y_j^2$ , they use the following estimator to recover  $f_\sigma(x)$ ,

$$\hat{f}_y(x) = \frac{1}{2\pi} \frac{1}{n} \sum_{j=1}^n \int_{-\infty}^{+\infty} \frac{\phi_K(th)}{\phi_k(t)} e^{-it(x-Z_j)} dt,$$

where  $\phi_K$  is the Fourier transform of a kernel function  $K$ ,  $\phi_k(t)$  is the characteristic function of  $\log \chi_1^2$  variable. Van Es, Spreij, and Van Zanten (2003) and Van Es, Spreij, and Van Zanten (2005) derive the convergence rate of the estimator, but a central limit theorem is missing.

If we assume the observed return sequence  $\{Z_j\}$ ,  $j = 1, \dots, n$  is generated by the SV model (23) with a strictly stationary,  $\alpha$ -mixing volatility process satisfies (12) and *i.i.d.* errors, a simple application of Theorem 2 will lead to the following corollary.

**Corollary 1** *In the stochastic volatility model (23), when the volatility process  $\{\sigma_j\}$ ,  $j = 1, \dots, n$  is  $\alpha$ -mixing with (12) satisfied;  $\varepsilon_{t_i}$ 's are i.i.d.  $N(0, 1)$ , independent with the volatility process; when  $\exp(1/h)/n \rightarrow 0$  as  $n \rightarrow \infty$  and  $h \rightarrow 0$ , it holds that*

$$\frac{\hat{f}_\sigma(x) - K_h * f_\sigma(x)}{\sqrt{\frac{1}{2\pi^2 n} \exp(\pi/h) f_y(x)}} \rightarrow^d N(0, 1).$$

Since the density  $f_u(x)$  can be estimated with the observed return sequence  $\{\log y_{t_i}^2\}$  consistently using classical kernel density estimator for any  $x$ , see e.g. Fan and Yao (2002). The above result can be used to construct pointwise confidence intervals for the kernel deconvolution density estimator.

## 5 Conclusion

In this paper, I have proved the asymptotic normality for kernel deconvolution estimator with logarithmic Chi-square noise. I consider both the case identical and independently distributed observations and strong mixing observations. The results are applied to prove the asymptotic normality of kernel deconvolution estimator for volatility density in stochastic volatility models.

## Acknowledgments

I am grateful for the Editor-in-Chief, Kerry Patterson and the two anonymous referees for helpful comments, which greatly improved the paper. The usual disclaimer applies.

## References

- Andersen, T. G., T. Bollerslev, F. X. Diebold, and P. Labys (2009). Parametric and nonparametric volatility measurement. *Handbook of financial econometrics 1*, 67–138.
- Butucea, C. (2004). Asymptotic normality of the integrated square error of a density estimator in the convolution model. *Sort* 28, 9–26.
- Carroll, R. and P. Hall (1988). Optimal rates of convergence for deconvolving a density. *Journal of the American Statistical Association* 83, 1184–1186.
- Comte, F. and V. Genon-Catalot (2006). Penalized projection estimator for volatility density. *Scandinavian Journal of Statistics* 33(4), 875–893.
- Delaigle, A. and I. Gijbels (2004). Practical bandwidth selection in deconvolution kernel density estimation. *Computational statistics & data analysis* 45(2), 249–267.
- Fan, J. (1991a). Asymptotic normality for deconvolution kernel density estimators. *Sankhya: The Indian Journal of Statistics, Series A* 53(1), 97–110.
- Fan, J. (1991b). On the optimal rates of convergence for nonparametric deconvolution problems. *Annals of Statistics* 19, 1257–1272.
- Fan, J. and Q. Yao (2002). *Nonlinear time series*, Volume 2. Springer.
- Fan, Y. and Y. Liu (1997). A note on asymptotic normality for deconvolution kernel density estimators. *Sankhya: The Indian Journal of Statistics, Series A* 59(1), 138–141.
- Kulik, R. (2008). Nonparametric deconvolution problem for dependent sequences. *Electronic Journal of Statistics* 2, 722–740.
- Masry, E. (1991). Multivariate probability density deconvolution for stationary random processes. *IEEE Transactions on Information Theory* 37, 1105–1115.

- Masry, E. (1993). Asymptotic normality for deconvolution estimators of multivariate densities of stationary processes. *Journal of Multivariate Analysis* 44(1), 47–68.
- Shephard, N. (Ed.) (2005). *Stochastic Volatility: Selected Readings*. Oxford University Press.
- Shephard, N. and T. G. Andersen (2009). Stochastic volatility: Origins and overview. In T. G. Andersen, R. A. Davis, J.-P. Kreiß, and T. Mikosch (Eds.), *Handbook of Financial Time Series*, pp. 233–254. Berlin: Springer.
- Stefanski, L. and R. Carroll (1990). Deconvoluting kernel density estimators. *Statistics* 21, 169–184.
- Taylor, S. J. (1982). Financial returns modelled by the product of two stochastic processes—a study of the daily sugar prices 1961–75. *Time series analysis: theory and practice* 1, 203–226.
- Todorov, V. and G. Tauchen (2012). Inverse realized laplace transforms for nonparametric volatility density estimation in jump-diffusions. *Journal of the American Statistical Association* 107(498), 622–635.
- Van Es, B., P. Spreij, and H. Van Zanten (2003). Nonparametric volatility density estimation. *Bernoulli* 9, 451–465.
- Van Es, B., P. Spreij, and H. Van Zanten (2005). Nonparametric volatility density estimation for discrete time models. *Nonparametric Statistics* 17, 237–251.
- Van Es, B. and H. Uh (2004). Asymptotic normality of nonparametric kernel type deconvolution density estimators: crossing the Cauchy boundary. *J. Nonparametr. Stat* 16, 261–277.
- Van Es, B. and H. Uh (2005). Asymptotic normality of kernel-type deconvolution estimators. *Scandinavian journal of statistics* 32(3), 467–483.