



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** Miqdad, A., Shehzad, A., Doran, T., Ferguson, B., Fleetcroft, R., Goddard, M., Goldblatt, P., Laudicella, M., Raine, R. & Cookson, R. (2016). How a universal health system reduces inequalities: lessons from England. *Journal of Epidemiology & Community Health*, 70, pp. 637-643. doi: 10.1136/jech-2015-206742

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/13038/>

**Link to published version:** <https://doi.org/10.1136/jech-2015-206742>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

City Research Online:

<http://openaccess.city.ac.uk/>

[publications@city.ac.uk](mailto:publications@city.ac.uk)

---

# **How a universal health system reduces inequalities: lessons from England**

Miqdad Asaria, Shehzad Ali, Tim Doran, Brian Ferguson, Robert Fleetcroft, Maria Goddard, Peter Goldblatt, Mauro Laudicella, Rosalind Raine, Richard Cookson

## **Corresponding author:**

Miqdad Asaria, Research Fellow, Centre for Health Economics, University of York, York YO10 5DD, [Miqdad.asaria@york.ac.uk](mailto:Miqdad.asaria@york.ac.uk), +44(0)1904 321973

## **Co-authors:**

Ali, S, Research Fellow, Department of Health Sciences, University of York, York YO10 5DD  
Doran, T, Professor of Health Policy, Department of Health Sciences, University of York, York YO10 5DD

Ferguson B, Director for Knowledge & Intelligence, Public Health England, York, YO10 5DG  
Fleetcroft R, Clinical Lecturer in General Practice, Norwich Medical School, University of East Anglia, Norwich NR4 7TJ

Goddard M, Professor of Health Economics and Director of the Centre for Health Economics, University of York, York YO10 5DD

Goldblatt P, Deputy Director, Institute of Health Equity, UCL, London WC1E 7HB

Laudicella M, School of Health Sciences, City University London, London EC1V 0HB

Raine R, Professor of Health Care Evaluation & Head of Department of Applied Health Research, UCL, London WC1E 7HB

Cookson R, Reader, Centre for Health Economics, University of York, York YO10 5DD

**Key words:** Health Inequalities, Primary Care, Access to Health Care, Quality of Health Care, Health Outcomes

**Word count:** 2,732 words

## **Abstract**

**Background:** Provision of universal coverage is essential for achieving equity in healthcare, but inequalities still exist in universal healthcare systems. Between 2004/5 and 2011/12 the English NHS, which has provided universal coverage since 1948, made sustained efforts to reduce health inequalities by strengthening primary care. We provide the first comprehensive assessment of trends in socioeconomic inequalities of primary care access, quality and outcomes during this period.

**Methods:** Whole-population small area longitudinal study, based on 32,482 neighbourhoods of approximately 1,500 people in England from 2004/5 to 2011/12. We measured slope indices of inequality in four indicators: (i) patients per family doctor (ii) primary care quality (iii) preventable emergency hospital admissions and (iv) mortality from conditions considered amenable to healthcare.

**Results:** Between 2004/5 and 2011/12 there were larger absolute improvements on all indicators in more deprived neighbourhoods. The modelled gap between the most and least deprived neighbourhoods in England decreased by: 193 patients per family doctor (95% confidence interval 173 to 213), 3.29 percentage points of primary care quality (3.13 to 3.45), 0.42 preventable hospitalisations per 1,000 people (0.29 to 0.55) and 0.23 amenable deaths per 1,000 people (0.15 to 0.31). By 2011/12 inequalities in primary care supply and quality were almost eliminated but socio-economic inequality was still associated with 158,396 preventable hospitalisations and 37,983 deaths amenable to healthcare.

**Conclusions:** Between 2004/5 and 2011/12 the NHS succeeded in substantially reducing socioeconomic inequalities in primary care access and quality but made only modest reductions in healthcare outcome inequalities.

## **WHAT THIS PAPER ADDS**

### *Section 1: What is already known on this topic*

- There are socioeconomic inequalities in primary care access, quality and outcomes in even in high income countries with universal healthcare systems
- Reducing these inequalities by strengthening primary care was a key priority for the NHS in England from 2004/5 to 2011/12, as part of the world's first cross-government strategy for reducing health inequality
- It is not known how far the NHS succeeded in addressing this priority since national trends in healthcare equity are still not routinely monitored

### *Section 2: What this study adds*

- This study presents the first comprehensive assessment of national trends in socioeconomic inequalities in primary care access, quality and outcomes in England from 2004/5 to 2011/12
- During this period, there were substantial reductions in socioeconomic inequalities in primary care supply and quality, but only modest reductions in preventable emergency hospitalisation and mortality amenable to healthcare
- We have developed a suite of indicators that could be used in other countries to monitor the contribution of healthcare services to tackling wider inequalities in community health

## **INTRODUCTION**

Equity is widely accepted by the medical professions as a fundamental element of quality, [1,2] and providing equitable care is a priority for most national healthcare systems. [3] Provision of universal coverage is a necessary, but not sufficient, requirement for achieving this goal. In the US, the Patient Protection and Affordable Care Act (PPACA) aims to provide near universal access to healthcare coverage and to improve quality and value [4]. Recent state-level expansions of healthcare coverage have improved access to care for disadvantaged populations [5] and have been associated with improvements in mortality for causes amenable to health care. [6] However, failure to address inequalities in care within the covered population will ultimately undermine wider programs to improve quality of care and patient outcomes [7].

In the UK, the National Health Service (NHS) has provided universal, comprehensive health care free at the point of delivery since 1948. Despite this, there are clear inequities in healthcare in the UK, and poorer access and worse patient outcomes remain strongly associated with social disadvantage. [8,9] Recognising this, in 2003 the UK Government made reducing health inequality a priority for the NHS in England, as part of a cross-governmental strategy with explicit national targets for reducing health inequality by 2010 [10] – the world’s first national strategy of this kind. [11] Strengthening primary care was central to these efforts, which included: (i) major investments in primary care supply and quality from 2004, including the world’s largest primary care pay-for-performance programme; [12] (ii) targeted investment in primary care supply in under-doctored areas of the country from 2008, [13] and (iii) national guidance and support for effective primary care interventions for chronic conditions in disadvantaged adults from 2007-9. [14]

It is not known how far the NHS contributed to reducing health inequalities during this key period because socioeconomic inequalities in primary care access, quality and outcomes have not been routinely monitored. [15] This hampers efforts to improve equity, since what is not measured may be marginalised. [16] National health inequalities targets introduced in the 2000s were limited from a healthcare quality perspective as they related to local government areas, thus masking important inequalities within these areas. They also focused on life expectancy and infant mortality, over which health care providers have little direct control since they are strongly influenced by social and economic factors (e.g. living and working conditions) and related lifestyle behaviours (e.g. smoking, diet and exercise).

In this paper we address these weaknesses by constructing a suite of four key summary measures relating to trends in socioeconomic inequality in healthcare access, quality and outcomes for which the health care system can plausibly be held to account. We present data describing trends in both absolute and relative inequality in these indicators at small area level, and provide the first comprehensive assessment of trends in healthcare equity performance during a key period of sustained effort by a national healthcare system to reduce socioeconomic inequalities in primary care access, quality and outcomes.

## **METHODS**

### **Data sources**

We extracted health data from four national administrative databases for financial years 2004/5 to 2011/12: i) the annual National Health Service General and Personal Medical Services workforce census (physician supply); ii) the Quality and Outcomes Framework (QOF) - the national primary care pay-for-performance programme (primary care quality); iii) hospital episode statistics (hospital admissions); iv) the Office for National Statistics (mortality). Data on physician supply and primary care quality were attributed from practice level to small area level using the NHS Attribution Data Set of GP-registered populations. Data on hospital activity and mortality were aggregated to small area level from individual level.

The basic geographical unit of analysis was the 2001 “lower super output area” (LSOA). There are 32,482 of these small area neighbourhoods, covering approximately 1,500 people each (minimum 1,000 and maximum 3,000). We measured the population size of each neighbourhood by age-sex group using mid-year population estimates from the ONS for years 2004 to 2011. We measured the socioeconomic status of each neighbourhood using the index of multiple deprivation (IMD 2010).

### **Indicators**

We aimed to provide a comprehensive assessment of socioeconomic inequalities of primary care access, quality and outcomes for which the NHS can be held accountable in its efforts to tackle health inequality. The indicator selection process included: reviewing existing indicators used by the NHS to monitor healthcare performance; consulting with health indicator experts about



technical feasibility and with clinical and policy experts about clinical and policy relevance; and a small-scale public consultation exercise. Four key indicators were selected:

### *1. Primary care supply*

We defined primary care supply as patients per full time equivalent general practitioner (GP), excluding registrars and retainers. In line with previous studies, we focused on full time equivalent (FTE) GP principals and salaried GPs, who make up the vast majority of the workforce. [17] Neighbourhood populations were adjusted for their relative needs for primary care using the workload adjustment aspect of the Carr-Hill formula for primary care resource allocation. [18] This adjustment takes into consideration the age and sex structure and IMD 2010 “health deprivation and disability” score of each LSOA.

### *2. Primary care quality*

We defined primary care quality using a modified version of the QOF-based public health impact score proposed by Ashworth and colleagues. [19] Our indicator is a weighted average of clinical process quality from 16 QOF indicators that were collected on a consistent basis throughout our study period. Weights were proportional to importance in terms of the estimated number of lives saved per 100,000 patients. We measured practices’ reported performance, which excludes patients reported as “exceptions” (and therefore considered not to be appropriate for the quality targets) [20]. In sensitivity analysis we included exception reported patients.

### *3. Preventable hospitalisation*

We defined preventable hospitalisation as the proportion of people with an emergency admission for a chronic ambulatory care sensitive condition – admissions that are potentially avoidable if

these chronic conditions are appropriately managed in primary care. [21] We focused on chronic rather than acute ambulatory care sensitive conditions, as the former are likely to be more sensitive to changes in primary care supply and quality. We used the same list of chronic ambulatory care sensitive conditions as the NHS Outcomes Framework (Indicator 2.3i). [22] We indirectly standardised each year of data for age and sex at LSOA level.

#### *4. Amenable mortality*

We defined amenable mortality as the proportion of people dying from causes considered amenable to health care. We used the list of causes of death from the NHS Outcomes Framework (Indicator 1.1). [23] We used a simple all-age mortality rate including deaths in those aged 75 and over. As with preventable hospitalisation we indirectly standardised amenable mortality for age and sex at LSOA level.

The two healthcare outcome indicators are widely used internationally to monitor the performance of whole healthcare systems, and are particularly useful for monitoring the performance of primary care and the co-ordination of care between primary and secondary services. [24,25] Full details of the indicator definitions and the standardisation processes are provided in the supplementary appendix.

#### **Analysis**

Our primary measures of inequality were the slope index of inequality (SII) and relative index of inequality (RII), both based on linear regression analysis at LSOA level. Each indicator was modelled as a linear function of LSOA level deprivation, entered as a continuous variable scaled

from 0 to 1. The SII is the coefficient in this regression; the RII is that coefficient divided by the mean. The SII can be interpreted as the modelled absolute gap between the most and least deprived small area, allowing for the whole socioeconomic gradient; the RII can be interpreted as the proportionate gap relative to the average.

We also computed the “inequity gap”, based on a counterfactual situation of full equality in which all neighbourhoods do as well as the least deprived neighbourhood in terms of modelled achievement on the indicator. For primary care supply, the “inequity gap” is calculated as the number of additional physicians required to achieve full equality. For primary care quality, it is the average deficit in quality attributable to socioeconomic inequality. For rates of preventable hospitalisation and amenable mortality it is the number of avoidable hospitalisations and deaths attributable to socioeconomic inequality.

Linear regression models were computed using pooled data for the first and last year, including interaction terms between year and deprivation to determine the magnitude of – and test for the statistical significance of – changes in inequality between the beginning and end of the analysis period.

## **RESULTS**

### **Inequalities in 2004/5**

There were clear and substantial socioeconomic gradients in all four indicators in 2004/5 (Figure 1), with less favourable primary care provision and health outcomes in more deprived areas. For primary care supply, there were fewer GPs relative to measured need (and therefore more

patients per GP) in deprived neighbourhoods than in less deprived neighbourhoods. This socioeconomic inequality was associated with a deficit of 1,008 GPs (924 to 1,093) nationally (Table 1). In other words, equalising GP provision in all neighbourhoods to the modelled level of GP provision in the least deprived neighbourhood would require an additional 1,008 GPs in relatively deprived neighbourhoods. Socioeconomic inequality was also associated with a deficit of 1.86 percentage points (1.79 to 1.94) in primary care quality, 160,397 (158,090 to 162,703) preventable hospitalisations, and 41,433 (39,899 to 42,966) amenable deaths.

### **Changes in inequality between 2004/5 and 2011/12**

All four indicators improved on average (i.e. inequalities reduced) between 2004/5 and 2011/12. Inequalities in primary care supply and quality decreased substantially, to the extent of being virtually eliminated by the end of the period, whereas changes in the social gradient in preventable hospitalisation and amenable mortality were less pronounced (Figure 1). By 2011/12 the supply of GPs had increased in all areas, with the greatest increases in the most deprived areas, leaving neighbourhoods in the middle of the deprivation range with the fewest GPs per patient. Socioeconomic inequality had been reduced to such an extent that deprived neighbourhoods had slightly more GPs relative to need than less deprived neighbourhoods, and socioeconomic inequality was associated with a surplus of 335 GPs (233 to 436), that is: equalising GP provision in all neighbourhoods to the level of the least deprived neighbourhood would require losing 335 GPs from relatively deprived neighbourhoods.

By 2011/12, socioeconomic inequality was also associated with an average deficit in primary care quality of 0.72 percentage points (0.69 to 0.77), 158,396 excess preventable hospitalisations

(155,995 to 160,797) and 37,983 excess amenable deaths (36,552 to 39,415). Looking more closely at the trends in inequality in the indicators over the period (Table 1 and Figure 2) there is a clear trend of decreasing inequality in both absolute and relative terms for both primary care supply and primary care quality. In contrast, preventable hospitalisation and amenable mortality show a mixed pattern of decreasing absolute inequality but increasing relative inequality.

## **DISCUSSION**

Our study presents the first comprehensive national picture of how far the English NHS succeeded in reducing socioeconomic inequalities in primary care supply, quality and outcomes from 2004/5 to 2011/12. During this period, primary care supply, quality and outcomes for the average patient all improved. We find that socioeconomic inequalities in both primary care supply relative to need and primary care quality decreased substantially in both absolute and relative terms. By the end of the period, inequality in primary care supply had been eliminated and inequality in primary care quality had been nearly eliminated. In contrast, inequality trends in preventable hospitalisation and amenable mortality were mixed, showing decreasing absolute inequality but increasing relative inequality. By 2011/12, deprived neighbourhoods had slightly better primary care supply than less deprived neighbourhoods (relative inequality -2%) and only slightly worse primary care quality (relative inequality 1%). However, there remained large inequalities in preventable hospitalisation (relative inequality 106%) and amenable mortality (relative inequality 57%).

### **Strengths and weaknesses of the study**

We used data on the entire population of England, including workload and quality data on virtually all primary care practices in England and outcomes data on virtually all individuals in

England. We used comprehensive indicators spanning the entire range of activities of the healthcare system, and inequality measures based on the entire socioeconomic gradient across all 32,482 small areas of England. We examined inequality in both absolute and relative terms, because absolute and relative inequality can change in opposite directions when the mean is changing over time. [26] One of our measures – the relative index of inequality – can also be compared between indicators measured on different scales to help assess the relative magnitude of different kinds of inequality.

However, our study does not include data on privately funded healthcare, which accounts for approximately 15% of total health expenditure in the UK. [27] We also lack detailed national data on changing patterns of multi-morbidity at small area level. One consequence is that our study may under-estimate additional needs for primary care in deprived neighbourhoods, which are likely to suffer from a greater burden of multi-morbidity. [28] We also cannot assess how far observed trends in preventable hospitalisation and amenable mortality are due to trends in multi-morbidity outside the control of the NHS. Another limitation is that the administrative health datasets do not contain information on individual socioeconomic characteristics. We therefore used the index of multiple deprivation, which assumes that individuals generally conform to the socioeconomic profile of their residential neighbourhood. Finally, our measure of primary care quality is based on indicators drawn from the UK primary care pay-for-performance scheme, which only captures part of clinical practice. [29] Under this scheme improvements in quality were most rapid in practices with low baseline performance, and these practices were concentrated in more deprived areas. [30] It is possible that aspects of primary care quality that

were not financially incentivised and monitored did not follow the same pattern, and inequalities in these may have persisted or even widened.

## **Findings**

The NHS succeeded in reducing inequality in primary care supply and quality from 2004/5 to 2011/12, eliminating the inequity in primary care supply and almost eliminating the inequity in primary care quality. These changes can partly be attributed to the substantial investments in primary care in the mid to late 2000s, including the QOF pay-for-performance programme from 2004/5 and provision of additional funding for new GP practices in “under-doctored” areas of the country from 2006. [31,32] However, the NHS did not have comparable success in reducing socioeconomic inequalities in healthcare outcomes. Although absolute inequalities in healthcare outcomes decreased slightly from 2004/5 to 2011/12, relative inequalities increased and substantial inequalities remained in 2011/12 in both preventable hospitalisation and amenable mortality. This is perhaps disappointing, given that this was a period of sustained large-scale expenditure growth in the English NHS [33] and that tackling health inequality was a high priority for the NHS. [34] It is possible that non-NHS factors were acting to increase socioeconomic inequalities in healthcare outcomes during this period – evidence suggests that socioeconomic inequalities increased between 2003 and 2008 for smoking, poor diet, physical inactivity and other unhealthy behaviours. [35] It is also possible that changes in primary care supply and quality have not yet been given sufficient time to substantially reduce inequalities in healthcare outcomes, or that the national pay-for-performance program over-emphasised management of existing chronic disease over primary prevention.

The main implication of our findings, however, is that reducing inequality in healthcare outcomes is more complex and challenging than reducing inequality of access to healthcare. [36] Socioeconomic inequalities in preventable hospitalisation and amenable mortality are not only due to inequalities in the supply of primary and hospital care. They are also attributable to socioeconomic-related differences in, and complex interactions between, (i) multi-morbidity, (ii) patient behaviours including healthcare seeking, self-care and lifestyle, (iii) informal social support networks, (iv) social care supply and quality, (v) primary care provider behaviour, (vi) secondary care provider behaviour, and (vii) the co-ordination of care between primary, secondary and social care providers. Reducing socioeconomic inequalities in healthcare outcomes is therefore likely to require complex interventions to improve the co-ordination of care between multiple actors within and outwith the healthcare system. There is a growing body of evidence about effective interventions to reduce preventable hospitalisation and amenable mortality, but little is known about how to reduce socioeconomic inequalities in these healthcare outcomes. [37,38] It is our hope that the indicators developed in this study can play a role in helping to develop the evidence base for reducing inequalities in healthcare outcomes, through application to equity monitoring at local, national and international levels.



**Table 1: Socioeconomic health care inequalities in England comparing 2004/5 with 2011/12**

Indicator	England Mean (95% CI)	RII (95% CI)	SII (95% CI)	Inequality Gap (95% CI)
<i>Primary Care Supply</i>				
2004	1814 (1814 to 1814)	0.09 (0.08 to 0.09)	156.1 (141.29 to 170.91)	1,008 (924 to 1,093)
2011	1689 (1689 to 1689)	-0.02 (-0.03 to -0.01)	-36.61 (-49.8 to -23.42)	-335 (-436 to -233)
Change 2011-2004	-125 (-125 to -125)	-0.11 (-0.12 to -0.1)	-192.71 (-212.55 to -172.87)	-1,343 (-1,473 to -1,213)
<i>Primary Care Quality</i>				
2004	76.91 (76.91 to 76.91)	0.05 (0.05 to 0.05)	3.73 (3.58 to 3.87)	1.86 (1.79 to 1.94)
2011	86.34 (86.34 to 86.34)	0.01 (0.00 to 0.01)	0.44 (0.37 to 0.51)	0.22 (0.18 to 0.26)
Change 2011-2004	9.44 (9.44 to 9.44)	-0.04 (-0.05 to -0.04)	-3.29 (-3.45 to -3.13)	-1.64 (-1.72 to -1.56)
<i>Preventable Hospitalisation</i>				
2004	6.43 (6.43 to 6.44)	1.01 (0.99 to 1.02)	6.48 (6.39 to 6.58)	160,397 (158,090 to 162,703)
2011	5.73 (5.73 to 5.74)	1.06 (1.04 to 1.07)	6.07 (5.97 to 6.16)	158,396 (155,995 to 160,797)
Change 2011-2004	-0.7 (-0.71 to -0.69)	0.05 (0.03 to 0.07)	-0.42 (-0.55 to -0.29)	-2,000 (-5,270 to 1,284)
<i>Amenable Mortality</i>				
2004	3.21 (3.21 to 3.22)	0.52 (0.5 to 0.54)	1.68 (1.62 to 1.74)	41,433 (39,899 to 42,966)
2011	2.53 (2.53 to 2.54)	0.57 (0.55 to 0.59)	1.45 (1.4 to 1.5)	37,983 (36,552 to 39,415)
Change 2011-2004	-0.68 (-0.69 to -0.67)	0.05 (0.02 to 0.08)	-0.23 (-0.31 to -0.15)	-3,449 (-5,516 to -1,375)

**Notes:** The England means and the SII indices are measured in terms of patients per physician, average primary care quality, preventable hospitalisation per 1,000 and amenable mortality per 1,000. The RII indices are the SII indices as a proportion of the England means. The inequality gaps refer to the number of GPs required to eliminate inequality, the average quality loss attributable to inequality, the total excess hospitalisations attributable to inequality and the total excess mortality attributable to inequality.

**Acknowledgements:** For helpful comments, we would like to thank Peter Smith, Hugh Gravelle, Rita Santos and Mark Dusheiko. NHS Attribution Data Set data on GP-registered populations was obtained under license from the Department of Health, GMS statistics data on GP supply and hospital episode statistics data were obtained under license from the Health and Social Care Information Centre, and mortality data were obtained under license from the Office of National Statistics.

**Competing interest declaration:** All authors have completed the Unified Competing Interest form at [www.icmje.org/coi\\_disclosure.pdf](http://www.icmje.org/coi_disclosure.pdf) (available on request from the corresponding author) and declare that MA, SA, TD, RF, and MG have nothing to declare. ML reports grants from the NIHR during the conduct of this study. PG reports grants from Department of Health, grants from Public Health England and grants from British Medical Association during the conduct of the study. RC reports grants from NIHR during the conduct of the study and is a member of the NHS Outcomes Framework Technical Advisory Group. RR was supported by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care (CLAHRC) North Thames at Bart's Health NHS Trust.

**Funding declaration:** This study is independent research supported by the National Institute for Health Research (HS&DR Project: 11/2004/39 - Developing indicators of change in NHS equity performance). The funding body did not play any role in study design; in the collection, analysis, and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health.

## REFERENCES

- 1 Institute of Medicine. *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington D.C.: : National Academy Press 2001.
- 2 Smedley BD, Stith AY, Nelson AR. *Unequal Treatment: Confronting Racial and Ethnic Disparities in Health Care*. The National Academies Press 2003.
- 3 Institute of Medicine. *Toward Health Equity and Patient-Centeredness: Integrating Health Literacy, Disparities Reduction, and Quality Improvement: Workshop Summary*. Washington, DC: National Academies Press; 2009.
- 4 The *Patient Protection and Affordable Care Act (PPACA)*, Pub. L. No. 111-148, 124 *Stat.* 119, 2010.
- 5 Pande A, Ross-Degnan D, Zaslavsky A, Salomon J. Effects of health-care reforms on coverage, access, and disparities: quasi-experimental analysis of evidence from Massachusetts. *Am J Prev Med.* 2011;41:1-8.
- 6 Sommers B, Long S, Baicker K. Changes in mortality after Massachusetts health care reform – a quasi-experimental study. *Annals of Internal Medicine* 2014;160:585-593.
- 7 Davis M, Walter J. Equality-in-quality in era of the Affordable Care Act. *The Journal of the American Medical Association* 2011;306(8):872-873.
- 8 Goddard M, Smith P. Equity of access to health care services: theory and evidence from the UK. *Soc Sci Med* 2001;53:1149–62. doi:10.1016/S0277-9536(00)00415-9

- 9 Dixon A, Le Grand J, Henderson J, *et al.* Is the British National Health Service equitable? The evidence on socioeconomic differences in utilization. *J Health Serv Res Policy* 2007;**12**:104–9. doi:10.1258/135581907780279549
- 10 Department of Health. Tackling health inequalities: A Programme for Action. 2003.
- 11 Mackenbach JP. Can we reduce health inequalities? An analysis of the English strategy (1997-2010). *J Epidemiol Community Health* 2011;**65**:568–75. doi:10.1136/jech.2010.128280
- 12 Roland M. Linking physicians’ pay to the quality of care--a major experiment in the United kingdom. *N Engl J Med* 2004;**351**:1448–54. doi:10.1056/NEJMhpr041294
- 13 Department of Health. Equitable Access to Primary Medical Care. 2007.<http://webarchive.nationalarchives.gov.uk/+/www.dh.gov.uk/en/Aboutus/Procurementandproposals/Procurement/ProcurementatPCTs/index.htm>
- 14 Office NA. Tackling inequalities in life expectancy in areas with the worst health and deprivation. *Prim Care* 2010.
- 15 Wenzl M, Mccuskee S, Mossialos E. Commissioning for equity in the NHS : rhetoric and practice. *Br Med Bull* 2015;1–13. doi:10.1093/bmb/ldv031
- 16 Fiscella K, Franks P, Gold MR, *et al.* Inequality in quality: addressing socioeconomic, racial, and ethnic disparities in health care. *JAMA* Published Online First: 2000. doi:10.1001/jama.283.19.2579

- 17 NHS GP Taskforce. Securing the future GP workforce. Delivering the mandate on GP expansion. GP taskforce final report. 2014.
- 18 NHS Employer and BMA. Review of the General Medical Services global sum formula. 2007.
- 19 Ashworth M, Schofield P, Doran T, *et al.* The Public Health Impact score: A new measure of public health effectiveness for general practices in England. *Br J Gen Pract* 2013;**63**:291–9. doi:10.3399/bjgp13X665260
- 20 Doran T, Kontopantelis E, Fullwood C, *et al.* Exempting dissenting patients from pay for performance schemes: retrospective analysis of exception reporting in the UK Quality and Outcomes Framework. *Bmj* 2012;**344**:e2405–e2405. doi:10.1136/bmj.e2405
- 21 Purdy S, Griffin T, Salisbury C, *et al.* Ambulatory care sensitive conditions: terminology and disease coding need to be more specific to aid policy makers and clinicians. *Public Health* 2009;**123**:169–73. doi:10.1016/j.puhe.2008.11.001
- 22 Health and Social Care Information Centre. NHS Outcomes Framework: Domain 2 Enhancing quality of life for people with long-term conditions - Indicator specifications version 1.5. 2015.
- 23 Health and Social Care Information Centre. NHS Outcomes Framework: Domain 1 Preventing people from dying prematurely - Indicator specifications version 1.5. 2015.
- 24 Mattke S, Epstein AM, Leatherman S. The OECD Health Care Quality Indicators Project: History and background. *Int J Qual Heal Care* 2006;**18**:1–4. doi:10.1093/intqhc/mzl019

- 25 Nolte E, McKee CM. In Amenable Mortality--Deaths Avoidable Through Health Care--Progress In The US Lags That Of Three European Countries. *Health Aff* 2012;**31**:2114–22. doi:10.1377/hlthaff.2011.0851
- 26 Asada Y. On the choice of absolute or relative inequality measures. *Milbank Q* 2010;**88**:616–22. doi:10.1111/j.1468-0009.2010.00614.x
- 27 OECD. OECD Health Data 2014. OECD Heal. Data. 2014.[http://stats.oecd.org/index.aspx?DataSetCode=HEALTH\\_STAT](http://stats.oecd.org/index.aspx?DataSetCode=HEALTH_STAT)
- 28 McLean G, Gunn J, Wyke S, *et al*. The influence of socioeconomic deprivation on multimorbidity at different ages: a cross-sectional study. *Br J Gen Pract* 2014;**64**:e440–7. doi:10.3399/bjgp14X680545
- 29 Roland M, Campbell S. Successes and Failures of Pay for Performance in the United Kingdom. *N Engl J Med* 2014;**370**:1944–9. doi:10.1056/NEJMhpr1316051
- 30 Doran T, Fullwood C, Kontopantelis E, *et al*. Effect of financial incentives on inequalities in the delivery of primary clinical care in England: analysis of clinical activity indicators for the quality and outcomes framework. *Lancet* 2008;**372**:728–36. doi:10.1016/S0140-6736(08)61123-X
- 31 Department of Health. Equitable Access to Primary Medical Care. 2007.<http://webarchive.nationalarchives.gov.uk/+/www.dh.gov.uk/en/Aboutus/Procurementandproposals/Procurement/ProcurementatPCTs/index.htm>
- 32 Mays N, Tan S. Evaluating Labour’s market reforms , 2002 – 10. 2012;**17**:1–6.


- 33 Department of Health. The NHS Plan: a plan for investment, a plan for reform. 2000.  
doi:10.1136/bmj.321.7257.315
- 34 Bleich SN, Jarlenski MP, Bell CN, *et al.* Health Inequalities: Trends, Progress, and Policy. *Annu Rev Public Health* 2012;**33**:7–40. doi:10.1146/annurev-publhealth-031811-124658
- 35 Buck D, Frosini F. Clustering of unhealthy behaviours over time. 2012.
- 36 Purdy S. Interventions to reduce unplanned hospital admissions: A series of systematic reviews. 2012.
- 37 Vest JR, Gamm LD, Oxford BA, *et al.* Determinants of preventable readmissions in the United States: a systematic review. *Implement Sci* 2010;**5**:88. doi:10.1186/1748-5908-5-88
- 38 Purdy S. Avoiding hospital admissions. *What does Res Evid say* 2010;:1–28.

### **Figure 1: Scatter plots of indicators in 2004/05 and 2011/12**


**Notes:** The black dots show deprivation decile groups of neighbourhoods (approximately 3,200 neighbourhoods per dot); the solid black line shows a linear regression through all 32,482 neighbourhoods; the shaded area shows the inequality gap; and the dashed red line shows the national average level for the indicator. \*inverted axis on primary care quality to ease comparisons with other indicators where decreasing implies improvement




### Primary Care Supply




### Primary Care Quality\*



### Preventable Hospitalisation



### Amenable Mortality



**figure 2: Inequality trends from 2004/5 to 2011/12**

**Notes:** \*inverted axis on primary care quality to ease comparisons with other indicators where decreasing implies improvement

