



City Research Online

## City, University of London Institutional Repository

---

**Citation:** Almond, M. H., O'Donoghue, M., Drey, N., Seneviratne, S., Lalvani, A., Wickremasinghe, M. & Kon, O. M. (2011). Interferon-gamma release assay (IGRA) conversion, reversion and implications for the diagnosis of latent tuberculosis infection using a multimodality approach: a retrospective, observational study within a central London TB centre. *Thorax*, 66(Suppl 4), A72. doi: 10.1136/thoraxjnl-2011-201054c.13

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/13826/>

**Link to published version:** <https://doi.org/10.1136/thoraxjnl-2011-201054c.13>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

City Research Online:

<http://openaccess.city.ac.uk/>

[publications@city.ac.uk](mailto:publications@city.ac.uk)

---

# **A comparison of a central London latent TB screening algorithm with national UK and US algorithms and the effect on LTBI diagnosis**

Marie O'Donoghue<sup>1</sup> Nicholas Drey<sup>2</sup> Mark Almond<sup>1</sup> Melissa Wickremasinghe<sup>1</sup> Suranjith Seneviratne<sup>1</sup>  
Ajit Lalvani<sup>3</sup> Onn Min Kon<sup>1</sup>

<sup>1</sup>Chest and Allergy Department, St Mary's Hospital, Imperial College Healthcare NHS Trust, London,

<sup>2</sup>School of Health Sciences, City University, London

<sup>3</sup>Tuberculosis Research Unit, National Heart and Lung Institute, Imperial College, London

## **Introduction**

There is continuing debate over the most effective strategy for the use of interferon gamma release assays (IGRAs) in clinical practice and this is reflected in the diversity of published recommendations. Practice in UK TB contact screening clinics reflects these widely divergent views.

The aims of our study were to:

1. Enumerate the prevalence of latent TB infection (LTBI) according to US, UK and local screening protocols
2. Explore which algorithm is the most effective in a central London TB clinic

## **Methods**

This was a cross sectional retrospective study comprising 558 consecutive adult contacts of TB cases that attended for screening tests for LTBI 2008-2010. The screening results were applied to patient management algorithms derived from US guidance (IGRA or tuberculin skin test but not both), UK NICE guidance (two step approach), and our local service (concurrent IGRA and TST). For the purpose of this analysis, LTBI was defined as the outcome predetermined by each algorithm.

## **Results**

In our cohort, NICE guidance had the highest completion rate (97%) with the most complex algorithm (local practice) at 87%. The diagnosis of LTBI had a variability of 5-26% dependent on which algorithm was used. NICE screens contacts aged >35 years with a chest x-ray and misses the immunodiagnosis of LTBI in 12-16% of this group. The NICE approach is also less sensitive in contacts aged <35 years because of the two step approach (IGRA is only performed following a positive TST) and the failure to rescreen contacts of pulmonary cases who are TST or IGRA negative at the initial screen. 11 (4.6%) of pulmonary TB contacts aged <35 years that would be discharged by NICE had a TST or IGRA conversion. The US IGRA alone algorithm misses a proportion (6.9%) of individuals; 11 (2%) had a strongly positive TST ( $\geq 20$ mm), and of these, 5 had a TST conversion varying 6-16mm. This approach

also potentially over diagnosed 8 (2%) of contacts who reverted on rescreen when initially discordant (TST negative / IGRA positive).

### **Conclusion**

The sensitivity of the screening depends not only on the performance of the tests but also what algorithm is used. As demonstrated in our local data, the overall effectiveness of any approach is also affected by how these impact on the patient journey and their ability to complete the screening process. These findings also highlight the impact of conversion and reversion phenomena in a clinical service.

Word count 399

