



City Research Online

City, University of London Institutional Repository

Citation: Santhi, K., Shafqat, K., Kyriacou, P. A. & Pal, S. K. (2007). Investigation of heart rate variability in patients undergoing hand surgery under axillary plexus block. British Journal of Anaesthesia, 98(2), 282P-283P. doi: 10.1093/bja/ael319

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/14701/>

Link to published version: <https://doi.org/10.1093/bja/ael319>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Investigation of heart rate variability in patients undergoing hand surgery under axillary plexus block.

K Shafqat^{*1}, S Kumari^{*2}, P A Kyriacou^{*1}, and S K Pal²

¹ School of Engineering and Mathematical Sciences, City University, London, UK

² St Andrew's Centre for Plastic Surgery & Burns, Broomfield Hospital, Chelmsford, Essex.

Heart Rate Variability (HRV) is the study of inter-beat variability of heart rate and can be used as a non-invasive technique to assess the autonomic influence on the heart ¹. The two most commonly used methods to analyse HRV are time domain and frequency domain analysis. In the frequency domain, three frequency bands can be distinguished in the spectrum of short term (2 to 5 minutes) HRV signals ¹. These components are termed as: a) High-Frequency (HF) 0.15 Hz to 0.4Hz; b) Low-Frequency (LF) band 0.04 Hz to 0.15 Hz; and c) Very Low-Frequency band (VLF) which is the band of less than 0.04-Hz frequencies. The HF region is mediated almost entirely by the vagal tone. The LF region is an indicator of sympathetic tone, but is also modulated by vagal activity. The physiological aspects behind VLF band are not well established and therefore in short term recording this band should be avoided ¹. Using this information HRV indices such as the ratio of LF/HF power or the fractional LF power has been used to describe sympathovagal balance ². The aim of this study was to investigate the effect of local anaesthetics in patients having brachial plexus block on HRV indices. We recruited 14 ASA I-II patients, 7 males and 7 females, aged 50.57 ± 20.72 years (mean weight 67 ± 15.33 Kg, mean height 1.63 ± 0.2 m) undergoing hand surgery under axillary block. All patients received a standard mixture of 30 ml lignocaine and 20 ml of 0.5% bupivacaine with adrenaline 1:200,000 for the block. Lead II ECG signal was collected from the patients. The monitoring started 30 minutes before the start of the block and continued 30 minutes after the surgery. For frequency domain analysis 5 minutes of data (overlapped by 50 %) was analysed using a non-parametric method of spectral analysis which has been used in various HRV studies ³. The analysis showed that in almost 80% of the cases the LF/HF ratio increases just after the application of the brachial block and then decreases considerably. The timing of the drop in the ratio value differs in patients, but in each case the drop occurs within an hour of start of the block. Some examples of such changes are presented in figure 1. This initial study has indicated that HRV parameters may change during the brachial plexus block using a mixture of lignocaine and bupivacaine with adrenaline.




Figure 1: Ratio LF/HF changes after the brachial block.

References

- 1) Task Force of the Euro. Society of Cardiology the N. American Society of Pacing Electrophysiology, *European Heart Journal* 1996; (18): 354-381.
- 2) J. J. Goldberger, *Am. J. Physiol.* 1999; (276): H1273-H1280.
- 3) D. Singh, K. Vinod, S. C. Saxena et al *Physiol. Meas.* 2004; (25): 721-735.