



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** Ashford, M., Olander, E. K. & Ayers, S. (2016). Finding Web-Based Anxiety Interventions on the World Wide Web: A Scoping Review. *JMIR Mental Health*, 3(2), e14. doi: 10.2196/mental.5349

This is the published version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/14842/>

**Link to published version:** <https://doi.org/10.2196/mental.5349>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

---



Review

# Finding Web-Based Anxiety Interventions on the World Wide Web: A Scoping Review

Miriam Thiel Ashford, BSc, MSc; Ellinor K Olander, PhD; Susan Ayers, PhD

Centre for Maternal and Child Health Research, School of Health Sciences, City University London, London, United Kingdom

**Corresponding Author:**

Miriam Thiel Ashford, BSc, MSc  
Centre for Maternal and Child Health Research  
School of Health Sciences  
City University London  
Northampton Square  
London, EC1V 0HB  
United Kingdom  
Phone: 44 20 7040 ext 5775  
Fax: 44 0207 040 5808  
Email: [miriam.ashford@city.ac.uk](mailto:miriam.ashford@city.ac.uk)

## Abstract

**Background:** One relatively new and increasingly popular approach of increasing access to treatment is Web-based intervention programs. The advantage of Web-based approaches is the accessibility, affordability, and anonymity of potentially evidence-based treatment. Despite much research evidence on the effectiveness of Web-based interventions for anxiety found in the literature, little is known about what is publically available for potential consumers on the Web.

**Objective:** Our aim was to explore what a consumer searching the Web for Web-based intervention options for anxiety-related issues might find. The objectives were to identify currently publically available Web-based intervention programs for anxiety and to synthesize and review these in terms of (1) website characteristics such as credibility and accessibility; (2) intervention program characteristics such as intervention focus, design, and presentation modes; (3) therapeutic elements employed; and (4) published evidence of efficacy.

**Methods:** Web keyword searches were carried out on three major search engines (Google, Bing, and Yahoo—UK platforms). For each search, the first 25 hyperlinks were screened for eligible programs. Included were programs that were designed for anxiety symptoms, currently publically accessible on the Web, had an online component, a structured treatment plan, and were available in English. Data were extracted for website characteristics, program characteristics, therapeutic characteristics, as well as empirical evidence. Programs were also evaluated using a 16-point rating tool.

**Results:** The search resulted in 34 programs that were eligible for review. A wide variety of programs for anxiety, including specific anxiety disorders, and anxiety in combination with stress, depression, or anger were identified and based predominantly on cognitive behavioral therapy techniques. The majority of websites were rated as credible, secure, and free of advertisement. The majority required users to register and/or to pay a program access fee. Half of the programs offered some form of paid therapist or professional support. Programs varied in treatment length and number of modules and employed a variety of presentation modes. Relatively few programs had published research evidence of the intervention's efficacy.

**Conclusions:** This review represents a snapshot of available Web-based intervention programs for anxiety that could be found by consumers in March 2015. The consumer is confronted with a diversity of programs, which makes it difficult to identify an appropriate program. Limited reports and existence of empirical evidence for efficacy make it even more challenging to identify credible and reliable programs. This highlights the need for consistent guidelines and standards on developing, providing, and evaluating Web-based interventions and platforms with reliable up-to-date information for professionals and consumers about the characteristics, quality, and accessibility of Web-based interventions.

(*JMIR Ment Health* 2016;3(2):e14) doi:[10.2196/mental.5349](https://doi.org/10.2196/mental.5349)

**KEYWORDS**

Anxiety; mental health; web-based interventions; internet; technology; consumer; access to health care

## Introduction

The National Comorbidity Survey Replication showed that 28.8% of people in the United States suffer from an anxiety disorder in their lifetime [1]. Reviews suggest that anxiety disorders are the most frequently occurring class of mental health disorders [2-4] and are considered chronic and disabling conditions worldwide [5]. Despite effective treatments being available, anxiety disorders are still widely underdiagnosed and undertreated [4,6]. The adverse effects of anxiety disorders on psychological and somatic health, as well as high economic costs [2,6-8] mean that treatment is a significant public health issue.

Lack of help-seeking behavior and perceived barriers to accessing treatment contribute to underdiagnosis and undertreatment. Generally, individuals with anxiety display a tendency not to seek help for their disorder [9,10]. Identified treatment barriers include lack of awareness of the presence of a disorder and available services, financial burden, and the stigma associated with disclosing mental health disorders [10,11].

Research has shown that many individuals use the Internet to find information or help for health-related topics [12], especially for topics that they experience as difficult to talk about [13,14]. A survey demonstrated that 18% of all surveyed Internet users had searched the Internet for mental health-related information, with higher prevalence for those who had a history of mental health issues and those who at the time stated that they were experiencing psychological distress [15]. Similarly, nationally representative surveys from the Pew Internet and American Life Project found that 26-39% of individuals who sought Web-based health information looked at mental health information [14,16]. When searching the Internet for mental health information, individuals may come across Web-based interventions.

A Web-based intervention has been defined as “a primarily self-guided intervention program that is executed by means of a prescriptive online program operated through a website.” (p. 5) [17]. Advantages of Web-based approaches include accessibility, affordability, and anonymity of mental health interventions [18,19]. Web-based interventions can be accessed anytime and anywhere from devices such as computers, laptops, tablets, and mobile phones and large audiences and rural areas can be reached in a cost-effective manner [20-22]. Web-based interventions also offer anonymity and privacy, which may attract individuals who experience difficulties with disclosing mental health disorders [23-24].

The efficacy of Web-based mental health intervention programs is well established. Meta-analyses of Web-based mental health interventions have shown that those interventions were as effective as face-to-face treatments and superior to control groups with substantial effect sizes [25,26]. With regard to anxiety disorders specifically, a meta-analysis concluded that computerized- and Internet-based cognitive behavioral therapy (CBT) for anxiety disorders had improved outcomes compared to waitlist and placebo assignments and these effects were equal to face-to-face treatment [27,28]. Another meta-analytic review concluded that computer-aided psychotherapy was as effective

as face-to-face therapy and that the effects did not differ across various anxiety disorders and types of delivery [29]. Similarly, a recent review reported moderate to large effect sizes for Internet-based CBT for a range of anxiety disorders ranging from 0.30 to 2.53 [30].

Despite an extensive body of literature evaluating the effectiveness of developed Web-based interventions, little research has examined the range and characteristics of publically available Web-based intervention programs for individuals with mental health issues. Research has started to identify, describe, and evaluate the range and characteristics of mental health mobile phone apps [31,32] and e-therapy or e-counselling services [33,34]. However, e-therapy is different from Web-based programs, as in e-therapy mental health professionals use text- or video-based formats (eg, email, chat, Skype) for delivering therapy. There is also a clinical online directory of Web-based mental health programs called Beacon available, which lists among others, intervention program websites for phobias, generalized anxiety disorder, social anxiety disorder, and panic disorder [35]. However, this website is not updated very often and is clinically directed rather than a systematic review of programs that are publically available. A few publically available Web-based programs were briefly discussed in a review; however, this was restricted to four programs available in Australia (FearFighter, Beating the Blues, Online Anxiety, CRUFAD) and included only a short overall description of the main program characteristics [36]. In addition, the programs were not identified by a systematic Web search. Recently, a scoping review has identified and evaluated currently available interactive Web-based interventions for depression [37]. However, to our knowledge no study has conducted a similar review for Web-based intervention programs for anxiety.

In summary, despite the clear advantages of Web-based anxiety interventions, there is only limited systematically identified and up-to-date information available on the characteristics of publically available Web-based interventions for anxiety and the quality of these services is currently unknown. This information would be helpful and important for consumers and practitioners interested in Web-based interventions for anxiety, as well as researchers developing and evaluating those interventions. Therefore, this study conducted a replicable Web search to identify freely available Web-based anxiety intervention programs and review these in terms of (1) website characteristics such as origin, accessibility, and credibility; (2) Web-based program characteristics, such as intervention focus, design, delivery, and features; (3) intervention characteristics such as the overall therapeutic approach and intervention features; as well as (4) published evidence of efficacy.

## Methods

### Search Strategy

Using the 3 most popular Web search engines, Google, Bing, and Yahoo [38,39], a keyword search for Web-based intervention programs for anxiety was performed in March 2015. UK versions of the search engines were used (.co.uk). Before starting the search, existing search history and cookies were deleted and future tracking and cookies were disabled in

the browser. A list of the 9 search term combinations used can be found in [Textbox 1](#). Primarily simple and lay keywords were used to simulate a Web search that was relatively likely to be conducted by an individual searching for Web-based programs. It has been shown that most individuals rarely consider more

than the first 20 links generated by a search engine [40]. As featured links placed at the top and bottom were also considered, we chose to assess the first 25 links. This resulted in 675 hyperlinks being screened (3 search engines  $\times$  9 search terms  $\times$  25 hyperlinks).

**Textbox 1.** Search terms used in Google, Bing, and Yahoo.

1. Internet therapy anxiety
2. Internet treatment anxiety
3. Internet cognitive behavioural therapy anxiety
4. Online therapy anxiety
5. Online treatment anxiety
6. Online cognitive behavioural therapy anxiety
7. Web therapy anxiety
8. Web treatment anxiety
9. Web cognitive behavioural therapy anxiety

## Program Identification

All 675 hyperlinks were screened for eligible Web-based programs for anxiety by the first author. The screening process consisted of two stages. The first stage involved screening all 675 hyperlinked websites to eliminate clearly irrelevant websites. All hyperlinks were screened and organized into 1 of the following 3 categories: websites with Web-based programs, websites linking to websites with Web-based programs, and websites with irrelevant content. Irrelevant content included, among others, e-counselling websites, mental health information websites, support groups/forums, online mental health screening/assessment, therapist or mental health clinic websites, scholarly articles, blogs, Facebook pages, Wikipedia, videos, and broken links.

For all websites categorized as “websites of Web-based programs” and “websites linking to websites of Web-based programs,” duplicates were removed. All remaining websites entered the second stage of screening and were screened according to the following criteria by the first author: (1) designed for anxiety symptoms (although they did not need to be focused on anxiety only), (2) currently publically accessible on the Internet (via registration, application, General Practitioner

(GP) referral), (3) online component, (4) structured treatment plan (information only), and (5) available in English. Programs were excluded if they were (1) not publically accessible; (2) Web-based counseling only (Skype, email, or instant message contact with a counselor only, with no structured program associated); (3) purely informational (psychoeducation only); or (4) exclusively part of a research study.

## Data Extraction

A data extraction form was created containing 4 main categories and 9 subcategories. [Table 1](#) provides an overview of the items in each main category and subcategories. The 4 main categories are based on the 4 specified study interests (website characteristics, program characteristics, intervention characteristics, and empirical evidence). The 9 subcategories of the 4 main categories were established by incorporating the 12 key facets of a framework designed for evaluating and reporting Internet intervention studies [41]. The extraction was undertaken by the first author in March 2015. Screenshots of all programs and websites were taken in case the program changed during the rating period. To ensure that the programs could be evaluated thoroughly, all program authors were contacted to request free access.

**Table 1.** Data extraction categories and subcategories.

Main Category	Subcategory	Item
Website characteristics	Origin	Country of origin
		Registration (yes/no—if yes, how?)
	Accessibility	Log-in available on website (yes/no)
		Access fee (yes/no—if yes, how much? Free trial available? Refund period? Length of subscription)
		Mobile phone rendering (yes/no)
	Credibility	Advertisements (yes/no—if yes, relevant vs irrelevant)
		Presented contact details (yes/no)
		Specified authorship (yes/no)
		Terms of use specified (yes/no)
		Privacy notice specified (yes/no)
Program characteristics	Intervention focus	Target anxiety issue
		Target audience
	Intervention design	Therapist support (yes/no—if yes, specify)
		Suggested or set treatment length
		Number of modules
	Intervention delivery	Presentation format
Intervention characteristics	Therapeutic approach	CBT <sup>a</sup> ; others (specify)
		Other therapeutic elements
	Intervention features	Worksheets (yes/no—if yes, specify format)
		Mood or symptom monitoring (yes/no)
		Diary (yes/no)
		Forum (yes/no)
Empirical evidence	Empirical evidence for program efficacy	Other features (yes/no—if yes, specify)
		Scrutinized program website for relevant information, contacted the author, and checked the Beacon <sup>b</sup> directory

<sup>a</sup>CBT: cognitive behavioral therapy

<sup>b</sup>Beacon: Australian clinical Web-based platform that describes different Web-based self-help treatment programs [49].

## Program Evaluation

Several validated and widely accepted scales are available to evaluate the methodology of published studies. However, there are currently no validated criteria available for evaluating actual Web-based interventions as found on the web. Renton et al [31] created a program scoring system to evaluate Web-based depression interventions. With permission from the authors, the

scale was adapted to fit the 4 specified study interests. The adapted version consists of 16 yes or no close-ended questions that are outlined in Table 2. Consistent with the scoring system used by Renton et al [31], a score of 1 was awarded if the answer was yes, and a score of 0 if the answer was no or the question could not be evaluated. Scores were converted into percentages, with higher scores indicating a larger number of met criteria on the scale.

**Table 2.** Program evaluation criteria.


Main Category	Question
Website characteristics	1. Was country of origin stated?
	2. Was a unique user name or password provided to users?
	3. Were the names and credentials of authors present?
	4. Were contact details provided?
	5. Were the Terms of Use specified?
	6. Was a Privacy Notice specified?
	7. Was evidence for the program provided to the user (ie, attrition data/success rate/completion rate/# of users in the program/testimonials)?
Program characteristics	8. Were the primary focus/goals/objectives of the intervention stated?
	9. Was the patient group or target mental health issue specified?
	10. Was the number of modules or time to complete each module stated?
	11. Was the intervention tailored to the user or was it generic for all users?
	12. Did the program offer multimedia content delivery (ie, a combination of text, video, graphics, and audio formats)?
	13. Was the program easy to navigate?
Intervention characteristics	14. Was the model of change (ie, type of therapy utilized) defined/stated?
	15. Was information on what is covered in the intervention modules provided (ie, names of modules or a short description)?
Empirical evidence	16. Has the program been empirically validated?

## Results

### Program Selection

A search log outlines the number of hits per search, as well as the number of included and excluded hyperlinks (see [Multimedia Appendix 1](#)). Most program websites were identified when the search terms “online treatment anxiety” (Program websites: n=14, websites with links to program websites: n=2) and “online cognitive behavioural therapy anxiety” (program websites: n=12, websites with links to program websites: n=4) were entered into Google. “Web therapy anxiety” entered in Yahoo did not identify any program websites. All 675 links were

assessed for inclusion. [Figure 1](#) displays the flowchart for the two-stage selection process of included programs and reasons for exclusion. In total, 176 of the 675 (25.3%) assessed hyperlinks led either directly to program websites (133/675, 19.7%) or to websites containing links to program websites (43/675, 6.4%). The first stage of the screening identified 35 potentially eligible program websites that were subsequently assessed for inclusion. Of those, 19 websites and 34 programs met the inclusion criteria. For 5 programs, the authors did not grant access and some aspects of those programs could therefore not be evaluated (Beating the Blues, Changing States, eCentreClinic, FearFighter, Social Anxiety Institute).

**Figure 1.** Flow diagram of program selection.

## Website Characteristics

### Country of Origin


Programs identified in this review originated from 4 different countries. The majority of programs were developed in Australia (16/34, 47%), followed by the UK (9/34, 28%) and the USA (8/34, 24%), and Sweden with one program.

### Accessibility

Out of the 34 programs evaluated, 25 (74%) had a compulsory online registration process to access the program, 5 (15%) required GP/clinician referral, 1 (3%) was accessible through either registration or GP referral, 1 (3%) through application, and 2 (6%) did not require registration to access the program (see Figure 2). Excluding the 2 programs that did not require registration, 29 (85%) had a log-in feature on their website. For

3 programs (9%), no log-in feature was found and it was unclear how users would log in after buying the program. For programs requiring registration, consumers had to enter personal information to set up a profile. Registration allowed tracking and saving of entered information. This was not possible for programs that required no registration.

Over half of programs (24/34, 71%) required an access fee, while 3 were free if signing up for a research trial. Costs varied from £14.99 (Changing States) for 1 module to £197 (FearFighter) for 9 modules and therapist support. Most programs that required a fee had either a free trial period, or a 100% refund period and were either weekly/monthly subscriptions or only valid for 1 to 6 months. Out of the 34 programs, 9 (28%) were accessible in a mobile phone version, while 3 (9%) could not be evaluated in this respect, as the authors did not grant access to the program.

**Figure 2.** Access to evaluated programs.


### Credibility

All programs specified authorship and all programs presented contact details either via a contact form and phone number (9/34, 28%), email address (7/34, 21%), email address and live chat (6/34, 18%), contact form (5/34, 15%), email address and mailing address (3/34, 9%), email address and phone number (2/34, 6%), phone number (1/34, 3%), or email address and contact form (1/34, 3%) (see [Figure 3](#)). Thirty out of 34 (88%) programs specified their terms of use and 32 (94%) had a privacy notice. All programs with a privacy notice also included

information about browser cookies, data collection, and data management. Thirty-one programs (91%) displayed no advertisements. One program's advertisement was deemed relevant (mental health self-help books) and the other two were deemed irrelevant (BBC news link and Google Ads).

### Intervention Program Characteristics

An overview of intervention program characteristics for each program can be found in [Table 3](#) and screenshots of all programs can be found in [Multimedia Appendix 2](#).

**Figure 3.** Methods of contacting the program owner.

**Table 3.** Intervention program characteristics of included Web-based intervention programs for anxiety.

Program (Ref#)	Target Anxiety Issue & Population	Therapist-Assisted	Structure & Length	Presentation Format	Therapeutic Approach	Intervention Features
AI-Therapy (#1)	Social anxiety	No	7 modules (1-2 modules per week)	- Text chapters with figures - Audio features with every chapter - Video features	CBT <sup>a</sup>	- Online worksheets - Online questionnaires - Symptom tracking - Email reminders - Knowledge quizzes - Personalized eBook
Beating the Blues (#2)	Anxiety & depression	No	8 sessions (over 8 weeks)	- Image slides with audio & video - Interactive slides	CBT <sup>a</sup>	- Worksheet printouts - Email reminders
Blues Begone (#3)	Anxiety & depression	No	30 modules (8 weeks)	- Text chapters with figures and images - Audio with every chapter - Cartoon videos	CBT <sup>a</sup>	- Worksheets - Symptom tracking - Diary
Changing States - The Stress and Anxiety Manager (#4)	Anxiety & stress	No	1 module divided in 4 main sections	- Slides with images, text, accompanied by audio - Notes for printing	Hypnotherapy & CBT <sup>a</sup>	- Relaxation technique audio files
CBT 7 Step Self Help Course (#5)	Anxiety, depression, & anger	Option of receiving paid email guidance and personalized formulation	7 modules	- Text chapters with figures - Audio features (need to be purchased separately)	CBT <sup>a</sup>	- PDF worksheet - Wiki
CCBT Limited – FearFighter (#6)	Panic and phobia	Via telephone (if purchased)	9 steps (recommended 9 weeks)	- Video text and image slides	CBT <sup>a</sup>	- Worksheet printouts - Progress monitoring - Emails with further tips at the end of each step
eCentreClinic - Mood Mechanic Course (#7)	Depression, social anxiety, panic attacks, & generalized worry; Australian adults aged 18 to 24	Weekly contact with clinician via email and telephone (depends on trial)	4 lessons (5 weeks)	- Text chapters and images	CBT <sup>a</sup>	- Online questionnaires and worksheets - Symptom tracking - Diary - Knowledge tests
eCouch - Anxiety & Worry Program (#8)	Anxiety & worry; Aged over 16	No	3 main sections (arm chair: 15 sections, tool kit, workbooks)	- Text chapters with figures and animated pictures - Audio features	CBT <sup>a</sup> & IPT <sup>b</sup>	- Online questionnaires and worksheets - Symptom tracking - Diary - Knowledge tests
eCouch - Social Anxiety Program (#9)	Social anxiety; Aged over 16	No	3 main sections (arm chair: 16 sections, tool kit, workbooks)	- Text chapters with figures and animated pictures - Audio features	CBT <sup>a</sup> & IPT <sup>b</sup>	- Online questionnaires and worksheets - Symptom tracking - Diary - Knowledge tests
Learn to Live (#10)	Social anxiety	No	8 lessons (8 weeks recommended)	- Animated slides with audio, images, and text - Videos - Slides require input from users	CBT <sup>a</sup>	- Online and printable worksheets - Forum - Symptom tracking - Online calendar - Questionnaires
Livanda - Free from Anxiety (#11)	Panic disorder, social phobia, & general anxiety	Through messaging system within the program (if paid for)	8-10 sections (12-15 weeks)	- Text chapters and slides - Audio features	CBT <sup>a</sup>	- Online worksheets - Symptom tracking

Program (Ref#)	Target Anxiety Issue & Population	Therapist-Assisted	Structure & Length	Presentation Format	Therapeutic Approach	Intervention Features
Living Life to the Full (#12)	Anxiety, stress, & life skills	User can designate a support practitioner	12 modules	- Text slides with figures - Audio with every slide	CBT <sup>a</sup>	- Alert emails for incomplete modules - Symptom tracking - PDF worksheets - Online books - Online questionnaires
Mental Health Online - Generalised Anxiety Disorder (#13)	Generalized anxiety disorder; Aged over 18	Weekly eTherapist emails, monitor progress, answer questions and provide support via email	12 modules (12 weeks)	- Text chapters with figures - Audio and video features	CBT <sup>a</sup>	- PDFs worksheets - Online worksheets - Symptom tracking - Diary
Mental Health Online - Social Anxiety Disorder (#14)	Social anxiety disorder; Aged over 18	Weekly eTherapist emails, monitor progress, answer questions and provide support via email	12 modules (12 weeks)	- Text chapters with figures - Audio and video features	CBT <sup>a</sup>	- PDF worksheets - Online worksheets - Symptom tracking - Diary
Mental Health Online - Panic Disorder with or without Agoraphobia (#15)	Panic disorder with or without agoraphobia; Aged over 18	Weekly eTherapist emails, monitor progress, answer questions and provide support via email	12 modules (12 weeks)	- Text chapters with figures - Audio and video features	CBT <sup>a</sup>	- PDF worksheets - Online worksheets - Symptom tracking - Diary
Mood Control (#16)	Anxiety & depression	No	12 modules (13 weeks)	- Video for every chapter with an introduction text	CBT <sup>a</sup>	- PDF worksheets - Online questionnaires - Symptom tracking - Forum - Bonus material (sessions for personal development and life change) - Additional worksheets
MoodGym (#17)	Anxiety & depression; Aged over 16	No	5 modules	- Text chapters with images	CBT <sup>a</sup> & IPT <sup>b</sup>	- Quizzes - Worksheets - Downloadable relaxation audio - Symptom tracking
myCompass (#18)	Anxiety, depression, & stress; Aged over 18; Mobile phone function for Australian residents only	No	12 modules (6-8 weeks)	- Text chapters with figures	CBT <sup>a</sup> , IPT <sup>b</sup> , & positive psychology	- PDF worksheets - Online worksheets - Symptom tracking - Diary - Wiki - SMS & email reminders - Real-life experience stories
Online Therapy – Anxiety (#19)	Anxiety; Aged over 18	Live support and email	8 sections (8 weeks)	- Text chapters with figures and images	CBT <sup>a</sup>	- Online worksheets - Online questionnaires - Symptom tracking - Diary - Forum - Chatroom for general help

Program (Ref#)	Target Anxiety Issue & Population	Therapist-Assisted	Structure & Length	Presentation Format	Therapeutic Approach	Intervention Features
Online Therapy - Generalized Anxiety Disorder (#20)	Generalized anxiety disorder; Aged over 18	Live support and email	8 sections (8 weeks)	- Text chapters with figures and images	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- Online worksheets</li> <li>- Online questionnaires</li> <li>- Symptom tracking</li> <li>- Diary</li> <li>- Forum</li> <li>- Chatroom for general help</li> </ul>
Online Therapy - Panic Attacks (#21)	Panic attacks; Aged over 18	Live support and email	8 sections (8 weeks)	- Text chapters with figures and images	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- Online worksheets</li> <li>- Online questionnaires</li> <li>- Symptom tracking</li> <li>- Diary</li> <li>- Forum</li> <li>- Chatroom for general help</li> </ul>
Online Therapy - Agoraphobia (#22)	Agoraphobia; Aged over 18	Live support and email	8 sections (8 weeks)	- Text chapters with figures and images	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- Online worksheets</li> <li>- Online questionnaires</li> <li>- Symptom tracking</li> <li>- Diary</li> <li>- Forum</li> <li>- Chatroom for general help</li> </ul>
Online Therapy - Social Anxiety (#23)	Social anxiety; Aged over 18	Live support and email	8 sections (8 weeks)	- Text chapters with figures and images	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- Online worksheets</li> <li>- Online questionnaires</li> <li>- Symptom tracking</li> <li>- Diary</li> <li>- Forum</li> <li>- Chatroom for general help</li> </ul>
Online Therapy - Speech Anxiety (#24)	Speech anxiety; Aged over 18	Live support and email	8 sections (8 weeks)	- Text chapters with figures and images	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- Online worksheets</li> <li>- Online questionnaires</li> <li>- Symptom tracking</li> <li>- Diary</li> <li>- Forum</li> <li>- Chatroom for general help</li> </ul>
Serenity Program - Anxiety Program (#25)	Stress, generalized anxiety disorder, social anxiety, & panic disorder; Aged over 18	No	9 modules (9 weeks)	<ul style="list-style-type: none"> <li>- Text slides with figures and animated images</li> <li>- Interactive content on slides</li> <li>- Audio feature</li> </ul>	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- PDF worksheets</li> </ul>
Social Anxiety Institute (#26)	Social anxiety	No	25 modules	<ul style="list-style-type: none"> <li>- Audio sessions</li> <li>- Video features</li> <li>- Hand-outs</li> </ul>	CBT <sup>a</sup>	None
This Way Up Clinic – Worry (#27)	Generalized anxiety disorder; Aged over 18	Supervised by clinician	6 modules (8 weeks)	- Comic slides	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- Online questionnaires</li> <li>- Downloadable homework</li> <li>- Symptom tracking</li> <li>- Recovery stories</li> <li>- Online calendar (set up email reminders)</li> <li>- Downloadable extra activities and information</li> </ul>

Program (Ref#)	Target Anxiety Issue & Population	Therapist-Assisted	Structure & Length	Presentation Format	Therapeutic Approach	Intervention Features
This Way Up Clinic -Worry and sadness (#28)	Depression & anxiety; Aged over 18	Supervised by clinician	6 modules (8 weeks)	- Comic slides	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- Online questionnaires</li> <li>- Downloadable homework</li> <li>- Symptom tracking</li> <li>- Recovery stories</li> <li>- Online calendar (set up email reminders)</li> <li>- Downloadable extra activities and information</li> </ul>
This Way Up Clinic – Panic (#29)	Panic/agoraphobia; Aged over 18	Supervised by clinician	6 modules (8 weeks)	- Comic slides	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- Online questionnaires</li> <li>- Downloadable homework</li> <li>- Symptom tracking</li> <li>- Recovery stories</li> <li>- Online calendar (set up email reminders)</li> <li>- Downloadable extra activities and information</li> </ul>
This Way Up Clinic - Shyness (#30)	Social phobia; Aged over 18	Supervised by clinician	6 modules (8 weeks)	- Comic slides	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- Online questionnaires</li> <li>- Downloadable homework</li> <li>- Symptom tracking</li> <li>- Recovery stories</li> <li>- Online calendar (set up email reminders)</li> <li>- Downloadable extra activities and information</li> </ul>
This Way Up Self-help - Shyness (#31)	Social phobia; Aged over 18	No	3 modules (3 weeks)	- Comic slides	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- Online questionnaires</li> <li>- Downloadable homework</li> <li>- Symptom tracking</li> <li>- Recovery stories</li> <li>- Online calendar (set up email reminders)</li> <li>- Downloadable extra activities and information</li> </ul>
This Way Up Self-help - Worry and Sadness (#32)	Depression & anxiety; Aged over 18	No	3 modules (3 weeks)	- Comic slides	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- Online questionnaires</li> <li>- Downloadable homework</li> <li>- Symptom tracking</li> <li>- Recovery stories</li> <li>- Online calendar (set up email reminders)</li> <li>- Downloadable extra activities and information</li> </ul>
This Way Up School -Overcoming Social Anxiety (#33)	Social anxiety; grade 11 and 12 high school	No	6 modules (6 weeks)	- Comic slides	CBT <sup>a</sup>	<ul style="list-style-type: none"> <li>- Online questionnaires</li> <li>- Downloadable homework</li> <li>- Symptom tracking</li> <li>- Recovery stories</li> <li>- Online calendar (set up email reminders)</li> <li>- Downloadable extra activities and information</li> </ul>

Program (Ref#)	Target Anxiety Issue & Population	Therapist-Assisted	Structure & Length	Presentation Format	Therapeutic Approach	Intervention Features
This Way Up School -Anxiety and Depression Prevention for Adolescents (#34)	Anxiety & depression; grade 9 to 11 high school	No	6 modules (6 weeks)	- Comic slides	CBT <sup>a</sup>	- Online questionnaires - Downloadable homework - Symptom tracking - Recovery stories - Online calendar (set up email reminders) - Downloadable extra activities and information

<sup>a</sup>CBT: Cognitive behavioral therapy


<sup>b</sup>IPT: Interpersonal Therapy

### Intervention Focus

Programs were designed for a range of issues including specific anxiety disorders; anxiety combined with depression and stress, or anger; various anxiety disorders combined; or anxiety in general. Figure 4 shows that the majority of programs were designed for social anxiety/phobia (9/34, 28%) or for mixed anxiety and depression (8/34, 24%). The remaining programs focused on anxiety in general (1/34, 3%); multiple anxiety disorders combined (3/34, 3%); anxiety mixed with stress (2/34,

6%); anxiety mixed with depression and stress or anger (2/34, 6%); or other specific anxiety disorders such as generalized anxiety disorder (3/34, 9%), panic attacks with or without agoraphobia (4/34, 12%), agoraphobia (1/34, 3%), and speech anxiety (1/34, 3%). Concerning the target audience, the majority of programs were designed for an adult population (aged over 16 or 18 years) (20/34, 59%), 2 were targeted at teenagers of high school age, 1 specifically for young adults aged 18-24 years, and 11 programs (32%) did not specify an age group, but based on content seemed to be designed for adults.


**Figure 4.** Intervention target of evaluated programs.



### Intervention Design

In total, 17 programs (50%) offered therapist support, either by email, instant messaging, or phone. See Figure 5 for a summary of the different forms of therapist support. Therapist support always required a fee; for the majority of these support programs (10/17, 59%), there was an option of paying only for the

self-guided version or paying extra for support. For one free program (Living Life to the Full), consumers could invite a professional to access their account and provide support within the program (support practitioner). The recommended length of the programs varied from 1 to 15 weeks (mean 8.85 weeks, SD 4.10) and the number of modules offered ranged from 1 to 30 (mean 9.38, SD 5.97).


**Figure 5.** Therapist support offered in evaluated programs.

### ***Modes of Therapy Presentation***

All programs used a combination of different modes of presentation (eg, text, images, audio, video, text entry-fields, and animation). Content was most frequently presented as text chapters with images or diagrams (23/34; 68%). Other identified modes of presentation included animated slides or pictures, comic slides, ebooks, and video sessions. In total, 15 programs (44%) incorporated audio components and 9 included video components (26%).

### ***General Therapeutic Approach***

All 34 programs claimed to be CBT-based and at least one cognitive and behavioral therapeutic element was employed for each program based on the examined module content. Some programs stated that they also incorporated other therapeutic approaches, such as interpersonal therapy (4/34, 13%), hypnotherapy (1/34, 3%), and positive psychology (1/34, 3%) (see Figure 6).

**Figure 6.** Therapeutic approaches used in evaluated programs.

### ***Other Therapeutic Elements***

Other popular therapeutic elements included psychoeducation modules, goal-setting features, features to create case conceptualizations for oneself, skills training exercises, various forms of relaxation exercises, mindfulness-based exercises, experience stories, sleep hygiene, and relapse prevention.

### ***Intervention Features***

All except for one program (33/34, 97%) provided the user with worksheets during the session or homework in PDF or online forms. Mood or symptom monitoring/tracking was part of the majority of programs (29/34, 85%). Most programs allowed the user to see results and access a result history either in a numerical or diagram format. In total, 12/34 programs (35%) offered an online diary and 9/34 programs (27%) incorporated a user forum. The review also revealed a great variety of other program features. One feature was the set-up of email or text message reminders for unfinished or future sessions (AI-Therapy, Beating the Blues, Living Life to the Full, myCompass) and an online treatment calendar to schedule the next session and set up alerts (Learn to Live, This Way Up programs). Other features included bonus material (eg, personal development offered in Mood Control), additional resources (ie, more worksheets to be used between sessions or after the end of treatment, offered in Mood Control, This Way Up), treatment items voted most useful by users, to-do-list maker,

personal note section, awards, commitment checks (Blues Begone), knowledge tests at the beginning and information about medication (AI-Therapy, e-couch), personalized eBooks (AI-Therapy), printable session recap and homework cards in wallet format (Mental Health Online), and a teammate function, which allowed the nomination of friends or family members for optional support (Learn to Live).

### **Empirical Evidence for Program Efficacy**

A summary of the types of published research evaluations for each program and the respective references can be found in [Table 4](#). For 3/34 programs (9%), indirect research evidence was identified. The two e-couch programs are based on the MoodGYM program, for which research evidence is available; however, the e-couch program's efficacy was not specifically empirically evaluated. The AI-Therapy program has only been evaluated for social anxiety in adults who stutter using a pre-post study without a control group. For 17/34 programs (50%), empirical studies evaluating efficacy or effectiveness were found. Studies ranged from case series and small to mid-sized pre-post interventions without comparison groups to controlled and randomized controlled trials (RCT). Both MoodGYM and This Way Up have been evaluated through 9 RCTs each. The efficacy of Beating the Blues was demonstrated by 2 RCTs, FearFighter by 2 RCTs, and myCompass by 1 RCT. For 14/34 programs (41%), no research evidence of the efficacy or effectiveness of the intervention was found.

**Table 4.** Types of research evaluations of included Web-based interventions

Program (Ref#)	Type of Research Evaluation Studies
AI-Therapy (#1)	<ul style="list-style-type: none"> <li>- Pre-post intervention for social anxiety in adults who stutter [42]</li> <li>- Case study [43]</li> <li>- Feasibility &amp; acceptability [44]</li> <li>- 2 RCTs<sup>a</sup>[45, 46]</li> <li>- Cost-effectiveness [47]</li> <li>- Pre-post intervention without comparison group [42, 49]</li> </ul>
Beating the Blues (#2)	<ul style="list-style-type: none"> <li>- Implementation [50- 52]</li> </ul>
Blues Begone (#3)	<ul style="list-style-type: none"> <li>- Pre-post intervention without comparison group [53]</li> </ul>
Changing States - The Stress and Anxiety Manager (#4)	Website: not specified; Beacon <sup>b</sup> : no research evidence
CBT 7 Step Self Help Course (#5)	Website: not specified; Beacon <sup>b</sup> : not reviewed <ul style="list-style-type: none"> <li>- Acceptability study [54]</li> <li>- Pre-post intervention pilot [55]</li> <li>- Case studies without comparison group [56]</li> <li>- Implementation study [57]</li> </ul>
CCBT Limited – FearFighter (#6)	<ul style="list-style-type: none"> <li>- 2 RCTs<sup>a</sup>[58, 59]</li> </ul>
eCentreClinic - Mood Mechanic Course (#7)	Website: nothing for this specific program; Beacon <sup>b</sup> : not reviewed
eCouch - Anxiety & Worry Program (#8, #9)	Adapted from MoodGYM
Learn to Live (#10)	Website: not specified; Beacon <sup>b</sup> : not reviewed
Livanda - Free from Anxiety (#11)	Website: not specified; Beacon <sup>b</sup> : no research evidence
Living Life to the Full (#12)	Website: not specified; Beacon <sup>b</sup> : no research evidence
Mental Health Online - Generalized Anxiety Disorder (#13, #14, #15)	<ul style="list-style-type: none"> <li>- Participant choice trial [60]</li> <li>- Implementation [61]</li> <li>- Predictors of pre-treatment attrition and treatment withdrawal [62]</li> </ul>
Mood Control (#16)	Website: not specified; Beacon <sup>b</sup> : no research evidence
MoodGym (#17)	<ul style="list-style-type: none"> <li>- Nine RCTs<sup>a</sup> [63- 71]</li> <li>- School and class-based trials [72, 73]</li> <li>- Implementation [74]</li> <li>- Program usage analysis [75]</li> <li>- Follow-up outcome analysis [76]</li> <li>- Compliance of community users and predictor of outcomes analysis [77]</li> </ul>
myCompass (#18)	<ul style="list-style-type: none"> <li>- RCT<sup>a</sup>[78, 79]</li> </ul>
Online Therapy (#19, #20, #21, #22, #23, #24)	Website: not specified; Beacon <sup>b</sup> : not reviewed
Serenity Program - Anxiety Program (#25)	<ul style="list-style-type: none"> <li>- Pilot pre-post treatment without comparison group [80]</li> </ul>
Social Anxiety Institute (#26)	Website: not specified; Beacon <sup>b</sup> : not reviewed
This Way Up Clinic (#27, #28, #29, #30) This Way Up Self-help (#31, #32)	Generalized anxiety disorder: <ul style="list-style-type: none"> <li>- 3 RCTs<sup>a</sup>[81- 83] - Implementation study [84]</li> </ul> Panic: - 1 RCT <sup>a</sup> [85] <ul style="list-style-type: none"> <li>- Pre-post intervention trial without comparison group [86] Social phobia:               <ul style="list-style-type: none"> <li>- 5 RCTs<sup>a</sup>[87- 91]</li> <li>- Implementation study [92]</li> </ul> </li> </ul>
This Way Up School (#33, #34) Anxiety	<ul style="list-style-type: none"> <li>- Cost-effectiveness, acceptability, and follow-up analysis [93]</li> </ul>

<sup>a</sup>RCT: randomized controlled trial<sup>b</sup>Beacon: Australian clinical online platform that describes different Web-based self-help treatment programs [35]

Results were examined for programs for which anxiety symptoms were evaluated in RCTs. Beating the Blues was found to lead to a significant reduction of anxiety both at the end of treatment and at 6 months' follow-up compared to treatment as usual [45]. For FearFighter, both the face-to-face and online program group had reduced anxiety at post-treatment and 1-month follow-up [59]. MoodGym was evaluated in adolescents and university students and levels of anxiety were found to be lower in the intervention group compared to the waitlist control group after the intervention [63,66]. In addition, combined face-to-face and online CBT was more effective in treating anxiety symptoms than either face-to-face or online as a standalone [70]. Compared to control subjects, participants in the myCompass intervention had reduced anxiety symptoms after the program and symptom scores remained at near normal levels at 3-month follow-up [78]. For This Way Up, a program targeted at generalized anxiety disorder, the intervention group participants showed significantly reduced symptoms of panic [81] and anxiety [82] at post-treatment compared to the control group. Symptom reduction was the same for technician and clinician-assisted versions of the treatment [83].

## Program Evaluation

Program evaluation scores for each program and each evaluation criteria can be found in [Multimedia Appendix 3](#). Scores for each program ranged from 69% (CBT 7 Step Self Help Course) to 100% (AI-Therapy) with an average score of 81% (SD 7%). Concerning the evaluation criteria, all program websites specified for which patient group or symptoms the program was designed, defined or stated the utilized model of change, presented program author names and credentials, and provided contact details. About half of the program websites had been empirically evaluated (19/34, 55.9%), specified which information was covered in the intervention modules (18/34, 52.9%), and provided evidence for the program to the user (eg, attrition data, success rate, completion rate, number of users in the program, testimonials) (14/34, 41.2%). Only 4 program websites (11.8%) specified whether the intervention was tailored to the user or generic for all users. This question could not be evaluated for one website (Social Anxiety Institute).

## Discussion

### Principal Findings

To our knowledge, this is the first review of publically available Web-based programs for anxiety that showcases what individuals seeking such treatment options might find if they search the Web. The review aimed at providing consumers, practitioners, and researchers with a summary of the availability, characteristics, and efficacy of currently freely available Web-based interventions for anxiety. The review identified a wide variety of programs for anxiety, specific anxiety disorders, or anxiety in combination with stress, depression, or anger with treatments based predominantly on CBT techniques. The majority of websites were found to be credible and accessible. Of the programs reviewed, the majority required that users register and/or pay a program access fee. Half of the programs offered some form of paid therapist or professional support. Programs varied in treatment length and number of modules

and employed a variety of presentation modes. Relatively few were evaluated in terms of efficacy. In particular, this review highlights two key issues: the large number and diversity of program formats and the lack of empirical evidence of efficacy for many of the identified programs. These will be discussed in more detail and results will be compared with a similar review of Web-based depression programs available on the Web [37].

First, the great variety and large number of identified programs for anxiety is noteworthy. Programs differed in their level of support, accessibility, and presentation. A similarly great variability among identified programs was also found for Web-based depression interventions [37]. Concerning accessibility, more than half of the programs required an access fee. Considering the high costs and waiting times for psychotherapy in many countries, paid Web-based programs may provide an affordable alternative. However, programs often could only be purchased for a limited period. Many users may not be able to finish the program in the allotted time, and being able to receive treatment at one's own pace might be an important reason for choosing Web-based treatment over face-to-face therapy.

Overall, most programs used a multimedia presentation for the intervention delivery. With the current rapid pace of advances in technology, more engaging ways of translating therapeutic techniques into interactive techniques could be created for Web-based interventions to distinguish them from traditional self-help material. Increased engagement through interactivity may increase adherence and effectiveness [94,95], especially when considering reports of low utilization and high dropout rates of Web-based interventions [19,96]. As individuals may differ in their preferred style of therapy and time and resources available for treatment, trying different programs and considering the access period is recommended before choosing a program.

The number of identified anxiety programs was similar to the number listed in the Beacon directory [35]. In total, 33 distinct programs were found in the directory. About half of those programs were also identified by this review and some of those identified in this review were not listed on Beacon. It is important to note here that the Beacon website is not updated very often; for example, some reviews of anxiety programs were last updated in 2009. The difference may also be a result of the keywords used and the way search engines are designed and work. Search engines are often referred to as "information gatekeepers," as they are able to include and exclude websites and influence the ranking of websites in the search results [39]. These results suggest that even though a multitude of Web-based programs exists, it may be difficult for interested consumers to identify and compare all options. Having specialized services like the Beacon directory and keeping them up-to-date is therefore important to provide consumers with knowledge about program differences, credibility, and effectiveness. This will in turn help consumers to be able to compare programs and choose the one most suitable for them. A review of Web-based depression interventions identified a similar number ( $n=32$ ) of programs on the Web and 12 of those programs were also included in this review [37]. Those were mostly programs that

offered interventions targeting both anxiety and depression issues.

To ensure that consumers access programs of appropriate quality and safety, national and/or international platforms are needed that provide consumers with reliable guidance on evidence-based and effective Web-based intervention options. For example, the E - Mental Health Strategy for Australia [97] outlines the development of an e-mental health portal that provides reliable information and accessible pathways for consumers and caregivers to navigate and use evidence-based Web-based mental health support. In addition, there is little consistent guidance on necessary quality standards, as well as legal and ethical issues regarding Web-based interventions for professionals. There is, for example, the Suggested Principles for the Online Provision of Mental Health Services by the International Society for Mental Health Online (ISMHO); however, this document mainly addresses online counselling and there are no guidelines specifically addressing Web-based programs. In the context of mental health apps, a review has also highlighted the need for standards and guidelines for developers to follow and frameworks for consumers to assess credibility and legitimacy [32].

Concerning the evidence base of the included programs, all were found to be based on CBT principles. This is consistent with prior reviews, which found that some form of CBT or other behavioral therapy was included in most Web-based interventions [98], as well as in publicly available Web-based intervention programs for depression [37]. In general, research evidence indicates that CBT is an effective treatment for anxiety disorders (eg, [99,100]). This suggests that all reviewed programs were to some extent developed using an evidence-based approach; however, this does not guarantee that the evidence-based approach used is necessarily effective in the program.

In this context, another major finding was that several programs did not provide any research evidence or provided only limited evidence of the efficacy of the treatment. This is similar to findings from the review of Web-based depression interventions, which showed that 63% had not been evaluated using RCTs [37]. This finding is interesting considering the numerous systematic reviews and meta-analyses of Web-based interventions [25,30,101]. However, some may currently be in the process of being evaluated and not yet published. In addition, the absence of evidence of efficacy in terms of RCTs for a particular intervention also does not necessarily mean that the intervention does not work, especially if it is based on evidence-based approaches such as CBT. For treatment efficacy, the predominant model has been “empirically supported treatments” [102]. However, recently it has been proposed that clinical treatment decisions should be based on the best available research evidence, a clinician’s expertise, and patient characteristics [103]. It has also been argued that RCTs evaluating interventions should focus on evaluating intervention principles rather than each actual implementation [104]. However, unlike therapists who require accreditation to practice an evidence-based approach such as CBT, no such accreditations currently exist for Web-based programs. Hence, any Web-based program can claim to be based on CBT, but may not fulfil all

requirements and therefore not work, which is especially problematic for programs requiring an access fee. Therefore, programs should ideally undergo appropriate empirical evaluation before being made available online. The development of an accreditation service for Web-based interventions may help improve this issue and enable consumers to make more informed decisions. It is also important to acknowledge competing interests within the eHealth space. Developers with a commercial focus may not be as concerned about treatment efficacy and researchers developing programs may not have the resources to sustain a publically available program. For programs for which published empirical studies were identified in this review, there was a large variety of study designs and quality of evidence. Only This Way Up programs, MoodGYM, myCompass, Beating the Blues, and FearFighter underwent rigorous evaluation through RCTs.

### Limitations

In regard to this study, a few limitations have to be noted. The representativeness and comprehensiveness of the search and identified programs may be affected by various characteristics of the Web, search engines, and search terms. First, the ranks of websites vary by location on commercial search engines. The search for this review was performed in the UK and it is likely that the same search in another country may have yielded different results. The Web and search engines are also dynamic. Results of search engines vary over time, meaning searches conducted several months before or after the current search could present a different set of programs. In addition, currently existing programs may change or be discontinued and new programs may be released. It is also possible that some individuals may not use the three search engines and would have therefore received different results. However, a considerable strength of this review is that the three most popular search engines were used.

Secondly, the first 25 hyperlinks from the search were included in this review. It is likely that more programs are available, which at the time did not have the page ranking to be identified by the search. This may especially be the case for recently created services [105]. However, it has been suggested that most people rarely consider more than the first 20 links [40]; thus the identified sample of this review is believed to be representative of what an average Internet user might discover when searching the Web for Web-based intervention options. Page ranking is also influenced by various search engine optimization techniques, algorithms of the search engines, as well as cookie settings of the browsers [106], and thereby impacts the results of the search. To combat this, we removed search engine histories and cookies were disabled on all three browsers. While this may not be a complete list of currently available Web-based programs for anxiety, it is a comprehensive snapshot of programs found in March 2015.

Third, the program evaluation scale used was an adapted version of the scoring system used by Renton et al [37]. However, the summative scores do not account for the fact that items within this scale may not be equivalent in terms of importance. Using different weightings based on importance would add great value to the rating. The development of such a scale was considered

beyond the scope of this review; however, it would be important to develop a weighted scale for similar future reviews.

Lastly, it is important to acknowledge that the definition of Web credibility is complex and consists of multiple dimensions [107]. In the case of this paper, only a limited number of credibility dimensions that focused on trustworthiness rather than expertise, were assessed.

## Conclusion

This review found that individuals searching for Web-based intervention programs for anxiety are presented with a large number and variety of potential programs to choose from. For consumers with limited knowledge about intervention quality criteria it may be challenging to choose an appropriate program. With the number of people using the Internet increasing, it is likely that more individuals will search for information about treatments options in general and, specifically, online interventions. It is therefore important for health professionals

working with mental health clients to be aware of the diversity of Web-based interventions and that not all have had their efficacy tested in robust research trials. Directories such as Beacon can assist clinicians, as well as individuals in this task; however, it is important to keep services like these up-to-date. There is a definite need for consistent guidelines and standards on developing and providing Web-based mental health intervention programs for professionals and a platform with reliable up-to-date information for professionals and consumers about the quality and accessibility of Web-based interventions. This review is the first to identify and review Web-based anxiety interventions available on the Web. Therefore, research is needed in reviewing and evaluating Web-based intervention programs for other mental health related issues. There is also a need to develop standardized evaluation scales for publically available Web-based intervention programs to facilitate the rating process and ensure its rigor. For future research, it may also be interesting to explore health professionals' and consumers' experiences and perceptions of those programs.

## Acknowledgments

We would like to thank all the authors of the interventions who provided us with free access to their Web-based interventions and additional information, such as relevant publications.

## Conflicts of Interest

None declared.

## Multimedia Appendix 1

Search log.

[[XLSX File \(Microsoft Excel File\), 490KB - jmir\\_v3i2e14\\_app1.xlsx](#)]

## Multimedia Appendix 2

Screenshots of programs.

[[PDF File \(Adobe PDF File\), 3MB - jmir\\_v3i2e14\\_app2.pdf](#)]

## Multimedia Appendix 3

Program evaluation scores.

[[XLSX File \(Microsoft Excel File\), 15KB - jmir\\_v3i2e14\\_app3.xlsx](#)]

## References

1. Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2005 Jun;62(6):617-627 [[FREE Full text](#)] [doi: [10.1001/archpsyc.62.6.617](#)] [Medline: [15939839](#)]
2. Somers JM, Goldner EM, Waraich P, Hsu L. Prevalence and incidence studies of anxiety disorders: A systematic review of the literature. *Can J Psychiatry* 2006 Feb;51(2):100-113. [Medline: [16989109](#)]
3. Kessler R, Ruscio A, Shear K, Wittchen H. Epidemiology of anxiety disorders. In: Stein MB, Steckler T, editors. *Behavioral Neurobiology of Anxiety and Its Treatment*. Berlin: Springer; 2010:978-983.
4. Wittchen HU, Jacobi F, Rehm J, Gustavsson A, Svensson M, Jönsson B, et al. The size and burden of mental disorders and other disorders of the brain in Europe 2010. *Eur Neuropsychopharmacol* 2011 Sep;21(9):655-679. [doi: [10.1016/j.euroneuro.2011.07.018](#)] [Medline: [21896369](#)]
5. Baxter AJ, Vos T, Scott KM, Ferrari AJ, Whiteford HA. The global burden of anxiety disorders in 2010. *Psychol Med* 2014 Aug;44(11):2363-2374. [doi: [10.1017/S0033291713003243](#)] [Medline: [24451993](#)]

6. Kroenke K, Spitzer RL, Williams JBW, Monahan PO, Löwe B. Anxiety disorders in primary care: Prevalence, impairment, comorbidity, and detection. *Ann Intern Med* 2007 Mar 6;146(5):317-325. [Medline: [17339617](#)]
7. Schonfeld WH, Verboncoeur CJ, Fifer SK, Lipschutz RC, Lubeck DP, Buesching DP. The functioning and well-being of patients with unrecognized anxiety disorders and major depressive disorder. *J Affect Disord* 1997 Apr;43(2):105-119. [Medline: [9165380](#)]
8. Living with Anxiety: Understanding the Role and Impact of Anxiety in Our Lives. London: Mental Health Foundation; 2014. URL: <http://www.mentalhealth.org.uk/publications/living-with-anxiety/> [accessed 2015-11-14] [WebCite Cache ID [6d1eEKytm](#)]
9. Pollard C, Henderson J, Frank M, Margolis RB. Help-seeking patterns of anxiety-disordered individuals in the general population. *J Anxiety Disord* 1989 Jan;3(3):131-138. [doi: [10.1016/0887-6185\(89\)90007-8](#)]
10. Mechanic D. Barriers to help-seeking, detection, and adequate treatment for anxiety and mood disorders: Implications for health care policy. *J Clin Psychiatry* 2007;68 Suppl 2:20-26. [Medline: [17288503](#)]
11. Collins KA, Westra HA, Dozois DJA, Burns DD. Gaps in accessing treatment for anxiety and depression: Challenges for the delivery of care. *Clin Psychol Rev* 2004 Sep;24(5):583-616. [doi: [10.1016/j.cpr.2004.06.001](#)] [Medline: [15325746](#)]
12. Powell J, Clarke A. The WWW of the World Wide Web: Who, What, and Why? *J Med Internet Res* 2002;4(1):e4 [FREE Full text] [doi: [10.2196/jmir.4.1.e4](#)] [Medline: [11956036](#)]
13. Kummervold PE, Gammon D, Bergvik S, Johnsen JK, Hasvold T, Rosenvinge JH. Social support in a wired world: Use of online mental health forums in Norway. *Nord J Psychiatry* 2002;56(1):59-65. [doi: [10.1080/08039480252803945](#)] [Medline: [11869468](#)]
14. Fox S, Rainie L. Vital Decisions: A Pew Internet Health Report.: Pew Research Center; 2002. URL: <http://www.pewinternet.org/2002/05/22/vital-decisions-a-pew-internet-health-report/> [accessed 2015-11-14] [WebCite Cache ID [6d1eWDQMg](#)]
15. Powell J, Clarke A. Internet information-seeking in mental health: Population survey. *Br J Psychiatry* 2006 Sep;189:273-277 [FREE Full text] [doi: [10.1192/bjp.bp.105.017319](#)] [Medline: [16946364](#)]
16. Fox S, Rainie L. The Online Health Care Revolution. 2000. URL: <http://www.pewinternet.org/2000/11/26/the-online-health-care-revolution/> [accessed 2015-11-14] [WebCite Cache ID [6d1ed5Yzt](#)]
17. Barak A, Klein B, Proudfoot JG. Defining internet-supported therapeutic interventions. *Ann Behav Med* 2009 Aug;38(1):4-17. [doi: [10.1007/s12160-009-9130-7](#)] [Medline: [19787305](#)]
18. Griffiths F, Lindenmeyer A, Powell J, Lowe P, Thorogood M. Why are health care interventions delivered over the internet? A systematic review of the published literature. *J Med Internet Res* 2006;8(2):e10 [FREE Full text] [doi: [10.2196/jmir.8.2.e10](#)] [Medline: [16867965](#)]
19. Mohr DC, Burns MN, Schueller SM, Clarke G, Klinkman M. Behavioral intervention technologies: Evidence review and recommendations for future research in mental health. *Gen Hosp Psychiatry* 2013 Aug;35(4):332-338 [FREE Full text] [doi: [10.1016/j.genhosppsych.2013.03.008](#)] [Medline: [23664503](#)]
20. Kaltenthaler E, Brazier J, De NE, Tumor I, Ferriter M, Beverley C, et al. Computerised cognitive behaviour therapy for depression and anxiety update: A systematic review and economic evaluation. *Health Technol Assess* 2006 Sep;10(33):iii, xi-ixiv, 1 [FREE Full text] [Medline: [16959169](#)]
21. Muñoz RF. Using evidence-based internet interventions to reduce health disparities worldwide. *J Med Internet Res* 2010;12(5):e60 [FREE Full text] [doi: [10.2196/jmir.1463](#)] [Medline: [21169162](#)]
22. Griffiths KM, Christensen H. Internet-based mental health programs: A powerful tool in the rural medical kit. *Aust J Rural Health* 2007 Apr;15(2):81-87. [doi: [10.1111/j.1440-1584.2007.00859.x](#)] [Medline: [17441815](#)]
23. Gega L, Marks I, Mataix-Cols D. Computer-aided CBT self-help for anxiety and depressive disorders: Experience of a London clinic and future directions. *J Clin Psychol* 2004 Feb;60(2):147-157. [doi: [10.1002/jclp.10241](#)] [Medline: [14724922](#)]
24. Rüsch N, Angermeyer MC, Corrigan PW. Mental illness stigma: Concepts, consequences, and initiatives to reduce stigma. *Eur Psychiatry* 2005 Dec;20(8):529-539. [doi: [10.1016/j.eurpsy.2005.04.004](#)] [Medline: [16171984](#)]
25. Barak A, Hen L, Boniel-Nissim M, Shapira N. A comprehensive review and a meta-analysis of the effectiveness of internet-based psychotherapeutic interventions. *J Technol Hum Serv* 2008 Jul 03;26(2-4):109-160. [doi: [10.1080/15228830802094429](#)]
26. Andrews G, Cuijpers P, Craske MG, McEvoy P, Titov N. Computer therapy for the anxiety and depressive disorders is effective, acceptable and practical health care: A meta-analysis. *PLoS One* 2010 Oct;5(10):e13196 [FREE Full text] [doi: [10.1371/journal.pone.0013196](#)] [Medline: [20967242](#)]
27. Reger MA, Gahm GA. A meta-analysis of the effects of internet- and computer-based cognitive-behavioral treatments for anxiety. *J Clin Psychol* 2009 Jan;65(1):53-75. [doi: [10.1002/jclp.20536](#)] [Medline: [19051274](#)]
28. Murean V, Montgomery GH, David D. Emotional outcomes and mechanisms of change in online cognitive-behavioral interventions: A quantitative meta-analysis of clinical controlled studies. *J Technol Hum Serv* 2012 Jan;30(1):1-13. [doi: [10.1080/15228835.2011.653290](#)]
29. Cuijpers P, Marks IM, van SA, Cavanagh K, Gega L, Andersson G. Computer-aided psychotherapy for anxiety disorders: A meta-analytic review. *Cogn Behav Ther* 2009;38(2):66-82. [doi: [10.1080/16506070802694776](#)] [Medline: [20183688](#)]

30. Mewton L, Smith J, Rossouw P, Andrews G. Current perspectives on Internet-delivered cognitive behavioral therapy for adults with anxiety and related disorders. *Psychol Res Behav Manag* 2014;7:37-46 [[FREE Full text](#)] [doi: [10.2147/PRBM.S40879](#)] [Medline: [24511246](#)]
31. Harrison AM, Goozee R. Psych-related iPhone apps. *J Ment Health* 2014 Feb;23(1):48-50. [doi: [10.3109/09638237.2013.869575](#)] [Medline: [24484194](#)]
32. Shen N, Levitan M, Johnson A, Bender JL, Hamilton-Page M, Jadad Alejandro Alex R, et al. Finding a depression app: A review and content analysis of the depression app marketplace. *JMIR Mhealth Uhealth* 2015 Feb;3(1):e16 [[FREE Full text](#)] [doi: [10.2196/mhealth.3713](#)] [Medline: [25689790](#)]
33. Heinlen K, Welfel E. The nature, scope, and ethics of psychologists' e-therapy web sites: What consumers find when surfing the Web. *Psychother Theory Res Pract Train* 2003;40(1):112-124. [doi: [10.1037/0033-3204.40.1.112](#)]
34. Recupero PR, Rainey SE. Characteristics of e-therapy web sites. *J Clin Psychiatry* 2006 Sep;67(9):1435-1440. [Medline: [17017831](#)]
35. Beacon 2.: Beacon 2015 URL: <https://beacon.anu.edu.au/> [accessed 2015-04-24] [[WebCite Cache ID 6Y1bEAONK](#)]
36. Andrews G, Titov N. Is internet treatment for depressive and anxiety disorders ready for prime time? *Med J Aust* 2010 Jun 7;192(11 Suppl):S45-S47. [Medline: [20528709](#)]
37. Renton T, Tang H, Ennis N, Cusimano MD, Bhalerao S, Schweizer TA, et al. Web-based intervention programs for depression: A scoping review and evaluation. *J Med Internet Res* 2014;16(9):e209 [[FREE Full text](#)] [doi: [10.2196/jmir.3147](#)] [Medline: [25249003](#)]
38. eBizMBA I. Top 15 Most Popular Search Engines. 2015 URL: <http://www.ebizmba.com/articles/search-engines> [accessed 2015-04-23] [[WebCite Cache ID 6Y1betZk5](#)]
39. Levene M. An Introduction to Search Engines and Web Navigation. Hoboken, NJ: John Wiley & Sons, Inc; 2010.
40. Eysenbach G, Powell J, Kuss O, Sa E. Empirical studies assessing the quality of health information for consumers on the world wide web: A systematic review. *JAMA* 2002;287(20):2691-2700. [Medline: [12020305](#)]
41. Proudfoot J, Klein B, Barak A, Carlbring P, Cuijpers P, Lange A, et al. Establishing guidelines for executing and reporting Internet intervention research. *Cogn Behav Ther* 2011;40(2):82-97. [doi: [10.1080/16506073.2011.573807](#)] [Medline: [25155812](#)]
42. Helgadóttir FD, Menzies RG, Onslow M, Packman A, O'Brian S. A standalone Internet cognitive behavior therapy treatment for social anxiety in adults who stutter: CBTpsych. *J Fluency Disord* 2014 Sep;41:47-54. [doi: [10.1016/j.jfludis.2014.04.001](#)] [Medline: [25173456](#)]
43. Helgadóttir FD, Menzies RG, Onslow M, Packman A, O'Brian S. Online CBT II: A Phase I trial of a standalone, online CBT treatment program for social anxiety in stuttering. *Behav Chang* 2009 Dec;26(4):254-270. [doi: [10.1375/bech.26.4.254](#)]
44. Proudfoot J, Swain S, Widmer S, Watkins E, Goldberg D, Marks I, et al. The development and beta-test of a computer-therapy program for anxiety and depression: Hurdles and lessons. *Comput Human Behav* 2003 May;19(3):277-289. [doi: [10.1016/S0747-5632\(02\)00062-6](#)]
45. Proudfoot J, Goldberg D, Mann A, Everitt B, Marks I, Gray JA. Computerized, interactive, multimedia cognitive-behavioural program for anxiety and depression in general practice. *Psychol Med* 2003 Feb;33(2):217-227. [Medline: [12622301](#)]
46. Proudfoot J, Ryden C, Everitt B, Shapiro DA, Goldberg D, Mann A, et al. Clinical efficacy of computerised cognitive-behavioural therapy for anxiety and depression in primary care: Randomised controlled trial. *Br J Psychiatry* 2004 Jul;185:46-54 [[FREE Full text](#)] [Medline: [15231555](#)]
47. McCrone P, Knapp M, Proudfoot J, Ryden C, Cavanagh K, Shapiro DA, et al. Cost-effectiveness of computerised cognitive-behavioural therapy for anxiety and depression in primary care: Randomised controlled trial. *Br J Psychiatry* 2004 Jul;185:55-62 [[FREE Full text](#)] [Medline: [15231556](#)]
48. Cavanagh K, Shapiro DA, Van Den Berg S, Swain S, Barkham M, Proudfoot J. The effectiveness of computerized cognitive behavioural therapy in routine care. *Br J Clin Psychol* 2006 Nov;45(Pt 4):499-514. [doi: [10.1348/014466505X84782](#)] [Medline: [17076960](#)]
49. Mitchell N, Dunn K. Pragmatic evaluation of the viability of CCBT self-help for depression in higher education. *Couns Psychother Res* 2007 Sep;7(3):144-150. [doi: [10.1080/14733140701565987](#)]
50. Fox E, Acton T, Wilding B, Corcoran S. Service development report: An assistant psychologist's perspective on the use of computerised cognitive behavioural therapy in a GP practice in Barnet. *Qual Prim Care* 2004;12:161-165.
51. van den Berg S, Shapiro DA, Bickerstaffe D, Cavanagh K. Computerized cognitive-behaviour therapy for anxiety and depression: A practical solution to the shortage of trained therapists. *J Psychiatr Ment Health Nurs* 2004 Oct;11(5):508-513. [doi: [10.1111/j.1365-2850.2004.00745.x](#)] [Medline: [15450016](#)]
52. Learmonth D, Rai S. Taking computerized CBT beyond primary care. *Br J Clin Psychol* 2008 Mar;47(Pt 1):111-118. [doi: [10.1348/014466507X248599](#)] [Medline: [17939879](#)]
53. Purves DG, Bennett M, Wellman N. An open trial in the NHS of Blues Begone: A new home based computerized CBT program. *Behav Cogn Psychother* 2009 Oct;37(5):541-551. [doi: [10.1017/S1352465809990282](#)] [Medline: [19703330](#)]
54. MacGregor AD, Hayward L, Peck DF, Wilkes P. Empirically grounded clinical interventions clients' and referrers' perceptions of computer-guided CBT (FearFighter). *Behav Cogn Psychother* 2009 Jan;37(1):1-9. [doi: [10.1017/S135246580800492X](#)] [Medline: [19364403](#)]

55. Kenwright M, Marks I, Marks IM, Gega L, Mataix-Cols D. Computer-aided self-help for phobia/panic via internet at home: A pilot study. *Br J Psychiatry* 2004 May;184:448-449 [[FREE Full text](#)] [Medline: [15123511](#)]
56. Gega L, Marks I, Mataix-Cols D. Computer-aided CBT self-help for anxiety and depressive disorders: Experience of a London clinic and future directions. *J Clin Psychol* 2004 Feb;60(2):147-157. [doi: [10.1002/jclp.10241](#)] [Medline: [14724922](#)]
57. Hayward L, MacGregor AD, Peck DF, Wilkes P. The feasibility and effectiveness of computer-guided CBT (FearFighter) in a rural area. *Behav Cogn Psychother* 2007 Apr 16;35(04):409. [doi: [10.1017/S1352465807003670](#)]
58. Schneider AJ, Mataix-Cols D, Marks IM, Bachofen M. Internet-guided self-help with or without exposure therapy for phobic and panic disorders. *Psychother Psychosom* 2005;74(3):154-164. [doi: [10.1159/000084000](#)] [Medline: [15832066](#)]
59. Marks IM, Kenwright M, McDonough M, Whittaker M, Mataix-Cols D. Saving clinicians' time by delegating routine aspects of therapy to a computer: A randomized controlled trial in phobia/panic disorder. *Psychol Med* 2004 Jan;34(1):9-17. [Medline: [14971623](#)]
60. Klein B, Meyer D, Austin DW, Kyrios M. Anxiety online: A virtual clinic: Preliminary outcomes following completion of five fully automated treatment programs for anxiety disorders and symptoms. *J Med Internet Res* 2011;13(4):e89 [[FREE Full text](#)] [doi: [10.2196/jmir.1918](#)] [Medline: [22057287](#)]
61. Al-Asadi AM, Klein B, Meyer D. Posttreatment attrition and its predictors, attrition bias, and treatment efficacy of the anxiety online programs. *J Med Internet Res* 2014;16(10):e232 [[FREE Full text](#)] [doi: [10.2196/jmir.3513](#)] [Medline: [25316533](#)]
62. Al-Asadi AM, Klein B, Meyer D. Pretreatment attrition and formal withdrawal during treatment and their predictors: An exploratory study of the anxiety online data. *J Med Internet Res* 2014;16(6):e152 [[FREE Full text](#)] [doi: [10.2196/jmir.2989](#)] [Medline: [24938311](#)]
63. Caelear AL, Christensen H, Mackinnon A, Griffiths KM, O'Kearney R. The YouthMood Project: A cluster randomized controlled trial of an online cognitive behavioral program with adolescents. *J Consult Clin Psychol* 2009 Dec;77(6):1021-1032. [doi: [10.1037/a0017391](#)] [Medline: [19968379](#)]
64. Christensen H, Griffiths KM, Jorm AF. Delivering interventions for depression by using the internet: Randomised controlled trial. *BMJ* 2004 Jan 31;328(7434):265 [[FREE Full text](#)] [doi: [10.1136/bmj.37945.566632.EE](#)] [Medline: [14742346](#)]
65. Christensen H, Griffiths KM, Mackinnon AJ, Brittliffe K. Online randomized controlled trial of brief and full cognitive behaviour therapy for depression. *Psychol Med* 2006 Dec;36(12):1737-1746. [doi: [10.1017/S0033291706008695](#)] [Medline: [16938144](#)]
66. Ellis L, Campbell A, Sethi S, O'Dea B. Comparative randomized trial of an online cognitive-behavioural therapy program and an online support group for depression and anxiety. *J Cyber Ther Rehabil* 2011;4(4):461-467.
67. Farrer L, Christensen H, Griffiths KM, Mackinnon A. Internet-based CBT for depression with and without telephone tracking in a national helpline: Randomised controlled trial. *PLoS One* 2011 Nov;6(11):e28099 [[FREE Full text](#)] [doi: [10.1371/journal.pone.0028099](#)] [Medline: [22140514](#)]
68. Hickie IB, Davenport TA, Luscombe GM, Moore M, Griffiths KM, Christensen H. Practitioner-supported delivery of internet-based cognitive behaviour therapy: Evaluation of the feasibility of conducting a cluster randomised trial. *Med J Aust* 2010 Jun 7;192(11 Suppl):S31-S35. [Medline: [20528705](#)]
69. Griffiths KM, Christensen H, Jorm AF, Evans K, Groves C. Effect of web-based depression literacy and cognitive-behavioural therapy interventions on stigmatising attitudes to depression: Randomised controlled trial. *Br J Psychiatry* 2004 Oct;185:342-349 [[FREE Full text](#)] [doi: [10.1192/bjp.185.4.342](#)] [Medline: [15458995](#)]
70. Sethi S, Campbell AJ, Ellis LA. The use of computerized self-help packages to treat adolescent depression and anxiety. *J Technol Hum Serv* 2010 Aug 31;28(3):144-160. [doi: [10.1080/15228835.2010.508317](#)]
71. Lintvedt OK, Griffiths KM, Sørensen K, Østvik AR, Wang CEA, Eisemann M, et al. Evaluating the effectiveness and efficacy of unguided internet-based self-help intervention for the prevention of depression: A randomized controlled trial. *Clin Psychol Psychother* 2013;20(1):10-27. [doi: [10.1002/cpp.770](#)] [Medline: [21887811](#)]
72. O'Kearney R, Gibson M, Christensen H, Griffiths KM. Effects of a cognitive-behavioural internet program on depression, vulnerability to depression and stigma in adolescent males: A school-based controlled trial. *Cogn Behav Ther* 2006;35(1):43-54. [doi: [10.1080/16506070500303456](#)] [Medline: [16500776](#)]
73. O'Kearney R, Kang K, Christensen H, Griffiths K. A controlled trial of a school-based Internet program for reducing depressive symptoms in adolescent girls. *Depress Anxiety* 2009;26(1):65-72. [doi: [10.1002/da.20507](#)] [Medline: [18828141](#)]
74. Christensen H, Griffiths KM, Korten AE, Brittliffe K, Groves C. A comparison of changes in anxiety and depression symptoms of spontaneous users and trial participants of a cognitive behavior therapy website. *J Med Internet Res* 2004 Dec 22;6(4):e46 [[FREE Full text](#)] [doi: [10.2196/jmir.6.4.e46](#)] [Medline: [15631970](#)]
75. Christensen H, Griffiths KM, Korten A. Web-based cognitive behavior therapy: Analysis of site usage and changes in depression and anxiety scores. *J Med Internet Res* 2002;4(1):e3 [[FREE Full text](#)] [doi: [10.2196/jmir.4.1.e3](#)] [Medline: [11956035](#)]
76. Mackinnon A, Griffiths KM, Christensen H. Comparative randomised trial of online cognitive-behavioural therapy and an information website for depression: 12-month outcomes. *Br J Psychiatry* 2008 Feb;192(2):130-134 [[FREE Full text](#)] [doi: [10.1192/bjp.bp.106.032078](#)] [Medline: [18245031](#)]

77. Christensen H, Griffiths K, Groves C, Korten A. Free range users and one hit wonders: community users of an Internet-based cognitive behaviour therapy program. *Aust N Z J Psychiatry* 2006;40(1):59-62. [doi: [10.1111/j.1440-1614.2006.01743.x](https://doi.org/10.1111/j.1440-1614.2006.01743.x)] [Medline: [16403040](#)]
78. Proudfoot J, Clarke J, Birch M, Whitton AE, Parker G, Manicavasagar V, et al. Impact of a mobile phone and web program on symptom and functional outcomes for people with mild-to-moderate depression, anxiety and stress: A randomised controlled trial. *BMC Psychiatry* 2013;13:312 [FREE Full text] [doi: [10.1186/1471-244X-13-312](https://doi.org/10.1186/1471-244X-13-312)] [Medline: [24237617](#)]
79. Clarke J, Proudfoot J, Birch M, Whitton AE, Parker G, Manicavasagar V, et al. Effects of mental health self-efficacy on outcomes of a mobile phone and web intervention for mild-to-moderate depression, anxiety and stress: Secondary analysis of a randomised controlled trial. *BMC Psychiatry* 2014;14:272 [FREE Full text] [doi: [10.1186/s12888-014-0272-1](https://doi.org/10.1186/s12888-014-0272-1)] [Medline: [25252853](#)]
80. Slegg G, Cottrell S, Nicholas H, Messenger L. A guided online anxiety self-help programme to aid a return to work: A pilot study. *J Occup Psychol Employ Disabil* 2009;11(2):63-77.
81. Titov N, Andrews G, Robinson E, Schwencke G, Johnston L, Solley K, et al. Clinician-assisted Internet-based treatment is effective for generalized anxiety disorder: Randomized controlled trial. *Aust NZ J Psychiatry* 2009 Jan;43(10):905-912. [doi: [10.1080/00048670903179269](https://doi.org/10.1080/00048670903179269)]
82. Titov N, Andrews G, Johnston L, Robinson E, Spence J. Transdiagnostic Internet treatment for anxiety disorders: A randomized controlled trial. *Behav Res Ther* 2010 Sep;48(9):890-899. [doi: [10.1016/j.brat.2010.05.014](https://doi.org/10.1016/j.brat.2010.05.014)] [Medline: [20561606](#)]
83. Robinson E, Titov N, Andrews G, McIntyre K, Schwencke G, Solley K. Internet treatment for generalized anxiety disorder: A randomized controlled trial comparing clinician vs. technician assistance. *PLoS One* 2010;5(6):e10942 [FREE Full text] [doi: [10.1371/journal.pone.0010942](https://doi.org/10.1371/journal.pone.0010942)] [Medline: [20532167](#)]
84. Mewton L, Wong N, Andrews G. The effectiveness of internet cognitive behavioural therapy for generalized anxiety disorder in clinical practice. *Depress Anxiety* 2012 Oct;29(10):843-849. [doi: [10.1002/da.21995](https://doi.org/10.1002/da.21995)] [Medline: [22949296](#)]
85. Wims E, Titov N, Andrews G, Choi I. Clinician-assisted Internet-based treatment is effective for panic: A randomized controlled trial. *Aust NZ J Psychiatry* 2010 Jul;44(7):599-607. [doi: [10.3109/00048671003614171](https://doi.org/10.3109/00048671003614171)] [Medline: [20560847](#)]
86. Wims E, Titov N, Andrews G. The Climate Panic program: An open trial of Internet-based treatment for panic disorder. *EJAP* 2008 Dec 23;4(2). [doi: [10.7790/ejap.v4i2.134](https://doi.org/10.7790/ejap.v4i2.134)]
87. Titov N, Andrews G, Schwencke G. Shyness 2: Treating social phobia online: Replication and extension. *Aust NZ J Psychiatry* 2008 Jul;42(7):595-605. [doi: [10.1080/00048670802119820](https://doi.org/10.1080/00048670802119820)] [Medline: [18612863](#)]
88. Titov N, Andrews G, Schwencke G, Solley K, Johnston L, Robinson E. An RCT comparing effect of two types of support on severity of symptoms for people completing Internet-based cognitive behaviour therapy for social phobia. *Aust NZ J Psychiatry* 2009 Jan;43(10):920-926. [doi: [10.1080/00048670903179228](https://doi.org/10.1080/00048670903179228)]
89. Titov N, Andrews G, Choi I, Schwencke G, Johnston L. Randomized controlled trial of web-based treatment of social phobia without clinician guidance. *Aust NZ J Psychiatry* 2009 Jan;43(10):913-919. [doi: [10.1080/00048670903179160](https://doi.org/10.1080/00048670903179160)]
90. Titov N, Andrews G, Schwencke G, Drobny J, Einstein D. Shyness 1: Distance treatment of social phobia over the Internet. *Aust NZ J Psychiatry* 2008 Jul;42(7):585-594. [doi: [10.1080/00048670802119762](https://doi.org/10.1080/00048670802119762)] [Medline: [18612862](#)]
91. Titov N, Andrews G, Choi I, Schwencke G, Mahoney A. Shyness 3: Randomized controlled trial of guided versus unguided Internet-based CBT for social phobia. *Aust NZ J Psychiatry* 2008 Dec;42(12):1030-1040. [doi: [10.1080/00048670802512107](https://doi.org/10.1080/00048670802512107)] [Medline: [19016091](#)]
92. Aydos L, Titov N, Andrews G. Shyness 5: The clinical effectiveness of Internet-based clinician-assisted treatment of social phobia. *Australas Psychiatry* 2009 Dec;17(6):488-492. [doi: [10.1080/10398560903284943](https://doi.org/10.1080/10398560903284943)] [Medline: [20001373](#)]
93. Titov N, Andrews G, Johnston L, Schwencke G, Choi I. Shyness programme: Longer term benefits, cost-effectiveness, and acceptability. *Aust NZ J Psychiatry* 2009 Jan;43(1):36-44. [doi: [10.1080/00048670802534424](https://doi.org/10.1080/00048670802534424)] [Medline: [19085526](#)]
94. Doherty G, Coyle D, Sharry J. Engagement with online mental health interventions: An exploratory clinical study of a treatment for depression. 2012 Presented at: SIGCHI Conference on Human Factors in Computing Systems CHI '12; 2012; New York, USA. [doi: [10.1145/2207676.2208602](https://doi.org/10.1145/2207676.2208602)]
95. Ritterband LM, Cox DJ, Gordon TL, Borowitz SM, Kovatchev BP, Walker LS, et al. Examining the added value of audio, graphics, and interactivity in an Internet intervention for pediatric encephalitis. *Children's Health Care* 2006 Mar;35(1):47-59. [doi: [10.1207/s15326888chc3501\\_5](https://doi.org/10.1207/s15326888chc3501_5)]
96. Christensen H, Griffiths KM, Farrer L. Adherence in internet interventions for anxiety and depression. *J Med Internet Res* 2009;11(2):e13 [FREE Full text] [doi: [10.2196/jmir.1194](https://doi.org/10.2196/jmir.1194)] [Medline: [19403466](#)]
97. Australian Government Department of Health and Ageing. E-Mental Health Strategy for Australia. 2012 URL: [http://www.health.gov.au/internet/main/publishing.nsf/Content/7C7B0BFEB985D0EBCA257BF0001BB0A6/\\$File/emstrat.pdf](http://www.health.gov.au/internet/main/publishing.nsf/Content/7C7B0BFEB985D0EBCA257BF0001BB0A6/$File/emstrat.pdf) [accessed 2015-11-14] [WebCite Cache ID 6d1e2IXTc]
98. Siemer CP, Fogel J, Van Voorhees BW. Telemental health and web-based applications in children and adolescents. *Child Adolesc Psychiatr Clin N Am* 2011 Jan;20(1):135-153 [FREE Full text] [doi: [10.1016/j.chc.2010.08.012](https://doi.org/10.1016/j.chc.2010.08.012)] [Medline: [21092918](#)]
99. Hofmann SG, Smits JAJ. Cognitive-behavioral therapy for adult anxiety disorders: A meta-analysis of randomized placebo-controlled trials. *J Clin Psychiatry* 2008 Apr;69(4):621-632 [FREE Full text] [Medline: [18363421](#)]

100. Stewart RE, Chambless DL. Cognitive-behavioral therapy for adult anxiety disorders in clinical practice: A meta-analysis of effectiveness studies. *J Consult Clin Psychol* 2009 Aug;77(4):595-606. [doi: [10.1037/a0016032](https://doi.org/10.1037/a0016032)] [Medline: [19634954](https://pubmed.ncbi.nlm.nih.gov/19634954/)]
101. Spek V, Cuijpers P, Nyklíček I, Riper H, Keyzer J, Pop V. Internet-based cognitive behaviour therapy for symptoms of depression and anxiety: A meta-analysis. *Psychol Med* 2007 Mar;37(3):319-328. [doi: [10.1017/S0033291706008944](https://doi.org/10.1017/S0033291706008944)] [Medline: [17112400](https://pubmed.ncbi.nlm.nih.gov/17112400/)]
102. Chambless DL, Hollon SD. Defining empirically supported therapies. *J Consult Clin Psychol* 1998 Feb;66(1):7-18. [Medline: [9489259](https://pubmed.ncbi.nlm.nih.gov/9489259/)]
103. Tolin DF, McKay D, Forman EM, Klonsky ED, Thombs BD. Empirically supported treatment: Recommendations for a new model. *Clin Psychol Sci Pract* 2015 Nov 09;22(4):317-338. [doi: [10.1111/cpsp.12122](https://doi.org/10.1111/cpsp.12122)]
104. Mohr DC, Schueller SM, Riley WT, Brown CH, Cuijpers P, Duan N, et al. Trials of intervention principles: Evaluation methods for evolving behavioral intervention technologies. *J Med Internet Res* 2015;17(7):e166 [FREE Full text] [doi: [10.2196/jmir.4391](https://doi.org/10.2196/jmir.4391)] [Medline: [26155878](https://pubmed.ncbi.nlm.nih.gov/26155878/)]
105. Cho J, Roy S. Impact of search engines on page popularity. 2004 Presented at: 13th Conference on World Wide Web - WWW; 2004; New York, USA p. 04. [doi: [10.1145/988672.988676](https://doi.org/10.1145/988672.988676)]
106. Evans MP. Analysing Google rankings through search engine optimization data. *Internet Res* 2007 Feb 06;17(1):21-37. [doi: [10.1108/10662240710730470](https://doi.org/10.1108/10662240710730470)]
107. Shah A, Ravana S, Hamid S, Ismail M. Information Research. 2015. Web Credibility Assessment: Affecting Factors and Assessment Techniques URL: <http://www.informationr.net/ir/20-1/paper663.html> [accessed 2015-03-15] [WebCite Cache ID 6X3QQGnRu]

## Abbreviations

**CBT:** Cognitive behavioral therapy  
**IPT:** Interpersonal Therapy  
**RCT:** Randomized controlled trial  
**GP:** General Practitioner

*Edited by J Torous; submitted 18.11.15; peer-reviewed by S Schueller, N Shen, J Bisslerbe; comments to author 11.12.15; revised version received 14.01.16; accepted 01.02.16; published 01.06.16*

*Please cite as:*

Ashford MT, Olander EK, Ayers S

Finding Web-Based Anxiety Interventions on the World Wide Web: A Scoping Review

*JMIR Ment Health* 2016;3(2):e14

URL: <http://mental.jmir.org/2016/2/e14/>

doi: [10.2196/mental.5349](https://doi.org/10.2196/mental.5349)

PMID: [27251763](https://pubmed.ncbi.nlm.nih.gov/27251763/)

©Miriam Thiel Ashford, Ellinor K Olander, Susan Ayers. Originally published in JMIR Mental Health (<http://mental.jmir.org>), 01.06.2016. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Mental Health, is properly cited. The complete bibliographic information, a link to the original publication on <http://mental.jmir.org>, as well as this copyright and license information must be included.