



City Research Online

City, University of London Institutional Repository

Citation: Chuang, J. & Tan, K. M. (2016). Canonical bases for Fock spaces and tensor products. *Advances in Mathematics*, 302, pp. 159-189. doi: 10.1016/j.aim.2016.07.008

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/15164/>

Link to published version: <https://doi.org/10.1016/j.aim.2016.07.008>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

CANONICAL BASES FOR FOCK SPACES AND TENSOR PRODUCTS

JOSEPH CHUANG AND KAI MENG TAN

ABSTRACT. We relate the canonical basis of the Fock space representation of the quantum affine algebra $U_q(\widehat{\mathfrak{gl}}_n)$, as defined by Leclerc and Thibon [LT], to the canonical basis of its restriction to $U_q(\mathfrak{sl}_n)$, regarded as a based module in the sense of Lusztig. More generally we consider the restriction to any Levi subalgebra. We deduce results on decomposition numbers and branching coefficients of Schur algebras over fields of positive characteristic, generalising those of Kleshchev [Kl] and of Tan and Teo [TT].

1. INTRODUCTION

The complete determination of the decomposition numbers of the symmetric groups and Schur algebras in positive characteristic p is a well-known and longstanding open problem, for which a complete solution does not seem to be forthcoming. Related to these decomposition numbers are the q -decomposition numbers arising from the canonical basis for the Fock space representation of the quantum affine algebra $U_q(\widehat{\mathfrak{gl}}_n)$. These q -decomposition numbers, as conjectured by Leclerc and Thibon [LT] and shown by Varagnolo and Vasserot [VV], are polynomials in q with nonnegative integer coefficients and when evaluated at $q = 1$ give the corresponding decomposition numbers for the v -Schur algebra in characteristic zero where v is a primitive n -th root of unity. As shown by James [Jam], when $n = p$, the decomposition matrix for the Schur algebra can be obtained by postmultiplying the decomposition matrix of the v -Schur algebra in characteristic zero by an adjustment matrix which has nonnegative integer entries and is unitriangular when the indexing set is suitably ordered. As such, the q -decomposition numbers provide a first approximation to the decomposition numbers of the Schur algebras.

James's Conjecture asserts that this first approximation is in fact an equality whenever the indexing partitions have p -weight less than p . Even though the conjecture is now known to be false in general [W], it has been proved in some cases, such as in Rouquier blocks [CT] and in blocks with p -weight less than 5 [F1, F2]. It has also been shown that the first

2010 *Mathematics Subject Classification*. 17B37, 20G43.

Supported by EPSRC grant GR/T00917 and Singapore MOE AcRF R-146-000-172-112.

approximation is in fact an equality in many cases irrespective of the p -weight of the indexing partitions; see for example [T, TT].

In [Kl], Kleshchev introduces the combinatorics of sign sequences, and uses it to describe the decomposition number $d_{\lambda\mu}$ when the partition λ is obtained from μ by moving one node. Subsequently, the present authors and Miyauchi [CMT2] provide closed formulas for the corresponding q -decomposition number $d_{\lambda\mu}(q)$ using the same combinatorics. More recently, the second author and Teo [TT] provide closed formulas for $d_{\lambda\mu}(q)$ when λ is obtained from μ by moving any number of nodes as long as all of them have the same n -residue, and show that, when $n = p$, $d_{\lambda\mu}(1) = d_{\lambda\mu}$. The astute reader of [TT] who is familiar with the work of Frenkel and Khovanov in [FK] will be struck by the uncanny similarity between the last closed formulas and those describing the canonical bases of tensor powers $V_2^{\otimes d}$ of the natural two-dimensional representation V_2 of the quantum enveloping algebra $U_q(\mathfrak{sl}_2)$, although the former is formulated using the combinatorics of sign sequences while the latter is described by graphical calculus. It is natural to attempt to find out the exact relationship between the latter canonical bases with that of the Fock space representation. This is the main motivation of our work appearing in this paper.

We briefly describe our results here. For our purposes it suffices to consider the subalgebra $\mathbf{U} = U_q(\widehat{\mathfrak{sl}}_n)$ of $U_q(\widehat{\mathfrak{gl}}_n)$. Let $\mathbf{n} = (n_1, \dots, n_r)$ be a tuple of positive integers such that $n_1 + \dots + n_r = n$, and let $\mathbf{U}(\mathbf{n})$ denote the subalgebra of \mathbf{U} isomorphic to $U_q(\mathfrak{sl}_{n_1}) \otimes \dots \otimes U_q(\mathfrak{sl}_{n_r})$. We show that the Fock space representation \mathcal{F}_s indexed by an integer s (see subsection 2.3 for formal definition), when restricted from \mathbf{U} to $\mathbf{U}(\mathbf{n})$, has a natural decomposition, which corresponds to partitioning the standard basis of \mathcal{F}_s into subsets in a certain way. Each summand $\mathcal{F}_{\mathbf{t}}$ of the restriction is isomorphic to a tensor product of irreducible $\mathbf{U}(\mathbf{n})$ -modules and hence, as a based module in the sense of Lusztig [Lu], is equipped with a canonical basis. We then show that a subset of the canonical basis of \mathcal{F}_s maps to this basis of $\mathcal{F}_{\mathbf{t}}$ under the natural projection map $\mathcal{F}_s \rightarrow \mathcal{F}_{\mathbf{t}}$ (Theorem 5.2). The difficulty in directly relating the canonical bases of \mathcal{F}_s and $\mathcal{F}_{\mathbf{t}}$ stems from the differing nature of the corresponding ‘bar involutions’ that fix basis vectors: in the Fock space, the involution is given by reversing q -wedges [LT], whereas for tensor products of based modules, it is defined through quasi- R -matrices [Lu].

Subsequently, the results in [FK] can be exploited to provide more information when $n_j = 2$ for some j . In particular, this establishes the exact relationship between the canonical basis of the Fock space and that of $V_2^{\otimes d}$. We also obtain closed formulas for some of the branching coefficients for \mathcal{F}_s .

We then turn our attention to the Schur algebras. Assuming the results of Kleshchev [Kl] describing the branching coefficients $[\text{Res}(L(\mu)) : L(\lambda)]$ when

λ is obtained from μ by removing a normal node, we obtain closed formulas for the decomposition numbers $d_{\lambda\mu}$ when λ is obtained from μ by moving some nodes whose p -residues are pairwise non-adjacent (Corollary 6.2), generalising the results of [TT] on the decomposition numbers. We also show that $[\text{Res}(L(\mu)) : L(\lambda)] = 0$ whenever λ is obtained from μ by removing a node and moving other nodes while preserving their p -residues, such that the p -residues of all these nodes are pairwise non-adjacent (Theorem 6.5).

We now indicate the layout of this paper. We begin in the next section with a short account of the background theory which we require. In section 3, we prove some preliminary results which we shall require in a general setting to deal simultaneously with the quantized Fock spaces and the Grothendieck group of the Schur algebras. In section 4, we describe the restriction of the quantized Fock space to $\mathbf{U}(\mathbf{n})$ as a direct sum of factors $\mathcal{F}_{\mathbf{t}}$ isomorphic to tensor products of exterior powers of the natural n_j -dimensional representations of the $U_q(\mathfrak{sl}_{n_j})$. In section 5, we relate the canonical basis of the quantized Fock space to that of $\mathcal{F}_{\mathbf{t}}$. In section 6, we apply an argument analogous to that used in section 5 to the Schur algebras and obtain closed formulas for the decomposition numbers mentioned above.

2. PRELIMINARIES

Denote $\mathbb{N} = \{1, 2, \dots\}$ and $\mathbb{N}_0 = \{0, 1, \dots\}$.

2.1. Partitions, β -numbers, abaci. A partition $\lambda = (\lambda_1, \lambda_2, \dots)$ is an infinite weakly decreasing sequence of non-negative integers such that $\lambda_k = 0$ for all large enough k . We write $|\lambda|$ for $\sum_{i=1}^{\infty} \lambda_i$, and denote the set of all partitions by \mathcal{P} .

For $\lambda \in \mathcal{P}$, define its Young diagram $[\lambda] = \{(i, j) \in \mathbb{N}^2 \mid j \leq \lambda_i\}$. The elements of \mathbb{N}^2 are usually called nodes in this context. A node (a, b) is to the left of a node (c, d) if $b < d$, in which case (c, d) is to the right of (a, b) . If $n \in \mathbb{N}$, the n -residue of a node (i, j) is the residue class of $j - i$ modulo n . For convenience, for $k \in \mathbb{Z}$, we will say a node has n -residue k if the n -residue of the node is the residue class of k modulo n .

A node $\mathbf{n} \in [\lambda]$ is removable if $[\lambda] \setminus \{\mathbf{n}\} = [\mu]$ for some $\mu \in \mathcal{P}$, in which case we also say that \mathbf{n} is an addable node of $[\mu]$ (or simply μ). The removable node is normal if it, as well as any of the removable or addable nodes with the same n -residue as and to the right of \mathbf{n} , has at least as many removable nodes as addable nodes of the same n -residue to its right.

For $(a, b) \in [\lambda]$, define

$$\mathfrak{h}_{a,b}(\lambda) = \{(i, j) \in [\lambda] \mid i \geq a, j \geq \max(b, \lambda_{i+1})\}.$$

This is a rim hook of $[\lambda]$ (or simply λ). Note that $[\lambda] \setminus \mathfrak{h}_{a,b}(\lambda) = [\nu]$ for some $\nu \in \mathcal{P}$; we say that ν is obtained from λ by unwrapping $\mathfrak{h}_{a,b}(\lambda)$ and λ is obtained from ν by wrapping $\mathfrak{h}_{a,b}(\lambda)$.

A subset B of \mathbb{Z} is a set of β -numbers if $|\mathbb{N}_0 \cap B| + |\mathbb{Z}_{<0} \setminus B|$ is finite. Denote the collection of all sets of β -numbers by \mathfrak{B} . If $B \in \mathfrak{B}$, then since it is bounded above, we can arrange its elements in decreasing order and obtain its associated β -sequence $\overline{B} = (\overline{B}_1, \overline{B}_2, \dots)$. Write $\mathfrak{s}(B) = |\mathbb{N}_0 \cap B| - |\mathbb{Z}_{<0} \setminus B|$, and for each $s \in \mathbb{Z}$, let $\mathfrak{B}_s = \{B \in \mathfrak{B} \mid \mathfrak{s}(B) = s\}$.

For each $\lambda \in \mathcal{P}$ and $s \in \mathbb{Z}$, define $\beta_s(\lambda) = \{\lambda_i + s - i \mid i \in \mathbb{N}\}$. Then $\beta_s(\lambda) \in \mathfrak{B}_s$. In fact, $\lambda \leftrightarrow \beta_s(\lambda)$ gives a one-to-one correspondence between \mathcal{P} and \mathfrak{B}_s . More generally, we have a bijection between $\mathcal{P} \times \mathbb{Z}$ and \mathfrak{B} given by $(\lambda, s) \leftrightarrow \beta_s(\lambda)$. For each $B \in \mathfrak{B}$, write $\text{Par}(B)$ for the partition such that $(\text{Par}(B), \mathfrak{s}(B)) \leftrightarrow B$.

The n -abacus was introduced by James to facilitate manipulations with rim hooks of size n (see, e.g. [JK]). It has n vertical runners, labelled $0, 1, \dots, n-1$ from left to right, and infinitely many rows, labelled by \mathbb{Z} in an ascending order from top down. The position on row i and runner j of the n -abacus is labelled $in + j$. Thus the positions on the n -abacus are labelled by integers running left to right and top to bottom, with position 0 in the leftmost runner. We may display any subset S of \mathbb{Z} on the n -abacus by placing a bead on position x for each $x \in S$. We do so especially for the elements $B \in \mathfrak{B}$. See Figure 2.1 for an example.




FIGURE 2.1. Displaying the partition $\lambda = (13, 12, 10, 8, 8, 8, 6, 5, 5, 3, 2, 1, 1)$ on a 9-abacus, with $s = 14$.

Moving a bead from position a on the n -abacus display of B to a vacant position b produces the display of $B' = B \setminus \{a\} \cup \{b\}$, and $\mathfrak{s}(B') = \mathfrak{s}(B)$. If $a > b$, then $\text{Par}(B')$ can be obtained from $\text{Par}(B)$ by unwrapping a rim hook of size $a - b$, namely $\mathfrak{h}_{x,y}(\text{Par}(B))$ where $a = \overline{B}_x$, $\overline{B}_{z+1} < b < \overline{B}_z$, and $y = z + b + 1 - \mathfrak{s}(B)$. Conversely, if $\mu \in \mathcal{P}$ is obtained from λ by unwrapping a rim hook of size c , then $\beta_s(\mu) = \beta_s(\lambda) \setminus \{x\} \cup \{y\}$ for some $x \in \beta_s(\lambda)$ and $y \notin \beta_s(\lambda)$ such that $x - y = c$.

In particular, moving a bead on runner i in the n -abacus display of B to its vacant preceding position corresponds to removing a removable node of n -residue i_s , where i_s is the residue class of $i - s$ modulo n , from $\text{Par}(B)$.

The n -core of λ is defined to be the partition obtained from λ by successively removing rim hooks of size n . This is the partition obtained by sliding the beads in the n -abacus display of λ up their respective runners as high up as possible, and is thus well-defined (i.e. independent of the order in which the rim hooks of size n are removed).

Let \geq denote the usual lexicographic ordering on the set of infinite sequences of integers, so that $(a_1, a_2, \dots) > (b_1, b_2, \dots)$ if and only if there exists $r \in \mathbb{N}$ such that $a_k = b_k$ for all $k < r$ and $a_r > b_r$. Then \geq restricts to a total order on \mathcal{P} and on $\{\overline{B} \mid B \in \mathfrak{B}\}$ and hence on \mathfrak{B} . It is easy to see that, for $\lambda, \mu \in \mathcal{P}$, the following statements are equivalent:

- (1) $\lambda \geq \mu$.
- (2) $\beta_s(\lambda) \geq \beta_s(\mu)$ for all $s \in \mathbb{Z}$.
- (3) $\beta_s(\lambda) \geq \beta_s(\mu)$ for some $s \in \mathbb{Z}$.

Note that, from our definition, it is possible for $\lambda \geq \mu$ when $|\lambda| \neq |\mu|$, and for $B \geq B'$ when $\mathfrak{s}(B) \neq \mathfrak{s}(B')$.

We now define a coarser order on \mathfrak{B} . Let $B, B' \in \mathfrak{B}$. We write $B \rightarrow_n B'$ if $B' = B \setminus \{a, b\} \cup \{a - in, b + in\}$ for some $a, b \in B$ and $i \in \mathbb{N}$ such that $a > b + in$ and $a - in, b + in \notin B$. The Jantzen order \geq_{J_n} on \mathfrak{B} is defined as follows: $B \geq_{J_n} B'$ if and only if there exist $B_0, \dots, B_t \in \mathfrak{B}$ ($t \in \mathbb{N}_0$) such that $B_0 = B$, $B_t = B'$, and $B_{i-1} \rightarrow_n B_i$ for all $i = 1, \dots, t$. We note that, if $\lambda, \mu \in \mathcal{P}$, then $\beta_s(\lambda) \rightarrow_n \beta_s(\mu)$ for some $s \in \mathbb{Z}$ if and only if $\beta_s(\lambda) \rightarrow_n \beta_s(\mu)$ for all $s \in \mathbb{Z}$. As such, we may define the relations \rightarrow_n and \geq_{J_n} on \mathcal{P} : if $\lambda, \mu \in \mathcal{P}$, then $\lambda \rightarrow_n \mu$ (resp. $\lambda \geq_{J_n} \mu$) if and only if $\beta_s(\lambda) \rightarrow_n \beta_s(\mu)$ (resp. $\beta_s(\lambda) \geq_{J_n} \beta_s(\mu)$) for some $s \in \mathbb{Z}$. Note that $\lambda \rightarrow_n \mu$ if and only if $\lambda \geq \mu$ and μ is obtained from λ by unwrapping a hook of size in from λ and then wrapping a hook of size in .

2.2. Quantized enveloping algebras. Let $\mathbf{U} = U_q(\widehat{\mathfrak{sl}}_n)$ be the quantized enveloping algebra of affine type $A_{n-1}^{(1)}$; it is the unital associative $\mathbb{C}(q)$ -algebra generated by e_i, f_i, K_i^\pm ($0 \leq i < n$), subject to the following relations:

$$\begin{aligned} K_i^+ K_i^- &= 1 = K_i^- K_i^+, & K_i^+ K_j^+ &= K_j^+ K_i^+, \\ K_i^+ e_j K_i^- &= q^{a_{ij}} e_j, & K_i^- f_j K_i^+ &= q^{-a_{ij}} f_j, \\ e_i f_j - f_j e_i &= \delta_{ij} \frac{K_i^+ - K_i^-}{q - q^{-1}}, \\ \sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_q e_i^{1-a_{ij}-k} e_j e_i^k &= 0 \quad (i \neq j), \end{aligned}$$

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_q f_i^{1-a_{ij}-k} f_j f_i^k = 0 \quad (i \neq j).$$

Here $A = (a_{ij})_{0 \leq i, j < n}$ is the Cartan matrix of type $A_{n-1}^{(1)}$, and $\begin{bmatrix} m \\ k \end{bmatrix}_q = \frac{[m]_q [m-1]_q \cdots [m-k+1]_q}{[k]_q [k-1]_q \cdots [1]_q}$ where $[i]_q = q^{1-i} + q^{3-i} + \cdots + q^{i-3} + q^{i-1}$ for all $i \in \mathbb{N}$.

For convenience, if $j \in \mathbb{Z}$ and $j \equiv_n \bar{j}$ with $\bar{j} \in \{0, 1, \dots, n-1\}$, we also write f_j for $f_{\bar{j}}$.

Fix a tuple (n_1, \dots, n_r) of positive integers such that $n_1 + \cdots + n_r = n$, and, for each $j = 1, \dots, r$, let

$$\sigma_j = \sum_{a=1}^j n_a.$$

Let $\mathbf{U}(\mathbf{n})$ be the corresponding Levi subalgebra of \mathbf{U} , isomorphic to $U_q(\mathfrak{sl}_{n_1}) \otimes \cdots \otimes U_q(\mathfrak{sl}_{n_r})$, the quantized enveloping algebra of finite type $A_{n_1-1} \times \cdots \times A_{n_r-1}$. So $\mathbf{U}(\mathbf{n})$ is the subalgebra of \mathbf{U} generated by the e_i , f_i and K_i^\pm such that $i \neq \sigma_k$ for all k with $0 \leq k \leq r-1$.

The assignments $\phi(e_i) = e_i$, $\phi(f_i) = f_i$, $\phi(K_i^\pm) = K_i^\mp$ define a $\mathbb{C}(q)$ -semilinear automorphism $\phi : \mathbf{U} \rightarrow \mathbf{U}$. (Here, and hereafter, a map $f : V \rightarrow W$ between $\mathbb{C}(q)$ -vector spaces is $\mathbb{C}(q)$ -semilinear if and only if $f(v_1 + a(q)v_2) = f(v_1) + a(q^{-1})f(v_2)$ for all $v_1, v_2 \in V$ and $a(q) \in \mathbb{C}(q)$.) Likewise, $\omega(e_i) = f_i$, $\omega(f_i) = e_i$, $\omega(K_i^\pm) = K_i^\pm$ can be extended to a $\mathbb{C}(q)$ -linear antiautomorphism of \mathbf{U} . The maps ϕ and ω restrict to an automorphism and an antiautomorphism of $\mathbf{U}(\mathbf{n})$, respectively.

We define a coproduct $\Delta : \mathbf{U} \rightarrow \mathbf{U} \otimes \mathbf{U}$ by

$$\begin{aligned} \Delta(e_i) &= e_i \otimes 1 + K_i^- \otimes e_i, \\ \Delta(f_i) &= f_i \otimes K_i^+ + 1 \otimes f_i, \\ \Delta(K_i^\pm) &= K_i^\pm \otimes K_i^\pm. \end{aligned}$$

It restricts to a coproduct on $\mathbf{U}(\mathbf{n})$. Note that Δ differs from the coproduct used in [Lu] and [Jan], which is $(\phi \otimes \phi) \circ \Delta \circ \phi$, and from that used in [Ka], [HK] and [Le], which is $(\omega \otimes \omega) \circ \Delta \circ \omega$.

Given two \mathbf{U} -modules (resp. $\mathbf{U}(\mathbf{n})$ -modules) M and N , we take the \mathbf{U} -module (resp. $\mathbf{U}(\mathbf{n})$ -module) structure on $M \otimes N$ to be given by the pullback along Δ .

2.3. Fock spaces. Let

$$\mathcal{F} := \bigoplus_{\lambda \in \mathcal{P}} \mathbb{C}(q)\lambda$$

be the $\mathbb{C}(q)$ -vector space with distinguished basis given by the set \mathcal{P} of partitions of all natural numbers.

In [LT], Leclerc and Thibon defined an $\mathbb{C}(q)$ -semilinear bar involution $x \mapsto \bar{x}$ on \mathcal{F} . They proved the existence of another distinguished basis

$\{G(\lambda) \mid \lambda \in \mathcal{P}\}$ of \mathcal{F} , called the canonical basis, which has the following characterization:

$$G(\lambda) - \lambda \in \sum_{\mu \in \mathcal{P}} q\mathbb{Z}[q]\mu, \quad \overline{G(\lambda)} = G(\lambda).$$

Let $\langle -, - \rangle$ be the symmetric bilinear form on \mathcal{F} with respect to which \mathcal{P} is orthonormal. For $\lambda, \mu \in \mathcal{P}$, define $d_{\lambda\mu}(q) \in \mathbb{C}(q)$ by

$$d_{\lambda\mu}(q) = \langle G(\mu), \lambda \rangle.$$

As the involution defined in [LT] depends on a fixed integer $n \geq 2$, $d_{\lambda\mu}(q)$ also implicitly depends on n . When we need to emphasize the role of n , we shall write $d_{\lambda\mu}^n(q)$ instead. For convenience, we further define $d_{\lambda\mu}^1(q) = \delta_{\lambda\mu}$.

We shall make use of the following remarkable properties of these q -decomposition numbers $d_{\lambda\mu}(q)$:

Theorem 2.1.

- (1) $d_{\mu\mu}(q) = 1$.
- (2) If $\lambda \neq \mu$, then $d_{\lambda\mu}(q) \in q\mathbb{N}_0[q]$.
- (3) If $d_{\lambda\mu}(q) \neq 0$, then $\mu \geq_{J_n} \lambda$.

Recall the collection \mathfrak{B} of all sets of β -numbers. Define $\mathcal{F}_{\mathbb{Z}}$ to be the $\mathbb{C}(q)$ -vector space with basis \mathfrak{B} . Following [H, MM], we define an action of $\mathbf{U} = U_q(\widehat{\mathfrak{sl}}_n)$ on $\mathcal{F}_{\mathbb{Z}}$ as follows. Let $B \in \mathfrak{B}$, and take $x \notin B$ such that $x - 1 \in B$. Let $C = B \setminus \{x - 1\} \cup \{x\}$, and

$$\begin{aligned} N_{>}(B, C) &= |\{y \in B \mid y > x, y \equiv_n x - 1\}| - |\{y \in B \mid y > x, y \equiv_n x\}|, \\ N_{<}(B, C) &= |\{y \notin B \mid y < x - 1, y \equiv_n x - 1\}| - |\{y \notin B \mid y < x - 1, y \equiv_n x\}|. \end{aligned}$$

Here, and hereafter, we write $a \equiv_n b$ for $a \equiv b \pmod{n}$. For $B \in \mathfrak{B}$ and $0 \leq i < n$, let

$$N_i(B) = |\{y \notin B \mid y - 1 \in B, y \equiv_n i\}| - |\{y \in B \mid y - 1 \notin B, y \equiv_n i\}|.$$

Then we have

$$e_i(C) = \sum_B q^{N_{<}(B, C)} C, \quad f_i(B) = \sum_C q^{N_{>}(B, C)} C, \quad K_i^+(B) = q^{N_i(B)},$$

where the first sum runs over all $B \in \mathfrak{B}$ such that $B = C \setminus \{x\} \cup \{x - 1\}$ for some $x \in C$, $x \equiv_n i$ and $x - 1 \notin C$, and second sum over all $C \in \mathfrak{B}$ such that $C = B \setminus \{x - 1\} \cup \{x\}$ for some $x - 1 \in B$, $x \equiv_n i$ and $x \notin B$.

Let $s \in \mathbb{Z}$, and write \mathcal{F}_s for the vector subspace of $\mathcal{F}_{\mathbb{Z}}$ with basis \mathfrak{B}_s ($= \{B \in \mathfrak{B} \mid \mathfrak{s}(B) = s\}$). It is easy to see that \mathcal{F}_s is invariant under the \mathbf{U} -action defined above. Furthermore, the bijection from \mathcal{P} to \mathfrak{B}_s defined by $\lambda \mapsto \beta_s(\lambda)$ induces a $\mathbb{C}(q)$ -linear isomorphism $\beta_s : \mathcal{F} \rightarrow \mathcal{F}_s$. Thus via this isomorphism, \mathcal{F}_s inherits the bar-involution from \mathcal{F} , while \mathcal{F} inherits an \mathbf{U} -action from \mathcal{F}_s . The action of e_i and f_i on \mathcal{F} via β_s may be described as follows. Let i_s be the residue class of $i - s$ modulo n . Suppose that $\lambda \in \mathcal{P}$

has an addable node \mathbf{n} of n -residue i_s , and let μ be the partition obtained by adding \mathbf{n} to λ . Let $N_{>}(\lambda, \mu)$ (resp. $N_{<}(\lambda, \mu)$) be the number of addable nodes of λ , of n -residue i_s and to the right (resp. left) of \mathbf{n} , minus the number of removable nodes of λ , of n -residue i_s and to the right (resp. left) of \mathbf{n} . For $\lambda \in \mathcal{P}$ and $0 \leq i < n$, let $N_i(\lambda)$ be the number of addable nodes of λ of n -residue i minus the number of removable nodes of λ of n -residue i . Then

$$e_i(\mu) = \sum_{\lambda} q^{-N_{<}(\lambda, \mu)} \lambda, \quad f_i(\lambda) = \sum_{\mu} q^{N_{>}(\lambda, \mu)} \mu, \quad K_i^+(\lambda) = q^{N_i(\lambda)} \lambda,$$

where the first sum runs over all partitions λ that can be obtained by removing a removable node of n -residue i_s from μ and the second sum runs over all partitions μ that can be obtained by adding an addable node of n -residue i_s to λ .

For each $\lambda \in \mathcal{P}$, write $G_s(\lambda)$ for $\beta_s(G(\lambda))$.

Theorem 2.2 ([LT]). *For all $u \in \mathbf{U}$ and $x \in \mathcal{F}_s$, we have*

$$\overline{ux} = \phi(u) \overline{x}.$$

In particular, for $0 \leq i < n$,

$$f_i(G_s(\lambda)) = \sum_{\mu \in \mathcal{P}} L_{\lambda}^{\mu}(q) G_s(\mu),$$

where $L_{\lambda}^{\mu}(q) = L_{\lambda}^{\mu}(q^{-1}) \in \mathbb{C}(q)$ for all $\mu \in \mathcal{P}$.

Lascoux and Thibon in fact extend the action of \mathbf{U} on \mathcal{F}_s to an action of the larger algebra $U_q(\widehat{\mathfrak{gl}}_n)$, and prove the theorem in this context. We will not require the extended action.

2.4. Tensor products of based modules. In this subsection we give an abbreviated introduction to Lusztig's theory of based modules over quantized enveloping algebras; see [Lu, Chapters 27-28] for full details, though note that since our coproduct is slightly different, we need to swap q and q^{-1} in Lusztig's account. The theory is valid in finite type; we consider specifically based modules of $\mathbf{U}(\mathbf{n})$, the subalgebra of \mathbf{U} defined in subsection 2.2.

A based $\mathbf{U}(\mathbf{n})$ -module M is a finite-dimensional weight $\mathbf{U}(\mathbf{n})$ -module equipped with a $\mathbb{C}(q)$ -basis B satisfying certain conditions, amongst which are the following:

- (1) each $b \in B$ is a weight vector.
- (2) for all $u \in \mathbf{U}(\mathbf{n})$ and $m \in M$, we have $\psi_M(um) = \phi(u)\psi_M(m)$, where ψ_M is the $C(q)$ -semilinear involution of M fixing every element of B .

Any highest weight irreducible $\mathbf{U}(\mathbf{n})$ -module, together with its canonical basis, is a based module.

Recall that $\mathbf{U}(\mathbf{n}) \cong U_q(\mathfrak{sl}_{n_1}) \otimes \cdots \otimes U_q(\mathfrak{sl}_{n_r})$. Given a based $U_q(\mathfrak{sl}_{n_j})$ -module $(M^{(j)}, B^{(j)})$ for each $j = 1, \dots, r$, we may form the 'external' tensor

product

$$M = M^{(1)} \otimes \cdots \otimes M^{(r)},$$

a $\mathbf{U}(\mathbf{n})$ -module. It is then clear that the basis $B = B^{(1)} \otimes \cdots \otimes B^{(r)}$ satisfies the two properties above, and in fact it is true that (M, B) is a based $\mathbf{U}(\mathbf{n})$ -module.

We now turn to the consideration of ‘internal’ tensor products. Let $(M_1, B_1), \dots, (M_d, B_d)$ be based modules of $\mathbf{U}(\mathbf{n})$. The coproduct on $\mathbf{U}(\mathbf{n})$ gives a $\mathbf{U}(\mathbf{n})$ -module structure $M = M_1 \otimes \cdots \otimes M_d$. However M together with the obvious basis $\mathbf{B}^\otimes := B_1 \otimes \cdots \otimes B_d$ is not necessarily a based module. In order to correct this deficiency, Lusztig defines an involution ψ_M of M , constructed out of the involutions ψ_{M_i} and the quasi- R -matrix of $\mathbf{U}(\mathbf{n})$, and shows that for each $\mathbf{b} = (b_1, \dots, b_d) \in \mathbf{B} := B_1 \times \cdots \times B_d$ there exists a unique element

$$\mathbf{b}^\diamond = b_1 \diamond \cdots \diamond b_d \in M$$

such that

$$\psi_M(\mathbf{b}^\diamond) = \mathbf{b}^\diamond \quad \text{and} \quad \mathbf{b}^\diamond - \mathbf{b}^\otimes \in \bigoplus_{\mathbf{c} \in \mathbf{B}} q\mathbb{Z}[q]\mathbf{c}^\otimes,$$

where $\mathbf{b}^\otimes = b_1 \otimes \cdots \otimes b_d$. He also proves that, writing

$$\mathbf{B}^\diamond = B_1 \diamond \cdots \diamond B_d := \{\mathbf{b}^\diamond \mid \mathbf{b} \in \mathbf{B}\},$$

(M, \mathbf{B}^\diamond) is a based $\mathbf{U}(\mathbf{n})$ -module.

In order to formulate some required additional properties of this canonical basis, we introduce a partial order on \mathbf{B} , a reverse lexicographic order, as follows: $\mathbf{b} = (b_1, \dots, b_d) > \mathbf{b}' = (b'_1, \dots, b'_d)$ if and only if there exists i such that $\text{wt}(b_j) = \text{wt}(b'_j)$ for all $i < j \leq d$ and $\text{wt}(b_i) < \text{wt}(b'_i)$.

Lemma 2.3. *Keep the notations above.*

(1) *We have, for all $\mathbf{b} \in \mathbf{B}$,*

$$\mathbf{b}^\diamond - \mathbf{b}^\otimes \in \bigoplus_{\mathbf{c} < \mathbf{b}} q\mathbb{Z}[q]\mathbf{c}^\otimes.$$

(2) *Let $\mathbf{b} = (b_1, \dots, b_d) \in \mathbf{B}$, and suppose that b_d is a highest weight vector in M_d . Writing $\mathbf{a} = (b_1, \dots, b_{d-1})$, we have*

$$\mathbf{b}^\diamond = \mathbf{a}^\diamond \otimes b_d.$$

Proof. This first part follows by iterated application of [Lu, Theorem 27.3.2(b)]. The second is checked by directly evaluating $\psi_M(\mathbf{a}^\diamond \otimes b_d)$; see the proof of [Lu, Lemma 28.2.6] \square

Finally we explain how to handle a mixture of external and internal tensor products of based modules. Suppose that for $j = 1, \dots, r$ we are given based $U_q(\mathfrak{sl}_{n_j})$ -modules

$$(M_1^{(j)}, B_1^{(j)}), \dots, (M_d^{(j)}, B_d^{(j)}).$$

Then, as described above, for each j we have a based $U_q(\mathfrak{sl}_{n_j})$ -module

$$M^{(j)} = M_1^{(j)} \otimes \cdots \otimes M_d^{(j)}$$

with basis

$$\mathbf{B}^{(j)\diamond} = B_1^{(j)} \diamond \cdots \diamond B_d^{(j)}.$$

We can then take the external tensor product, obtaining a based $\mathbf{U}(\mathbf{n})$ -module

$$M' = M^{(1)} \otimes \cdots \otimes M^{(r)}$$

with basis

$$\mathbf{B}^{\diamond\diamond} = \mathbf{B}^{(1)\diamond} \otimes \cdots \otimes \mathbf{B}^{(r)\diamond}.$$

On the other hand, we may take external tensor products first, before taking internal tensor products. For each $i = 1, \dots, d$, we have the based $\mathbf{U}(\mathbf{n})$ -module

$$M_i = M_i^{(1)} \otimes \cdots \otimes M_i^{(r)}.$$

with basis

$$\mathbf{B}_i^{\otimes} = B_i^{(1)} \otimes \cdots \otimes B_i^{(r)}.$$

The internal tensor product

$$M'' = M_1 \otimes \cdots \otimes M_d$$

is then a based $\mathbf{U}(\mathbf{n})$ -module with basis

$$\mathbf{B}^{\otimes\diamond} = \mathbf{B}_1^{\otimes} \diamond \cdots \diamond \mathbf{B}_d^{\otimes}.$$

Thus, to each $\mathbf{b} \in \mathbf{B} = \prod_{1 \leq i \leq d, 1 \leq j \leq r} B_i^{(j)}$, we have associated a distinguished basis element $\mathbf{b}^{\diamond\diamond} \in \mathbf{B}^{\diamond\diamond} \subseteq M'$ and another distinguished basis element $\mathbf{b}^{\otimes\diamond} \in \mathbf{B}^{\otimes\diamond} \subseteq M''$.

Proposition 2.4. *Keep the notation as above. The obvious correspondence between M' and M'' gives an isomorphism of based $\mathbf{U}(\mathbf{n})$ -modules. Furthermore $\mathbf{b}^{\diamond\diamond} \leftrightarrow \mathbf{b}^{\otimes\diamond}$ for each $\mathbf{b} \in \mathbf{B}$ under this correspondence.*

Proof. Since the coproduct of $\mathbf{U}(\mathbf{n})$ is simply the tensor product of the coproducts of its factors $\mathbf{U}_q(\mathfrak{sl}_{n_j})$, the natural identification of vector spaces

$$M = M^{(1)} \otimes \cdots \otimes M^{(r)} = \bigotimes_{\substack{1 \leq i \leq d \\ 1 \leq j \leq r}} M_i^{(j)} = M_1 \otimes \cdots \otimes M_d = M''$$

is an isomorphism of $\mathbf{U}(\mathbf{n})$ -modules. Under this identification $\mathbf{b}^{\diamond\diamond} = \mathbf{b}^{\otimes\diamond}$ for all $\mathbf{b} \in \mathbf{B}$, as they are characterized in terms of involutions that coincide because the quasi- R -matrix of $\mathbf{U}(\mathbf{n})$ is the tensor product of the quasi- R -matrices of its factors $U_q(\mathfrak{sl}_{n_j})$. \square

2.5. Schur algebras. Let $\mathbf{S}_m = S(m, m)$ be the Schur algebra of degree m over a field \mathbb{F} of characteristic $p > 0$. This is a quasi-hereditary algebra with indexing set $\{\lambda \in \mathcal{P} \mid |\lambda| = m\}$, together with the order opposite to

the lexicographic order; here we are following the notation conventions in [Jam]. Denote by $L(\lambda)$, $\Delta(\lambda)$ and $P(\lambda)$ the simple, standard and projective indecomposable modules associated to λ . By Brauer-Humphrey's reciprocity the decomposition number $d_{\lambda\mu} := [\Delta(\lambda) : L(\mu)]$ may be expressed as the multiplicity $[P(\mu) : \Delta(\lambda)]$ of $\Delta(\lambda)$ in a Δ -filtration of $P(\mu)$; even though the filtrations are not unique, the multiplicities are. These numbers enjoy the following well-known properties:

Proposition 2.5. *Let $\lambda, \mu \in \mathcal{P}$. Then*

- (1) $d_{\lambda\lambda} = 1$;
- (2) $d_{\lambda\mu} \neq 0$ only if $\lambda \leq_{J_p} \mu$ (see [TT, Theorem 2.5(iv)]);
- (3) $d_{\lambda\mu} \geq d_{\lambda\mu}(1)$ (by [VV] and [Jam]).

For each residue class i modulo p , one can define functors

$$i\text{-Res}_m : \mathbf{S}_{m+1}\text{-mod} \rightarrow \mathbf{S}_m\text{-mod},$$

$$i\text{-Ind}_m : \mathbf{S}_m\text{-mod} \rightarrow \mathbf{S}_{m+1}\text{-mod}.$$

They are a biadjoint pair of exact functors, sending projectives to projectives. In particular, if $\lambda \in \mathcal{P}$ with $|\lambda| = m$, then

$$i\text{-Ind}_m(P(\lambda)) \cong \bigoplus_{\substack{\mu \in \mathcal{P} \\ |\mu|=m+1}} P(\mu)^{\oplus L_\lambda^\mu},$$

where each $L_\lambda^\mu \in \mathbb{N}_0$. In fact,

$$L_\lambda^\mu = \dim_{\mathbb{F}}(\text{Hom}_{\mathbf{S}_{m+1}}(i\text{-Ind}_m(P(\lambda)), L(\mu))) = \dim_{\mathbb{F}}(\text{Hom}_{\mathbf{S}_m}(P(\lambda), i\text{-Res}_m(L(\mu)))),$$

so that L_λ^μ is the multiplicity $[i\text{-Res}_m(L(\mu)) : L(\lambda)]$ of $L(\lambda)$ as a composition factor of $i\text{-Res}_m(L(\mu))$. Kleshchev describes some of these multiplicities.

Theorem 2.6 ([KL, Theorem 9.3]). *Let $\mu \in \mathcal{P}$ with $|\mu| = m$. Suppose that μ has a normal node \mathbf{n} of p -residue i , and let λ be the partition obtained when this node is removed from μ . Then*

$$[i\text{-Res}_m(L(\mu)) : L(\lambda)] =$$

$$1 + \text{number of normal nodes of } \mu \text{ of } p\text{-residue } i \text{ and to the right of } \mathbf{n}.$$

Let $i\text{-Res} = \bigoplus_m i\text{-Res}_m$ and $i\text{-Ind} = \bigoplus_m i\text{-Ind}_m$. For convenience, for $k \in \mathbb{Z}$, we also write $k\text{-Res}$ and $k\text{-Ind}$, which mean $\bar{k}\text{-Res}$ and $\bar{k}\text{-Ind}$ respectively, where \bar{k} is the residue class of k modulo p .

The effects of $i\text{-Res}$ and $i\text{-Ind}$ on the standard module $\Delta(\lambda)$ can be easily described in the Grothendieck group $K_0(\bigoplus_m \mathbf{S}_m\text{-mod})$ as follows:

$$\begin{aligned} [i\text{-Res}(\Delta(\lambda))] &= \sum_{\tau} [\Delta(\tau)]; \\ [i\text{-Ind}(\Delta(\lambda))] &= \sum_{\mu} [\Delta(\mu)], \end{aligned}$$

where the first sum runs over all partitions τ that can be obtained by removing a removable node of p -residue i from λ , while the second sum runs over all partitions μ that can be obtained by adding an addable node of p -residue i to λ .

It is natural to identify $K_0(\bigoplus_m \mathbf{S}_m\text{-mod})$ with the classical (non-quantized) Fock space $\mathcal{F}_c = \bigoplus_{\lambda \in \mathcal{P}} \mathbb{Z}\lambda$ under the correspondence $[\Delta(\lambda)] \leftrightarrow \lambda$.

Recall the quantized Fock space \mathcal{F} introduced in [subsection 2.3](#). Let $A = \mathbb{Z}[q, q^{-1}]$, and let \mathcal{F}_A be the free A -submodule of \mathcal{F} with basis \mathcal{P} . Then $G(\lambda) \in \mathcal{F}_A$ for all $\lambda \in \mathcal{P}$ by [Theorem 2.1](#). Similarly, let $\mathcal{F}_{s,A}$ denote the free A -submodule of \mathcal{F}_s with basis \mathfrak{B}_s for each $s \in \mathbb{Z}$. We have a surjective \mathbb{Z} -linear map $\varepsilon_{q=1} : \mathcal{F}_{\mathbb{Z},A} \rightarrow \mathcal{F}_c$ defined by $a(q)B \mapsto a(1)\text{Par}(B)$ for all $a(q) \in A$ and $B \in \mathfrak{B}$.

Let $n = p$, and let \mathbf{U}_A be the A -subalgebra of \mathbf{U} generated by $\{e_i, f_i, K_i^\pm \mid i = 0, 1, \dots, p-1\}$. Then for each $s \in \mathbb{Z}$, $\mathcal{F}_{s,A}$ is a \mathbf{U}_A -submodule of \mathcal{F}_s . Under the above identification of $K_0(\bigoplus_m \mathbf{S}_m\text{-mod})$ with \mathcal{F}_c , we have that $\varepsilon_{q=1}$, when restricted to $\mathcal{F}_{s,A}$, intertwines the Chevalley generators e_i and f_i with the $(i-s)$ -Res and $(i-s)$ -Ind functors. More precisely, we have

$$\varepsilon_{q=1} \circ f_i = (i-s)\text{-Ind} \circ \varepsilon_{q=1} \quad \text{and} \quad \varepsilon_{q=1} \circ e_i = (i-s)\text{-Res} \circ \varepsilon_{q=1}.$$

3. GENERAL SETUP

In order to deal with the quantized Fock spaces and the classical non-quantized Fock space simultaneously, we consider the following general setup.

Let R be an integral domain, and let \mathfrak{F} be the free R -module with R -basis $\{s(\lambda) \mid \lambda \in \mathcal{P}\}$, indexed by the set \mathcal{P} of all partitions. Let $\langle \cdot, \cdot \rangle$ denote the symmetric R -bilinear form on \mathfrak{F} with respect to which $\{s(\lambda) \mid \lambda \in \mathcal{P}\}$ is orthonormal.

For each $\lambda \in \mathcal{P}$, let $H(\lambda) \in \mathfrak{F}$ be such that $\langle H(\lambda), s(\mu) \rangle \neq 0$ only if $\lambda \geq_{J_n} \mu$, and $\langle H(\lambda), s(\lambda) \rangle = 1$. Then $\{H(\lambda) \mid \lambda \in \mathcal{P}\}$ is another R -basis of \mathfrak{F} . We note the following easy lemma:

Lemma 3.1. *Let $x = \sum_{\mu \in \mathcal{P}} a_\mu H(\mu) \in \mathfrak{F}$, where $a_\mu \in R$ for all $\mu \in \mathcal{P}$. Let $\nu \in \mathcal{P}$ be such that $a_\mu = 0$ whenever $\mu >_{J_n} \nu$. Then $\langle x, s(\nu) \rangle = a_\nu$.*

Proof. We have $a_\mu = 0$ if $\mu >_{J_n} \nu$, while $\langle H(\mu), s(\nu) \rangle = 0$ if $\mu \not\geq_{J_n} \nu$, so that

$$\langle x, s(\nu) \rangle = \left\langle \sum_{\mu \in \mathcal{P}} a_\mu H(\mu), s(\nu) \right\rangle = \langle a_\nu H(\nu), s(\nu) \rangle = a_\nu.$$

□

Fix $n \in \mathbb{N}$ and let \mathfrak{r} be a residue class modulo n . Let $\mathfrak{f}_{\mathfrak{r}} : \mathfrak{F} \rightarrow \mathfrak{F}$ be an R -linear map such that $\langle \mathfrak{f}_{\mathfrak{r}}(s(\lambda)), s(\mu) \rangle \neq 0$ only if λ can be obtained from μ by removing a removable node of n -residue \mathfrak{r} , and equals 1 if λ is obtained by removing the rightmost such node which also lies to the right of all addable nodes of n -residue \mathfrak{r} .

Proposition 3.2. *Let $\lambda \in \mathcal{P}$, and suppose that its rightmost addable node of n -residue \mathfrak{r} lies to the right of all its removable nodes of n -residue \mathfrak{r} , and let μ be the partition obtained by adding this node. Then*

$$\mathfrak{f}_{\mathfrak{r}}(H(\lambda)) = H(\mu) + \sum_{\substack{\nu \in \mathcal{P} \\ |\nu| = |\mu| \\ \nu < \mu}} a_{\nu} H(\nu),$$

with $a_{\nu} \in R$ for all ν .

Proof. Note first that when we remove from μ its rightmost removable node of n -residue \mathfrak{r} , we obtain λ .

Since $\{H(\nu) \mid \nu \in \mathcal{P}\}$ is a basis for \mathfrak{F} , we have $\mathfrak{f}_{\mathfrak{r}}(H(\lambda)) = \sum_{\nu \in \mathcal{P}} a_{\nu} H(\nu)$, where $a_{\nu} \in R$ for all $\nu \in \mathcal{P}$. Since $\langle H(\rho), s(\sigma) \rangle \neq 0$ only if $\rho \geq_{J_n} \sigma$, and the latter only if $|\rho| = |\sigma|$, we see that $a_{\nu} = 0$ unless $|\nu| = |\lambda| + 1 = |\mu|$. Thus, we need to show that $a_{\mu} = 1$, and $a_{\nu} \neq 0$ only if $\nu \leq \mu$.

Let $\sigma \in \mathcal{P}$ be maximal (with respect to \geq) subject to $a_{\sigma} \neq 0$. Then, by [Lemma 3.1](#), we have

$$\begin{aligned} (0 \neq) a_{\sigma} &= \langle \mathfrak{f}_{\mathfrak{r}}(H(\lambda)), s(\sigma) \rangle = \langle \mathfrak{f}_{\mathfrak{r}} \left(\sum_{\rho \in \mathcal{P}} \langle H(\lambda), s(\rho) \rangle s(\rho) \right), s(\sigma) \rangle \\ &= \sum_{\rho \in \mathcal{P}} \langle H(\lambda), s(\rho) \rangle \langle \mathfrak{f}_{\mathfrak{r}}(s(\rho)), s(\sigma) \rangle. \end{aligned}$$

Thus there exists $\rho \in \mathcal{P}$ such that $\langle H(\lambda), s(\rho) \rangle, \langle \mathfrak{f}_{\mathfrak{r}}(s(\rho)), \sigma \rangle \neq 0$; in particular, $\rho \leq \lambda$, $|\rho| = |\lambda|$, and ρ can be obtained from σ by removing a removable node of n -residue \mathfrak{r} . This also shows that $|\sigma| = |\mu|$.

If $\rho = \lambda$, then σ can be obtained by adding an addable node of n -residue \mathfrak{r} to λ , so that since μ is obtained by adding the rightmost addable node of λ of n -residue \mathfrak{r} , we have $\mu \geq \sigma$ (in fact, $\mu \geq_{J_n} \sigma$).

If $\rho \neq \lambda$, then $\rho < \lambda$. Let $a, b \in \mathbb{N}$ be such that $\mu_i = \lambda_i + \delta_{ai}$, $\sigma_i = \rho_i + \delta_{bi}$ for all $i \in \mathbb{N}$. Also, let $k \in \mathbb{N}$ be such that $\rho_i = \lambda_i$ for all $i < k$ and $\rho_k < \lambda_k$. We consider the cases $a < k$, $a = k$ and $a > k$ separately.

$a < k$: Note first that in this case $a \leq b$: otherwise $\lambda_i = \rho_i$ for all $i \leq b$ while $\sigma_b = \rho_b + 1$, so that $(b, \rho_b + 1) = (b, \lambda_b + 1)$ is an addable node of ρ and λ of n -residue \mathfrak{r} lying to the right of $(a, \lambda_a + 1)$, contradicting the latter being the rightmost addable node of λ of n -residue \mathfrak{r} . Thus $\mu_i = \lambda_i = \rho_i = \sigma_i$ for all $i < a$, and $\mu_a = \lambda_a + 1 = \rho_a + 1 = \sigma_a + 1 - \delta_{ba}$. Consequently either $\mu_a > \sigma_a$ or $a = b$; in the latter case $\mu_i = \lambda_i + \delta_{ai} = \rho_i + \delta_{bi} = \sigma_i$ for all $i < k$, and $\mu_k = \lambda_k > \rho_k = \sigma_k$. Either way, we have $\mu > \sigma$.

$a = k$: In this case, we have $\mu_i = \lambda_i = \rho_i$ for all $i < a = k$, and $\mu_a = \lambda_a + 1 = \lambda_k + 1 > \rho_k + 1 = \rho_a + 1 = \sigma_a + 1 - \delta_{ba} \geq \sigma_a$. Thus $\mu > \sigma$.

$a > k$: In this case, we have $\mu_i = \lambda_i = \rho_i$ for all $i < k$, and $\mu_k = \lambda_k > \rho_k = \sigma_k - \delta_{bk}$. Thus either $\mu_k > \sigma_k$ or both $b = k$ and $\mu_k = \sigma_k$. In the latter case, the node $(k, \sigma_k) = (k, \mu_k)$ is a removable node of σ whose removal produces ρ , so that it has n -residue \mathfrak{r} ; but this cannot be a removable node of μ since (a, μ_a) is its rightmost removable node of n -residue \mathfrak{r} . As such, $(k+1, \mu_k) \in [\mu] \setminus [\sigma]$. This gives $\mu_{k+1} > \sigma_{k+1}$. Thus, once again, either way, we get $\mu > \sigma$.

We thus conclude that $\mu \geq \sigma$ always. To finish off the proof, it suffices to show that $a_\mu = 1$. By Lemma 3.1, we have, as above,

$$a_\mu = \langle \mathfrak{f}_{\mathfrak{r}}(H(\lambda)), s(\mu) \rangle = \sum_{\rho \in \mathcal{P}} \langle H(\lambda), s(\rho) \rangle \langle \mathfrak{f}_{\mathfrak{r}}(s(\rho)), s(\mu) \rangle.$$

For each ρ such that $\langle \mathfrak{f}_{\mathfrak{r}}(s(\rho)), s(\mu) \rangle \neq 0$, ρ must be obtained from μ by removing a removable node of n -residue \mathfrak{r} . Since λ is obtained by removing from μ its rightmost removable node of n -residue \mathfrak{r} , we see that $\lambda \leq \rho$ (in fact, $\lambda \leq_{J_n} \rho$), so that $\langle H(\lambda), s(\rho) \rangle = 0$ unless $\lambda = \rho$. Thus,

$$a_\mu = \langle H(\lambda), s(\lambda) \rangle \langle \mathfrak{f}_{\mathfrak{r}}(s(\lambda)), s(\mu) \rangle = 1.$$

□

Remark. Proposition 3.2 in fact holds even when $H(\lambda)$ satisfies the weaker condition that $\langle H(\lambda), s(\mu) \rangle \neq 0$ only if $\lambda > \mu$ and $|\lambda| = |\mu|$. The proof is similar, uses a modified version of Lemma 3.1. We leave it to the reader as an easy exercise.

Applying Proposition 3.2 to $\mathcal{F}_c (\cong K_0(\bigoplus_m \mathbf{S}_m\text{-mod}))$ and $\mathcal{F}_{A,s}$ in the place of \mathfrak{F} , and $i\text{-Ind}$ and f_{s+i} in the place of \mathfrak{f} , we get the following immediately corollary.

Corollary 3.3. *Let $\lambda \in \mathcal{P}$, and suppose that its rightmost addable node of n -residue i lies to the right of all its removable nodes of n -residue i , and let μ be the partition obtained by adding this node. Then*

(1) *when $n = p$,*

$$i\text{-Ind}(P(\lambda)) \cong P(\mu) \oplus \bigoplus_{\substack{\nu \in \mathcal{P} \\ |\nu| = |\mu| \\ \nu < \mu}} P(\nu)^{\oplus [i\text{-Res}(L(\nu)) : L(\lambda)]},$$

(2)

$$f_{s+i}(G_s(\lambda)) = G_s(\mu) + \sum_{\substack{\nu \in \mathcal{P} \\ |\nu| = |\mu| \\ \nu < \mu}} L_\lambda^\nu(q) G_s(\nu),$$

where $L_\lambda^\nu(q) \in \mathbb{Z}[q, q^{-1}]$ for all ν .

Suppose further that $\mathcal{P} = \bigcup_{\mathbf{t} \in \mathbf{T}} \mathcal{P}_{\mathbf{t}}$ can be partitioned into subsets which are indexed by a set \mathbf{T} , and that the indexing set \mathbf{T} can be partially ordered

by \geq so that if $\lambda \in \mathcal{P}_{\mathbf{t}}$, $\mu \in \mathcal{P}_{\mathbf{s}}$ and $\lambda \geq_{J_n} \mu$, then $\mathbf{t} \geq \mathbf{s}$. For each $\mathbf{t} \in \mathbf{T}$, let $\mathfrak{F}_{\mathbf{t}}$ be the free R -submodule of \mathfrak{F} with R -basis $\{s(\lambda) \mid \lambda \in \mathcal{P}_{\mathbf{t}}\}$. Then $\mathfrak{F} = \bigoplus_{\mathbf{t} \in \mathbf{T}} \mathfrak{F}_{\mathbf{t}}$; let $\pi_{\mathbf{t}} : \mathfrak{F} \rightarrow \mathfrak{F}$ denote the natural projection onto $\mathfrak{F}_{\mathbf{t}}$ via this decomposition. We have, as immediate consequences of such an order on \mathbf{T} , the following: for $\mathbf{t} \in \mathbf{T}$ and $\lambda \in \mathcal{P}_{\mathbf{t}}$,

- $H(\lambda) \in \bigoplus_{\mathbf{s} \leq \mathbf{t}} \mathfrak{F}_{\mathbf{s}}$;
- if $\mathbf{s} \not\leq \mathbf{t}$, then $\pi_{\mathbf{s}}(H(\lambda)) = 0$.

Lemma 3.4. *Let $\xi : \mathfrak{F} \rightarrow \mathfrak{F}$ be an R -linear map satisfying $\xi(\mathfrak{F}_{\mathbf{s}}) \subseteq \mathfrak{F}_{\mathbf{s}}$ for all $\mathbf{s} \in \mathbf{T}$. Let $\mathbf{t} \in \mathbf{T}$ and $\lambda \in \mathcal{P}_{\mathbf{t}}$. Suppose that $\xi(H(\lambda)) = \sum_{\mu \in \mathcal{P}} a_{\mu} H(\mu)$, where $a_{\mu} \in R$ for all $\mu \in \mathcal{P}$. Then*

- (1) $a_{\mu} = 0$ for all $\mu \in \mathcal{P}_{\mathbf{s}}$ with $\mathbf{s} \not\leq \mathbf{t}$;
- (2) $\xi(\pi_{\mathbf{t}}(H(\lambda))) = \pi_{\mathbf{t}}(\xi(H(\lambda))) = \sum_{\mu \in \mathcal{P}_{\mathbf{t}}} a_{\mu} \pi_{\mathbf{t}}(H(\mu))$.

Proof. Since $H(\lambda) \in \bigoplus_{\mathbf{s} \leq \mathbf{t}} \mathfrak{F}_{\mathbf{s}}$, we have $\xi(H(\lambda)) \in \bigoplus_{\mathbf{s} \leq \mathbf{t}} \mathfrak{F}_{\mathbf{s}}$. If there exists $\nu \in \mathcal{P}_{\mathbf{s}}$ with $\mathbf{s} \not\leq \mathbf{t}$ such that $a_{\nu} \neq 0$, choose ν to be maximal with respect to \geq_{J_n} . Then by [Lemma 3.1](#),

$$a_{\nu} = \langle \xi(H(\lambda)), s(\nu) \rangle = 0,$$

since $\xi(H(\lambda)) \in \bigoplus_{\mathbf{s} \leq \mathbf{t}} \mathfrak{F}_{\mathbf{s}}$, a contradiction. Thus, (1) holds, and hence $\xi(H(\lambda)) = \sum_{\mathbf{s} \leq \mathbf{t}} \sum_{\mu \in \mathcal{P}_{\mathbf{s}}} a_{\mu} H(\mu)$, so that

$$\begin{aligned} \pi_{\mathbf{t}}(\xi(H(\lambda))) &= \pi_{\mathbf{t}}\left(\sum_{\mathbf{s} \leq \mathbf{t}} \sum_{\mu \in \mathcal{P}_{\mathbf{s}}} a_{\mu} H(\mu)\right) \\ &= \sum_{\mathbf{s} \leq \mathbf{t}} \sum_{\mu \in \mathcal{P}_{\mathbf{s}}} a_{\mu} \pi_{\mathbf{t}}(H(\mu)) \\ &= \sum_{\mu \in \mathcal{P}_{\mathbf{t}}} a_{\mu} \pi_{\mathbf{t}}(H(\mu)), \end{aligned}$$

proving the second equality of (2). The first equality of (2) follows from the fact that ξ and $\pi_{\mathbf{t}}$ commute, since each $\mathfrak{F}_{\mathbf{s}}$ is invariant under ξ . \square

4. RESTRICTION TO LEVI SUBALGEBRA

Fix $n \in \mathbb{N}$ and a tuple $\mathbf{n} = (n_1, \dots, n_r)$ of positive integers such that $n_1 + \dots + n_r = n$, and for each $j = 0, 1, \dots, r$, let $\sigma_j = \sum_{i=1}^j n_i$.

We wish to consider the restrictions of the action of \mathbf{U} on the Fock spaces $\mathcal{F}_{\mathbf{s}}$ to the Levi subalgebra $\mathbf{U}(\mathbf{n})$, as defined in [subsection 2.2](#). The restrictions have natural direct sum decompositions; to better understand the summands, consider the example pictured in [Figure 4.1](#).

In view of the description of the action in terms of the abacus, it is natural to split the n -abacus into sections, each of which is represented by a gray region in [Figure 4.1](#). Formally, for each $i \in \mathbb{Z}$ and $j \in \{1, \dots, r\}$, let

$$\mathbb{Z}_{i,j}^{\mathbf{n}} = \{x \in \mathbb{Z} \mid \sigma_{j-1} \leq x - in < \sigma_j\}.$$




FIGURE 4.1. A 9-abacus display of the partition $\lambda = (13, 12, 10, 8, 8, 8, 6, 5, 5, 3, 2, 1, 1)$ and the corresponding vector in $(\wedge^3 V_4 \otimes V_2 \otimes V_3) \otimes (\wedge^2 V_4 \otimes V_2 \otimes \wedge^2 V_3) \otimes (\wedge^2 V_4 \otimes \wedge^2 V_3)$

Given a subset B of \mathbb{Z} , we write

$$X_{i,j}^{\mathbf{n}}(B) = (B \cap \mathbb{Z}_{i,j}^{\mathbf{n}}) - in - \sigma_{j-1} = \{x - in - \sigma_{j-1} \mid x \in B \cap \mathbb{Z}_{i,j}^{\mathbf{n}}\}.$$

Thus $X_{i,j}^{\mathbf{n}}(B)$ describes, in a normalized form, the beads lying in the section $\mathbb{Z}_{i,j}^{\mathbf{n}}$ in the n -abacus display of B . In addition, let $\mathbf{t}_B^{\mathbf{n}} : \mathbb{Z} \times \{1, \dots, r\} \rightarrow \mathbb{N}_0$ be defined by $\mathbf{t}_B^{\mathbf{n}}(i, j) = |B \cap \mathbb{Z}_{i,j}^{\mathbf{n}}| = |X_{i,j}^{\mathbf{n}}(B)|$, so that it describes the number of beads lying in each section in the n -abacus display of B . Note that B is a set of β -numbers if and only if

$$\sum_{\substack{i \geq 0 \\ 1 \leq j \leq r}} \mathbf{t}_B^{\mathbf{n}}(i, j) + \sum_{\substack{i < 0 \\ 1 \leq j \leq r}} (n_j - \mathbf{t}_B^{\mathbf{n}}(i, j)) < \infty,$$

in which case

$$\sum_{\substack{i \geq 0 \\ 1 \leq j \leq r}} \mathbf{t}_B^{\mathbf{n}}(i, j) - \sum_{\substack{i < 0 \\ 1 \leq j \leq r}} (n_j - \mathbf{t}_B^{\mathbf{n}}(i, j)) = \mathfrak{s}(B).$$

Let $\mathbf{T}_{\mathbf{n}}$ be the set of functions $\mathbf{t} : \mathbb{Z} \times \{1, \dots, r\} \rightarrow \mathbb{N}_0$ such that $\mathbf{t}(i, j) \leq n_j$ for all $i \in \mathbb{Z}$ and that $\sum_{\substack{i \geq 0 \\ 1 \leq j \leq r}} \mathbf{t}(i, j) + \sum_{\substack{i < 0 \\ 1 \leq j \leq r}} (n_j - \mathbf{t}(i, j)) < \infty$. For each $\mathbf{t} \in \mathbf{T}_{\mathbf{n}}$, let

$$\mathfrak{s}(\mathbf{t}) = \sum_{\substack{i \geq 0 \\ 1 \leq j \leq r}} \mathbf{t}(i, j) - \sum_{\substack{i < 0 \\ 1 \leq j \leq r}} (n_j - \mathbf{t}(i, j)).$$

Then for each $B \in \mathfrak{B}$, $\mathfrak{s}(B) = \mathfrak{s}(\mathbf{t}_B^{\mathbf{n}})$. For each $s \in \mathbb{Z}$, let $\mathbf{T}_{\mathbf{n},s} = \{\mathbf{t} \in \mathbf{T}_{\mathbf{n}} \mid \mathfrak{s}(\mathbf{t}) = s\}$.

We have a surjection from \mathfrak{B} to $\mathbf{T}_{\mathbf{n}}$ defined by $B \mapsto \mathbf{t}_B^{\mathbf{n}}$, and this restricts to give a surjection from \mathfrak{B}_s to $\mathbf{T}_{\mathbf{n},s}$ for each $s \in \mathbb{Z}$. Given $\mathbf{t} \in \mathbf{T}_{\mathbf{n}}$, let $\mathfrak{B}_{\mathbf{t}} = \{B \in \mathfrak{B} \mid \mathbf{t}_B^{\mathbf{n}} = \mathbf{t}\}$. Then the $\mathfrak{B}_{\mathbf{t}}$'s partition \mathfrak{B}_s as \mathbf{t} runs over $\mathbf{T}_{\mathbf{n},s}$.

Let $\mathbf{t} \in \mathbf{T}_{\mathbf{n}}$ and write $s = \mathfrak{s}(\mathbf{t})$. We now describe a characterisation of $B \in \mathfrak{B}_{\mathbf{t}}$ in terms of $\text{Par}(B)$. Define $B_{\mathbf{t}} \subseteq \mathbb{Z}$ as follows: $X_{i,j}^{\mathbf{n}}(B_{\mathbf{t}}) = \{0, \dots, \mathbf{t}(i, j) - 1\}$. Then $B_{\mathbf{t}} \in \mathfrak{B}_{\mathbf{t}}$, and write $\lambda_{\mathbf{t}}$ for $\text{Par}(B_{\mathbf{t}})$.

Proposition 4.1. *Let $B \in \mathfrak{B}_s$. Then $B \in \mathfrak{B}_{\mathbf{t}}$ if and only if $\text{Par}(B)$ and $\lambda_{\mathbf{t}}$ have the same sets of nodes of n -residue $\sigma_j - s$ for all $j = 0, \dots, r-1$.*

Proof. Note that $B_{\mathbf{t}}$ is the unique element in $\mathfrak{B}_{\mathbf{t}}$ such that $|\text{Par}(B_{\mathbf{t}})|$ is the least. We prove by induction on $|\text{Par}(B)|$. If $|\text{Par}(B)| \leq |\lambda_{\mathbf{t}}|$, then $B \in \mathfrak{B}_{\mathbf{t}}$ if and only if $B = B_{\mathbf{t}}$. On the other hand, $\lambda_{\mathbf{t}}$ is characterised by its sets of nodes of n -residue $\sigma_j - s$ for all $j = 0, \dots, r-1$ and that all its removable nodes lie among these. As such, any partition μ having the same sets of nodes of n -residue $\sigma_j - s$ as $\lambda_{\mathbf{t}}$ for all $j = 0, \dots, r-1$ must have $[\mu] \supseteq [\lambda]$ as otherwise λ would have a removable node not in $[\mu]$. Thus, if $|\text{Par}(B)| \leq |\lambda_{\mathbf{t}}|$, then $\text{Par}(B)$ has the same sets of nodes of n -residue $\sigma_j - s$ as $\lambda_{\mathbf{t}}$ for all $j = 0, \dots, r-1$ if and only if $\text{Par}(B) = \lambda_{\mathbf{t}}$, or equivalently $B = B_{\mathbf{t}}$. Thus the base case for the induction holds.

Suppose that $|\text{Par}(B)| > |\lambda_{\mathbf{t}}|$. If $B \in \mathfrak{B}_{\mathbf{t}}$, then there exists $x \in B$, $x-1 \notin B$ and $x \not\equiv_n \sigma_j$ for all j . Let $C = B \cup \{x-1\} \setminus \{x\}$. Then $C \in \mathfrak{B}_{\mathbf{t}}$, and $\text{Par}(C)$ is obtained from $\text{Par}(B)$ by removing a node of n -residue $x-s$. By induction, $\text{Par}(C)$ has the same sets of nodes of n -residue $\sigma_j - s$ as $\lambda_{\mathbf{t}}$ for all $j = 0, \dots, r-1$, and thus, so does $\text{Par}(B)$. On the other hand, if $\text{Par}(B)$ has the same sets of nodes of n -residue $\sigma_j - s$ as $\lambda_{\mathbf{t}}$ for all $j = 0, \dots, r-1$, then since $|\text{Par}(B)| > |\lambda_{\mathbf{t}}|$, we see that $\text{Par}(B)$ must have some removable node of n -residue not equal to $\sigma_j - s$ for all $j = 0, \dots, r-1$. Let μ be the partition obtained by removing this node. Then $\mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}} = \mathbf{t}_B^{\mathbf{n}}$, and μ has the same sets of nodes of n -residue $\sigma_j - s$ as $\lambda_{\mathbf{t}}$ for all $j = 0, \dots, r-1$, so that by induction, $\mathbf{t} = \mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}} = \mathbf{t}_B^{\mathbf{n}}$, and we are done. \square

We get the following immediate corollary.

Corollary 4.2. *Let $\lambda, \mu \in \mathcal{P}$ and $s \in \mathbb{Z}$. Then $\mathbf{t}_{\beta_s(\lambda)}^{\mathbf{n}} = \mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}$ if and only if λ and μ have the same sets of nodes of n -residue $\sigma_j - s$ for all $j = 0, \dots, r-1$.*

For each $\mathbf{t} \in \mathbf{T}_{\mathbf{n}}$, let $\mathcal{F}_{\mathbf{t}}$ be the $\mathbb{C}(q)$ -vector space with basis $\mathfrak{B}_{\mathbf{t}}$. Then $\mathcal{F}_{\mathbf{t}}$ is a vector subspace of \mathcal{F}_s if $\mathfrak{s}(\mathbf{t}) = s$, and is invariant under the action of $\mathbf{U}(\mathbf{n})$. Moreover, $\mathcal{F}_s = \bigoplus_{\mathbf{t} \in \mathbf{T}_{\mathbf{n},s}} \mathcal{F}_{\mathbf{t}}$.

In the special case where $\mathbf{n} = (n)$, we use the following notational conventions:

- $\mathbf{T}_{(n)} = \{\mathbf{t} : \mathbb{Z} \rightarrow \{0, 1, \dots, n\} \mid \sum_{i \geq 0} \mathbf{t}(i) + \sum_{i < 0} (n - \mathbf{t}(i)) < \infty\}$.
- If $\mathbf{t} \in \mathbf{T}_{(n)}$, then $\mathfrak{s}(\mathbf{t}) = \sum_{i \geq 0} \mathbf{t}(i) - \sum_{i < 0} (n - \mathbf{t}(i))$.
- If $B \subseteq \mathbb{Z}$, then $X_i^{(n)}(B) = \{x - in \mid x \in B, in \leq x < (i+1)n\}$ for all $i \in \mathbb{Z}$, and $\mathbf{t}_B^{(n)} : \mathbb{Z} \rightarrow \{0, 1, \dots, n\}$ sends i to $|X_i^{(n)}(B)|$.

We now proceed to identify $\mathcal{F}_{\mathbf{t}}$ as a $\mathbf{U}(\mathbf{n})$ -module. Let V_m denote the natural module for $U_q(\mathfrak{sl}_m) = \langle e_1, \dots, e_{m-1}; f_1, \dots, f_{m-1}; K_1^{\pm}, \dots, K_{m-1}^{\pm} \rangle$. It has distinguished basis $v_{m,0}, \dots, v_{m,m-1}$, with action given by

$$e_i v_{m,j} = \delta_{i,j} v_{m,j-1}, \quad f_i v_{m,j} = \delta_{i-1,j} v_{m,j+1}$$

and

$$K_i^+ v_{m,j} = \begin{cases} qv_{m,j} & \text{if } j = i - 1 \\ q^{-1}v_{m,j} & \text{if } j = i \\ v_{m,j} & \text{otherwise.} \end{cases}$$

The exterior power $\bigwedge^d V_m$ is an irreducible $U_q(\mathfrak{sl}_m)$ -module; its canonical basis is given by $\{v_{m,X} := v_{m,i_1} \wedge \dots \wedge v_{m,i_d} \mid X = \{i_1 < \dots < i_d\} \in \mathfrak{P}_{m,d}\}$, where here and hereafter $\mathfrak{P}_{m,d}$ denotes the set of d -element subsets of $\{0, 1, \dots, m-1\}$. So

$$\begin{aligned} e_i v_{m,X} &= \begin{cases} v_{m, X \cup \{i-1\} \setminus \{i\}} & \text{if } i \in X \text{ and } i-1 \notin X \\ 0 & \text{otherwise,} \end{cases} \\ f_i v_{m,X} &= \begin{cases} v_{m, X \cup \{i\} \setminus \{i-1\}} & \text{if } i-1 \in X \text{ and } i \notin X \\ 0 & \text{otherwise,} \end{cases} \\ K_i^+ v_{m,X} &= \begin{cases} qv_{m,X} & \text{if } i-1 \in X \text{ and } i \notin X \\ q^{-1}v_{m,X} & \text{if } i \in X \text{ and } i-1 \notin X \\ v_{m,X} & \text{otherwise.} \end{cases} \end{aligned}$$

It follows that for any tuple $\mathbf{d} = (d_1, \dots, d_r)$ of integers with $0 \leq d_j \leq n_j$ for all j , $\bigwedge_{\mathbf{n}}^{\mathbf{d}}(V) := \bigwedge^{d_1} V_{n_1} \otimes \dots \otimes \bigwedge^{d_r} V_{n_r}$ is an irreducible $\mathbf{U}(\mathbf{n})$ -module with canonical basis

$$C_{\mathbf{d}} = \{v_{n_1, X_1} \otimes \dots \otimes v_{n_r, X_r} \mid X_j \in \mathfrak{P}_{n_j, d_j} \ \forall j\}.$$

Lemma 4.3. *Let $\mathbf{t} \in \mathbf{T}_{\mathbf{n}}$. Write $\bigwedge^{\mathbf{t}}(V)$ for $\bigotimes_{i \in \mathbb{Z}, 1 \leq j \leq r} \bigwedge^{\mathbf{t}(i,j)} V_{n_j}$.*

- (1) *The set $\left\{ \bigotimes_{i \in \mathbb{Z}, 1 \leq j \leq r} v_{n_i, X_{i,j}} \mid X_{i,j} \in \mathfrak{P}_{n_j, \mathbf{t}(i,j)} \ \forall i, j \right\}$ is a basis for $\bigwedge^{\mathbf{t}}(V)$.*
- (2) *The map $\mathfrak{B}_{\mathbf{t}} \rightarrow \prod_{i \in \mathbb{Z}, 1 \leq j \leq r} \mathfrak{P}_{n_j, \mathbf{t}(i,j)}$ defined by $B \mapsto (X_{i,j}^{\mathbf{n}}(B))_{i \in \mathbb{Z}, 1 \leq j \leq r}$ for all $B \in \mathfrak{B}_{\mathbf{t}}$ is bijective.*
- (3) *The $\mathbb{C}(q)$ -linear map $\Theta_{\mathbf{t}} : \mathcal{F}_{\mathbf{t}} \rightarrow \bigwedge^{\mathbf{t}}(V)$ defined by $B \mapsto \bigotimes_{i \in \mathbb{Z}, 1 \leq j \leq r} v_{n_j, X_{i,j}^{\mathbf{n}}(B)}$ for all $B \in \mathfrak{B}_{\mathbf{t}}$ is bijective, and is an isomorphism of $\mathbf{U}(\mathbf{n})$ -modules if $\bigwedge^{\mathbf{t}}(V)$ is viewed as*

$$\dots \otimes \bigwedge_{\mathbf{n}}^{\mathbf{t}_{-1}}(V) \otimes \bigwedge_{\mathbf{n}}^{\mathbf{t}_0}(V) \otimes \bigwedge_{\mathbf{n}}^{\mathbf{t}_1}(V) \otimes \dots,$$

where $\mathbf{t}_i = (\mathbf{t}(i, 1), \dots, \mathbf{t}(i, r))$ for all $i \in \mathbb{Z}$.

Note that when $i \gg 0$ or $i \ll 0$, the component $\bigwedge^{\mathbf{t}(i,j)} V_{n_j}$, and hence $\bigwedge_{\mathbf{n}}^{\mathbf{t}_i}(V)$, is canonically isomorphic to the trivial module $\mathbb{C}(q)$, so the tensor product $\bigwedge^{\mathbf{t}}(V)$ is essentially a finite one.

Proof. This is straightforward to check. The reader may find [Figure 4.1](#) useful. \square

Each of the irreducible $\mathbf{U}(\mathbf{n})$ -modules $\bigwedge_{\mathbf{n}}^{\mathbf{d}}(V)$ with its canonical basis $C_{\mathbf{d}} = \{v_{\mathbf{X}} = v_{n_1, X_1} \otimes \cdots \otimes v_{n_r, X_r} \mid \mathbf{X} = (X_j)_{1 \leq j \leq r} \in \prod_{j=1}^r \mathfrak{P}_{n_j, d_j}\}$ is a based module. Hence, by Lusztig's theory, the tensor product $\bigwedge^{\mathbf{t}}(V) = \cdots \otimes \bigwedge_{\mathbf{n}}^{\mathbf{t}_{-1}}(V) \otimes \bigwedge_{\mathbf{n}}^{\mathbf{t}_0}(V) \otimes \bigwedge_{\mathbf{n}}^{\mathbf{t}_1}(V) \otimes \cdots$ is a based module with basis

$$\mathbf{C}^{\diamond} = \cdots \diamond C_{\mathbf{t}_{-1}} \diamond C_{\mathbf{t}_0} \diamond C_{\mathbf{t}_1} \diamond \cdots,$$

while $\mathbf{C}^{\otimes} = \cdots \otimes C_{\mathbf{t}_{-1}} \otimes C_{\mathbf{t}_0} \otimes C_{\mathbf{t}_1} \otimes \cdots$ is the basis given in Lemma 4.3(1). For each $B \in \mathfrak{B}_{\mathbf{t}}$, let $\mathbf{X}_i(B) = (X_{i,j}^{\mathbf{n}}(B))_{1 \leq j \leq r} \in \prod_{j=1}^r \mathfrak{P}_{n_j, \mathbf{t}(i,j)}$ for each $i \in \mathbb{Z}$, and let

$$\begin{aligned} v_B^{\otimes} &= \cdots \otimes v_{\mathbf{X}_{-1}(B)} \otimes v_{\mathbf{X}_0(B)} \otimes v_{\mathbf{X}_1(B)} \otimes \cdots \in \mathbf{C}^{\otimes}, \\ v_B^{\diamond} &= \cdots \diamond v_{\mathbf{X}_{-1}(B)} \diamond v_{\mathbf{X}_0(B)} \diamond v_{\mathbf{X}_1(B)} \diamond \cdots \in \mathbf{C}^{\diamond}. \end{aligned}$$

Then $\mathbf{C}^{\otimes} = \{v_B^{\otimes} \mid B \in \mathfrak{B}_{\mathbf{t}}\} = \{\Theta_{\mathbf{t}}(B) \mid B \in \mathfrak{B}_{\mathbf{t}}\}$ by Lemma 4.3(3). Also, $\mathbf{C}^{\diamond} = \{v_B^{\diamond} \mid B \in \mathfrak{B}_{\mathbf{t}}\}$.

For each $B \in \mathfrak{B}_{\mathbf{t}}$, let $G_{\mathbf{t}}(B) = \Theta_{\mathbf{t}}^{-1}(v_B^{\diamond})$. Denote by $\psi_{\mathbf{t}}$ the $\mathbb{C}(q)$ -semilinear involution of $\mathcal{F}_{\mathbf{t}}$ fixing $G_{\mathbf{t}}(B)$ for each $B \in \mathfrak{B}_{\mathbf{t}}$.

Let $s = \mathfrak{s}(\mathbf{t})$. Recall that $\beta_s : \mathcal{P} \rightarrow \mathfrak{B}_s$ is a bijection, with inverse Par . Let $\mathcal{P}_{\mathbf{t}} = \text{Par}(\mathfrak{B}_{\mathbf{t}})$, so that the following three statements are equivalent:

$$(1) \mu \in \mathcal{P}_{\mathbf{t}}; \quad (2) \beta_s(\mu) \in \mathfrak{B}_{\mathbf{t}}; \quad (3) \mathbf{t}_{\beta_s(\mu)} = \mathbf{t}.$$

For each $\mu \in \mathcal{P}_{\mathbf{t}}$, we write $G_{\mathbf{t}}(\mu)$ for $G_{\mathbf{t}}(\beta_s(\mu))$. In addition define $d_{\lambda\mu}^{\mathbf{t}}(q) \in \mathbb{C}(q)$ for $\lambda, \mu \in \mathcal{P}_{\mathbf{t}}$ by

$$G_{\mathbf{t}}(\mu) = \sum_{\lambda \in \mathcal{P}_{\mathbf{t}}} d_{\lambda\mu}^{\mathbf{t}}(q) \beta_s(\lambda).$$

From Lemma 2.3(1) we deduce the following.

Lemma 4.4. *Let $\lambda, \mu \in \mathcal{P}_{\mathbf{t}}$.*

- (1) $d_{\mu\mu}^{\mathbf{t}}(q) = 1$.
- (2) *If $\lambda \neq \mu$, then $d_{\lambda\mu}^{\mathbf{t}}(q) \in q\mathbb{Z}[q]$.*
- (3) *If $d_{\lambda\mu}^{\mathbf{t}}(q) \neq 0$, then $\mu \geq \lambda$.*
- (4) *If $m \in \mathcal{F}_{\mathbf{t}}$ satisfies $\psi_{\mathbf{t}}(m) = m$ and $m - \beta_s(\mu) \in \sum_{B \in \mathfrak{B}_{\mathbf{t}}} q\mathbb{Z}[q]B$, then $m = G_{\mathbf{t}}(\mu)$.*

Proposition 4.5. *Let $\mathbf{t} \in \mathbf{T}_{\mathbf{n}}$. For each $j = 1, \dots, r$, define $\mathbf{t}_j : \mathbb{Z} \rightarrow \mathbb{N}_0$ by $\mathbf{t}_j(i) = \mathbf{t}(i, j)$.*

- (1) *For each $j = 1, \dots, r$, $\mathbf{t}_j \in \mathbf{T}_{(n_j)}$. In addition, $\sum_{j=1}^r \mathfrak{s}(\mathbf{t}_j) = \mathfrak{s}(\mathbf{t})$.*
- (2) *The $\mathbb{C}(q)$ -linear map $\Phi_{\mathbf{t}} : \mathcal{F}_{\mathbf{t}} \rightarrow \mathcal{F}_{\mathbf{t}_1} \otimes \cdots \otimes \mathcal{F}_{\mathbf{t}_r}$ defined by $\Phi_{\mathbf{t}}(B) = B_1 \otimes \cdots \otimes B_r$ for all $B \in \mathfrak{B}_{\mathbf{t}}$, where $B_j = \{x + in_j \mid i \in \mathbb{Z}, x \in X_{i,j}^{\mathbf{n}}(B)\}$ for all j , is a well-defined isomorphism of $\mathbf{U}(\mathbf{n})$ -modules (where we identify $\mathbf{U}(\mathbf{n})$ with $U_q(\mathfrak{sl}_{n_1}) \otimes \cdots \otimes U_q(\mathfrak{sl}_{n_r})$ in the obvious way), and*

$$\Phi_{\mathbf{t}}(G_{\mathbf{t}}(B)) = G_{\mathbf{t}_1}(B_1) \otimes \cdots \otimes G_{\mathbf{t}_r}(B_r)$$

for all $B \in \mathfrak{B}_{\mathbf{t}}$.

Proof. We show only the last equality; all other assertions can be easily verified.

By Lemma 4.3(2), we have, for each $j = 1, \dots, r$, a bijective $\mathbb{C}(q)$ -linear map (actually an isomorphism of $U_q(\mathfrak{sl}_{n_j})$ -modules) $\Theta_{\mathbf{t}_j} : \mathcal{F}_{\mathbf{t}_j} \rightarrow \bigotimes_{i \in \mathbb{Z}} \wedge^{\mathbf{t}_j(i)} V_{n_j}$ sending B_j to $\bigotimes_{i \in \mathbb{Z}} v_{n_j, X_i^{(n_j)}(B_j)}$ for all $B_j \in \mathfrak{B}_{\mathbf{t}_j}$. Since

$$\begin{aligned} ((\Theta_{\mathbf{t}_1}^{-1} \otimes \dots \otimes \Theta_{\mathbf{t}_r}^{-1}) \circ \Theta_{\mathbf{t}})(B) &= (\Theta_{\mathbf{t}_1}^{-1} \otimes \dots \otimes \Theta_{\mathbf{t}_r}^{-1}) \left(\bigotimes_{\substack{i \in \mathbb{Z} \\ 1 \leq j \leq r}} v_{n_j, X_{i,j}^{\mathbf{n}}(B)} \right) \\ &= \bigotimes_{j=1}^r \Theta_{\mathbf{t}_j}^{-1} \left(\bigotimes_{i \in \mathbb{Z}} v_{n_j, X_i^{(n_j)}(B_j)} \right) = \bigotimes_{j=1}^r B_j = \Phi_{\mathbf{t}}(B) \end{aligned}$$

for all $B \in \mathfrak{B}_{\mathbf{t}}$, we see that $\Phi_{\mathbf{t}} = (\Theta_{\mathbf{t}_1}^{-1} \otimes \dots \otimes \Theta_{\mathbf{t}_r}^{-1}) \circ \Theta_{\mathbf{t}}$. Thus

$$\begin{aligned} \Phi_{\mathbf{t}}(G_{\mathbf{t}}(B)) &= (\Theta_{\mathbf{t}_1}^{-1} \otimes \dots \otimes \Theta_{\mathbf{t}_r}^{-1})(\Theta_{\mathbf{t}}(G_{\mathbf{t}}(B))) \\ &= (\Theta_{\mathbf{t}_1}^{-1} \otimes \dots \otimes \Theta_{\mathbf{t}_r}^{-1})(v_B^{\diamond}) \\ &= (\Theta_{\mathbf{t}_1}^{-1} \otimes \dots \otimes \Theta_{\mathbf{t}_r}^{-1}) \left(\bigotimes_{i \in \mathbb{Z}} \left(\bigotimes_{j=1}^r v_{n_j, X_{i,j}^{\mathbf{n}}(B)} \right) \right) \\ &= (\Theta_{\mathbf{t}_1}^{-1} \otimes \dots \otimes \Theta_{\mathbf{t}_r}^{-1}) \left(\bigotimes_{j=1}^r \left(\bigotimes_{i \in \mathbb{Z}} v_{n_j, X_i^{(n_j)}(B_j)} \right) \right) \\ &= \bigotimes_{j=1}^r \Theta_{\mathbf{t}_j}^{-1} \left(\bigotimes_{i \in \mathbb{Z}} v_{n_j, X_i^{(n_j)}(B_j)} \right) \\ &= \bigotimes_{j=1}^r G_{\mathbf{t}_j}(B_j), \end{aligned}$$

using Proposition 2.4 for the fourth equality. \square

5. COMPARISON OF CANONICAL BASES

Fix $n \in \mathbb{N}$ and a tuple $\mathbf{n} = (n_1, \dots, n_r)$ of positive integers such that $n_1 + \dots + n_r = n$. Recall our subalgebra $\mathbf{U}(\mathbf{n})$ of \mathbf{U} and our set $\mathbf{T}_{\mathbf{n}}$ defined in the last section. We define a partial order \geq on $\mathbf{T}_{\mathbf{n}}$ as follows: for $\mathbf{t}, \mathbf{t}' \in \mathbf{T}_{\mathbf{n}}$, we have $\mathbf{t} > \mathbf{t}'$ if and only if $\mathfrak{s}(\mathbf{t}) = \mathfrak{s}(\mathbf{t}')$ and there exists $(i_0, j_0) \in \mathbb{Z} \times \{1, \dots, r\}$ such that $\mathbf{t}(i_0, j_0) > \mathbf{t}'(i_0, j_0)$ and $\mathbf{t}(i, j) = \mathbf{t}'(i, j)$ for all (i, j) with $i > i_0$ or both $i = i_0$ and $j > j_0$. Note that \geq restricts to a total order on $\mathbf{T}_{\mathbf{n},s}$ for each $s \in \mathbb{Z}$.

Lemma 5.1. *Let $\lambda, \mu \in \mathcal{P}$. If $\lambda \geq_{J_n} \mu$, then $\mathbf{t}_{\beta_s(\lambda)}^{\mathbf{n}} \geq \mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}$ for all $s \in \mathbb{Z}$.*

Proof. Note that $B \rightarrow_n C$ implies $\mathbf{t}_B^n \geq \mathbf{t}_C^n$. \square

For $\mathbf{t} \in \mathbf{T}_{\mathbf{n},s}$ let $\pi_{\mathbf{t}} : \mathcal{F}_s \rightarrow \mathcal{F}_{\mathbf{t}}$ be the natural projection defined by $\pi_{\mathbf{t}}(B) = B$ if $B \in \mathfrak{B}_{\mathbf{t}}$ and zero otherwise.

Since, for each $s \in \mathbb{Z}$, $\mathcal{P} = \bigcup_{\mathbf{t} \in \mathbf{T}_{\mathbf{n},s}} \mathcal{P}_{\mathbf{t}}$ is a partition of \mathcal{P} , and the order \geq on $\mathbf{T}_{\mathbf{n},s}$ defined above respects the Jantzen order \geq_{J_n} by Lemma 5.1, we can apply Lemma 3.4 with $H(\lambda) = G(\lambda)$ and ξ an element of $\mathbf{U}(\mathbf{n})$, or $H(\lambda) = [P(\lambda)]$ and ξ an i -Ind functor.

Theorem 5.2. *Let $s \in \mathbb{Z}$ and $\mathbf{t} \in \mathbf{T}_{\mathbf{n},s}$. If $\mu \in \mathcal{P}_{\mathbf{t}}$, then $\pi_{\mathbf{t}}(G_s(\mu)) = G_{\mathbf{t}}(\mu)$. Equivalently $d_{\lambda\mu}(q) = d_{\lambda\mu}^{\mathbf{t}}(q)$ for all $\lambda, \mu \in \mathcal{P}_{\mathbf{t}}$.*

Proof. We prove by induction on $|\mu|$ the following equivalent statement: if $\mu \in \mathcal{P}$, then $\pi_{\mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}}(G_s(\mu)) = G_{\mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}}(\mu)$ for all $s \in \mathbb{Z}$. If $\mu = \emptyset$, then $G(\mu) = \mu$ by Theorem 2.1(1,3), while $G_{\mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}}(\mu) = \beta_s(\mu)$ by Lemma 4.4(1,3), so that the result holds. So let $\mu \neq \emptyset$ and assume that the statement holds for all partitions $\tilde{\mu}$ such that $|\tilde{\mu}| < |\mu|$, or both $|\tilde{\mu}| = |\mu|$ and $\tilde{\mu} < \mu$. Let $s \in \mathbb{Z}$ and let $\mathbf{t} = \mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}$. Let $c = \max\{i \in \mathbb{Z} \mid \mathbf{t}(i, j) > 0 \text{ for some } j\}$, so that row c is the bottommost row that contains some bead in the n -abacus display of $\beta_s(\mu)$. There are two cases to consider.

If $X_{c,j}^{\mathbf{n}}(\beta_s(\mu)) = \{0, \dots, \mathbf{t}(c, j) - 1\}$ for all j (i.e. the beads occurring in each section of row c in the n -abacus display of $\beta_s(\mu)$ occupy the leftmost positions), then $v_{n_1, X_{c,1}^{\mathbf{n}}(\beta_s(\mu))} \otimes \dots \otimes v_{n_r, X_{c,r}^{\mathbf{n}}(\beta_s(\mu))}$ is the highest weight vector for the irreducible $\mathbf{U}(\mathbf{n})$ -module $\bigwedge^{\mathbf{t}(c,1)}(V_{n_1}) \otimes \dots \otimes \bigwedge^{\mathbf{t}(c,r)}(V_{n_r})$.

Let $\lambda \in \mathcal{P}_{\mathbf{t}}$. Note that if $\lambda > \mu$, then $d_{\lambda\mu}(q) = 0$, by Theorem 2.1(3), and $d_{\lambda\mu}^{\mathbf{t}}(q) = 0$ by Lemma 4.4(3). So we may assume that $\lambda \leq \mu$. In this case, the beads in each section of row c of the n -abacus display of $\beta_s(\lambda)$ must be occupying the leftmost positions too, i.e. $X_{c,j}^{\mathbf{n}}(\beta_s(\lambda)) = X_{c,j}^{\mathbf{n}}(\beta_s(\mu))$ for all j . Let $\mathbf{t}' \in \mathbf{T}_{\mathbf{n}}$ such that

$$\mathbf{t}'(i, j) = \begin{cases} \mathbf{t}(i, j), & \text{if } i \neq c; \\ 0, & \text{if } i = c. \end{cases}$$

Write $s' = \mathfrak{s}(\mathbf{t}')$. Let $\tilde{\lambda}, \tilde{\mu} \in \mathcal{P}$ such that $\beta_{s'}(\tilde{\lambda})$ and $\beta_{s'}(\tilde{\mu})$ are obtained by removing the beads in row c from the n -abacus displays of $\beta_s(\lambda)$ and $\beta_s(\mu)$ respectively. Then $\tilde{\lambda}, \tilde{\mu} \in \mathcal{P}_{\mathbf{t}'}$ and are the partitions obtained from λ and μ respectively by removing the first $\sum_{j=1}^r \mathbf{t}(c, j)$ parts, so that $|\tilde{\mu}| < |\mu|$. We have

$$d_{\lambda\mu}(q) = d_{\tilde{\lambda}\tilde{\mu}}(q) = d_{\tilde{\lambda}\tilde{\mu}}^{\mathbf{t}'}(q) = d_{\lambda\mu}^{\mathbf{t}}(q),$$

where the first equality is ‘row removal’ [CMT1, Theorem 1(1)], the second by induction and the last by Lemma 2.3(2).

We now proceed to the second case. In this case, there is a bead in row c of the n -abacus display of $\beta_s(\mu)$ having a vacant preceding position that is in the same section as the bead. Formally, there exists $k \in \beta_s(\mu)$ such that

$\lfloor \frac{k}{n} \rfloor = c$, $k-1 \notin \beta_s(\mu)$, and $f_k \in \mathbf{U}(\mathbf{n})$. Let $\lambda = \text{Par}(\beta_s(\mu) \cup \{k-1\} \setminus \{k\})$. Then $\lambda \in \mathcal{P}_{\mathbf{t}}$, and we have, by [Corollary 3.3\(2\)](#),

$$f_k(G_s(\lambda)) = G_s(\mu) + \sum_{\substack{\nu \in \mathcal{P} \\ |\nu| = |\mu| \\ \nu < \mu}} L_{\lambda}^{\nu}(q) G_s(\nu),$$

with $L_{\lambda}^{\nu}(q) = L_{\lambda}^{\nu}(q^{-1})$ for all ν by [Theorem 2.2](#). Applying [Lemma 3.4](#) with $\xi = f_k$, we get

$$f_k(\pi_{\mathbf{t}}(G_s(\lambda))) = \pi_{\mathbf{t}}(f_k(G_s(\lambda))) = \pi_{\mathbf{t}}(G_s(\mu)) + \sum_{\substack{\nu \in \mathcal{P}_{\mathbf{t}} \\ |\nu| = |\mu| \\ \nu < \mu}} L_{\lambda}^{\nu}(q) \pi_{\mathbf{t}}(G_s(\nu)).$$

Applying the induction hypothesis to λ and ν , we get

$$\pi_{\mathbf{t}}(G_s(\mu)) = f_k(G_{\mathbf{t}}(\lambda)) - \sum_{\substack{\nu \in \mathcal{P}_{\mathbf{t}} \\ |\nu| = |\mu| \\ \nu < \mu}} L_{\lambda}^{\nu}(q) G_{\mathbf{t}}(\nu).$$

Thus, $\pi_{\mathbf{t}}(G_s(\mu))$ is $\psi_{\mathbf{t}}$ -invariant. Furthermore, since $G(\mu) - \mu \in \bigoplus_{\nu \in \mathcal{P}} q\mathbb{Z}[q]\nu$, we see that

$$G_s(\mu) - \beta_s(\mu) = \beta_s(G(\mu) - \mu) \in \bigoplus_{\nu \in \mathcal{P}} q\mathbb{Z}[q]\beta_s(\nu) = \bigoplus_{B \in \mathfrak{B}_s} q\mathbb{Z}[q]B,$$

so that

$$\pi_{\mathbf{t}}(G_s(\mu)) - \beta_s(\mu) = \pi_{\mathbf{t}}(G_s(\mu) - \beta_s(\mu)) \in \bigoplus_{B \in \mathfrak{B}_{\mathbf{t}}} q\mathbb{Z}[q]B.$$

Hence $\pi_{\mathbf{t}}(G_s(\mu)) = G_{\mathbf{t}}(\mu)$ by [Lemma 4.4\(4\)](#), as desired. \square

Corollary 5.3. *Let $\mathbf{t} \in \mathbf{T}_{\mathbf{n}}$, and let $\lambda, \mu \in \mathcal{P}_{\mathbf{t}}$. Then $d_{\lambda\mu}^{\mathbf{t}}(q) \neq 0$ only if $\mu \geq_{J_n} \lambda$.*

Proof. This follows immediately from [Theorem 5.2](#) and [Theorem 2.1\(3\)](#). \square

Corollary 5.4. *Let $s \in \mathbb{Z}$ and $\mathbf{t} \in \mathbf{T}_{\mathbf{n},s}$. Let $\lambda \in \mathcal{P}_{\mathbf{t}}$, and let $\xi \in \mathbf{U}(\mathbf{n})$. Suppose that $\xi(G_s(\lambda)) = \sum_{\mu \in \mathcal{P}} a_{\mu}(q) G_s(\mu)$. Then*

$$\xi(G_{\mathbf{t}}(\lambda)) = \sum_{\mu \in \mathcal{P}_{\mathbf{t}}} a_{\mu}(q) G_{\mathbf{t}}(\mu).$$

Proof. This follows immediately from [Lemma 3.4](#) and [Theorem 5.2](#). \square

The next result may be regarded as a ‘runner removal’ theorem.

Theorem 5.5. *Let $s \in \mathbb{Z}$ and $\mathbf{t} \in \mathbf{T}_{\mathbf{n},s}$. Let $\lambda, \mu \in \mathcal{P}_{\mathbf{t}}$. For each $j = 1, \dots, r$, let $s_j = \sum_{i \geq 0} \mathbf{t}(i, j) + \sum_{i < 0} (\mathbf{t}(i, j) - n_j)$, and let $\lambda^{(j)}, \mu^{(j)} \in \mathcal{P}$ such that*

$$\begin{aligned} \beta_{s_j}(\lambda^{(j)}) &= \{x + in_j \mid i \in \mathbb{Z}, x \in X_{i,j}^{\mathbf{n}}(\beta_s(\lambda))\}, \\ \beta_{s_j}(\mu^{(j)}) &= \{x + in_j \mid i \in \mathbb{Z}, x \in X_{i,j}^{\mathbf{n}}(\beta_s(\mu))\}. \end{aligned}$$

(Thus, $\lambda^{(j)}$ is the partition read off from the n_j -abacus consisting of runners i for $\sigma_{j-1} \leq i < \sigma_j$ in the n -abacus display of $\beta_s(\lambda)$. Similarly for $\mu^{(j)}$.) Then

$$d_{\lambda\mu}^n(q) = \prod_{j=1}^r d_{\lambda^{(j)}\mu^{(j)}}^{n_j}(q).$$

Proof. For each $j = 1, \dots, r$, define $\mathbf{t}_j : \mathbb{Z} \rightarrow \mathbb{N}_0$ by $\mathbf{t}_j(i) = \mathbf{t}(i, j)$ for all $i \in \mathbb{Z}$. By [Proposition 4.5](#), we have

$$\Phi_{\mathbf{t}}(\beta_s(\mu)) = \bigotimes_{j=1}^r \beta_{s_j}(\mu^{(j)}) \quad \text{and} \quad \Phi_{\mathbf{t}}(G_{\mathbf{t}}(\mu)) = \bigotimes_{j=1}^r G_{\mathbf{t}_j}(\mu^{(j)})$$

for all $\mu \in \mathcal{P}_{\mathbf{t}}$. Thus, $d_{\lambda\mu}^{\mathbf{t}}(q) = \prod_{j=1}^r d_{\lambda^{(j)}\mu^{(j)}}^{\mathbf{t}_j}(q)$. Now apply [Theorem 5.2](#). \square

We end this section with a discussion of the case where $\mathbf{n} = (n_1, \dots, n_r)$ has $n_j = 2$ for some j . We first look at the special case where $\mathbf{n} = (2)$. In this case, since $\bigwedge^1 V_2 = V_2$ and $\bigwedge^0 V_2 = \bigwedge^2 V_2 = \mathbb{C}(q)$, for each $\mathbf{t} \in \mathbf{T}_{\mathbf{n}}$, $\mathcal{F}_{\mathbf{t}}$ is isomorphic to a tensor power $V_2^{\otimes d}$ as $U_q(\mathfrak{sl}_2)$ -modules. The latter has been studied extensively by Frenkel and Khovanov in [\[FK\]](#). In particular, they provide closed formulas for v_B^\diamond in terms of graphical calculus combinatorics for each $B \in \mathfrak{B}_{\mathbf{t}}$, so that there are closed formulas for $d_{\lambda\mu}^{\mathbf{t}}(q)$ for $\lambda, \mu \in \mathcal{P}_{\mathbf{t}}$. They also describe $f(v_B^\diamond)$, which we translate to our language as follows:

Theorem 5.6 ([\[FK\]](#)). *Let $\mathbf{n} = (2)$, and let $\lambda \in \mathcal{P}$. Let $s \in \mathbb{Z}$ and write $\mathbf{t} = \mathbf{t}_{\beta_s(\lambda)}^{\mathbf{n}}$. Suppose that*

$$f(G_{\mathbf{t}}(\lambda)) = \sum_{\mu \in \mathcal{P}_{\mathbf{t}}} L_{\lambda}^{\mu}(q) G_{\mathbf{t}}(\mu).$$

Then

$$L_{\lambda}^{\mu}(q) = \begin{cases} [1 + n_{\lambda\mu}]_q, & \text{if } \lambda \text{ can be obtained from } \mu \\ & \text{by removing a normal node } \mathbf{n}_{\lambda\mu} \text{ of 2-residue } s+1; \\ 0, & \text{otherwise.} \end{cases}$$

Here $n_{\lambda\mu}$ is the number of normal nodes of μ with 2-residue $s+1$ and to the right of $\mathbf{n}_{\lambda\mu}$.

Proof. We briefly describe our translation of the results in [\[FK\]](#). The basis vector $v_{2,1}$ of V_2 , represented by an up arrow in the diagrams in [\[FK\]](#), corresponds to a removable node of 2-residue $s+1$, while the other basis vector $v_{2,0}$, represented by a down arrow in the diagrams in [\[FK\]](#), corresponds to an addable node of 2-residue $s+1$. An arc connecting the up and down arrows induces a pairing of the corresponding removable node with the addable node. We should point out that the description in [\[FK, p.445-7\]](#) is given in terms of the basis dual to $\{G_{\mathbf{t}}(\lambda)\}$ with respect to an inner product for which e and f are adjoint operators, and so we use their description of the action of e rather than that of f . \square

Corollary 5.7. *Let $s \in \mathbb{Z}$, and $\lambda \in \mathcal{P}$. Write \mathbf{t} for $\mathbf{t}_{\beta_s(\lambda)}^{\mathbf{n}}$. Suppose that $\mathbf{n} = (n_1, \dots, n_r)$ has $n_d = 2$ for some d , and let $k = \sum_{j=1}^{d-1} n_j + 1$. Let*

$$f_k(G_{\mathbf{t}}(\lambda)) = \sum_{\mu \in \mathcal{P}_{\mathbf{t}}} L_{\lambda}^{\mu}(q) G_{\mathbf{t}}(\mu).$$

Then

$$L_{\lambda}^{\mu}(q) = \begin{cases} [1 + n_{\lambda\mu}]_q, & \text{if } \lambda \text{ can be obtained from } \mu \\ & \text{by removing a normal node } \mathbf{n}_{\lambda\mu} \text{ of } n\text{-residue } k - s; \\ 0, & \text{otherwise.} \end{cases}$$

Here $n_{\lambda\mu}$ is the number of normal nodes of μ with n -residue $k - s$ and to the right of $\mathbf{n}_{\lambda\mu}$.

Proof. By [Proposition 4.5](#) and adopting the notations there with $\lambda = \text{Par}(B)$ and $\lambda^{(i)} = \text{Par}(B_i)$, we have

$$\begin{aligned} f_k(G_{\mathbf{t}}(\lambda)) &= \Phi_{\mathbf{t}}^{-1}(G_{\mathbf{t}_1}(\lambda^{(1)}) \otimes \dots \otimes G_{\mathbf{t}_{d-1}}(\lambda^{(d-1)}) \otimes \\ &\quad f(G_{\mathbf{t}_d}(\lambda^{(d)}) \otimes G_{\mathbf{t}_{d+1}}(\lambda^{(d+1)}) \otimes \dots \otimes G_{\mathbf{t}_r}(\lambda^{(r)})) \\ &= \Phi_{\mathbf{t}}^{-1}(G_{\mathbf{t}_1}(\lambda^{(1)}) \otimes \dots \otimes G_{\mathbf{t}_{d-1}}(\lambda^{(d-1)}) \otimes \\ &\quad \sum_{\mu^{(d)} \in \mathcal{P}_{\mathbf{t}_d}} L_{\lambda^{(d)}}^{\mu^{(d)}}(q) (G_{\mathbf{t}_d}(\mu^{(d)}) \otimes G_{\mathbf{t}_{d+1}}(\lambda^{(d+1)}) \otimes \dots \otimes G_{\mathbf{t}_r}(\lambda^{(r)})) \end{aligned}$$

The result thus follows from [Theorem 5.6](#). \square

Corollary 5.8. *Let $\lambda, \mu \in \mathcal{P}$. Suppose that there exists a residue class i modulo n such that λ and μ has the same sets of nodes of n -residue $i - 1$ and $i + 1$. Let $s \in \mathbb{Z}$. Writing $f_{s+i}(G_s(\lambda)) = \sum_{\nu \in \mathcal{P}} L_s^{\nu}(q) G_s(\nu)$, we have*

$$L_s^{\mu}(q) = \begin{cases} [1 + n_{\lambda\mu}]_q, & \text{if } \lambda \text{ can be obtained from } \mu \\ & \text{by removing a normal node } \mathbf{n}_{\lambda\mu} \text{ of } n\text{-residue } i; \\ 0, & \text{otherwise.} \end{cases}$$

Here, $n_{\lambda\mu}$ is the number of normal nodes of μ with n -residue i and to the right of $\mathbf{n}_{\lambda\mu}$.

Proof. Let $\mathbf{n} = (2, n - 2)$, and let $x = 1 - i$. By [Corollary 4.2](#), our conditions on λ and μ force $\mathbf{t}_{\beta_x(\lambda)}^{\mathbf{n}} = \mathbf{t}_{\beta_x(\mu)}^{\mathbf{n}}$. Thus, by [Corollary 5.4](#) and [Corollary 5.7](#) (with $k = 1$), we see that the result holds for $L_x^{\nu}(x)$. To deal with a general s , let $\theta_{ab} : \mathcal{F}_a \rightarrow \mathcal{F}_b$ ($a, b \in \mathbb{Z}$) be the $\mathbb{C}(q)$ -linear isomorphism defined by $\theta_{ab}(\beta_a(\nu)) = \beta_b(\nu)$ for all $\nu \in \mathcal{P}$. Then $\theta_{ab}(G_a(\lambda)) = G_b(\nu)$ for all $\nu \in \mathcal{P}$. Furthermore θ_{ab} intertwines f_r with f_{r+b-a} for all residue class r modulo n . Thus, applying θ_{xs} to $f_1(G_x(\lambda)) = \sum_{\nu \in \mathcal{P}} L_x^{\nu}(q) G_x(\nu)$, we get $f_{s+i}(G_s(\lambda)) = \sum_{\nu \in \mathcal{P}} L_s^{\nu}(q) G_s(\nu)$, and the result follows. \square

6. DECOMPOSITION NUMBERS

Recall that we identified the Grothendieck group $K_0(\bigoplus_m \mathbf{S}_m\text{-mod})$ with the classical (non-quantized) Fock space \mathcal{F}_c . Recall also the A -lattice $\mathcal{F}_{\mathbb{Z},A}$ (where $A = \mathbb{Z}[q, q^{-1}]$) and the \mathbb{Z} -linear map $\varepsilon_{q=1} : \mathcal{F}_{\mathbb{Z},A} \rightarrow \mathcal{F}_c$ which intertwines the Chevalley generators of \mathbf{U}_A^- with the i -Ind functors.

We assume in this section that $n = p$, the characteristic over which the Schur algebras \mathbf{S}_m are defined. Let $\mathbf{n} = (n_1, \dots, n_r)$ be a tuple of positive integers summing to p . For each $\mathbf{t} \in \mathbf{T}_{\mathbf{n}}$, define $\varrho_{\mathbf{t}} : \mathcal{F}_c \rightarrow \mathcal{F}_c$ to be the natural projection such that $\varrho_{\mathbf{t}}(\lambda) = \lambda$ if $\lambda \in \mathcal{P}_{\mathbf{t}}$ and zero otherwise, and we write $\mathcal{F}_{c,\mathbf{t}}$ for the image of $\varrho_{\mathbf{t}}$. In addition, if M is a module of a Schur algebra, we write $[M]_{\mathbf{t}}$ for $\varrho_{\mathbf{t}}[M]$.

Theorem 6.1. *Let $\mathbf{n} = (n_1, \dots, n_r)$ be a tuple of positive integers summing to p . Suppose that $n_j \in \{1, 2\}$ for all j , so that $\mathbf{U}(\mathbf{n})$ is isomorphic to a tensor product of copies of $U_q(\mathfrak{sl}_2)$. Let $\mathbf{t} \in \mathbf{T}_{\mathbf{n}}$. Then for all $\mu \in \mathcal{P}_{\mathbf{t}}$, we have $[P(\mu)]_{\mathbf{t}} = \varepsilon_{q=1}(G_{\mathbf{t}}(\mu))$. Equivalently, $d_{\lambda\mu} = d_{\lambda\mu}^{\mathbf{t}}(1)$ for all $\lambda, \mu \in \mathcal{P}_{\mathbf{t}}$.*

Proof. We follow an inductive argument reminiscent of the proof of [Theorem 5.2](#). Once again, we prove the following equivalent statement: if $\mu \in \mathcal{P}$, then $[P(\mu)]_{\mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}} = \varepsilon_{q=1}(G_{\mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}}(\mu))$ for all $s \in \mathbb{Z}$. If $\mu = \emptyset$ the result is clear. So let $\mu \neq \emptyset$ and assume the statement holds for all partitions $\tilde{\mu}$ such that $|\tilde{\mu}| < |\mu|$, or both $|\tilde{\mu}| = |\mu|$ and $\tilde{\mu} < \mu$. Let $s \in \mathbb{Z}$ and let $\mathbf{t} = \mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}$. Let $c = \max\{i \in \mathbb{Z} \mid \mathbf{t}(i, j) > 0 \text{ for some } j\}$. There are two cases to consider.

If $X_{c,j}^{\mathbf{n}}(\beta_s(\mu)) = \{0, \dots, \mathbf{t}(c, j) - 1\}$ for all j (i.e. the beads occurring in each section of row b in the n -abacus display of B occupy the leftmost positions), then $v_{n_1, X_{c,1}^{\mathbf{n}}(\beta_s(\mu))} \otimes \dots \otimes v_{n_r, X_{c,r}^{\mathbf{n}}(\beta_s(\mu))}$ is the highest weight vector for the irreducible $\mathbf{U}(\mathbf{n})$ -module $\bigwedge^{\mathbf{t}(c,1)}(V_{n_1}) \otimes \dots \otimes \bigwedge^{\mathbf{t}(c,r)}(V_{n_r})$.

Let $\lambda \in \mathcal{P}_{\mathbf{t}}$. Note that if $\lambda > \mu$, then $d_{\lambda\mu} = 0$ by [Proposition 2.5\(2\)](#), and $d_{\lambda\mu}^{\mathbf{t}}(q) = 0$ by [Lemma 4.4\(3\)](#). So we may assume that $\lambda \leq \mu$. In this case, $X_{c,j}^{\mathbf{n}}(\beta_s(\lambda)) = X_{c,j}^{\mathbf{n}}(\beta_s(\mu))$ for all j . Let $\mathbf{t}' \in \mathbf{T}_{\mathbf{n}}$ such that

$$\mathbf{t}'(i, j) = \begin{cases} \mathbf{t}(i, j), & \text{if } i \neq c; \\ 0, & \text{if } i = c. \end{cases}$$

Write $s' = \mathfrak{s}(\mathbf{t}')$. Let $\tilde{\lambda}, \tilde{\mu} \in \mathcal{P}$ such that $\beta_{s'}(\tilde{\lambda})$ and $\beta_{s'}(\tilde{\mu})$ may be obtained by removing the beads in row c from the n -abacus displays of $\beta_s(\lambda)$ and $\beta_s(\mu)$ respectively. Then $\tilde{\lambda}, \tilde{\mu} \in \mathcal{P}_{\mathbf{t}'}$ and are the partitions obtained from λ and μ respectively by removing the first $s - s'$ parts, so that $|\tilde{\mu}| < |\mu|$. We have

$$d_{\lambda\mu} = d_{\tilde{\lambda}\tilde{\mu}} = d_{\tilde{\lambda}\tilde{\mu}}^{\mathbf{t}'}(1) = d_{\lambda\mu}^{\mathbf{t}}(1),$$

where the first equality is ‘row removal’ [[Jam](#), Theorem 6.18], the second by induction and the last by [Lemma 2.3\(2\)](#).

We now proceed to the second case. In this case, there is a bead in row c of the n -abacus display of $\beta_s(\mu)$ having a vacant preceding position that is in the same section as the bead. Formally, there exists $k \in \beta_s(\mu)$ such that $\lfloor \frac{k}{n} \rfloor = c$, $k-1 \notin \beta_s(\mu)$, and $f_k \in \mathbf{U}(\mathbf{n})$. Let λ be the partition such that $\beta_s(\lambda) = \beta_s(\mu) \cup \{k-1\} \setminus \{k\}$. Then $\lambda \in \mathcal{P}_{\mathbf{t}}$, and we have, by [Corollary 3.3](#) and [Corollary 5.4](#),

$$\begin{aligned} f_k(G_{\mathbf{t}}(\lambda)) &= G_{\mathbf{t}}(\mu) + \sum_{\substack{\nu \in \mathcal{P}_{\mathbf{t}} \\ |\nu|=|\mu| \\ \nu < \mu}} L_{\lambda}^{\nu}(q) G_{\mathbf{t}}(\nu) \\ [(k-s)\text{-Ind}(P(\lambda))] &= [P(\mu)] + \sum_{\substack{\nu \in \mathcal{P}_{\mathbf{t}} \\ |\nu|=|\mu| \\ \nu < \mu}} L_{\lambda}^{\nu}[P(\nu)], \end{aligned}$$

and $L_{\lambda}^{\nu} = [(k-s)\text{-Res}(L(\nu)) : L(\lambda)] \in \mathbb{N}_0$ for all ν . Note that $\mathcal{F}_{c,\mathbf{s}}$ is invariant under $(k-s)\text{-Ind}$ for all $\mathbf{s} \in \mathfrak{B}_s$. Thus applying $\varrho_{\mathbf{t}}$ to the last equation above, we get by [Lemma 3.4](#)

$$(k-s)\text{-Ind}([P(\lambda)]_{\mathbf{t}}) = [P(\mu)]_{\mathbf{t}} + \sum_{\substack{\nu \in \mathcal{P}_{\mathbf{t}} \\ |\nu|=|\mu| \\ \nu < \mu}} L_{\lambda}^{\nu}[P(\nu)]_{\mathbf{t}}.$$

Hence

$$\begin{aligned} [P(\mu)]_{\mathbf{t}} - \varepsilon_{q=1}(G_{\mathbf{t}}(\mu)) &= (k-s)\text{-Ind}([P(\lambda)]_{\mathbf{t}}) - \varepsilon_{q=1}(f_k(G_{\mathbf{t}}(\lambda))) \\ &\quad + \sum_{\substack{\nu \in \mathcal{P}_{\mathbf{t}} \\ |\nu|=|\mu| \\ \nu < \mu}} (L_{\lambda}^{\nu}(1)\varepsilon_{q=1}(G_{\mathbf{t}}(\nu)) - L_{\lambda}^{\nu}[P(\nu)]_{\mathbf{t}}) \\ &= \sum_{\substack{\nu \in \mathcal{P}_{\mathbf{t}} \\ |\nu|=|\mu| \\ \nu < \mu}} (L_{\lambda}^{\nu}(1) - L_{\lambda}^{\nu}) \varepsilon_{q=1}(G_{\mathbf{t}}(\nu)), \end{aligned} \tag{*}$$

using the fact that $\varepsilon_{q=1}$ intertwines f_k and $(k-s)\text{-Ind}$, and applying induction hypothesis to λ and ν .

By [Theorem 2.6](#) and [Corollary 5.7](#), we see that $L_{\lambda}^{\nu}(1) \leq L_{\lambda}^{\nu}$ for all $\nu \in \mathcal{P}_{\mathbf{t}}$. If $L_{\lambda}^{\nu}(1) \neq L_{\lambda}^{\nu}$ for some $\nu \in \mathcal{P}_{\mathbf{t}}$, let $\rho \in \mathcal{P}_{\mathbf{t}}$ be maximal with respect to \geq_{J_p} such that $L_{\lambda}^{\rho}(1) \neq L_{\lambda}^{\rho}$. Then $L_{\lambda}^{\rho}(1) < L_{\lambda}^{\rho}$, and, for all $\nu \in \mathcal{P}_{\mathbf{t}}$, $L_{\lambda}^{\nu}(1) - L_{\lambda}^{\nu} = 0$ if $\nu >_{J_p} \rho$ while $d_{\rho\nu}(q) = 0$ if $\nu \not\geq_{J_p} \rho$ by [Theorem 2.1\(3\)](#). Thus, comparing the coefficient of $\beta_s(\rho)$ in both sides of (*), we get by [Theorem 5.2](#)

$$d_{\rho\mu} - d_{\rho\mu}(1) = \sum_{\substack{\nu \in \mathcal{P}_{\mathbf{t}} \\ |\nu|=|\mu| \\ \nu < \mu}} (L_{\lambda}^{\nu}(1) - L_{\lambda}^{\nu}) d_{\rho\nu}(1) = L_{\lambda}^{\rho}(1) - L_{\lambda}^{\rho}.$$

But $d_{\rho\mu} \geq d_{\rho\mu}(1)$ by [Proposition 2.5\(3\)](#) while $L_{\lambda}^{\rho}(1) < L_{\lambda}^{\rho}$, giving us a contradiction. Therefore, $L_{\lambda}^{\nu}(1) = L_{\lambda}^{\nu}$ for all $\nu \in \mathcal{P}_{\mathbf{t}}$, and we get from (*) that $[P(\mu)]_{\mathbf{t}} = \varepsilon_{q=1}(G_{\mathbf{t}}(\mu))$, as desired. \square

Corollary 6.2. *Let $\mathbf{n} = (n_1, \dots, n_r)$ be a tuple of positive integers summing to p , with $n_j \in \{1, 2\}$ for all j . Then $d_{\lambda\mu} = d_{\lambda\mu}(1)$ for all $\lambda, \mu \in \mathcal{P}_{\mathbf{t}}$ and $\mathbf{t} \in \mathbf{T}_{\mathbf{n}}$.*

Proof. This follows immediately from [Theorem 5.2](#) and [Theorem 6.1](#). \square

We note that if $\mathbf{n} = (n_1, \dots, n_r)$ is a tuple of positive integers summing to p , with $n_j \in \{1, 2\}$ for all j , then $d_{\lambda\mu}^{\mathbf{t}}(q)$, and hence $d_{\lambda\mu}(q)$ and $d_{\lambda\mu}$ by virtue of [Theorem 6.1](#) and [Corollary 6.2](#), can be described by the closed formulas found by Frenkel and Khovanov using [Theorem 5.5](#) for all $\mathbf{t} \in \mathbf{T}_{\mathbf{n}}$, $\lambda, \mu \in \mathcal{P}_{\mathbf{t}}$.

Remark. [Corollary 6.2](#) does not hold for arbitrary \mathbf{n} , even under James's hypothesis that the size of the partitions λ and μ is strictly less than p^2 [[Jam](#)]. Williamson [[W](#)] produces counterexamples in which $\lambda, \mu \in \mathcal{P}_{\mathbf{t}}$ are partitions of $p \binom{N}{2} < p^2$, and where $\mathbf{n} = (p)$ and \mathbf{t} is given by

$$\mathbf{t}(i) = \begin{cases} p & \text{if } i \leq 0 \\ 1 & \text{if } 1 \leq i \leq N \\ 0 & \text{if } i \geq N + 1. \end{cases}$$

Two residue classes a, b modulo p are *adjacent* if and only if $a - b \equiv_p \pm 1$.

Lemma 6.3. *Let $\lambda, \mu \in \mathcal{P}$, and let I be the set of p -residues of the nodes in $([\lambda] \setminus [\mu]) \cup ([\mu] \setminus [\lambda])$. Suppose that no two elements of I are adjacent. Then there exist $s \in \{0, 1\}$ and $\mathbf{n} = (n_1, \dots, n_r)$, with $n_j \in \{1, 2\}$ for all j and $\sum_{j=1}^r n_j = p$, such that $\mathbf{t}_{\beta_s(\lambda)}^{\mathbf{n}} = \mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}$ and $I = \{\sum_{j=1}^i n_j - s - 1 \mid n_i = 2\}$.*

Proof. Let $s = 1$ if $0 \in I$ and $s = 0$ if $0 \notin I$. Let $\mathbf{n} = (n_1, \dots, n_r)$ be a sequence of positive integers summing to p such that $\{\sum_{j=1}^i n_j - s \mid s \leq i \leq r - 1 + s\} = \{0, 1, \dots, p - 1\} \setminus I$. Since the elements of I are pairwise non-adjacent, we see that $n_j \leq 2$ for all j . Furthermore, $\mathbf{t}_{\beta_s(\lambda)}^{\mathbf{n}} = \mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}$ by [Corollary 4.2](#) and $I = \{\sum_{j=1}^i n_j - s - 1 \mid n_i = 2\}$. \square

If $\mu \in \mathcal{P}$, and λ is obtained from μ by removing a removable node of p -residue i and adding an addable node of p -residue i , we say that λ is obtained from μ by *moving* a node of p -residue i .

Theorem 6.4. *Let $\mu \in \mathcal{P}$ and suppose that λ is obtained from μ by moving some nodes whose p -residues are pairwise non-adjacent. Let I be the subset of residue classes modulo p which occur as p -residues of the nodes moved to obtain λ . For each $i \in I$, let $\lambda(i)$ be the partition obtained from μ by moving only those nodes with p -residue i . Then*

$$d_{\lambda\mu} = \prod_{i \in I} d_{\lambda(i), \mu}.$$

Proof. By [Lemma 6.3](#), there exist $s \in \{0, 1\}$ and $\mathbf{n} = (n_1, \dots, n_r)$, with $n_j \in \{1, 2\}$ for all j and $\sum_{j=1}^r n_j = p$, such that $\mathbf{t}_{\beta_s(\lambda)}^{\mathbf{n}} = \mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}$. Furthermore,

by [Corollary 4.2](#), $\mathbf{t}_{\beta_s(\lambda)}^{\mathbf{n}} = \mathbf{t}_{\beta_s(\lambda)(i)}^{\mathbf{n}} = \mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}$ for all $i \in I$. By [Theorem 5.5](#) and adopting the notations there, we have

$$d_{\lambda\mu}^p(q) = \prod_{j=1}^r d_{\lambda^{(j)}\mu^{(j)}}^{n_j}(q) = \prod_{j: n_j=2} d_{\lambda^{(j)}\mu^{(j)}}^2(q),$$

since $\lambda^{(j)} = \mu^{(j)}$ if $n_j = 1$. For each j such that $n_j = 2$, let $i_j = \sum_{i=1}^j n_i - s - 1$. Then

$$d_{\lambda^{(i_j)}\mu}^p(q) = d_{\lambda^{(j)}\mu^{(j)}}^2(q) \left(\prod_{k \neq j} d_{\mu^{(k)}\mu^{(k)}}^{n_k}(q) \right) = d_{\lambda^{(j)}\mu^{(j)}}^2(q),$$

by [Theorem 5.5](#) and [Theorem 2.1\(1\)](#). Thus,

$$d_{\lambda\mu}^p = d_{\lambda\mu}^p(1) = \prod_{j: n_j=2} d_{\lambda^{(j)}\mu^{(j)}}^2(1) = \prod_{j: n_j=2} d_{\lambda^{(i_j)}\mu}^p(1) = \prod_{i \in I} d_{\lambda^{(i)}\mu}^p(1) = \prod_{i \in I} d_{\lambda(i)\mu}$$

by [Lemma 6.3](#) and [Corollary 6.2](#). \square

Theorem 6.5. *Let $\mu \in \mathcal{P}$ and let λ be the partition obtained from μ by removing a removable node \mathbf{n} of p -residue i and moving m other nodes, such that the p -residues of these nodes (including \mathbf{n}) are pairwise non-adjacent. Then*

$$[i\text{-Res}(L(\mu)) : L(\lambda)] = 0,$$

unless $m = 0$ and \mathbf{n} is a normal node of μ .

Proof. By [Lemma 6.3](#), there exist $s \in \{0, 1\}$ and $\mathbf{n} = (n_1, \dots, n_r)$ with $n_j \in \{1, 2\}$ for all j and $\sum_{j=1}^r n_j = p$ such that $\mathbf{t}_{\beta_s(\lambda)}^{\mathbf{n}} = \mathbf{t}_{\beta_s(\mu)}^{\mathbf{n}}$. Let $\mathbf{t} = \mathbf{t}_{\beta_s(\lambda)}^{\mathbf{n}}$. Applying $\varrho_{\mathbf{t}}$ to $i\text{-Ind}([P(\lambda)]) = \bigoplus_{\nu \in \mathcal{P}} L_{\lambda}^{\nu}[P(\nu)]$, where $L_{\lambda}^{\nu} = [i\text{-Res}(L(\nu)) : L(\lambda)]$ for all $\nu \in \mathcal{P}$, we get by [Lemma 3.4\(2\)](#)

$$i\text{-Ind}([P(\lambda)]_{\mathbf{t}}) = \bigoplus_{\nu \in \mathcal{P}_{\mathbf{t}}} L_{\lambda}^{\nu}[P(\nu)]_{\mathbf{t}}.$$

On the other hand, applying $\varepsilon_{q=1}$ to $f_{s+i}(G_{\mathbf{t}}(\lambda)) = \sum_{\nu \in \mathcal{P}_{\mathbf{t}}} L_{\lambda}^{\nu}(q)G_{\mathbf{t}}(\nu)$, where $L_{\lambda}^{\nu}(q) \in A$ for all $\nu \in \mathcal{P}_{\mathbf{t}}$, we get by [Theorem 6.1](#)

$$i\text{-Ind}([P(\lambda)]_{\mathbf{t}}) = \bigoplus_{\nu \in \mathcal{P}_{\mathbf{t}}} L_{\lambda}^{\nu}(1)[P(\nu)]_{\mathbf{t}}.$$

Thus, $L_{\lambda}^{\nu}(1) = L_{\lambda}^{\nu}$ for all $\nu \in \mathcal{P}_{\mathbf{t}}$. The statement now follows from [Corollary 5.7](#). \square

Remark.

- (1) Exactly the same proofs of [Theorem 6.4](#) and [Theorem 6.5](#) in fact prove the respective results hold in a slightly more general setting: whenever any two nodes in the symmetric difference of $[\lambda]$ and $[\mu]$ do not have adjacent p -residues. But the additional information is actually trivial—they can easily be seen to hold with our current knowledge of decomposition numbers and branching coefficients of Schur algebras.

- (2) We have deliberately left out in [Theorem 6.5](#) the closed formula for $[i\text{-Res}(L(\mu)) : L(\lambda)]$ when λ is obtained from μ by removing a normal node of residue i (which is one plus the number of normal nodes of μ of residue i and to the right of the normal node removed to obtain λ , see [Theorem 2.6](#)) so as not to give a false impression that we have an alternative and independent proof of [Theorem 2.6](#).

REFERENCES

- [CMT1] J. Chuang, H. Miyachi, K. M. Tan. Row and column removal in the q -deformed Fock space. *J. Algebra*, 254:84–91, 2002.
- [CMT2] J. Chuang, H. Miyachi, K. M. Tan. Kleshchev’s decomposition numbers and branching coefficients in the Fock space. *Trans. Amer. Math. Soc.*, 360:1179–1191, 2008.
- [CT] J. Chuang, K. M. Tan. Filtrations in Rouquier blocks of symmetric groups and Schur algebras. *Proc. London Math. Soc.* (3), 86:685–706, 2003.
- [F1] M. Fayers. Decomposition numbers for weight three blocks of symmetric groups and Iwahori-Hecke algebras. *Trans. Amer. Math. Soc.* 360:1341–1376, 2008.
- [F2] M. Fayers. James’s Conjecture holds for weight four blocks of Iwahori-Hecke algebras. *J. Algebra* 317:593–633, 2007.
- [FK] I. B. Frenkel, M. Khovanov. Canonical bases in tensor products and graphical calculus for $U_q(\mathfrak{sl}_2)$. *Duke Math J.*, 87:409–480, 1997.
- [H] T. Hayashi. q -analogues of Clifford and Weyl algebras—spinor and oscillator representations of quantum enveloping algebras. *Comm. Math. Phys.*, 127:129–144, 1990.
- [HK] J. Hong, S.-J. Kang. Introduction to quantum groups and crystal bases. Graduate Studies in Mathematics, 42. American Mathematical Society, Providence, RI, 2002.
- [Jam] G. D. James. The decomposition matrices of $GL_n(q)$ for $n \leq 10$. *Proc. London Math. Soc.* (3), 60(2):225–265, 1990.
- [JK] G. D. James, A. Kerber. The representation theory of the symmetric group. With a foreword by P. M. Cohn. With an introduction by Gilbert de B. Robinson. Encyclopedia of Mathematics and its Applications, 16. Addison-Wesley Publishing Co., Reading, Mass., 1981.
- [Jan] J. C. Jantzen. Lectures on quantum groups. Graduate Studies in Mathematics, 6. American Mathematical Society, Providence, RI, 1996.
- [Ka] Kashiwara, M. On crystal bases of the Q -analogue of universal enveloping algebras. *Duke Math. J.* 63: 465–516, 1991.
- [Kl] A. Kleshchev. On decomposition numbers and branching coefficients for symmetric and special linear groups. *Proc. London Math. Soc.* (3), 75:497–558, 1997.
- [Le] B. Leclerc. Fock space representations of $U_q(\widehat{\mathfrak{sl}}_n)$. Geometric methods in representation theory. I, 343–385, Sémin. Congr., 24-I, Soc. Math. France, Paris, 2012.
- [LT] B. Leclerc, J.-Y. Thibon. Canonical bases of q -deformed Fock spaces. *Internat. Math. Res. Notices*, 9:447–456, 1996.
- [Lu] G. Lusztig. Introduction to quantum groups. Reprint of the 1994 edition. Modern Birkhuser Classics. Birkhuser/Springer, New York, 2010.
- [MM] K. Misra, T. Miwa. Crystal base for the basic representation of $U_q(\mathfrak{sl}(n))$. *Comm. Math. Phys.*, 134:79–88, 1990.
- [T] K. M. Tan. Beyond Rouquier partitions. *J. Algebra* 321:248–263, 2009.

- [TT] K. M. Tan, W. H. Teo. Sign sequences and decomposition numbers. *Trans. Amer. Math. Soc.* 365:6385–6401, 2013.
- [VV] M. Varagnolo and E. Vasserot. On the decomposition matrices of the quantized Schur algebra. *Duke Math. J.*, 100(2):267–297, 1999.
- [W] G. Williamson. Schubert calculus and torsion explosion. Preprint, [arxiv:1309.5055](https://arxiv.org/abs/1309.5055), 2015.

DEPARTMENT OF MATHEMATICS, CITY UNIVERSITY LONDON, NORTHAMPTON SQUARE,
LONDON EC1V 0HB, UNITED KINGDOM. Email: j.chuang@city.ac.uk

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, BLOCK S17,
10 LOWER KENT RIDGE ROAD, SINGAPORE 119076. Email: tankm@nus.edu.sg