



City Research Online

City, University of London Institutional Repository

Citation: Gen, J., Correa, F. & Fring, A. (2017). Time-delay and reality conditions for complex solitons. *Journal of Mathematical Physics*, 58(3), 32901. doi: 10.1063/1.4978864

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/17162/>

Link to published version: <https://doi.org/10.1063/1.4978864>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Time-delay and reality conditions for complex solitons

Julia Cen[•] and Francisco Correa^{†,‡} and Andreas Fring[•]

• *Department of Mathematics, City University London,
Northampton Square, London EC1V 0HB, UK*

† *Instituto de Ciencias Físicas y Matemáticas, Universidad Austral de Chile,
Casilla 567, Valdivia, Chile*

‡ *Institut für Theoretische Physik and Riemann Center for Geometry and Physics,
Leibniz Universität Hannover, Appelstraße 2, 30167 Hannover, Germany*

E-mail: julia.cen.1@city.ac.uk, francisco.correa@itp.uni-hannover.de, a.fring@city.ac.uk

ABSTRACT: We compute lateral displacements and time-delays for a scattering processes of complex multi-soliton solutions of the Korteweg de-Vries equation. The resulting expressions are employed to explain the precise distinction between solutions obtained from different techniques, Hirota's direct method and a superposition principle based on Bäcklund transformations. Moreover they explain the internal structures of degenerate compound multi-solitons previously constructed. Their individual one-soliton constituents are time-delayed when scattered amongst each other. We present generic formulae for these time-dependent displacements. By recalling Gardner's transformation method for conserved charges, we argue that the structure of the asymptotic behaviour resulting from the integrability of the model together with its \mathcal{PT} -symmetry ensure the reality of all of these charges, including in particular the mass, the momentum and the energy.

1. Introduction

It is one of the defining features of classical multi-soliton solutions to nonlinear integrable equations that individual one-soliton contributions maintain their overall shape before and after a scattering event. The only net effect is that they are delayed or advanced in time as a result of the scattering with other solitons when compared to the undisturbed motion of a single one-soliton [1, 2, 3]. Besides providing a more detailed picture on the motion of classical solitons, following ideas of Wigner and Eisenbud [4], the concrete values of the delay times are also important for the quantization of the theory as they can be related to quantum mechanical scattering matrices in a semi-classical approximation.

We present here a detailed analysis of the delay times for complex soliton solutions to the Korteweg de-Vries equation (KdV) previously reported in [5, 6]. We use our results

for a variety of purposes. On a technical level the explicit expressions allow to clarify the precise distinctions between solutions obtained from different types of solution methods, in particular Hirota's direct method and Bäcklund transformations. Moreover, the time-delays also shed new on the degenerate multi-soliton solutions constructed in [6], especially the internal structure of compound multi-solitons can be explained in detail when using the expression for the time-delays or some approximate asymptotic formulae.

In the quantum mechanical context it is well understood which role \mathcal{PT} -symmetries, or better antilinear symmetries [7], play in order to explain the reality of the energy eigen-spectrum [8, 9, 10]. For nonlinear integrable wave equations and their \mathcal{PT} -symmetric deformations [11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21] there is still some uncertainty about the precise reasoning. For the model discussed here we argue that physical quantities based on one-soliton solutions, \mathcal{PT} -symmetric or not, can always be made \mathcal{PT} -symmetric and in this case that feature alone guarantees their reality. The integrability of the model then ensures that asymptotically any multi-soliton solution separates into a collection of one-soliton solutions, possibly time-delayed, of which each contributes only a real value to an overall conserved charge. We recall the structure of all of these charges constructed from Gardner's transformation and then use it to show that \mathcal{PT} -symmetry and integrability ensure the reality of all conserved charges.

Our manuscript is organized as follows: Starting from some general definitions and properties of conserved quantities and complex one-soliton solutions we compute in section 2 the time-delays in nondegenerate and degenerate two and three-soliton solutions. We provide closed expressions for the time-dependent displacements for degenerate N -solitons for any N . In section 3 we present the reasoning that ensures the reality of all conserved quantities. Our conclusions and an outlook into future work and open issues is presented in section 4. In general the detailed computations are omitted in section 2, but in order to illustrate the working we present some sample computations in an appendix.

2. Time-delays for multi-soliton solutions

2.1 Generalities

Following [1, 2, 3] the classical time-delay of a scattering process is defined as follows: We consider the trajectories of a particle, or a soliton for that matter, with velocity v before and after the collision as $x = vt + x^{(i)}$ and $x = vt + x^{(f)}$, respectively. The *lateral displacement* resulting from the scattering event is then defined as the difference between these two trajectories, that is simply

$$\Delta_x := x^{(f)} - x^{(i)}, \quad (2.1)$$

so that the *time-delay* is naturally defined as

$$\Delta_t := t^{(f)} - t^{(i)} = -\frac{x^{(f)}}{v} + \frac{x^{(i)}}{v} = -\frac{\Delta_x}{v}. \quad (2.2)$$

Negative and positive time-delays are interpreted as attractive and repulsive forces, respectively. In a multi-particle scattering process of particles, or solitons, of type k the

corresponding lateral displacements and time-delays $(\Delta_x)_k$ and $(\Delta_t)_k$, respectively, have to satisfy certain consistence conditions [3]. Demanding for instance that the centre of mass coordinate

$$X = \frac{\sum_k m_k x_k}{\sum_k m_k} \quad (2.3)$$

remains the same before and after the collision, i.e. $X^{(i)} = X^{(f)}$, immediately implies that

$$\sum_k m_k (\Delta_x)_k = 0. \quad (2.4)$$

Furthermore, given that $m\Delta_x = -mv\Delta_t = -p\Delta_t$ yields

$$\sum_k p_k (\Delta_t)_k = 0, \quad (2.5)$$

where p_k is the momentum of a particle of type k . We will use the relations (2.4) and (2.5) for consistency checks.

2.2 Complex KdV multi-soliton scattering

Let us now see how the above applies to the scattering of multi-solitons that are solutions of the KdV equation for the complex field $u(x, t) = p(x, t) + iq(x, t)$ with $p(x, t), q(x, t) \in \mathbb{R}$

$$u_t + 6uu_x + u_{xxx} = 0. \quad (2.6)$$

Separating (2.6) into its real and imaginary part one may view it of course as set of coupled equations for the real fields $p(x, t)$ and $q(x, t)$. In the limits $(pq)_x \rightarrow pq_x$ and $q_{xxx} \rightarrow 0$ they reduce to some well studied systems, the Hirota-Satsuma [22] and Ito equations [23], respectively. The total mass, momentum and energy associated to the solution $u(x, t)$ are defined as

$$m = \int_{-\infty}^{\infty} u dx, \quad p = \int_{-\infty}^{\infty} u^2 dx, \quad E = \int_{-\infty}^{\infty} (2u^3 - u_x^2) dx, \quad (2.7)$$

respectively. See section 3 for a derivation of these expressions. We have to establish that these quantities are real as they are meant to be physical, i.e. observable.

2.2.1 Properties of the one-soliton solutions

First we need to compute complex solutions to the KdV equation. We recall that they may be constructed for instance from Hirota's direct method [24]. Defining the quantities $\eta_{\mu;\alpha} := \alpha x - \alpha^3 t + \mu$, we consider the τ -function for a one-soliton

$$\tau_{\mu;\alpha}(x, t) = 1 + e^{\eta_{\mu;\alpha}}, \quad (2.8)$$

from which the corresponding solution to the complex KdV equation is obtained as $u(x, t) = 2[\ln \tau(x, t)]_{xx}$. Taking the value for μ in a form that respects the \mathcal{PT} -symmetry of the solutions, i.e. purely imaginary $\mu = i\theta$ with $\theta \in \mathbb{R}$, we obtain [5]

$$u_{i\theta;\alpha}(x, t) = \frac{\alpha^2 + \alpha^2 \cos \theta \cosh(\alpha x - \alpha^3 t)}{[\cos \theta + \cosh(\alpha x - \alpha^3 t)]^2} - i \frac{\alpha^2 \sin \theta \sinh(\alpha x - \alpha^3 t)}{[\cos \theta + \cosh(\alpha x - \alpha^3 t)]^2}. \quad (2.9)$$

We may restrict ourselves to this choice, because \mathcal{PT} -symmetry breaking choices, such as $\mu = \kappa + i\theta$ with $\kappa \in \mathbb{R}$, can be converted to the form (2.9) by simple shifts in x or t , $u_{\kappa+i\theta;\alpha}(x, t) = u_{i\theta;\alpha}(x + \kappa/\alpha, t) = u_{i\theta;\alpha}(x, t - \kappa/\alpha^3)$. The former may then be absorbed in the limits of integrals when computing physical quantities and the latter may be neglected when considering conserved charges. Using this solution we compute the quantities as defined in (2.7). For the mass of the complex one-soliton we obtain always a real value

$$m_\alpha = \int_{-\infty}^{\infty} u_{i\theta;\alpha}(x, t) dx = \frac{\alpha \sinh(\alpha x - \alpha^3 t) + i\alpha \sin \theta}{\cos \theta + \cosh(\alpha x - \alpha^3 t)} \Big|_{x=-\infty}^{\infty} = 2\alpha. \quad (2.10)$$

Thus only $p(x, t)$ component contributes to the mass of the soliton $u(x, t)$, so that $q(x, t)$ may be viewed as a massless soliton. Likewise, the momentum of the one-soliton turns out to be always real

$$p_\alpha = \int_{-\infty}^{\infty} u_{i\theta;\alpha}^2 dx = \frac{\alpha^3 \sinh \eta_{0;\alpha} [5 + 6 \cos \theta \cosh \eta_{0;\alpha} + \cosh(2\eta_{0;\alpha})]}{6 (\cos \theta + \cosh \eta_{0;\alpha})^3} \Big|_{x=-\infty}^{\infty} + i \frac{\alpha^3 \sin \theta (5 + \cos(2\theta) + 6 \cos \theta \cosh(\eta_{0;\alpha}))}{6 (\cos \theta + \cosh \eta_{0;\alpha})^3} \Big|_{x=-\infty}^{\infty} = \frac{2}{3} \alpha^3, \quad (2.11)$$

and the value for the energy was reported in [5] to be real as well

$$E_\alpha = \int_{-\infty}^{\infty} [2u_{i\theta;\alpha}^3 - (u_{i\theta;\alpha})_x^2] dx = \frac{2}{5} \alpha^5. \quad (2.12)$$

In comparison with [5] we have rescaled the energy by a factor of -2 for reason that become apparent in section 3.

We now follow [3] by choosing a reference frame that tracks a distinct point on the soliton, such the crest or trough of the wave. Tracking the one-soliton solution by keeping the traveling wave coordinate $x - \alpha^2 t$ at a fixed value, we obtain for the real part of the solution (2.9) the constant values

$$p_{i\theta;\alpha} [t\alpha^2, t] = \frac{\alpha^2}{2} \sec^2 \left(\frac{\theta}{2} \right) =: \hat{P}_\alpha(\theta), \quad (2.13)$$

$$p_{i\theta;\alpha} \left[t\alpha^2 \pm \frac{1}{\alpha} \Delta_r(\theta), t \right] = -\frac{\alpha^2}{4} \cot^2(\theta) =: \check{P}_\alpha(\theta), \quad (2.14)$$

corresponding to a maximum and two minima, respectively, with shift function

$$\Delta_r(\theta) := \operatorname{arccosh}(\cos \theta - 2 \sec \theta). \quad (2.15)$$

Notice that the minima only emerge when $\theta \in ((4n+1)\pi/2, (4n+3)\pi/2)$ with $n \in \mathbb{Z}$ as otherwise the argument of the $\operatorname{arccosh}$ in (2.15) is smaller 1. For the imaginary part we define the shift function

$$\Delta_i(\theta) := \operatorname{arccosh} \left[\frac{1}{2} \cos \theta + \frac{\sqrt{2}}{4} \sqrt{17 + \cos(2\theta)} \right], \quad (2.16)$$

and compute the minimal and maximal values

$$q_{i\theta;\alpha} \left[t\alpha^2 \pm \frac{1}{\alpha} \Delta_i(\theta), t \right] = \mp \frac{8\alpha^2 \sin \theta \sqrt{5 + \cos(2\theta) + \sqrt{2} \cos \theta \sqrt{17 + \cos(2\theta)}}}{\left[6 \cos \theta + \sqrt{2} \sqrt{17 + \cos(2\theta)} \right]^2} =: \mp Q_\alpha(\theta). \quad (2.17)$$

With these shifts the real part of the one-soliton solution is simply fixed to remain on the crest of the wave as time evolves in (2.13) or on either of the two minima in (2.14) when they exist. For the imaginary part we have the option to track either the crest or trough as specified in (2.17).

We summarize these features in figure 1.




Figure 1: \mathcal{PT} -symmetric one-soliton solution (2.9) of the KdV equation (2.6) with $\alpha = 6/5$ and $\theta = 6/5\pi$ at time $t = 20$.

2.2.2 Properties of nondegenerate two-soliton solutions

Next we consider the complex two-soliton solution. Abbreviating the reoccurring constant $\varkappa(\alpha, \beta) := (\alpha - \beta)^2 / (\alpha + \beta)^2$, the two-soliton τ -function is compactly expressed as

$$\tau_{\mu,\nu;\alpha,\beta}(x, t) = 1 + e^{\eta_{\mu;\alpha}} + e^{\eta_{\nu;\beta}} + \varkappa(\alpha, \beta) e^{\eta_{\mu;\alpha} + \eta_{\nu;\beta}}, \quad (2.18)$$

from which, by using again the transformation $u(x, t) = 2[\ln \tau(x, t)]_{xx}$, we compute the solution

$$u_{i\theta,i\phi;\alpha,\beta} = \frac{2[\alpha^2 e^{\eta_{i\theta;\alpha}} + \beta^2 e^{\eta_{i\phi;\beta}} + \varkappa(\alpha, \beta)(\alpha^2 e^{\eta_{i\theta;\alpha} + 2\eta_{i\phi;\beta}} + \beta^2 e^{2\eta_{i\theta;\alpha} + \eta_{i\phi;\beta}}) + 2(\alpha - \beta)^2 e^{\eta_{i\theta;\alpha} + \eta_{i\phi;\beta}}]}{\tau_{i\theta,i\phi;\alpha,\beta}^2}. \quad (2.19)$$

We now have to track the one-soliton contributions within the solution (2.19) and according to our definitions (2.1) and (2.2) we need to compare the values in the infinite past with the one in the infinite future in order to find the lateral displacements and time-delays. From (2.13) we can read off which frame we have to choose. Tracking the maxima for the real part of the two-soliton solution (2.19), we compute the asymptotic values

$$\begin{aligned} \lim_{t \rightarrow -\infty} p_{i\theta,i\phi;\alpha,\beta} [t\alpha^2, t] &= \lim_{t \rightarrow +\infty} p_{i\theta,i\phi;\alpha,\beta} [t\alpha^2 + \delta_{\alpha}^{\alpha,\beta}, t] = \hat{P}_{\alpha}(\theta), \\ \lim_{t \rightarrow +\infty} p_{i\theta,i\phi;\alpha,\beta} [t\beta^2, t] &= \lim_{t \rightarrow -\infty} p_{i\theta,i\phi;\alpha,\beta} [t\beta^2 + \delta_{\beta}^{\alpha,\beta}, t] = \hat{P}_{\beta}(\phi), \end{aligned} \quad (2.20)$$

where for definiteness we have taken the ordering $\alpha > \beta$ and furthermore abbreviated the quantity

$$\delta_x^{y,z} := \frac{2}{x} \ln \left(\frac{y+z}{y-z} \right), \quad (2.21)$$

for conciseness. See appendix A for the details of this computation. According to our definitions (2.1) and (2.2) we can now read off the lateral displacements and time-delays from the asymptotic values in (2.20) by comparing the infinite future and the infinite past. For the soliton with velocity α^2 we find

$$(\Delta_x)_\alpha = \delta_\alpha^{\alpha,\beta}, \quad \text{and} \quad (\Delta_t)_\alpha = -\frac{1}{\alpha^2} \delta_\alpha^{\alpha,\beta}, \quad (2.22)$$

and for the soliton with velocity β^2 we identify

$$(\Delta_x)_\beta = -\delta_\beta^{\alpha,\beta}, \quad \text{and} \quad (\Delta_t)_\beta = \frac{1}{\beta^2} \delta_\beta^{\alpha,\beta}. \quad (2.23)$$




Figure 2: Lateral displacements for the complex \mathcal{PT} -symmetric two-soliton KdV solution (2.19) with $\alpha = 3/2$, $\beta = 1$, $\theta = \pi/3$ and $\phi = \pi/4$. The plots in the negative and positive regime of x correspond to the time taken to be $t = -20$ and $t = 20$, respectively.

Using the values for the masses and momenta computed in (2.10) and (2.11), we verify that the quantities (2.22) and (2.23) indeed satisfy the consistency relations (2.4) and (2.5), since

$$m_\alpha(\Delta_x)_\alpha + m_\beta(\Delta_x)_\beta = 2\alpha\delta_\alpha^{\alpha,\beta} - 2\beta\delta_\beta^{\alpha,\beta} = 0, \quad (2.24)$$

$$p_\alpha(\Delta_t)_\alpha + p_\beta(\Delta_t)_\beta = -\frac{2}{3}\alpha^3\frac{1}{\alpha^2}\delta_\alpha^{\alpha,\beta} + \frac{2}{3}\beta^3\frac{1}{\beta^2}\delta_\beta^{\alpha,\beta} = 0. \quad (2.25)$$

We may of course also track any other point and should obtain the same values for the displacement and delays. For instance, tracking the minima for the real part we compute

$$\begin{aligned} \lim_{t \rightarrow -\infty} p_{i\theta, i\phi; \alpha, \beta} \left[t\alpha^2 \pm \frac{\Delta_r(\theta)}{\alpha}, t \right] &= \lim_{t \rightarrow +\infty} p_{i\theta, i\phi; \alpha, \beta} \left[t\alpha^2 \pm \frac{\Delta_r(\theta)}{\alpha} + \delta_\alpha^{\alpha,\beta}, t \right] = \check{P}_\alpha(\theta), \\ \lim_{t \rightarrow +\infty} p_{i\theta, i\phi; \alpha, \beta} \left[t\beta^2 \pm \frac{\Delta_r(\phi)}{\beta}, t \right] &= \lim_{t \rightarrow -\infty} p_{i\theta, i\phi; \alpha, \beta} \left[t\beta^2 \pm \frac{\Delta_r(\phi)}{\beta} + \delta_\beta^{\alpha,\beta}, t \right] = \check{P}_\beta(\phi), \end{aligned} \quad (2.26)$$

or tracking the minimum or maximum for the imaginary part we obtain

$$\begin{aligned} \lim_{t \rightarrow -\infty} q_{i\theta, i\phi; \alpha, \beta} \left[t\alpha^2 \pm \frac{\Delta_i(\theta)}{\alpha}, t \right] &= \lim_{t \rightarrow +\infty} q_{i\theta, i\phi; \alpha, \beta} \left[t\alpha^2 \pm \frac{\Delta_i(\theta)}{\alpha} + \delta_{\alpha}^{\alpha, \beta}, t \right] = \mp Q_{\alpha}(\theta), \\ \lim_{t \rightarrow +\infty} q_{i\theta, i\phi; \alpha, \beta} \left[t\beta^2 \pm \frac{\Delta_i(\phi)}{\beta}, t \right] &= \lim_{t \rightarrow -\infty} q_{i\theta, i\phi; \alpha, \beta} \left[t\beta^2 \pm \frac{\Delta_i(\phi)}{\beta} + \delta_{\beta}^{\alpha, \beta}, t \right] = \mp Q_{\beta}(\phi). \end{aligned} \quad (2.27)$$

We depict these features in figure 2.

We also remark here that the time-delays are important to clarify the precise relation between the solutions obtained from the Hirota method (2.19) and those constructed via Bäcklund transformations [5]

$$u_{\mu, \nu; \alpha, \beta}^B(x, t) = \frac{\alpha^2 - \beta^2}{2} \frac{\beta^2 \operatorname{sech} \left[\frac{1}{2}(\beta x - \beta^3 t + \nu) \right]^2 - \alpha^2 \operatorname{sech} \left[\frac{1}{2}(\alpha x - \alpha^3 t + \mu) \right]^2}{\left[\alpha \tanh \left[\frac{1}{2}(\alpha x - \alpha^3 t + \mu) \right] - \beta \tanh \left[\frac{1}{2}(\beta x - \beta^3 t + \nu) \right] \right]^2}. \quad (2.28)$$

Tracking the maxima of the real part of (2.28) we compute the asymptotic values

$$\begin{aligned} \lim_{t \rightarrow +\infty} p_{i\theta, i\phi; \alpha, \beta}^B \left[\alpha^2 t + \frac{\delta_{\alpha}^{\alpha, \beta}}{2} \right] &= \lim_{t \rightarrow -\infty} p_{i\theta, i\phi; \alpha, \beta}^B \left[\alpha^2 t - \frac{\delta_{\alpha}^{\alpha, \beta}}{2} \right] = \hat{P}_{\alpha}(\theta + \pi) \\ \lim_{t \rightarrow +\infty} p_{i\theta, i\phi; \alpha, \beta}^B \left[\beta^2 t - \frac{\delta_{\beta}^{\alpha, \beta}}{2} \right] &= \lim_{t \rightarrow -\infty} p_{i\theta, i\phi; \alpha, \beta}^B \left[\beta^2 t + \frac{\delta_{\beta}^{\alpha, \beta}}{2} \right] = \hat{P}_{\beta}(\phi) \end{aligned} \quad (2.29)$$




Figure 3: Complex Hirota two-soliton KdV solution (2.19) versus two-soliton KdV solution obtained from Bäcklund transformations (2.28) for $\alpha = 1.2$, $\beta = 0.8$, $\theta = \pi/3$ and $\phi = \pi/4$. The plots in the negative and positive regime of x correspond to the time taken to be $t = -20$ and $t = 20$, respectively.

Comparing the real parts of the one-soliton contribution within the Bäcklund two-soliton solution with those in the Hirota solution, we find that the slower and faster one are delayed by the amount $\delta_{\beta}^{\alpha, \beta}/2$ and $\delta_{\alpha}^{\alpha, \beta}/2$, respectively. Thus the two types of solutions are not simply shifted by an overall amount, but each of the individual one-soliton contributions shifted by a different amount relative to each other. Furthermore, comparing (2.26) and (2.29) we observe that we require a shift in θ by π in the Hirota solution in order to obtain the same qualitative features in both solution, in the sense of matching amplitudes and occurrence of minima. Overall we obtain the same total delays (2.22) from any of the solutions.

Tracking the minima and maxima for the imaginary part we obtain

$$\begin{aligned} \lim_{t \rightarrow +\infty} q_{i\theta, i\phi; \alpha, \beta}^B \left[\alpha^2 t + \frac{\delta_{\alpha, \beta}^{\alpha, \beta}}{2} \pm \frac{\Delta_i(\theta + \pi)}{\alpha} \right] &= \lim_{t \rightarrow -\infty} q_{i\theta, i\phi; \alpha, \beta}^B \left[\alpha^2 t - \frac{\delta_{\alpha, \beta}^{\alpha, \beta}}{2} \pm \frac{\Delta_i(\theta + \pi)}{\alpha} \right] = \mp Q_{\alpha}(\theta + \pi) \\ \lim_{t \rightarrow +\infty} q_{i\theta, i\phi; \alpha, \beta}^B \left[\beta^2 t - \frac{\delta_{\beta}^{\alpha, \beta}}{2} \pm \frac{1}{\beta} \Delta_i(\phi) \right] &= \lim_{t \rightarrow -\infty} q_{i\theta, i\phi; \alpha, \beta}^B \left[\beta^2 t + \frac{\delta_{\beta}^{\alpha, \beta}}{2} \pm \frac{1}{\beta} \Delta_i(\phi) \right] = \mp Q_{\beta}(\phi). \end{aligned} \quad (2.30)$$

As expected, we observe from this that the imaginary parts of the soliton solutions are displaced by the same amount as the real parts, but for the faster soliton we also acquired an overall minus sign.

These features are displayed in figure 3.

We notice that the time-delay caused by the scattering between the faster and the slower soliton is the same in both solutions, i.e. they are preserved quantities.

2.2.3 Properties of degenerate two-soliton solutions

As discussed in [6], the degenerate solutions for which some of the energies are the same are quite special. In general the limit $\beta \rightarrow \alpha$ to degenerate energies in a multi-soliton solution is divergent. However, as discussed in detail in [6], in a complex setting, when tuning certain parameters it can be performed consistently, leading to the solution

$$u_{i\theta, i\phi; \alpha, \alpha}(x, t) = \frac{2\alpha^2 [(\alpha x - 3\alpha^3 t + i\phi) \sinh(\eta_{i\theta; \alpha}) - 2 \cosh(\eta_{i\theta; \alpha}) - 2]}{[\alpha x - 3\alpha^3 t + i\phi + \sinh(\eta_{i\theta; \alpha})]^2}. \quad (2.31)$$

Defining the time-dependent displacement

$$\Delta(t) := \frac{1}{\alpha} \ln(4\alpha^3 |t|), \quad (2.32)$$

we track the maximum for the real part of the degenerate solution (2.31) as

$$\lim_{t \rightarrow +\infty} p_{i\theta, i\phi; \alpha, \alpha} [t\alpha^2 - \Delta(t), t] = \lim_{t \rightarrow -\infty} p_{i\theta, i\phi; \alpha, \alpha} [t\alpha^2 + \Delta(t), t] = \hat{P}_{\alpha}(\theta), \quad (2.33)$$

$$\lim_{t \rightarrow +\infty} p_{i\theta, i\phi; \alpha, \alpha} [t\alpha^2 + \Delta(t), t] = \lim_{t \rightarrow -\infty} p_{i\theta, i\phi; \alpha, \alpha} [t\alpha^2 - \Delta(t), t] = \hat{P}_{\alpha}(\theta + \pi), \quad (2.34)$$

See appendix A for a derivation of these asymptotic expressions. Comparing (2.33) with (2.13) we observe that the time-dependent shift is tracking the maximum of the one-soliton solution. The second maximum (2.34) corresponds to the one-soliton solution (2.13) with $\theta \rightarrow \theta + \pi$, which relates the sech^2 to the csch^2 solution. We expect these solutions to emerge as in the real case they correspond to the two independent solutions from which the degenerate one (2.31) was constructed in [6] when using Wronskians involving Jordan states. Moreover, regarding the internal structure of the degenerate two-soliton we deduce that the one-solitons with amplitude $\hat{P}_{\alpha}(\theta)$ and $\hat{P}_{\alpha}(\theta + \pi)$ are laterally displaced by $-\Delta(t)$ and $\Delta(t)$, respectively, as a result of the scattering process. When $t \rightarrow \pm\infty$ the displacements tend to infinity as we somehow expect from the displacement (2.22) which diverges when $\beta \rightarrow \alpha$.

As we have seen, the real part of one of the one-soliton solutions also develops two minima, for which we compute the limits

$$\lim_{t \rightarrow \sigma\infty} p_{i\theta, i\phi; \alpha, \alpha} \left[t\alpha^2 - \sigma\Delta(t) \pm \frac{1}{\alpha} \Delta_r(\theta), t \right] = \check{P}_{\alpha}(\theta). \quad (2.35)$$

where σ can be $+1$ or -1 . Since the two one-solitons are relatively shifted to each other by $\theta \rightarrow \theta + \pi$ it follows from the remarks after (2.15) that these minima can only emerge in one of the two solitons.

For the imaginary part we compute the eight limits

$$\lim_{t \rightarrow \sigma\infty} q_{i\theta, i\phi; \alpha, \alpha} \left[t\alpha^2 - \sigma\Delta(t) \pm \frac{1}{\alpha}\Delta_i(\theta), t \right] = \mp Q_\alpha(\theta), \quad (2.36)$$

$$\lim_{t \rightarrow \sigma\infty} q_{i\theta, i\phi; \alpha, \alpha} \left[t\alpha^2 + \sigma\Delta(t) \pm \frac{1}{\alpha}\Delta_i(\theta + \pi), t \right] = \pm Q_\alpha(\theta + \pi). \quad (2.37)$$

We also observe from the imaginary part that the overall time-delays are $\pm 2\Delta(t)$.

2.2.4 Properties of nondegenerate three-soliton solutions

Let us now consider the three-soliton solution for which the τ -function reads

$$\begin{aligned} \tau_{\mu, \nu, \rho; \alpha, \beta, \gamma}(x, t) &= 1 + e^{\eta_{\mu; \alpha}} + e^{\eta_{\nu; \beta}} + e^{\eta_{\rho; \gamma}} + \varkappa(\alpha, \beta)e^{\eta_{\mu; \alpha} + \eta_{\nu; \beta}} + \varkappa(\alpha, \gamma)e^{\eta_{\mu; \alpha} + \eta_{\rho; \gamma}} \\ &\quad + \varkappa(\beta, \gamma)e^{\eta_{\nu; \beta} + \eta_{\rho; \gamma}} + \varkappa(\alpha, \beta)\varkappa(\alpha, \gamma)\varkappa(\beta, \gamma)e^{\eta_{\mu; \alpha} + \eta_{\nu; \beta} + \eta_{\rho; \gamma}}, \end{aligned} \quad (2.38)$$

leading to the three-soliton solution

$$u_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma}(x, t) = 2 \left[\ln \tau_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma}(x, t) \right]_{xx}, \quad (2.39)$$

which we do not report here explicitly. Assuming the ordering $\alpha > \beta > \gamma$ we track the maxima for the real parts and compute the asymptotic values

$$\begin{aligned} \lim_{t \rightarrow -\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma} [t\alpha^2, t] &= \lim_{t \rightarrow +\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma} [t\alpha^2 + \delta_\alpha^{\alpha, \beta} + \delta_\alpha^{\alpha, \gamma}, t] = \hat{P}_\alpha(\theta), \\ \lim_{t \rightarrow -\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma} [t\beta^2 + \delta_\beta^{\alpha, \beta}, t] &= \lim_{t \rightarrow +\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma} [t\beta^2 + \delta_\beta^{\beta, \gamma}, t] = \hat{P}_\beta(\phi), \\ \lim_{t \rightarrow -\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma} [t\gamma^2 + \delta_\gamma^{\alpha, \gamma} + \delta_\gamma^{\beta, \gamma}, t] &= \lim_{t \rightarrow +\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma} [t\gamma^2, t] = \hat{P}_\gamma(\vartheta). \end{aligned} \quad (2.40)$$

When tracking the minima or maxima in the imaginary part we obtain

$$\begin{aligned} \lim_{t \rightarrow -\infty} q_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma} \left[t\alpha^2 \pm \frac{\Delta_i(\theta)}{\alpha}, t \right] &= \lim_{t \rightarrow \infty} q_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma} \left[t\alpha^2 + \delta_\alpha^{\alpha, \beta} + \delta_\alpha^{\alpha, \gamma} \pm \frac{\Delta_i(\theta)}{\alpha}, t \right] = \mp Q_\alpha(\theta), \\ \lim_{t \rightarrow -\infty} q_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma} \left[t\beta^2 + \delta_\beta^{\alpha, \beta} \pm \frac{\Delta_i(\phi)}{\beta}, t \right] &= \lim_{t \rightarrow \infty} q_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma} \left[t\beta^2 + \delta_\beta^{\beta, \gamma} \pm \frac{\Delta_i(\phi)}{\beta}, t \right] = \mp Q_\beta(\phi), \\ \lim_{t \rightarrow -\infty} q_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma} \left[t\gamma^2 + \delta_\gamma^{\alpha, \gamma} + \delta_\gamma^{\beta, \gamma} \pm \frac{\Delta_i(\vartheta)}{\gamma}, t \right] &= \lim_{t \rightarrow \infty} q_{i\theta, i\phi, i\vartheta; \alpha, \beta, \gamma} \left[t\gamma^2 \pm \frac{\Delta_i(\vartheta)}{\gamma}, t \right] = \mp Q_\gamma(\vartheta). \end{aligned}$$

Similarly as in the previous section for the two-soliton solution we read off the lateral displacements from these expressions as

$$(\Delta_x)_\alpha = \delta_\alpha^{\alpha, \beta} + \delta_\alpha^{\alpha, \gamma}, \quad (\Delta_x)_\beta = \delta_\beta^{\beta, \gamma} - \delta_\beta^{\alpha, \beta}, \quad (\Delta_x)_\gamma = -\delta_\gamma^{\alpha, \gamma} - \delta_\gamma^{\beta, \gamma}. \quad (2.41)$$

The corresponding time-delays are

$$(\Delta_t)_\alpha = -\frac{1}{\alpha^2} \left(\delta_\alpha^{\alpha, \beta} + \delta_\alpha^{\alpha, \gamma} \right), \quad (\Delta_t)_\beta = \frac{1}{\beta^2} \left(\delta_\beta^{\alpha, \beta} - \delta_\beta^{\beta, \gamma} \right), \quad (\Delta_t)_\gamma = \frac{1}{\gamma^2} \left(\delta_\gamma^{\alpha, \gamma} + \delta_\gamma^{\beta, \gamma} \right). \quad (2.42)$$

Once again we may use the values for the soliton mass (2.10) and momentum (2.11) to confirm that these quantities satisfy the consistency relation (2.4) and (2.5). As the values in (2.42) are simply the sums of the scattering of two solitons, this confirms the well known factorization property in integrable systems stating that a multiple scattering process can always be understood as consecutive scattering of two particles for which any ordering is equivalent [25]. In the quantized version of the model this property is reflected in the Yang-Baxter and bootstrap equations.

2.2.5 Properties of degenerate three-soliton solutions

Let us now consider the degenerate three-soliton solution for which the limit $\beta, \gamma \rightarrow \alpha$ is carried out. In [6] a solution to this scenario was reported as

$$u_{i\theta, i\phi, i\vartheta; \alpha, \alpha, \alpha}(x, t) = 2 \left\{ \ln \left[\left(1 + \left(\eta_{i\vartheta; \alpha}^{(3)} \right)^2 + \cosh \eta_{i\vartheta; \alpha}^{(1)} \right) \sinh \frac{\eta_{i\theta; \alpha}^{(1)}}{2} - \eta_{i\phi; \alpha}^{(9)} \cosh \frac{\eta_{i\theta; \alpha}^{(1)}}{2} \right] \right\}_{xx}, \quad (2.43)$$

with $\eta_{\mu; \alpha}^{(\lambda)} := \alpha x - \lambda \alpha^3 t + \mu$. In this case we find the three maxima for the real parts in the limits

$$\lim_{t \rightarrow \pm\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \alpha, \alpha} [t\alpha^2, t] = \hat{P}_\alpha(\theta + \pi), \quad (2.44)$$

$$\lim_{t \rightarrow \pm\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \alpha, \alpha} [t\alpha^2 \pm \bar{\Delta}(t), t] = \hat{P}_\alpha(\theta), \quad (2.45)$$

where the time-dependent displacement is defined as

$$\bar{\Delta}(t) := \frac{1}{\alpha} \ln (8\alpha^6 t^2). \quad (2.46)$$

Thus we find the center soliton converging to the sech^2 -one-soliton solution and the two outer ones to the csch^2 -one-soliton solution. The outer ones keep moving away from the center as $|t|$ increases. Similarly as for the degenerate two-soliton, the two individual outer solitons with amplitudes $\hat{P}_\alpha(\theta)$ are time-dependently displaced by the different amounts $\pm 2\bar{\Delta}(t)$. This might not be obvious from the limits (2.45) as these one-solitons are identical, but they have actually exchanged their position. The one-soliton in the center with amplitude $\hat{P}_\alpha(\theta + \pi)$ is not displaced or time-delayed and simply travels identically to a one-soliton solution. The real part of the solutions possess also minima in the regimes for θ as specified after (2.15). For those we compute

$$\lim_{t \rightarrow \pm\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \alpha, \alpha} \left[t\alpha^2 \pm \frac{1}{\alpha} \Delta_r(\theta + \pi), t \right] = \check{P}_\alpha(\theta), \quad (2.47)$$

$$\lim_{t \rightarrow \pm\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \alpha, \alpha} \left[t\alpha^2 \pm \bar{\Delta}(t) \pm \frac{1}{\alpha} \Delta_r(\theta), t \right] = \check{P}_\alpha(\theta). \quad (2.48)$$

For the imaginary parts we evaluate

$$\lim_{t \rightarrow \sigma\infty} q_{i\theta, i\phi, i\vartheta; \alpha, \alpha, \alpha} \left[t\alpha^2 \pm \frac{1}{\alpha} \Delta_i(\theta + \pi), t \right] = \mp Q_\alpha(\theta + \pi), \quad (2.49)$$

$$\lim_{t \rightarrow \sigma\infty} q_{i\theta, i\phi, i\vartheta; \alpha, \alpha, \alpha} \left[t\alpha^2 + \bar{\Delta}(t) \pm \frac{1}{\alpha} \Delta_i(\theta), t \right] = \mp Q_\alpha(\theta), \quad (2.50)$$

$$\lim_{t \rightarrow \sigma\infty} q_{i\theta, i\phi, i\vartheta; \alpha, \alpha, \alpha} \left[t\alpha^2 - \bar{\Delta}(t) \pm \frac{1}{\alpha} \Delta_i(\theta), t \right] = \mp Q_\alpha(\theta). \quad (2.51)$$

Using these limits we deduce the same values for the displacements as from the real part.

Next we consider the degenerate three-soliton solution for which only the limit $\beta \rightarrow \alpha$ is carried out, such that only two of the contributions are degenerate. We recall a solution for this from [6]

$$u_{i\theta, i\phi, i\vartheta; \alpha, \alpha, \gamma}(x, t) = 2 \left\{ \ln \left[\cosh \frac{\eta_{i\vartheta; \gamma}^{(1)}}{2} \left[\frac{\alpha^2 + \gamma^2}{8} \sinh \left(\eta_{i\theta; \alpha}^{(1)} \right) - \frac{\alpha^2 - \gamma^2}{8} \eta_{i\phi; \alpha}^{(3)} \right] \right. \right. \\ \left. \left. - \frac{\alpha\gamma}{2} \cosh^2 \left(\frac{\eta_{i\theta; \alpha}^{(1)}}{2} \right) \sinh \left(\frac{\eta_{i\vartheta; \gamma}^{(1)}}{2} \right) \right] \right\}_{xx}. \quad (2.52)$$

For the degenerated compound soliton the two maxima of the real part have the properties

$$\lim_{t \rightarrow \sigma\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \alpha, \gamma} \left[t\alpha^2 + \sigma\Delta(t) + \sigma\frac{\delta_\alpha^{\alpha, \gamma}}{2}, t \right] = \hat{P}_\alpha(\theta), \quad (2.53)$$

$$\lim_{t \rightarrow \sigma\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \alpha, \gamma} \left[t\alpha^2 - \sigma\Delta(t) + \sigma\frac{\delta_\alpha^{\alpha, \gamma}}{2}, t \right] = \hat{P}_\alpha(\theta + \pi), \quad (2.54)$$

and for the non-degenerated one-soliton contribution we compute

$$\lim_{t \rightarrow \sigma\infty} p_{i\theta, i\phi, i\vartheta; \alpha, \alpha, \gamma} [t\gamma^2 - \sigma\delta_\gamma^{\alpha, \gamma}, t] = \hat{P}_\gamma(\vartheta). \quad (2.55)$$

The degenerate one-solitons with amplitudes $\hat{P}_\alpha(\theta)$ and $\hat{P}_\alpha(\theta + \pi)$ are now time-dependently displaced due to the scattering amongst each other and in addition displaced by a constant due to the scattering by

$$(\Delta_x)_\alpha^\theta = 2\Delta(t) + \delta_\alpha^{\alpha, \gamma} \quad \text{and} \quad (\Delta_x)_\alpha^{\theta+\pi} = -2\Delta(t) + \delta_\alpha^{\alpha, \gamma} \quad (2.56)$$

respectively. From (2.55) we deduce

$$(\Delta_x)_\gamma = -2\delta_\gamma^{\alpha, \gamma} \quad (2.57)$$

the constant displacement for the one-soliton with amplitude $\hat{P}_\gamma(\vartheta)$. Our consistency equation (2.4) is satisfied as

$$m_\alpha(\Delta_x)_\alpha^\theta + m_\alpha(\Delta_x)_\alpha^{\theta+\pi} + m_\gamma(\Delta_x)_\gamma = 4\alpha\delta_\alpha^{\alpha, \gamma} - 4\gamma\delta_\gamma^{\alpha, \gamma} = 0. \quad (2.58)$$

We can argue similarly for the time-delays. Thus while the two individual degenerate contributions are time-dependently displaced, there are in addition nonvanishing constant contributions as a result of the scattering with the remaining non-degenerate one-soliton. In a similar fashion as above, these features are confirmed when tracking the minima in the real part or the minima and maxima in the imaginary part.

These features are summarized in figure 4.




Figure 4: Time-delays for a complex \mathcal{PT} -symmetric three-soliton KdV solution with a compound two-soliton with $\alpha = 6/5$, $\gamma = 4/5$, $\theta = \pi/3$ and $\vartheta = \phi = \pi/4$. The plots in the negative and positive regime of x correspond to the time taken to be $t = -30$ and $t = 30$, respectively.

2.2.6 Properties of degenerate multi-soliton solutions

We have seen in sections 2.2.3 and 2.2.5 that the individual one-soliton constituents within the degenerate two and three-soliton solutions (2.31) and (2.43) are shifted relative to each other by the time-dependent displacements (2.32) and (2.46), respectively. As these shifts are logarithmic in time the change is very slow and when confined to some finite regions they may be viewed as a compound N -soliton as advocated in [6]. This qualitative behaviour remains the same for degenerate N -soliton solutions for any N , albeit the formulae for the time-dependent displacements (2.32) and (2.46) need to be generalized.

Using the notation

$$\lim_{\alpha_2, \dots, \alpha_N \rightarrow \alpha_1 = \alpha} u_{i\theta_1 = i\theta, \dots, i\theta_N; \alpha_1, \dots, \alpha_N}(x, t) = p_{i\theta, \dots, i\theta_N; N\alpha}(x, t) + iq_{i\theta, \dots, i\theta_N; N\alpha}(x, t) \quad (2.59)$$

We compute the asymptotic limits for N even and N odd separately. For the even case we compute

$$\lim_{t \rightarrow \sigma\infty} p_{i\theta, \dots, i\theta_{2n}; 2n\alpha} [t\alpha^2 + \sigma\Delta_{n, \ell, 1}(t), t] = \hat{P}_\alpha \left(\theta + \frac{1 - (-1)^{n+\ell+1}}{2} \pi \right) \quad (2.60)$$

$$\lim_{t \rightarrow \sigma\infty} p_{i\theta, \dots, i\theta_{2n}; 2n\alpha} [t\alpha^2 - \sigma\Delta_{n, \ell, 1}(t), t] = \hat{P}_\alpha \left(\theta + \frac{1 - (-1)^{n+\ell}}{2} \pi \right) \quad (2.61)$$

for $n = 1, 2, \dots$, $\ell = 1, 2, \dots, n$ and for the odd case we obtain

$$\lim_{t \rightarrow \sigma\infty} p_{i\theta, \dots, i\theta_{2n+1}; (2n+1)\alpha} [t\alpha^2 \pm \Delta_{n, \ell, 0}(t), t] = \hat{P}_\alpha \left(\theta + \frac{1 - (-1)^{n+\ell}}{2} \pi \right) \quad (2.62)$$

for $n = 0, 1, 2, \dots$, $\ell = 0, 1, 2, \dots, n$. The time-dependent displacement takes on the general form

$$\Delta_{n, \ell, \kappa}(t) = \frac{1}{\alpha} \ln \left[\frac{(n - \ell)!}{(n + \ell - \kappa)!} (4|t|\alpha^3)^{2\ell - \kappa} \right]. \quad (2.63)$$

As in the previous cases we could also track the minima in the real part when they are present or the minima and maxima in the imaginary part, which leads to the same expressions for the time-dependent displacement (2.63).

3. Reality conditions for conserved charges

We will now argue that \mathcal{PT} -symmetry **together** with integrability will guarantee that **all** conserved charges in the model will be real. In order to see the structure of all of the charges we briefly recall how they can be constructed from the so-called Gardner transformation [26, 27, 28]. The central idea is to expand the KdV-field $u(x, t)$ in terms of a new field $w(x, t)$

$$u(x, t) = w(x, t) + \varepsilon w_x(x, t) - \varepsilon^2 w^2(x, t), \quad (3.1)$$

for some deformation parameter $\varepsilon \in \mathbb{R}$. The substitution of $u(x, t)$ into the KdV equation (2.6) yields

$$(1 + \varepsilon \partial_x - 2\varepsilon^2 w) [w_t + (w_{xx} + 3w^2 - 2\varepsilon^2 w^3)_x] = 0. \quad (3.2)$$

Since the last bracket is in form of a conservation law and needs to vanish by itself, one concludes that $\int_{-\infty}^{\infty} w(x, t) dx = \text{const.}$ Expanding the new field as

$$w(x, t) = \sum_{n=0}^{\infty} \varepsilon^n w_n(x, t) \quad (3.3)$$

then implies that also the quantities $I_n := \int_{-\infty}^{\infty} w_{2n-2}(x, t) dx$ are conserved. We may then use the relation (3.1) to construct the charge densities in a recursive manner

$$w_n = u \delta_{n,0} - (w_{n-1})_x + \sum_{k=0}^{n-2} w_k w_{n-k-2}. \quad (3.4)$$

Solving (3.4) recursively, by taking $w_n = 0$ for $n < 0$, we obtain easily the well known expressions for the first charge densities

$$w_0 = u, \quad (3.5)$$

$$w_1 = -(w_0)_x = -u_x, \quad (3.6)$$

$$w_2 = -(w_1)_x + w_0^2 = u_{xx} + u^2, \quad (3.7)$$

$$w_3 = -(w_2)_x + 2w_0 w_1 = -u_{xxx} - 2(u^2)_x, \quad (3.8)$$

$$w_4 = -(w_3)_x + 2w_0 w_2 + w_1^2 = u_{xxxx} + 6(uu_x)_x + 2u^3 - u_x^2. \quad (3.9)$$

The expressions simplify substantially when we drop surface terms and we recover the first three charges reported in (2.7). For the energy to be part of this general series is the reason why we rescaled it as compared to [5, 6].

For the charges constructed from the one-soliton solution (2.9) we obtain real expressions

$$I_n = \int_{-\infty}^{\infty} w_{2n-2}(x, t) dx = \frac{2}{2n-1} \alpha^{2n-1} \quad \text{and} \quad I_{n/2} = 0. \quad (3.10)$$

The reality of all charges build on one-soliton solutions is guaranteed by \mathcal{PT} -symmetry alone: When realizing the \mathcal{PT} -symmetry as $\mathcal{PT}: u \rightarrow u, x \rightarrow -x, t \rightarrow -t, i \rightarrow -i$ it is easily seen from (3.4) that the charge densities transform as $w_n \rightarrow (-1)^n w_n$. This mean

when $u(x, t)$ is \mathcal{PT} -symmetric so are the even graded charge densities $w_{2n}(x, t)$. Changing the argument of the functional dependence to the traveling wave coordinate $\zeta_\alpha = x - \alpha^2 t$ this means we can separate $w_{2n}(\zeta_\alpha)$ into a \mathcal{PT} -even and \mathcal{PT} -odd part $w_{2n}^e(\zeta_\alpha) \in \mathbb{R}$ and $w_{2n}^o(\zeta_\alpha) \in \mathbb{R}$, respectively, as $w_{2n}(\zeta_\alpha) = w_{2n}^e(\zeta_\alpha) + iw_{2n}^o(\zeta_\alpha)$, which allows us to conclude

$$I_n(\alpha) = \int_{-\infty}^{\infty} w_{2n-2}(x, t) dx = \int_{-\infty}^{\infty} [w_{2n-2}^e(\zeta_\alpha) + iw_{2n-2}^o(\zeta_\alpha)] d\zeta_\alpha = \int_{-\infty}^{\infty} w_{2n-2}^e(\zeta_\alpha) d\zeta_\alpha \in \mathbb{R}. \quad (3.11)$$

It is easily seen that the previous argument applies directly to the charges build from the solution $u_{i\theta; \alpha}(x, t)$ in (2.9), i.e. the real part and imaginary part are even and odd in ζ_α , respectively. When the parameter μ has a nonvanishing real part the \mathcal{PT} -symmetry is broken, but it can be restored by absorbing the real part by a shift either in t or x as argued in [5].

In order to ensure the same for the multi-soliton solutions we use the fact that the multi-soliton solutions separate asymptotically into single solitons with distinct support. As the charges are conserved in time we may compute I_n at any time. In the asymptotic regime any charge build from an N -soliton $u_{i\theta_1, \dots, i\theta_N; \alpha_1, \dots, \alpha_N}^{(N)}$ decomposes into the sum of charges build on the one-soliton solutions.

$$I_n(\alpha_1, \dots, \alpha_N) = \int_{-\infty}^{\infty} \left(w_{i\theta_1, \dots, i\theta_N; \alpha_1, \dots, \alpha_N}^{(N)} \right)_{2n-2}(x, t) dx, \quad (3.12)$$

$$= \int_{-\infty}^{\infty} \sum_{k=1}^N \left[\left(w_{i\theta_k; \alpha_k}^{(1)} \right)_{2n-2}(\zeta_{\alpha_k}) \right] d\zeta_{\alpha_k}, \quad (3.13)$$

$$= \sum_{k=1}^N I_n(\alpha_k), \quad (3.14)$$

$$= \frac{2}{2n-1} \sum_{k=1}^N \alpha_k^{2n-1}. \quad (3.15)$$

We used here the decomposition of the N -soliton into a sum of one-solitons in the asymptotic regime $u_{i\theta_1, \dots, i\theta_N; \alpha_1, \dots, \alpha_N}^{(N)} = \sum_{k=1}^N \left(u_{i\theta_k; \alpha_k}^{(1)} \right)$, which we have seen in detail above. Since each of the one-solitons is well localized we always have $u_{i\theta_k; \alpha_k}^{(1)} \cdot u_{i\theta_l; \alpha_l}^{(1)} = 0$ when $k \neq l$, which implies that

$$\left[u_{i\theta_1, \dots, i\theta_N; \alpha_1, \dots, \alpha_N}^{(N)} \right]^m = \left[\sum_{k=1}^N \left(u_{i\theta_k; \alpha_k}^{(1)} \right) \right]^m = \sum_{k=1}^N \left(u_{i\theta_k; \alpha_k}^{(1)} \right)^m. \quad (3.16)$$

As all the derivatives are finite and the support is the same as for the u s, this also implies

$$\left[\left(u_{i\theta_1, \dots, i\theta_N; \alpha_1, \dots, \alpha_N}^{(N)} \right)_{nx} \right]^m = \left[\sum_{k=1}^N \left(u_{i\theta_k; \alpha_k}^{(1)} \right)_{nx} \right]^m = \sum_{k=1}^N \left(u_{i\theta_k; \alpha_k}^{(1)} \right)_{nx}^m, \quad (3.17)$$

and similarly for mixed terms involving different types of derivatives. As all charge densities are made up from u and its derivatives we obtain

$$\left(w_{i\theta_1, \dots, i\theta_N; \alpha_1, \dots, \alpha_N}^{(N)} \right)_{2n-2} = \sum_{k=1}^N \left(w_{i\theta_k; \alpha_k}^{(1)} \right)_{2n-2} \quad (3.18)$$

in the asymptotic regime, which is used in the step from (3.12) to (3.13). In the remaining two steps (3.14) and (3.15) we use (3.10).

Thus \mathcal{PT} -symmetry **and** integrability guarantee the reality of **all** charges.

4. Conclusions

We have explicitly computed lateral displacements and time-delays for complex two and three-soliton solutions of the KdV equation. Our solutions satisfy the consistency equations (2.4) and (2.5) resulting from the preservation of the centre of mass coordinate. The expressions for the time-delay of the three-soliton scattering, being the sum of the delays of two two-soliton time-delays, confirm on classical level the standard factorization property of the scattering matrix for integrable systems that allows to treat any multiple scattering process as a succession of two particle scatterings. The imaginary part in our solutions may be thought of as a massless soliton partaking in the scattering process.

We used our expressions for three different purposes: Firstly we made the relation between solutions obtained from Hirota's direct method on one hand and those constructed from a superposition principle based on Bäcklund transformations precise. They differ by non-identical lateral displacements in each of their one-soliton constituents and additional shifts by π in the shift parameters. Overall they lead to the same values of the time-delays as they are preserved quantities. Secondly we elaborated on the internal structure of compound soliton solutions within degenerate multi-soliton solutions. We found that the degenerate one-soliton contributions are displaced relative to each other by a time-dependent shift. When scattered with any nondegenerate one-soliton constituent they are all displaced by the same amount. With (2.63) we presented a generic formula for the relative time-dependent displacement valid for any degenerate N -soliton solution. Thirdly we clarified the role \mathcal{PT} -symmetry plays in guaranteeing the reality of conserved charges, the energy being one of them. It turned out that \mathcal{PT} -symmetry is solely responsible for the reality of any charge based on one-soliton solutions. For charges constructed from multi-soliton solutions we need to invoke integrability having the effect of separating asymptotically the multi-solitons into single solitons to ensure the reality these charges.

As our approach is entirely model independent, it would naturally be interesting to apply it to other complex integrable systems. Furthermore, it would be very interesting to employ the expressions obtained for a semi-classical quantization. In particular the role played here by the massless soliton might shed some new light on some old results [29].

Acknowledgments: FC would like to thank the Alexander von Humboldt Foundation (grant number CHL 1153844 STP) for financial support and City University London for kind hospitality.

A. Sample time-delay computations

Most of the computations are rather cumbersome, so that it suffices to present a few samples. Let us for instance derive the values for the shifts in (2.20). For simplicity we take the real part of $u_{i\theta, i\phi, \alpha, \beta}(x, t)$ in (2.19) at specific values of θ and ϕ . Taking $\theta = \phi = \pi/2$ it acquires the relatively simple form

$$4 \frac{\left[\alpha e^{\alpha^3 t + 2\beta^3 t + \alpha x} + \beta e^{2\alpha^3 t + \beta^3 t + \beta x} + \alpha \chi(\alpha, \beta) e^{\alpha^3 t + \alpha x + 2\beta x} + \beta \chi(\alpha, \beta) e^{\beta^3 t + 2\alpha x + \beta x} \right]^2}{\left[e^{2t(\alpha^3 + \beta^3)} + \frac{8\alpha\beta}{(\alpha + \beta)^2} e^{t(\alpha^3 + \beta^3) + x(\alpha + \beta)} + e^{2\alpha^3 t + 2\beta x} + e^{2\beta^3 t + 2\alpha x} + \chi^2(\alpha, \beta) e^{2x(\alpha + \beta)} \right]^2}. \quad (\text{A.1})$$

Tracking the soliton with speed α^2 in the multi-soliton solution and trying to match it with the one-soliton, we may try to solve the equivalence relation

$$u_{i\pi, i\pi; \alpha, \beta}(t\alpha^2 + \Delta, t) \sim u_{i\pi; \alpha}(t\alpha^2, t), \quad (\text{A.2})$$

in the asymptotic regimes $t \rightarrow \pm\infty$ for some as yet unknown constant Δ . The right hand side of (A.2) is easily computed from (2.9) to α^2 . Replacing x by $t\alpha^2 + \Delta$ in (A.1) and identifying $e^{4t(\alpha^3 + \alpha^2\beta)}$ as the dominant term in the numerator and denominator in the large t regime we read off the corresponding coefficients. Thus for large t the equivalence relation (A.2) becomes

$$\frac{4\alpha^2 (\alpha^2 - \beta^2)^4 e^{2\alpha\Delta}}{2 (\alpha^2 - \beta^2)^4 e^{2\alpha\Delta} + (\alpha - \beta)^8 e^{4\alpha\Delta} + (\alpha + \beta)^8} = \alpha^2. \quad (\text{A.3})$$

Solving this equation for Δ leads to the time-displacement $\Delta = 2/\alpha \ln[(\alpha + \beta)/(\alpha - \beta)]$ for $t \rightarrow \infty$, which corresponds to the shift reported in (2.20) for generic values of θ and ϕ .

Next we present a sample computation for a degenerate multi-soliton solution for which the lateral displacement becomes time dependent rather than just being constant. We derive the asymptotic relation (2.33). Taking x to be $t\alpha^2 + \Delta$ in the degenerate two-soliton solution (2.31) for some constant Δ , we observe that the limits $t \rightarrow \pm\infty$ always yield zero and we will not be able to obtain a finite value such as the maximum of the one-soliton $\hat{P}_\alpha(\theta)$. Hence we are forced to include a time-dependence into Δ . This mildest dependence we may introduce is a logarithmic one. Taking therefore as an Ansatz $\Delta(t) = -1/\alpha \ln(\kappa |t|)$ for some unknown constant κ we compute

$$\frac{4\alpha^2 e^{i\theta} \kappa |t| [\kappa^2 t^2 (2\alpha^3 t - \nu - 2 + \ln(\kappa |t|)) - e^{2i\theta} (2\alpha^3 t + 2 - \nu + \ln(\kappa |t|)) - 4e^{i\theta} \kappa |t|]}{[e^{2i\theta} - \kappa^2 t^2 - 2e^{i\theta} \kappa |t| (2\alpha^3 t - \nu + \ln(\kappa |t|))]^2}. \quad (\text{A.4})$$

For large $|t|$ we can now find matching powers in t in the numerator and denominator. We replace now $|t|$ by σt and take σ to be ± 1 depending on whether t is negative or positive. The leading order terms are proportional to t^4 . Neglecting all other terms, the expression in (A.4) reduces to

$$\frac{8\alpha^5 e^{i\theta} \kappa \sigma}{(\kappa + 4\sigma\alpha^3 e^{i\theta})^2}. \quad (\text{A.5})$$

For $\kappa = 4\alpha^3$ this equals $\hat{P}_\alpha(\theta)$ and $\hat{P}_\alpha(\theta + \pi)$ for $\sigma = 1$ and $\sigma = -1$, respectively. Thus we have derived (2.33) and (2.34) with the lateral displacement taking on the form (2.32).

References

- [1] J. Rubinstein, Sine-Gordon Equation, *Journal of Mathematical Physics* **11**, 258–266 (1970).
- [2] R. Jackiw and G. Woo, Semiclassical scattering of quantized nonlinear waves, *Physical Review D* **12**, 1643 (1975).
- [3] A. Fring, P. Johnson, M. Kneipp, and D. I. Olive, Vertex operators and soliton time delays in affine Toda field theory, *Nuclear Physics B* **430**, 597–614 (1994).

- [4] E. P. Wigner and L. Eisenbud, Higher angular momenta and long range interaction in resonance reactions, *Physical Review* **72**(1), 29 (1947).
- [5] J. Cen and A. Fring, Complex solitons with real energies, arXiv preprint arXiv:1602.05465, accepted for publication in *Journal of Physics A: Mathematical and Theoretical* (2016).
- [6] F. Correa and A. Fring, Regularized degenerate multi-solitons, arXiv preprint arXiv:1605.06371 (2016).
- [7] E. Wigner, Normal form of antiunitary operators, *J. Math. Phys.* **1**, 409–413 (1960).
- [8] C. M. Bender and S. Boettcher, Real Spectra in Non-Hermitian Hamiltonians Having PT Symmetry, *Phys. Rev. Lett.* **80**, 5243–5246 (1998).
- [9] C. M. Bender, Making sense of non-Hermitian Hamiltonians, *Rept. Prog. Phys.* **70**, 947–1018 (2007).
- [10] A. Mostafazadeh, Pseudo-Hermitian Representation of Quantum Mechanics, *Int. J. Geom. Meth. Mod. Phys.* **7**, 1191–1306 (2010).
- [11] C. M. Bender, D. C. Brody, J. Chen, and E. Furlan, \mathcal{PT} -symmetric extension of the Korteweg-de Vries equation, *J. Phys. A* **40**, F153–F160 (2007).
- [12] A. Fring, \mathcal{PT} -Symmetric deformations of the Korteweg-de Vries equation, *J. Phys. A* **40**, 4215–4224 (2007).
- [13] C. M. Bender and J. Feinberg, Does the complex deformation of the Riemann equation exhibit shocks?, *J. Phys. A* **41**, 244004(8) (2008).
- [14] B. Bagchi and A. Fring, \mathcal{PT} -symmetric extensions of the supersymmetric Korteweg-De Vries equation, *J. Phys. A* **41**, 392004(9) (2008).
- [15] P. E. G. Assis and A. Fring, Integrable models from \mathcal{PT} -symmetric deformations, *J. of Phys. A* **42**, 105206 (2009).
- [16] C. M. Bender, F. Cooper, A. Khare, B. Mihaila, and A. Saxena, Compactons in PT-symmetric generalized Korteweg-de Vries Equations, *Pramana Journal of Physics* **73**, 375–385 (2009).
- [17] P. E. G. Assis and A. Fring, Compactons versus solitons, *Pramana Journal of Physics* **74**, 857–865 (2010).
- [18] A. Cavaglia, A. Fring, and B. Bagchi, \mathcal{PT} -symmetry breaking in complex nonlinear wave equations and their deformations, *J. Phys. A* **44**, 325201(42) (2011).
- [19] Z. Yan, Complex \mathcal{PT} -symmetric extensions of the nonlinear ultra-short light pulse model, *J. Phys. A* **45**, 444035 (2012).
- [20] A. Cavaglia and A. Fring, \mathcal{PT} -symmetrically deformed shock waves, *J. Phys. A* **45**, 444010(16) (2012).
- [21] P. E. G. Assis, PT-symmetry in quasi-integrable models, *Journal of Physics A: Mathematical and Theoretical* **49**(24), 245201 (2016).
- [22] R. Hirota and J. Satsuma, Soliton solutions of a coupled Korteweg-de Vries equation, *Phys. Lett. A* **85**, 407–408 (1981).
- [23] M. Ito, Symmetries and conservation laws of a coupled nonlinear wave equation, *Phys. Lett. A* **91**, 335–338 (1982).

- [24] R. Hirota, Exact Solution of the Korteweg-de Vries Equation for Multiple Collisions of Solitons, *Phys. Rev. Lett.* **27**, 1192 – 1194 (1971).
- [25] T. Hollowood, Solitons in affine Toda field theory, *Nucl. Phys.* **B384**, 523–540 (1992).
- [26] R. M. Miura, C. S. Gardner, and M. D. Kruskal, Korteweg-de Vries equation and generalizations. II. Existence of conservation laws and constants of motion, *Journal of Mathematical physics* **9**(8), 1204–1209 (1968).
- [27] R. M. Miura, The Korteweg-de Vries Equation: A Survey of Results, *SIAM Review* **18**, 412–459 (1976).
- [28] B. A. Kupershmidt, On the nature of the Gardner transformation, *Journal of Mathematical Physics* **22**(3), 449–451 (1981).
- [29] A. B. Zamolodchikov and A. B. Zamolodchikov, Massless factorized scattering and sigma models with topological terms, *Nuclear Physics B* **379**(3), 602–623 (1992).