



City Research Online

City, University of London Institutional Repository

Citation: Flinton, D. M., Singh, M. K. & Haria, K. (2018). Readability of internet based patient information for Radiotherapy Patients. *Journal of Radiotherapy in Practice*, 17(2), pp. 142-150. doi: 10.1017/s1460396917000620

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/18872/>

Link to published version: <https://doi.org/10.1017/s1460396917000620>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

David Flinton, EdD, MSc, BSc

Manrita Singh, BSc

Krupesh Haria, BSc

Readability of Internet based patient information for Radiotherapy Patients.

Abstract

Background. Information is key to patient informed choice and the internet is currently a major source of health information for adults in the UK. In order for the users to make use of the information it must be presented in a way that the user can understand. This depends on a number of factors one being that the document is written at the right level to be understood by the reader, readability. Aim. The aim of this study was to assess the readability of radiotherapy related documents on the internet and compare their levels to published norms. Method. An internet search was undertaken using Google, to identify UK based literature. Once identified documents were downloaded into Word and cleaned of punctuation other than that at the end of the sentence, documents were then analysed by the software package Readability Studio. Results and Conclusions. Documents tended to be written at too high a reading level, but the reading level had improved from a similar study conducted in 2006. The level of readability appears to show a relationship to the use of passive voice, which was very variable in the sample collected and reduction in the use of passive voice could help with the readability of the information.

Introduction

Information is essential to inform patient choice and can aid in improving patients' experiences. Patients require information about their diagnosis, treatment options and other issues surrounding their treatment if choices are to be informed and guidance followed. Health professionals have a responsibility to provide this information which can take a number of different forms. A common medium used to provide patients with information used widely within the NHS and other institutions to supplement verbal information is that of printed information leaflets (PILs) available within the departments and now ever more increasingly as on-line material via the internet. The internet has seen a significant increase in its use as a source of information with over 80% of the adult population using the internet in 2016 compared with approximately 35% in 2006¹. The use of the internet is different at different life stages, however one of the main uses of the internet is that of finding healthcare information^{1,2} which both the employed (45%) and retired (39%) user groups are more effective at when compared with students (32%)².

Using information which can be read can offer several advantages over verbal communication as patients are frequently distressed and may not fully comprehend or remember information provided in a face to face meeting. It is also possible to review the information once provided, which allows the document to be used as a reference source throughout their treatment, giving patients a greater awareness of what to expect and allowing them to make more informed choices. Written information is a key element for the patient, therefore, it is important that information is easy to understand. The NHS information standard principles state that "each product is in plain language".³ The National Cancer Patient Experience Survey⁴ stated that of patients who received information about the type of cancer that they had 28% did not find the information easy to understand. This approximately corresponds with the findings of both the Organisation for Economic Co-operation and Development (OECD) report⁵ and skills for life survey 2015⁶. The OECD report stated that approximately 9 million working aged adults in England (over a quarter of adults aged 16-65) have low literacy or numeracy skills or both. In terms of literacy, this would mean that they would have difficulty with simple written information. The Skills for Life survey reported that half of businesses stated that they were aware of problems in basic literacy amongst some of their employees, and more than 40% of businesses in the past year have had to provide remedial training in basic skills for at least some of their adult employees. In most countries, younger people (16-24 years of age) have better literacy levels than those nearing retirement, however, in the UK, figures for both groups are very similar⁵. Because of this and other factors that affect reading, the UK Government recommends a reading age of 9 for web based materials⁷.

Several key issues need to be considered when reviewing whether written information is effective. Firstly, authors need to know what information is required by the patient group, and whether everything relevant is included and correct. Once the information has been quality assured, it is essential that the document is constructed in such a way as to be easily understood by the intended patient group. How effective any written communication will be is dependent on a number of factors, including how legible and readable the target patient group finds the document.

Legibility is a measure of how well the text can be viewed or read. Legibility does not consider how well the reader can interpret the information, but merely how apparent the written information is. Legibility can be affected by a number of factors such as structure and design of the written information and includes numerous factors such as, the font, contrast between the text and the background, the size of the font, letter and word spacing or how far away the screen is; it is also affected by age⁸.

Readability can be defined as how easy the text is to comprehend due to the style of writing. A common way of measuring readability is to use readability formulae that were first developed in the USA in the early twentieth century. Readability formulas are simple algorithms that aid in the objective comparison of text. Most readability formulas look at the complexity of the words used within the document (semantic) and the sentence length (syntactic) elements of the work^{9,10}.

The formulas have the advantage of being a quick method of predicting the approximate readability of a document with evidence suggesting that they are related to the speed of reading, the probability that the document will be read and the knowledge of content after the article has been read¹¹. They are, however, not without fault and have been subject to much criticism. Although there are validation studies most readability formulae are not based on any particular theory of reading, but rather on observed correlations¹². However, the main criticisms are reserved for the fact they fail to account for many other factors known to affect readability, such as whether the material makes sense, the style of writing, prior knowledge of the reader, the appropriateness of the vocabulary or the design features that may hinder or help the reader^{13,14}.

The aim of this study was to look at literature aimed at patients undergoing radiotherapy, specifically to the information about treatment and advice on side-effects. Due to large variation of cancers, the study was limited to the web pages offering advice on prostate and

breast cancer aiding direct comparison of the documents. The rationale behind using these two cancers was that they represent the largest patient groups and the evidence collected could be compared with a similar study undertaken in 2010¹⁵.

Method

An online search was conducted using Google to source patient information pages linked to radiotherapy treatments of either the prostate or breast. In order to source patient information pages general search terms were used, for example, terms such as “breast”, “prostate,” “radiotherapy,” “treatment” and “information”. Data collection of information ended in January 2016. Once suitable internet pages had been sourced the pages were copied into Microsoft Word files; the resulting documents were then cleaned; a process that involved removing headings, headers footers, titles, copyright information and contact details. Also, any punctuation within the sentence such as, “thirty-two”, or “C.T.” were removed. This process was undertaken in order to improve the consistency of the readability statistics generated by each document, which may vary by up to 2 reading grades due to the way each different formula treats punctuation¹⁶.

Files were then imported into Readability Studio 2015, Oleander Software Ltd, (Vandalia, USA) for readability analysis. This study utilised well established formulae authors had previously used to calculate readability within healthcare settings: the four formulae were, Flesch Kincaid which is the most commonly used test in healthcare documents (57.42%)¹⁷, Simple Measure Of Gobbledygook (SMOG) which Wang¹⁶ states appears best suited for health care applications and both the New Dale-Chall and Gunning FOG measures.

Readability Studio produces both the reading age and reading grade for all four tests. This article will utilise the reading grade, which corresponds to the US grade level of education, as this is the more commonly quoted figure used in publications. Conversion between reading grade and reading age is relatively straight forward with the reading age equalling the reading grade level +5¹⁸ for example the average age of students in the 8th grade being 13 years old. Details of what the tests look for in a document to work out readability score can be found in Table 1. The fifth formula utilised, the Flesch Reading Ease scale does not report a reading age or grade, but rather grades text on a 0–100 scale with scores of 0–30 corresponding to very difficult; 30–50, difficult; 50–60, fairly difficult; 60–70, standard; 70–80, fairly easy; 80–90, easy; and 90–100, very easy. The Flesch Reading Ease scale was also included as again it is a commonly used test and it is commonly available, being included in many word processing software packages such as Microsoft Word.

Some studies have shown that an excessive use of passive voice construction can be associated with reduced levels of readability. Passive voice is when the focus of the sentence is on the action, not who or what is performing the action, and it is recommended that its use should be minimised to less than 10%¹⁹. The amount of passive voice used in the text is also calculated by the Readability Studio software and this metric was also recorded for each document. {Table 1}

Statistical analysis was performed using IBM SPSS 22, the significance level was set at 0.05.

Results Readability

48 separate information sources were found and analysed, which were divided equally between breast and prostate information documents. The vast majority 43 (89.6%) of the information websites were from hospitals posting advice online, 5 (10.4%) came from support groups such as Cancer Research UK.

The Flesch Reading Ease scale rated 26 (54%) of the articles as being of a standard level, 20 (42%) as difficult and 2 (4%) as very difficult with an average score for all documents of 59.9 (95% Confidence Interval (CI) = 58.2-61.6). Collectively, when averaged using the four readability scales, 48 internet pages were found to be written at a mean reading grade of 9.39, s.d. 0.52, with very comparable results for both prostate documents, mean 9.84, standard deviation (s.d.) 0.75, and breast documents, mean 9.83, s.d. 0.67. The individual documents had an average reading grade of between 8.4 and 11.5. Figure 1 shows the article scores. {Insert Figure 1}

There was considerable variation in length of the information, the mean length being 2,240 words with a range between 535 and 5,108. The length of the document had no correlation to the readability of the text $r = -0.041$, $p = 0.713$.

Passive voice

Overall the mean use of passive voice was 12.4%, s.d. 4.5, ranging from 2.78% to 20.23% with only 15 documents (31.3%) having a passive voice being less than 10% of the time. {Figure 2} There was a medium positive correlation between the use of passive voice in the documents and their readability score $r = 0.389$, $p = 0.006$. {Figure 3}

Comparison with data from 10 years ago

A sample of 85 patient information leaflets were collected in 2006 from seventeen radiotherapy departments for readability analysis. The results were presented at the Radiotherapy in

Practice conference and published in 2008¹⁵. The sample examined included 20 breast and 15 prostate information leaflets. These leaflets were re-examined using Readability Studio using the same parameters as those used for the more recent documents to try and establish if the readability of the documents had changed over time. Like the documents sourced in the more recent data collection, there was a considerable range of document size ranging between 332 and 4,264 with a mean of 1,287 words.

The overall average scores for readability were 9.98 in 2006 compared to 9.39 in 2016. This change in readability scores was significant, $t=3.063$, $p=0.03$. An improvement in the use of passive voice was also observed decreasing from 15.01% in 2006 to 12.64% in 2016 although this difference did not reach significance, $t=1.629$, $p=0.107$. {Figure 4 & 5}

Discussion

The internet provides anonymous, convenient access to a wealth of healthcare information with an ever-increasing reliance on the web for health information^{20,21}. Over 70% of patients report that the information they collect from the Internet influences their treatment decisions²². In order to ensure the information is readable by their target audience, the American Medical Association and the National Institutes of Health recommend that patient education resources should be written at no higher than the sixth-grade level. The International Patient Decision Aid Standards Collaboration²³ recommends health documents be written at grade 8 equivalent or less and even go so far as to recommend which readability test be used, SMOG or Fry²⁴. The average reading grade of the documents in this study was 9.39, which is slightly higher than that recommended, by the above institutions and well above the target figure of the reading age of a 9 year old (grade 4) suggested by the Government⁷.

No document was written at a reading grade of less than 8, but four documents (8.3%) did have a reading grade between 8 and 9. The reading grade reported here although higher than that recommended, is better than that found in many other studies. One study looking at patient education materials available on the European Society of Radiology (ESR) website found articles to have a mean grade level of 13.0 ± 1.6 with a range from 10.8 to 17.2²⁵. Similarly, three other studies^{26,27,28} found that compliance to the guidelines were poor. Fitzsimmons²⁶ found 60–89% of online Parkinson's disease information webpages were written above the 12th grade level, and Trivedi²⁷ noted or reported that medicinal labels had an average reading age of a 16 year old (reading grade 11). Finally Weiss²⁸ who looked at online lung cancer information found an average reading grade of 11.2. No material from any study complied with the American maximum recommended sixth-grade level. A further study looking at readability of laryngeal cancer patient information leaflets²⁹ and reported a Flesch

Reading Ease score of 48.2 difficult. The results in the current study were again better, the mean score reported being 59.9 fairly difficult, which was very close to the cut-off point of them being classed as a standard document or plain English which starts at 60.

There was no significant relationship between document length and readability, suggesting that reading difficulty is independent of the word count.

Despite the fact that the online information did not reach the recommended reading grade, it must be acknowledged that there had been a significant improvement in the readability of patient information in respect to the articles examined in 2006¹⁵.

One issue with readability formulae is that they ignore vocabulary and tend to assume a strong negative correlation between word length and readability. For example, “iff” meaning “if and only if” only has one syllable and therefore has a low semantic measure in many of the readability scales outlined in Table 1, despite many people probably not knowing what the word means, whereas “wheelbarrow” would give a larger semantic measure despite most people being able to read it and understand what it means. Our analysis utilised four readability scales to determine the level of readability of the text to try and overcome the limitations present in the metric of any one formula. A number of different formulae could have been used, Readability Studio being able to produce readability outputs for over 30 such tests. The justification outlined earlier was the common use of these tests in other readability publications in healthcare settings, however, a different selection of tests would have resulted in a slightly different readability score. Finally writing that utilises technical language does tend to have a slightly raised level of difficulty³⁰ and technical words such as radiotherapy and physiotherapy could raise the readability level of the document and it is difficult to think of alternatives to make the writing easier, however, the use of supraclavicular, and telangiectasia, terms present in seven documents examined as part of this study would be definite cases where alternative, simpler terms, could and should have been used.

The average use of passive voice within the documents was 12.4%, which was in keeping with that found by Harwood & Harrison³¹ (12.8%), but higher than that found by Pothier³² who reported average passive voice figures of 8.31%. The figure is also slightly higher than that recommended by the Plain English Campaign of 10%. As can be seen in Figure 2, there is considerable variance in the amount of passive voice used, with many documents falling into the acceptable category (48%). The information present on web pages will probably have been written by healthcare professionals with degrees, where the use of the third person and passive voice is promoted in many forms of written work. The moderation of passive voice

within the documents could lead to a further improvement in the readability level of many of the documents and comparison with the 2006 data did show a marked, but non-significant reduction in the use of passive voice so perhaps this is an area of writing style that is improving.

Conclusions

The results of this study revealed a difference among websites of both reading level and the levels of use of the passive voice with most online material being written above the proposed target reading grade of 8, however the results were noticeably better than those obtained in 2006 and that reported in other publications of online material. Some authors could improve their writing style and improve the readability of the documents produced, by more careful selection of terms and by restricting the use of passive voice, the level of which was very variable in the documents. The correlation between the two indices, reading age and passive voice probably reflects the increased syntactic difficulty of the documents that tends to occur due to the longer sentence structure when writing in the passive voice and the benefit in terms of making the documents simpler to read can be gained from looking at this aspect of the writing.

Acknowledgements

None

References

1. Office for National Statistics. (2016) Statistical Bulletin. Internet Access – households and individuals: 2016. What the internet is used for and types of purchases made by adults (aged 16 or over). Accessed online 20th August 2017.
2. Dutton, W., Blank, G. (2011) Next Generation Users: The Internet in Britain 2011. Oxford Internet Institute. University of Oxford. Oxford. England.
3. NHS England. The information standard Principles.
<https://www.england.nhs.uk/tis/about/the-info-standard/> Accessed online 22nd August 2017.
4. NHS England (2016) The National Cancer Patient Experience Survey 2015. Summary of key national and local results.
https://www.england.nhs.uk/statistics/wpcontent/uploads/sites/2/2016/06/20160713-CPES-2015-Presentation_revised.pdf Accessed online 22nd August 2017.
5. Kuczera, M., Field, S., Windisch, H.C. (2016) Building Skills for All: A Review of England. OECD Skills Studies. OECD, Paris, France.
6. Confederation of British Industry (CBI) (2015) Inspiring Growth. CBI/Pearson Education and Skills Survey 2015. CBI, London, UK.
7. Government Digital Services (2016) Content design: planning, writing and managing content. <https://www.gov.uk/guidance/content-design/writing-for-gov-uk> Accessed online 22nd August 2017.
8. Wolfe, B., Dobres, J., Kosovicheva, A., Rosenholtz, R., Reimer, B. (2016) AgeRelated differences in the legibility of degraded text. Cognitive Research: Principles and Implications. 1:22, 1-13.
9. Waller, E. (2011) What makes a good document? The criteria we use. Technical Paper 2. Simplification Centre. University of Reading, Reading, UK.
10. Kang, T., Elhadad, N., Weng, C. (2015) Initial Readability Assessment of Clinical Trial Eligibility Criteria. AMIA Annu Symp Proc. 687–696.

11. Janan, D., Wray, D. (2012) Readability: The limitations of an approach through formulae. Paper presented at the British Educational Research Association Annual Conference, University of Manchester, Manchester England.
12. Crossley, S.A., Skalicky, S., Dascalu, M., McNamara, D.S., Kyle, K. (2017) Predicting Text Comprehension, Processing, and Familiarity in Adult Readers: New Approaches to Readability Formulas. *Discourse Processes*. 54:5–6, 340–359.
13. Bailin, A., Grafstein, A. (2001) "The linguistic assumptions underlying readability formulae: a critique." *Language & Communication*. 21: 285–301.
14. Doak, C.C., Doak, L.G., Root, J.H. (1996). *Teaching Patients with Low Literacy Skills*. 2nd Edition. Lippincott Williams and Wilkins. Philadelphia, USA.
15. Flinton D. (2008) Readability and Legibility of Printed Information Leaflets in Radiotherapy. *Journal of Radiotherapy in Practice*. 7, 186-187.
16. Zouh, S., Jeong, H., Green, P.A. (2017) How Consistent Are the Best-Known Readability Equations in Estimating the Readability of Design Standards? *IEEE Transactions on Professional Communication*. 60:1, 97-111.
17. Wang, L-W., Miller, M.J., Schmitt, M.R., Wen, F.K. (2013) "Assessing readability formula differences with written health information materials: Application, results, and recommendations." *Research in Social and Administrative Pharmacy*. 9: (5), 503– 516.
18. Johnson, K. (1998) Readability. <http://www.timetabler.com/readable.pdf> Accessed online 22nd August 2017.
19. Plain English Campaign (2001) How to write medical information in plain English. Plain English Campaign. <http://www.plainenglish.co.uk/files/medicalguide.pdf> Accessed online 20th August 2017.
20. Blair, J. (2004) Assessing the value of the internet in health improvement. *Nursing Times*. 100: 35, 28–30.

21. Penson, R.T, Benson, R.C., Parles, K., Chabner, B.A., Lynch Jr, T.J. (2002) "Virtual Connections: Internet Health Care." *The Oncologist*. 7: 555-568.
22. Hansberry, D.R., Agarwal, N., Baker, S.R. (2015) "Health Literacy and Online Educational Resources: An Opportunity to Educate Patients." *American Journal of Roentgenology*. 204, 111-116.
23. International Patient Decision Aid Standards (IPDAS) Collaboration (2005) Criteria for Judging the quality of Patient Decision Aids.
http://ipdas.ohri.ca/IPDAS_checklist.pdf Accessed online 20th August 2017.
24. Smith, S. (2010) Guide to Appraising Health Information. Patient Information Forum.
<https://www.pifonline.org.uk/wp-content/uploads/2014/11/PIF-GuideAppraising-Health-Information-2010.pdf> Accessed online 22nd August 2017.
25. Hansberry, D.R, Ann John, A., John, E., Agarwal, N., Gonzales, S.F., Baker, S.R. (2014) A Critical Review of the Readability of Online Patient Education Resources from RadiologyInfo.Org. *American Journal of Roentgenology*. 202: (3), 566-575.
26. Fitzsimmons, P.R., Michael, B.D., Hulley, J.L., Scott, G.O. (2010) A readability assessment of online Parkinson's disease information. *J R Coll Physicians Edinb* 2010; 40: 292–296.
27. Trivedi, H., Trivedi, A., Hannan, M.F. (2014) Readability and comprehensibility of over-the-counter medication labels. *Renal Failure*. 36: (3), 473–477.
28. Weiss, K.D., Vargas, C.R., Ho., O.A., Chuang, D.J., Weiss, J., Lee, B.T. (2016) Readability analysis of online resources related to lung cancer. *Journal of Surgical Research*. 206: (1), 90-97.
29. Narwani, V., Nalamada, K., Lee, M., Kothari, P., Lakhani, R. (2015) Readability and quality assessment of internet-based patient education materials related to laryngeal cancer. *Head & Neck*. 28: (4), 601-605
30. Janan, D., Wray, D. (2012) Readability: The limitations of an approach through formulae. Paper presented at the British Educational Research Association Annual Conference.
<http://www.leeds.ac.uk/educol/documents/213296.pdf> Accessed 24th May 2017.

31. Harwood, A., Harrison, J.E. (2004) How readable are orthodontic patient information leaflets? *Journal of Orthodontics* 31: (3), 210-219.
32. Pothier, L., Day, R., Harris, C., Pothier, D.D. (2008) Readability statistics of patient information leaflets in a Speech and Language Therapy Department. *International Journal of Language and Communication Disorders*. 43: (6), 712–72.

Figure 1. Reading grade of web documents downloaded.




Figure 2. Passive voice usage in web documents downloaded.




Figure 3. Relationship between Passive voice use and readability grade.




Figure 4. Readability of 2006 documents compared to 2016 documents.




Figure 5. Comparison in the use of passive voice in 2006 and 2016.




Table 1. Semantic and Syntactic measures used in Readability formulae.

	Semantic Measure	Syntactic Measure
Flesch-Kincaid*	Total Syllables/Total words	Total words/total sentences
Dale-Chall	% of words on a word list	Average number of words per sentence
Gunning Fog	Words of more than 2 syllables	Average sentence length
SMOG	Number of 3+ syllable words	Average number of sentences

* The Flesch-Kincaid score is a conversion of the Flesch Reading Ease score.