



City Research Online

City, University of London Institutional Repository

Citation: Gardulf, A., Abolhassani, H., Gustafson, R., Eriksson, L. E. & Hammarström, L. (2018). Predictive markers for humoral influenza vaccine response in patients with common variable immunodeficiency (CVID). *Journal of Allergy and Clinical Immunology*, 142(6), 1922-1931.e2. doi: 10.1016/j.jaci.2018.02.052

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/19584/>

Link to published version: <https://doi.org/10.1016/j.jaci.2018.02.052>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Predictive markers for humoral influenza vaccine response in patients with common variable immunodeficiency (CVID)

Ann Gardulf, RN, PhD^{1,2}, Hassan Abolhassani, MD, PhD^{1,2,7}, Rolf Gustafson, MD, PhD³,
Lars E. Eriksson, RN, MSc, PhD^{4,5,6}, Lennart Hammarström, MD, PhD^{1,2}

¹*Division of Clinical Immunology and Transfusion Medicine, Department of Laboratory Medicine, Karolinska Institutet, Campus Flemingsberg, Huddinge, SE-141 86 Stockholm, Sweden*

²*Jeffrey Modell Diagnostic and Research Center for Primary Immunodeficiencies, Karolinska University Hospital, Campus Flemingsberg, Huddinge, SE-141 86 Stockholm, Sweden.*

³*The Immunodeficiency Unit, Department of Infectious Diseases, Karolinska University Hospital, Huddinge, SE-141 86 Stockholm, Sweden .*

⁴*Department of Learning, Informatics, Management and Ethics, Karolinska Institutet, SE-171 77 Stockholm, Sweden.*

⁵*Division of Nursing, School of Health Sciences, City, University of London, London EC1V 0HB, UK.*

⁶*Department of Infectious Diseases, Karolinska University Hospital, Huddinge, SE-141 86 Stockholm, Sweden.*

⁷*Research Center for Immunodeficiencies, Pediatrics Center of Excellence, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran*

Corresponding author: Ann Gardulf

Address: Department of Laboratory Medicine, Division of Clinical Immunology and Transfusion Medicine, Alfred Nobels allé 8, Karolinska Institutet, Huddinge, SE-141 86 Stockholm, Sweden

25 **E-mail:** ann.gardulf@ki.se.

26

27 **Running title:** Humoral response to A(H1N1) pandemic influenza vaccine in CVID

28

29 **Statement of financial sources:** The study was funded by the Jeffrey Modell Foundation,
30 New York; the Swedish Research Council, Stockholm, Sweden; and research funds from
31 Karolinska Institutet, Stockholm, Sweden.

32

33 **Conflict of interests:** The authors declare that there was no financial or personal relationship
34 with third parties whose interests could be positively or negatively influence the content.

Abstract

Background: A subgroup of patients with common variable immunodeficiencies (CVID) responds to vaccination. The aim of the study was to try to identify predictive markers for those who developed a humoral immune response after influenza vaccination.

Methods: 48 patients with CVID (29 females, 19 males, mean age 59.4 years) were vaccinated with the A(H1N1) influenza vaccine Pandemrix® and boosted after one month. Blood samples were collected prior to each vaccination and two months later. Patients with a 4-fold titer increase of the hemagglutinin inhibition test ($\geq 1:40$) were considered responders and compared to non-responders for clinical, immunological and genetic markers.

Results: Eight (16.7%) patients responded to the vaccination. A significantly higher proportion of the responders, who showed a Euroclass SmB⁺Tr^{norm}21^{norm} profile ($p=0.03$) with a post-germinal center B cell pattern ($p=0.04$) in blood, suffered from enteropathies ($p=0.04$) as compared to non-responders. Bronchiectasis on the other hand, was exclusively found among non-responders ($n=7$), as was autoimmune cytopenia ($n=5$). Non-responders with a Euroclass SmB⁺21^{low}Tr^{norm} profile ($p=0.02$), had a significantly higher prevalence of progressive antibody deficiency ($p=0.048$) and, at diagnosis, a higher mean serum IgM level ($p=0.03$), a lower mean serum IgG1 level ($p=0.007$), an expansion of absolute counts of cytotoxic CD8⁺ T-cells ($p=0.033$) and an increased proportion of memory CD8⁺ T-cells ($p=0.044$) in blood. CVID associated HLA markers were not detected in non-responders ($p=0.03$).

Conclusion: About one-fifth of the CVID patients achieved protective antibody levels after A(H1N1) vaccination and selected clinical and immunological markers were identified that may predict a positive outcome of influenza vaccination.

Key messages:

- A subgroup of patients with CVID develops a humoral immunological response from influenza vaccination.
- Selected clinical and immunological markers may help to identify a possible positive influenza vaccination outcome and are enteropathy, Euroclass SmB^{norm}Tr^{norm}21^{norm} and post-germinal center B-cell pattern.
- HLA CVID genetic predisposing markers, pronounced IL-12 production and Th1 polarity were observed only in non-responders.
- Despite the fact that not all patients with CVID will develop protective antibody levels after influenza vaccination it is concluded that they should be offered this prophylactic measure due to the potential severity of influenza and risk for bacterial complications.

Capsule Summary: One-fifth of the patients with CVID vaccinated against pandemic influenza A(H1N1) converted to a $\geq 1:40$ titer of specific antibodies against the antigen and selected clinical and immunological predictive markers were identified in this subgroup.

Keywords: Common variable immunodeficiency, CVID, specific antibody deficiency, vaccination, influenza, pandemic influenza, immune response, A(H1N1), Pandemrix

Abbreviations:

CVID: Common variable immunodeficiency

DNA: Deoxyribonucleic acid

ELISA: Enzyme-linked immunosorbent assay

HA: Hemagglutinin

HI: Haemagglutination inhibition

HLA: Human leukocyte antigen

- 83 **Igs:** Immunoglobulins
- 84 **IFN- γ :** Interferon gamma
- 85 **IL:** Interleukin
- 86 **PBS:** Phosphate-buffered saline
- 87 **PCR:** Polymerase chain reaction
- 88 **PID:** Primary immunodeficiency disorder
- 89 **PHA:** Phytohemagglutinin
- 90 **RDE:** Receptor-destroying enzymes
- 91 **WHO:** World Health Organization
- 92

93 Introduction

94 Common variable immunodeficiency (CVID) is a heterogeneous primary
95 immunodeficiency disorder (PID) characterized by low serum concentrations of
96 immunoglobulins (Igs) and impaired specific antibody production.¹⁻³ Specific antibody
97 deficiency is a diagnostic criterion in most of the standard clinical guidelines and constitutes
98 the absence of natural Igs (e.g. isohemagglutinins) or poor response to novel protein or
99 polysaccharide antigens (e.g. in vaccines).^{2, 4, 5}

100 Approximately 10-20% of clinically diagnosed patients with CVID have a residual
101 response to vaccination against protein antigens and, to a lesser extent, against polysaccharide
102 antigens.⁶⁻¹² The heterogeneity in specific antibody defects in CVID patients may be due to
103 the association of this complex disease with several different genetic defects.^{13, 14} Although it
104 is uncertain whether vaccination with killed/inactivated vaccines will protect an individual
105 with CVID, the use of these vaccines is nevertheless recommended for diagnostic purposes.^{10,}
106 ¹⁵ Moreover, the use of killed/inactivated vaccines to patients with PID is indeed
107 recommended to follow the same schemes as in general populations.¹⁶

108 Current evidence suggests that a subgroup of patients with CVID is not only capable of
109 re-stimulation *in vitro* for production of class-switched Igs but may show residual antibody
110 production *in vivo*.¹⁷⁻¹⁹ Characterization of this subgroup of patients regarding their clinical,
111 immunological and genetic profile could potentially help developing targeted therapy for
112 these patients^{18, 20} and provide evidence-based advises regarding administering influenza
113 vaccine to patients with CVID.

114 To our knowledge, the immunological response to the same antigen as used in the current
115 study - A(H1N1) pandemic influenza vaccine (antigen X179a 2009, Pandemrix[®]) - has so far
116 only been investigated in three patients with CVID and one patient with X-linked
117 agammaglobulinemia in a 3-months long follow-up study.¹⁹ Two of the patients with CVID
118 responded to the vaccination by a >4-fold rise in haemagglutination inhibition (HI) antibodies

119 and all three showed a Th1-cell response. It was concluded that some patients with CVID
120 might produce an influenza-specific humoral immune response, but that this should be
121 confirmed in larger studies.

122 The aim of the study was to try to identify clinical, immunological and genetic predictive
123 markers for those patients with CVID who developed a humoral immune response after
124 influenza vaccination. This was performed by investigating to what extent a group of patients
125 with CVID responded to vaccination by producing protective antibodies against the
126 glycopeptide pandemic influenza A(H1N1) antigen.

Material and methods

The present study had a prospective design, monitoring patients with CVID before and up to one year after vaccination with the pandemic influenza A(H1N1) vaccine (Pandemrix[®], GlaxoSmithKline, Belgium).

Patients

Patients with CVID cared for and receiving Ig replacement therapy at the Immunodeficiency Unit at the Karolinska University Hospital in Stockholm, Sweden, and who had not already been vaccinated against the A(H1N1) pandemic virus were invited to participate in the study carried out during 2010-2011. None of the patients used any steroids or any other immuno-suppressive therapy during the study period and 3 months prior to study start. All patients had been diagnosed as having CVID based on the diagnostic criteria established by the European Society for Immunodeficiencies (<http://esid.org/WorkingParties/Registry/Diagnosis-criteria>) and the American Academy of Allergy, Asthma and Immunology practice parameter for the diagnosis and management of PID.^{4, 21} Before study start, all patients were re-evaluated for fulfilling the diagnostic criteria of CVID and secondary causes of dysgammaglobulinemia were ruled out.²² None of the invited patients presented with protein losing enteropathy, nor with pre-study protective A(H1N1) pandemic virus antibody levels ($\geq 1:40$ titer).

Fifty-seven patients gave their informed consent and volunteered to participate in the study by receiving the first dose of the vaccine. Nine patients (five women and four men, mean age 48.8 ± 20.3) dropped out of the study; seven at the first follow-up visit and two patients at the second follow-up visit. Totally 48 patients (60.4% females) participated in all appointments and were included in the final vaccine response analyses. Forty-seven of the patients received regular weekly subcutaneous (minimum 100 mg/kg/week) and one patient received monthly intravenous (400 mg/kg/month) Ig replacement therapy before and during

the entire study period. Different brands were used but each patient received the same brand and batch of Ig at least three months before and at least until the last blood sampling.

An evaluation document was used to summarize the demographic information of each patient including age at diagnosis, family history and history of previous vaccinations shown as specific antibody responses, and clinical, immunological and genetic data.

Study overview

The A(H1N1) pandemic influenza vaccine Pandemrix[®], a monovalent X179a 2009 vaccine, was injected intramuscularly at study start and boosted once after one month. In each vaccination dose, 3.75 µg of hemagglutinin (HA) of the X179a antigen was administered. At study start, concentrations of total IgG, IgG subclasses, IgA and IgM were determined in fresh serum samples taken immediately prior to the next Ig substitution. Specific antibody titers for antibodies against the pandemic influenza A(H1N1) antigen were measured in samples collected prior to the first and the second vaccination and also two months after the second dose. Follow-up visits due to the vaccine study were scheduled for the patients at regular intervals until one year after study start for the recording of any influenza-like symptoms and/or incidental side effects of the vaccination.

Investigation and definition of vaccination response

One volume of serum was treated with four volumes of neuraminidase receptor destroying enzyme (RDE, Seiken, Japan) in phosphate-buffered saline (PBS) at 37°C overnight before inactivation at 56°C for 30 minutes to prevent non-specific inhibition. Serial two-fold dilutions of RDE treated sera were then incubated with eight haemagglutinating units of the X179a pandemic influenza A(H1N1) antigens for one hour, followed by the addition of 0.7% ([volume of solute/volume of solution] ×100%) turkey erythrocytes. After 30 minutes of incubation, the HA inhibition titers were read as the reciprocal of the highest

dilution at which 50% HA was inhibited. Each serum sample was run in duplicate in two independent experiments and the titers are presented as a mean of these two experiments for each patient.

A positive response to the pandemic influenza A(H1N1) antigen was defined as a 4-fold titer increase of the HA inhibition test ($\geq 1:40$).²³ Patients with such increase are hereinafter referred to as responders.

Clinical and immunological phenotyping and immunological assays

The clinical phenotyping of the 48 patients was based on the suggested division into distinct clinical phenotypes by Chapel *et al.*²⁴ and included patients with infections but no other disease-related complications or patients with infections and autoimmunity, infections and enteropathy and/or infections and polyclonal lymphocytic infiltration. Patients with ≥ 3 clinical findings were classified as overlapping phenotypes. Clinical data were collected from the medical and nursing records and covered the time from diagnosis to the study start.

Thirty-eight of the patients had previously been classified based on two main classifications for B-cell subsets including the Paris, Freiburg, EUROclass²⁵ and the B-cell pattern classification.²⁶ In 27 of the patients, ionizing radiation sensitivity assays had previously been performed on primary fibroblasts irradiated with different doses and serial dilutions.²⁷ This is a method used for classification of patients with CVID and defective DNA repair machinery required for variable (V), diversity (D) and joining (J) genes rearrangement and class-switch recombination.²⁸⁻³¹

Peripheral blood mononuclear cells from all 48 patients were stimulated for 48 hours with phytohemagglutinin (PHA) and cytokine production of interferon gamma (IFN- γ), interleukin 2 (IL-2), IL-5, IL-10 and IL-12 were measured by enzyme-linked immunosorbent assay (ELISA) using a method described previously.³²

205 ***Screening of positional candidate genetic markers***

206 HLA-A, -B, -DQ and -DR alleles were determined using low-resolution DNA-based
 207 typing (polymerase chain reaction [PCR]/sequence specific oligonucleotide probe).³³ The
 208 PCR amplification of tumour necrosis factor receptor superfamily member 13 B and C
 209 (*TNFRSF13B* and *TNFRSF13C*) and their Sanger sequencing were performed using primers
 210 and conditions as previously described.³⁴ Of note, selected monogenic forms of PID were
 211 excluded in studied patients according to the CVID diagnostic criteria using a targeted gene
 212 panel sequencing (covering 260 known monogenic diseases) including *BTK*, *BLNK*, *CD79A*,
 213 *CD79B*, *IGHM*, *IGLL1*, *TCF3*, *CD19*, *CD20*, *CD21*, *CD81*, *AICDA*, *UNG*, *INO80*, *MSH6*,
 214 *CARD11*, *NFKB1*, *NFKB2*, *PI3KCD*, *PIK3R1*, *PTEN*, *DOCK2*, *IKAROS*, *IRF2BP2*, *MOGS*,
 215 *TWEAK*, *IL21*, *IL21R*, *LRBA* and *CTLA4* genes. Moreover, whole exome sequencing in 17
 216 patients were also performed, but neither non-responders nor responders had a confirmed
 217 candidate monogenic disease.

219 ***Ethical considerations***

220 The World Health Organization declared in June 2009 that the outbreak of A(H1N1)
 221 influenza fulfilled the criteria of a pandemic situation. As a consequence, the National Board
 222 of Health and Welfare in Sweden recommended the general population to be vaccinated and
 223 certain risk groups, including individuals with an impaired immune system, to be prioritized.
 224 The pandemic influenza A(H1N1) vaccine had not previously been used in a mass vaccination
 225 situation, but the potential severity of the disease motivated patients with CVID to be offered
 226 the new vaccine.

227 Based on the Ethical Review of Research Involving Humans³⁵ the vaccination including
 228 blood sampling and follow-up schedule were approved by the regional Ethical Committee
 229 (approval number 2009/1646-31-3). The patients were given oral and written information

about the study. The principle of volunteering was emphasized and informed written consent to participate including a one-year follow-up was obtained from all patients.

Statistical analyses

Statistical analyses were performed using SPSS (version 21.0.0, Statistics software, SPSS, Chicago, Illinois) and R statistical systems (version 3.4.1., R Foundation for Statistical Computing, Vienna, Austria). The one-sample Kolmogorov-Smirnov test was applied to estimate whether the data distribution was normal and based on the findings of this evaluation independent T-test (in normal distributions) and Mann-Whitney U test (in skewed distributions) were used to compare continuous variables between responders and non-responders. Differences in categorical variables between responders and non-responders were examined using χ^2 tests and Fisher's exact tests (the latter when variables had a low frequency). Pearson's correlation coefficient analysis was used to investigate the relationship between specific antibody responses to different antigens. A *p*-value of 0.05 or less was considered statistically significant.

Results

Three months after the first vaccine dose and two months after the second dose, eight (16.7%) patients had reached a $\geq 1:40$ titer of specific antibodies against the pandemic influenza A(H1N1) antigen (responders) whereas the remaining 40 patients (83.3%) were considered as non-responders. The production of specific antibodies against the antigen is depicted in **Figure 1**. Four out of eight responders reached protective levels already after the first vaccine dose. Comparing the four early responders with the four late responders, no significant differences could be found between their current age (57.0 ± 14.7 vs. 54.2 ± 11.7 years) or age at CVID diagnosis (41.5 ± 10.0 vs. 39 ± 16.8 years).

During the study period, none of the patients showed any clinical symptoms of having been infected with the ongoing pandemic influenza A(H1N1) and during the one-year follow-up period, no influenza-like symptoms and/or negative side-effects other than local reactions on the injection sites were reported.

Table 1 summarizes the demographic and essential immunological data of responders and non-responders. Response to the A(H1N1) pandemic vaccine was independent of the gender of the patients. There were also no significant differences in the mean age at diagnosis, mean age at study start, mean years from onset of infections to diagnosis or the mean follow-up time after the CVID diagnosis between responders and non-responders, respectively. Nine patients had a familiar form of CVID, all belonging to the non-responder group. In the total group, 14 individuals (29.1%) had evolved from an IgA deficiency (n=12) and/or IgG subclass deficiency (n=2) to a CVID diagnosis during the course of their disease. All these 14 patients were non-responders constituting 35.0% of this group ($p=0.048$, compared to absence of this progressive PID form in responders) (**Table 1**).

Regarding the Ig replacement therapy there was no difference in the distribution of different Ig brands used between responders and non-responders (data not shown).

Clinical phenotyping

Before diagnosis, all 48 patients had experienced recurrent upper respiratory tract infections and 44 had also been diagnosed as recurrently suffer from lower respiratory tract infections (bronchitis and/or pneumonia). Clinical phenotyping of the 48 patients revealed that 14 (29.2%) of them presented with infections without any other disease-related complications, 14 (29.2%) by infections and autoimmunity, four (8.3%) by infections and enteropathy, and three (6.2%) by infections and polyclonal lymphocytic infiltration. Thirteen patients (27.1%) also manifested overlapping phenotypes (**Figure 2, Table S2**).

When dividing the clinical phenotyping between responders and non-responders the proportion of patients with infections without any other disease-related complications was 12.5% among the responders and 32.5% among the non-responders ($p=0.12$) (**Figure 2, Table S2**). Lower respiratory tract infections were documented in five responders (62.5%) and in 33 non-responders (82.5%, $p=0.20$). None of the responders showed signs of bronchiectasis, while seven non-responders (17.5%; $p=0.10$) suffered from this condition.

The phenotype infections and enteropathy was significantly higher in responders (50% vs. 0%, $p<0.001$). Although infections and autoimmunity was present in about the same proportion in both groups (25% in responders and 30% in non-responders, n.s.) (**Figure 2, Table S2**), autoimmune cytopenia was exclusively observed among the non-responders ($n=5$).

Malignancies were recorded in 12 patients (30%) in non-responders, mainly due to thymoma and lymphoma and in two patients (25%) among the responders (one breast cancer and one colon cancer) ($p=0.3882$) (**Table S2**).

No significant difference was observed regarding the proportion of patients with IgE mediated atopic disorders: four responders (50%) and nine non-responders (22.5%, $p=0.12$) (**Table S2**).

Immunological phenotyping and classification

At the time of the COVID diagnosis, a significantly higher mean serum level of IgM (34.8 ± 29.6 vs. 17.0 ± 16.0 mg/dl; $p=0.03$) and a significantly lower mean serum level of IgG1 (122.4 ± 96.3 vs. 219.2 ± 198.3 mg/dl; $p=0.007$) were noted in non-responders as compared to responders. Lymphocyte subset analyses showed an expansion of absolute counts of cytotoxic T-cells ($p=0.033$) as well as an increased proportion of memory CD8⁺ T-cells ($p=0.044$) in the non-responders as compared to the responders. Furthermore, there was a tendency of a higher number of NK-cells among the non-responders as compared to the responders ($p=0.06$). There was no significant difference between responders and non-responders regarding absolute B-cell counts, although the non-responders presented a decreased number of plasmablast and an increased CD21^{low} percentage ($p=0.007$ and $p=0.041$, respectively) (**Table 2**). Radiosensitivity was only documented in the group of non-responders (4/20 tested non-responders, 20%, vs. 0/7 tested responders, $p=0.09$).

Although a positive correlation was observed in responders with increment in the level of protective antibodies after the vaccination with the pandemic influenza A(H1N1) antigen regarding the production of other specific antibodies against protein and polysaccharide antigens ($r=0.75$, $p=0.08$), no statistically significant differences were found between responders and non-responders. Details of the humoral immune response to other antigens are presented in **Table S1**.

Immunological classification of the patients revealed that the most frequent immune profile in the responders was Euroclass SmB⁻Tr^{norm}21^{norm} ($p=0.03$) and post-germinal center B-cell pattern (normal naïve, transitional, marginal and memory B-cell subsets) ($p=0.04$) as compared to the non-responders, whereas the most frequent immune profile of non-responders was Euroclass SmB⁻21^{low}Tr^{norm} ($p=0.02$ compared to the responders) (**Figure 3**).

The PHA induced cytokine production in vitro in responders and non-responders are presented in **Figure 4**. A significant difference was found in mean IL-12 levels; in non-responders $1,081.2 \pm 651.3$ pg/ml and for responders 283.3 ± 256.5 pg/ml ($p=0.007$).

Genetic markers associated with CVID and pandemic influenza A(H1N1) antibody production

Four patients with *TNFRSF13B* (10%) and one patient with *TNFRSF13C* (2.5%) CVID susceptibility variants were found among the non-responders while only WT *TNFRSF13* genes were identified among the responders (12.5% vs 0%, $p=0.38$). HLA markers associated with CVID were detected in 16/40 (40%) of the non-responders; HLA-DR3-DQ2 in six patients, HLA-A1-B8 in six patients and HLA-A2-B44 in four patients. HLA markers associated with CVID were not found in any of the responders (40% vs 0%, $p=0.03$).

Discussion

The outbreak of the pandemic influenza A(H1N1) and the recommendation to specifically vaccinate individuals with an impaired immune system did not only stress the immediate medical need to offer this new vaccine with a A(H1N1) clade selected by WHO to patients with CVID, but it also opened up a possibility for a scientific evaluation of the vaccination in this group of individuals. The background to the study was the perpetual discussion whether patients with CVID should routinely be offered vaccination against seasonal influenza. Only our study and the study by Pedersen *et al.*¹⁹ have presented data from the use of the specific antigen X179a to individuals with CVID. Our study was designed as an evaluation within the group of patients and therefore no healthy controls were included. However, it has been shown that between 67-98.3% of healthy adults produce protective levels of antibodies against the influenza A(H1N1) vaccine Pandemrix® 21days after a single dose of 3.75 µg of the vaccine.^{36, 37}

Cross-reactive A(H1N1) antibodies are present in healthy populations³⁸ and consequently, antibodies against the pandemic influenza A(H1N1) antigen may be present in Ig preparations.³⁹ All participants in the study were therefore tested for pre-existing A(H1N1) antibodies before entering the study. One 73-year old male had a HA inhibition titer of 1:10 at the study start but this patient did not respond to the vaccination. All patients continued with the same brand and batch of Ig during the study period.

The WHO declaration in 2009 of a pandemic situation in 74 countries and territories rapidly led to a demand for the vaccine that was greater than the supply. On a European level it was considered important to offer the vaccine rapidly to as many individuals as possible. For this reason it was decided by the authorities to start the vaccination campaign with an antigen dose of 3.75 µg to the entire population but immunodeficient patients were to be offered a second dose after one month. It has later been shown that a single dose of 3.75 µg of Pandemrix® with the antigen X179a brings about immunity to protective level in healthy

adults and elderly.⁴⁰ In the study by Pedersen *et al.* about the immunological response to the antigen X179a in CVID¹⁹, the intention was to give two doses of 3.75 µg to three patients with CVID but the patients were accidentally given a double dose of Pandemrix® at study start. Two of the patients were then given a second dose of 3.75 µg after three weeks whilst the third patient declined being given the second dose. The two patients receiving two doses (7.5 µg + 3.75 µg) responded with a >4-fold humoral response while the patient receiving one dose (7.5 µg) had a HI titer below $\geq 1:40$. In our study, four of the responders developed protective levels of antibodies against the X179a antigen already after one dose of 3.75 µg, while the four others needed two doses to produce protective levels of antibodies (**Figure 1**). The four early responders were not older at diagnosis or study start than the four late responders, which could have explained the result as older patients might have been in contact with cross-reactive antigens earlier. Based on the current study and the study by Pedersen *et al.*¹⁹, it seems reasonable to assume that two doses of 3.75 µg of the antigen X179a would be required to obtain humoral immunity in patients with CVID. Whether these patients should be offered one or two doses of inactivated seasonal influenza vaccine has never been addressed in the four groups of patients with CVID where the outcome of this type of vaccine has been investigated^{9, 17, 41, 42} and any answer to this question is therefore not available. However, Eibl & Wolf¹⁶ suggest that the primary immunization of inactivated influenza vaccine should follow the same scheme as for healthy individuals, but that more frequent booster immunizations might be necessary depending on the assessment of vaccination response in PID.

In the influenza, polysaccharide or protein vaccine studies that have been conducted in patients with CVID^{7, 9, 19, 41-44} they have shown more frequent and earlier decline in antibody responses against polysaccharides compared to proteins, suggesting a preservation of T cell-dependent specific antibody response in a subgroup of patients.^{7, 19, 43} In a previous study, Zhan *et al.*⁴¹ reported generation of specific IgG-secreting memory B cells post seasonal

influenza vaccines in 50% of the studied CVID patients. In our study, we found that about one-fifth (16.7%) of the patients responded by producing protective levels of antibodies against the pandemic influenza A(H1N1) antigen. Consistent with other reports, we also observed a functional classification of CVID patients, according to the level of the response to the vaccine.^{43, 45}

There are to date no studies regarding the clinical phenotypes of patients who seroconvert or not after influenza vaccination.⁴⁶ The finding among the responders of a significantly higher proportion of enteropathies, a phenotype that usually presents due to increased apoptotic bodies in the gastrointestinal crypts⁴⁷, may suggest that the immunological phenotype of the responders with normal B cell subsets could be a result of an increased apoptosis of the long-term plasmacells. In line with this notion these patients can produce normal antibodies against specific antigens during a short-term, but these antibody-producing plasmacells subsequently disappear. The increased susceptibility to apoptosis in the gastrointestinal crypts could be a trigger of chronic inflammation. Regarding the severe lower respiratory tract infections and the bronchiectasis among the non-responders it appears that the non-responders are less likely to mount an immune response against bacteria and consequently an effective elimination in the lower respiratory tract, resulting in persistent tissue damage in the lungs.

The capability to respond to certain vaccines *in vivo*, strongly suggest that some patients with CVID can produce class-switched isotypes, as has been shown by cytokine stimulation *in vitro*.⁴⁸ Although CD27⁺ IgD⁻ isotype-switched memory B cells are generally reduced in CVID patients⁴⁹, our findings confirm that responders had a higher level of plasmablasts and lower counts of CD21^{low} B cells (compatible with a Euroclass SmB⁻Tr^{norm}21^{norm} and post-germinal center B-cell pattern). In contrast, we found that the non-responders at diagnosis presented with lower mean serum level of IgG1 and a higher mean serum level of IgM, highlighting a severe defective class switching recombination in this group of patients.

Moreover, radiosensitivity was only found in non-responders but this test was performed only in 27 patients and the difference was not statically significant. These findings emphasize the role of the class-switching capability for developing a vaccine response. In two studies by Goldacker *et al.*⁷ and Chovancova *et al.*¹³, respectively, they presented B cell subset analysis; both reported that group II of the Freiburg classification or the EUROclass group smB+, which represent patients with CVID with nearly normal numbers of class-switched memory B cells, constitutes patient with measurable antibody responses.

Stimulating peripheral blood mononuclear cells with PHA enhanced the production of IL-12 at significantly higher mean level in non-responders as compared to responders, suggesting a significant immune regulation in this group toward activation of cellular immunity. Up-regulation of the IL-12 as an initial factor in a subgroup of patients with CVID skews the immune response away from Ig production towards a polarized Th1-type chronic inflammation.^{50, 51} Increased expression of the IL-12 β 1 receptor has also been reported in these patients.⁵² Although the IFN- γ level, another marker of Th1-response, did not differ between the two groups, Ig replacement may alter the serum pattern of this cytokine but not the IL-12 level.⁵³ Failure of an immune deviation from a systemic Th1 response to a Th2 immune response may explain the mechanism responsible for the numerically lower proportion of IgE mediated atopic diseases observed in the non-responders as compared to responders.⁵⁴

Humoral response against influenza vaccine has been shown to be the major source of protection against infections and individuals with annual injections have a broader antibody recognition profile after pathogenic confrontation.⁵⁵ However, influenza-specific CD8+ T cells can be generated after vaccination targeted to conserved viral proteins (nucleoprotein and the matrix protein) to provide heterosubtypic immunity. Since long-term annual vaccination in the presence of normal humoral immunity may interfere the induction of heterosubtypic immunity, it can be hypothesized that virus-specific CD8+ T cell responses are

more pronounced in patients with antibody deficiency.⁵⁶ As the current study only focused on humoral immunity, further evaluation of the cellular immunity in these patients may be of considerable interest.

The presence of the familiar form of CVID only in non-responders, taken together with the finding of predisposing HLA and non-HLA factors, suggest a strong association of CVID pathogenetic in this group of patients.⁵⁷ Moreover, the progressive primary antibody deficiency was exclusively found in the non-responders, a phenomenon that has previously been linked to genetic susceptibility markers in patients with CVID, in particular HLA A1, B8, DR3 and DQ2.^{58, 59} Investigation of non-genetic etiologies among responders should be prioritized, including epigenome and microbiome assays. The essential role of specific HLA haplotype in response to vaccine has been investigated also in healthy individuals; particularly CVID HLA susceptibility markers such as DQ2-DR3 phenotypes have been linked with non-responsiveness to hepatitis B antigen in vaccine in the normal population.^{60, 61} Specific antibody response to vaccines has also been investigated in patients with *TNFRSF13B* mutated alleles and transmembrane activator and CAML interactor (*TACI*) knockout mice due to the role of *TACI* in signaling for induction of Ig class switching. Most of the animal studies have showed consistent defective responses to vaccination with T independent antigens.⁶² However, humans with heterozygous *TNFRSF13B* mutations show a wide range of ability to respond to vaccine from absent to some antibody production^{63, 64} even within the same family with same mutation⁶⁵, suggesting a presence of other genetic modifier rather than *TACI*.

To summarize, we identified protective increment in the level of antibodies against the pandemic influenza A(H1N1) antigen in eight (16.7%) patients of our study population including totally 48 individuals. This is to date the largest study performed investigating influenza vaccine response in patients with CVID. We found that a positive outcome of the influenza vaccination might be expected in patients with certain identified specific B cell patterns. The responders were sporadic cases without genetic susceptibility markers, but with

normal class switching recombination and DNA repair machinery. Therefore they could still produce residual specific antibodies against the current antigen, and they presented with low rate of severe lower respiratory tract infections and no infectious complications such as bronchiectasis. Instead, these patients more often presented with enteropathy.

Predictive markers for patients with CVID who will respond to influenza vaccine were found to be enteropathy, Euroclass SmB^{norm}Tr^{norm}21^{norm} and post-germinal center B-cell pattern. Despite the fact that not all patients with CVID developed protective antibody levels after two doses of the vaccine, it is concluded that patients with CVID should be offered vaccination also against seasonal influenza^{9, 16, 17, 42, 46} due to the potential severity of the infection and risk for bacterial complications. Adverse events are not a major issue and inactivated influenza vaccine can safely be given to patients with CVID.

474 **Acknowledgments**

475 We would like to express our sincere gratitude to the participating patients and staff at the
476 Immunodeficiency Unit, Department of Infectious Diseases, Karolinska University Hospital,
477 Huddinge, Stockholm, Sweden.

478 The study was funded by the Jeffrey Modell Foundation, New York; the Swedish
479 Research Council, Stockholm, Sweden; and research funds from Karolinska Institutet,
480 Stockholm, Sweden.

481

482 **Author contributions**

483 (1) The conception and design of the study

484 (2) Acquisition of data,

485 (3) Analysis and interpretation of data,

486 (4) Drafting the article

487 (5) Revising it critically for important intellectual content,

488 (6) Final approval of the version to be submitted

489

490 AG(2,3,4,5,6), HA (3,4,5,6), RG(2,4,5,6), LE(3,5,6) and LH, (1,3,5,6).

References

1. Jolles S. The variable in common variable immunodeficiency: a disease of complex phenotypes. *The Journal of Allergy and Clinical Immunology: In Practice* 2013; 1:545-56.
2. Yong PF, Thaventhiran JE, Grimbacher B. "A Rose is a Rose is a Rose," but CVID is Not CVID: Common Variable Immune Deficiency (CVID), What do we Know in 2011? *Advances in immunology* 2011; 111:47.
3. Abolhassani H, Parvaneh N, Rezaei N, Hammarstrom L, Aghamohammadi A. Genetic defects in B-cell development and their clinical consequences. *J Investig Allergol Clin Immunol* 2014; 24:6-22; quiz 2 p following
4. Bonilla FA, Barlan I, Chapel H, Costa-Carvalho BT, Cunningham-Rundles C, de la Morena MT, et al. International Consensus Document (ICON): Common Variable Immunodeficiency Disorders. *J Allergy Clin Immunol Pract* 2016; 4:38-59.
5. Fischer MB, Wolfram W, Binder CJ, Bohmig GA, Wahrmann M, Eibl MM, et al. Surface Plasmon Resonance Analysis Shows an IgG-Isotype-Specific Defect in ABO Blood Group Antibody Formation in Patients with Common Variable Immunodeficiency. *Front Immunol* 2015; 6:211.
6. Yarchoan R, Schneider HS, Wray BB, Nelson DL. Specific anti-influenza virus antibody production in vitro by lymphocytes from a subset of patients with hypogammaglobulinemia. *J Clin Invest* 1983; 71:1720-7.
7. Goldacker S, Draeger R, Warnatz K, Huzly D, Salzer U, Thiel J, et al. Active vaccination in patients with common variable immunodeficiency (CVID). *Clin Immunol* 2007; 124:294-303.
8. Sanchez-Ramon S, de Gracia J, Garcia-Alonso A, Rodriguez Molina JJ, Melero J, de Andres A, et al. Multicenter study for the evaluation of the antibody response against salmonella typhi Vi vaccination (EMPATHY) for the diagnosis of Anti-

- 517 polysaccharide antibody production deficiency in patients with primary
 518 immunodeficiency. Clin Immunol 2016; 169:80-4.
- 519 9. Hanitsch LG, Lobel M, Mieves JF, Bauer S, Babel N, Schweiger B, et al. Cellular and
 520 humoral influenza-specific immune response upon vaccination in patients with
 521 common variable immunodeficiency and unclassified antibody deficiency. Vaccine
 522 2016; 34:2417-23.
- 523 10. Hong R, Agrawal S, Gollapudi S, Gupta S. Impaired pneumovax-23-induced
 524 monocyte-derived cytokine production in patients with common variable
 525 immunodeficiency. J Clin Immunol 2010; 30:435-41.
- 526 11. Rezaei N, Siadat SD, Aghamohammadi A, Moin M, Pourpak Z, Norouzian D, et al.
 527 Serum bactericidal antibody response 1 year after meningococcal polysaccharide
 528 vaccination of patients with common variable immunodeficiency. Clin Vaccine
 529 Immunol 2010; 17:524-8.
- 530 12. Janssen WJ, Nierkens S, Sanders EA, Boes M, van Montfrans JM. Antigen-specific
 531 IgA titres after 23-valent pneumococcal vaccine indicate transient antibody deficiency
 532 disease in children. Vaccine 2015; 33:6320-6.
- 533 13. Chovancova Z, Vlkova M, Litzman J, Lokaj J, Thon V. Antibody forming cells and
 534 plasmablasts in peripheral blood in CVID patients after vaccination. Vaccine 2011;
 535 29:4142-50.
- 536 14. Carsetti R, Rosado MM, Donnanno S, Guazzi V, Soresina A, Meini A, et al. The loss
 537 of IgM memory B cells correlates with clinical disease in common variable
 538 immunodeficiency. J Allergy Clin Immunol 2005; 115:412-7.
- 539 15. Rezaei N, Aghamohammadi A, Read RC. Response to polysaccharide vaccination
 540 amongst pediatric patients with common variable immunodeficiency correlates with
 541 clinical disease. Iran J Allergy Asthma Immunol 2008; 7:231-4.

- 542 16. Eibl MM, Wolf HM. Vaccination in patients with primary immune deficiency,
543 secondary immune deficiency and autoimmunity with immune regulatory
544 abnormalities. *Immunotherapy* 2015; 7:1273-92.
- 545 17. van Assen S, de Haan A, Holvast A, Horst G, Gorter L, Westra J, et al. Cell-mediated
546 immune responses to inactivated trivalent influenza-vaccination are decreased in
547 patients with common variable immunodeficiency. *Clin Immunol* 2011; 141:161-8.
- 548 18. Ballow M. Vaccines in the assessment of patients for immune deficiency. *J Allergy*
549 *Clin Immunol* 2012; 130:283-4 e5.
- 550 19. Pedersen G, Halstensen A, Sjursen H, Naess A, Kristoffersen EK, Cox RJ. Pandemic
551 influenza vaccination elicits influenza-specific CD4+ Th1-cell responses in
552 hypogammaglobulinaemic patients: four case reports. *Scand J Immunol* 2011; 74:210-
553 8.
- 554 20. Wolf HM, Thon V, Litzman J, Eibl MM. Detection of impaired IgG antibody
555 formation facilitates the decision on early immunoglobulin replacement in
556 hypogammaglobulinemic patients. *Front Immunol* 2015; 6:32.
- 557 21. Conley ME, Notarangelo LD, Etzioni A. Diagnostic criteria for primary
558 immunodeficiencies. *Clinical immunology* 1999; 93:190-7.
- 559 22. Bogaert DJ, Dullaers M, Lambrecht BN, Vermaelen KY, De Baere E, Haerynck F.
560 Genes associated with common variable immunodeficiency: one diagnosis to rule
561 them all? *J Med Genet* 2016; 53:575-90.
- 562 23. Products, EAftEoM, Note for guidance on harmonization of requirements for
563 influenza vaccines (CPMP/BWP/214/96), 1997. Available on January 11, 2018,
564 at http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2009/09/WC500003945.pdf.
565
- 566 24. Chapel H, Lucas M, Patel S, Lee M, Cunningham-Rundles C, Resnick E, et al.
567 Confirmation and improvement of criteria for clinical phenotyping in common

- variable immunodeficiency disorders in replicate cohorts. *J Allergy Clin Immunol* 2012; 130:1197-8 e9.
25. Wehr C, Kivioja T, Schmitt C, Ferry B, Witte T, Eren E, et al. The EUROclass trial: defining subgroups in common variable immunodeficiency. *Blood* 2008; 111:77-85.
26. Driessen GJ, van Zelm MC, van Hagen PM, Hartwig NG, Trip M, Warris A, et al. B-cell replication history and somatic hypermutation status identify distinct pathophysiologic backgrounds in common variable immunodeficiency. *Blood* 2011; 118:6814-23.
27. Peron S, Pan-Hammarstrom Q, Imai K, Du L, Taubenheim N, Sanal O, et al. A primary immunodeficiency characterized by defective immunoglobulin class switch recombination and impaired DNA repair. *J Exp Med* 2007; 204:1207-16.
28. Vorechovsky I, Scott D, Haeney MR, Webster DA. Chromosomal radiosensitivity in common variable immune deficiency. *Mutat Res* 1993; 290:255-64.
29. Palanduz S, Palanduz A, Yalcin I, Somer A, Ones U, Ustek D, et al. In vitro chromosomal radiosensitivity in common variable immune deficiency. *Clin Immunol Immunopathol* 1998; 86:180-2.
30. Aghamohammadi A, Moin M, Kouhi A, Mohagheghi MA, Shirazi A, Rezaei N, et al. Chromosomal radiosensitivity in patients with common variable immunodeficiency. *Immunobiology* 2008; 213:447-54.
31. Offer SM, Pan-Hammarstrom Q, Hammarstrom L, Harris RS. Unique DNA repair gene variations and potential associations with the primary antibody deficiency syndromes IgAD and CVID. *PLoS One* 2010; 5:e12260.
32. Agarwal S, Smereka P, Harpaz N, Cunningham-Rundles C, Mayer L. Characterization of immunologic defects in patients with common variable immunodeficiency (CVID) with intestinal disease. *Inflamm Bowel Dis* 2011; 17:251-9.

- 593 33. International MHC, Autoimmunity Genetics N, Rioux JD, Goyette P, Vyse TJ,
594 Hammarstrom L, et al. Mapping of multiple susceptibility variants within the MHC
595 region for 7 immune-mediated diseases. Proc Natl Acad Sci U S A 2009; 106:18680-
596 5.
- 597 34. Salzer U, Neumann C, Thiel J, Woellner C, Pan-Hammarstrom Q, Lougaris V, et al.
598 Screening of functional and positional candidate genes in families with common
599 variable immunodeficiency. BMC Immunol 2008; 9:3.
- 600 35. Lag (2003:460) om etikprövning av forskning som avser människor. Svensk
601 författningssamling SFS 2003:460, Utbildningsdepartementet, Stockholm, Sverige. In
602 Swedish. (The Swedish Act (2003:460) on the Ethical Review of Research Involving
603 Humans. The Swedish Code of Statutes SFS 2003:460, The Ministry for Higher
604 Education and Research. Retrieved August 15, 2017, from
605 [https://www.riksdagen.se/sv/dokument-lagar/dokument/svensk-](https://www.riksdagen.se/sv/dokument-lagar/dokument/svensk-forfattningssamling/lag-2003460-om-etikprovning-av-forskning-som_sfs-2003-460)
606 [forfattningssamling/lag-2003460-om-etikprovning-av-forskning-som_sfs-2003-460](https://www.riksdagen.se/sv/dokument-lagar/dokument/svensk-forfattningssamling/lag-2003460-om-etikprovning-av-forskning-som_sfs-2003-460).
- 607 36. Manuel O, Pascual M, Hoschler K, Giulieri S, Alves D, Ellefsen K, et al. Humoral
608 response to the influenza A H1N1/09 monovalent AS03-adjuvanted vaccine in
609 immunocompromised patients. Clin Infect Dis 2011; 52:248-56.
- 610 37. European Medicines Agency. Pandemrix. Annex I. Summary of product
611 characteristics. European Medicines Agency, London, United Kingdom. Available on
612 January 11, 2018,
613 at [http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-](http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/000832/WC500038121.pdf)
614 [_Product_Information/human/000832/WC500038121.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/000832/WC500038121.pdf).
- 615 38. Ikonen N, Strengell M, Kinnunen L, Osterlund P, Pirhonen J, Broman M, et al. High
616 frequency of cross-reacting antibodies against 2009 pandemic influenza A(H1N1)
617 virus among the elderly in Finland. Euro Surveill 2010; 15.

- 618 39. Hong DK, Tremoulet AH, Burns JC, Lewis DB. Cross-reactive neutralizing antibody
619 against pandemic 2009 H1N1 influenza a virus in intravenous immunoglobulin
620 preparations. *Pediatr Infect Dis J* 2011; 30:67-9.
- 621 40. European Medicines Agency. Pandemrix. EMA/388182/2016. European Medicines
622 Agency, London, United Kingdom, 2016. Available on January 11, 2018,
623 at [http://www.ema.europa.eu/docs/en_GB/document_library/EPAR -](http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Summary_for_the_public/human/000832/WC500038122.pdf)
624 [_Summary_for_the_public/human/000832/WC500038122.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Summary_for_the_public/human/000832/WC500038122.pdf).
- 625 41. Zhan W, Yue F, Liu J, Lee J, Ho J, Moir S, et al. Response to influenza immunization
626 in patients with common variable immunodeficiency. *Allergy Asthma Clin Immunol*
627 2011; 7:A28.
- 628 42. van Assen S, Holvast A, Telgt DS, Benne CA, de Haan A, Westra J, et al. Patients
629 with humoral primary immunodeficiency do not develop protective anti-influenza
630 antibody titers after vaccination with trivalent subunit influenza vaccine. *Clin*
631 *Immunol* 2010; 136:228-35.
- 632 43. Al-Herz W, McGeady SJ. Antibody response in common variable immunodeficiency.
633 *Ann Allergy Asthma Immunol* 2003; 90:244-7.
- 634 44. Baldassin MPM, Grumach A, De Oliveira AKB, Kalil J. B cell subtypes' kinetics over
635 a 6 monthxs period in CVID patients submitted to influenza and H1N1 immunization.
636 *World Allergy Organization Journal* 2015; 8:A226.
- 637 45. Ko J, Radigan L, Cunningham-Rundles C. Immune competence and switched memory
638 B cells in common variable immunodeficiency. *Clin Immunol* 2005; 116:37-41.
- 639 46. Mieves JF, Wittke K, Freitag H, Volk HD, Scheibenbogen C, Hanitsch LG. Influenza
640 Vaccination in Patients with Common Variable Immunodeficiency (CVID). *Curr*
641 *Allergy Asthma Rep* 2017; 17:78.

- 642 47. Choi SW, Seo KW, Oh JS, Lee JS, Cha HJ. Successful treatment of enteropathy
643 associated with common variable immunodeficiency. *Korean J Intern Med* 2014;
644 29:256-8.
- 645 48. Borte S, Pan-Hammarstrom Q, Liu C, Sack U, Borte M, Wagner U, et al. Interleukin-
646 21 restores immunoglobulin production ex vivo in patients with common variable
647 immunodeficiency and selective IgA deficiency. *Blood* 2009; 114:4089-98.
- 648 49. Sanchez-Ramon S, Radigan L, Yu JE, Bard S, Cunningham-Rundles C. Memory B
649 cells in common variable immunodeficiency: clinical associations and sex differences.
650 *Clin Immunol* 2008; 128:314-21.
- 651 50. Cambronero R, Sewell WA, North ME, Webster AD, Farrant J. Up-regulation of IL-
652 12 in monocytes: a fundamental defect in common variable immunodeficiency. *J*
653 *Immunol* 2000; 164:488-94.
- 654 51. Martinez-Pomar N, Raga S, Ferrer J, Pons J, Munoz-Saa I, Julia MR, et al. Elevated
655 serum interleukin (IL)-12p40 levels in common variable immunodeficiency disease
656 and decreased peripheral blood dendritic cells: analysis of IL-12p40 and interferon-
657 gamma gene. *Clin Exp Immunol* 2006; 144:233-8.
- 658 52. McQuaid A, Tormey VJ, Trafford B, Webster AD, Bofill M. Evidence for increased
659 expression of regulatory cytokine receptors interleukin-12R and interleukin-18R in
660 common variable immunodeficiency. *Clin Exp Immunol* 2003; 134:321-7.
- 661 53. Ibanez C, Sune P, Fierro A, Rodriguez S, Lopez M, Alvarez A, et al. Modulating
662 effects of intravenous immunoglobulins on serum cytokine levels in patients with
663 primary hypogammaglobulinemia. *BioDrugs* 2005; 19:59-65.
- 664 54. Hirahara K, Nakayama T. CD4+ T-cell subsets in inflammatory diseases: beyond the
665 Th1/Th2 paradigm. *Int Immunol* 2016; 28:163-71.
- 666 55. McCaw JM, McVernon J, McBryde ES, D. MJ. Influenza: accounting for prior
667 immunity. *Science*; 325:1071.

- 668 56. Bodewes R, Fraaij P, Geelhoed-Mieras M, van Baalen C, Tiddens H, van Rossum A,
669 et al. Annual Vaccination against Influenza Virus Hampers Development of Virus-
670 Specific CD8+ T Cell Immunity in Children. *Journal Virology* 2011; 85:11995-2000.
- 671 57. Li J, Jorgensen SF, Maggadottir SM, Bakay M, Warnatz K, Glessner J, et al.
672 Association of CLEC16A with human common variable immunodeficiency disorder
673 and role in murine B cells. *Nat Commun* 2015; 6:6804.
- 674 58. Aghamohammadi A, Mohammadi J, Parvaneh N, Rezaei N, Moin M, Espanol T, et al.
675 Progression of selective IgA deficiency to common variable immunodeficiency. *Int*
676 *Arch Allergy Immunol* 2008; 147:87-92.
- 677 59. Cheraghi T, Aghamohammadi A, Mirminachi B, Keihanian T, Hedayat E,
678 Abolhassani H, et al. Prediction of the evolution of common variable
679 immunodeficiency: HLA typing for patients with selective IgA deficiency. *J Investig*
680 *Allergol Clin Immunol* 2014; 24:198-200.
- 681 60. Desombere I, Van der Wielen M, Van Damme P, Stoffel M, De Clercq N, Goilav C, et
682 al. Immune response of HLA DQ2 positive subjects, vaccinated with HBsAg/AS04, a
683 hepatitis B vaccine with a novel adjuvant. *Vaccine* 2002; 20:2597-602.
- 684 61. Li ZK, Nie JJ, Li J, Zhuang H. The effect of HLA on immunological response to
685 hepatitis B vaccine in healthy people: a meta-analysis. *Vaccine* 2013; 31:4355-61.
- 686 62. von Bulow GU, van Deursen JM, Bram RJ. Regulation of the T-independent humoral
687 response by TACI. *Immunity* 2001; 14:573-82.
- 688 63. Martinez-Gallo M, Radigan L, Almejun MB, Martinez-Pomar N, Matamoros N,
689 Cunningham-Rundles C. TACI mutations and impaired B-cell function in subjects
690 with CVID and healthy heterozygotes. *J Allergy Clin Immunol* 2013; 131:468-76.
- 691 64. Romberg N, Chamberlain N, Saadoun D, Gentile M, Kinnunen T, Ng YS, et al.
692 CVID-associated TACI mutations affect autoreactive B cell selection and activation. *J*
693 *Clin Invest* 2013; 123:4283-93.

- 694 65. Koopmans W, Woon ST, Brooks AE, Dunbar PR, Browett P, Ameratunga R. Clinical
695 variability of family members with the C104R mutation in transmembrane activator
696 and calcium modulator and cyclophilin ligand interactor (TACI). J Clin Immunol
697 2013; 33:68-73.
- 698

699 **Table 1-** Demographic data of 48 patients with CVID including eight responders ($\geq 1:40$ titer) and 40
 700 non-responders to the A(H1N1) pandemic influenza vaccine.

Parameters	Total group	Responders	Non-responders	<i>p</i> -value*
Number	48	8	40	-
Gender (F/M)	29/19	5/3	24/16	0.62
Mean age at diagnosis of CVID (years)	44.2 \pm 18.0	40.5 \pm 15.4	44.9 \pm 18.5	0.26
Mean age at study start (years)	57.7 \pm 14.9	54.3 \pm 16.2	58.3 \pm 17.4	0.60
Mean years from onset of infections to diagnosis (years)	12.9 \pm 10.6	9.4 \pm 8.2	13.6 \pm 9.7	0.44
Mean follow-up after diagnosis (years)	13.4 \pm 8.8	13.8 \pm 8.0	13.4 \pm 9.1	0.72
Familial cases (%)	9(18.7)	0	9(22.5)	0.16
Progressive primary antibody ^a deficiency (%)	14(29.1)	0	14(35)	0.048

701

702 ^a Progression of other types of primary antibody deficiency including IgA deficiency and IgG subclass
 703 deficiency to CVID.

704 * *p* values <0.05 were regarded significant and are **bolded**

705 **Table 2-** Immunologic data of 48 patients with CVID at diagnosis including 8 responders and 40 non-
 706 responders to A(H1N1) pandemic vaccine.

Parameters	Normal range	Total CVID	Responders	Non-responders	<i>p</i> -value*
IgM (mg/dl)	50-370	31.7±9.7	17.0±16.0	34.8±29.6	0.03
IgA (mg/dl)	80-380	9.2±8.4	9.5±8.4	9.2±7.0	0.76
IgG (mg/dl)	600-1500	209.2±173.4	243.7±206.1	201.7±167.8	0.20
IgG1 (mg/dl)	280-800	141.3±125.7	219.2±198.3	122.4±96.3	0.007
IgG2 (mg/dl)	115-570	49.3±37.0	48.0±36.1	49.7±37.0	0.74
IgG3 (mg/dl)	24-125	19.9±18.1	19.3±16.5	22.3±21.4	0.30
White Blood cells (cell/ul)	3500-8800	5828.5±2395.2	5971.4±1729.8	5791.4±2565.9	0.49
Lymphocyte (cell/ul)	1130-2720	1364.4±607.2	1237.3±318.0	1397.4±662.6	0.10
T cells (cell/ul)	780-2070	1126.6±489.8	1032.8±264.8	1150.9±534.1	0.056
CD4⁺ T cells (cell/ul)	490-1340	516.4±244.3	504.2±187.6	519.6±260.0	0.46
Naive CD4 (% of CD4⁺ T cells)	11-35	13.0±8.0	16.1±9.5	12.1±7.4	0.24
Memory CD4 (% of CD4⁺ T cells)	31-74	70.4±14.8	70.5±14.5	70.4±15.2	0.8
CD8⁺ T cells (cell/ul)	190-800	625.4±342.3	530.5±230.4	650.0±365.2	0.033
Naive CD8 (% of CD8⁺ T cells)	27-69	35.4±12.1	32.8±14.0	36.1±11.7	0.42
Memory CD8 (% of CD8⁺ T cells)	12-50	35.5±2.0	33.2±9.7	43.1±16.4	0.044
CD4:CD8	1-4	1.1±0.9	1.0±0.4	1.1±1.0	0.28
NK cells (cell/ul)	70-420	94.1±61.6	78.5±23.4	98.1±67.9	0.06
B cell (cell/ul)	90-400	123.7±94.9	114.2±91.2	126.2±94.5	0.56
Naïve B (% of CD19⁺ T cells)	47-84	70.2±21.4	71.6±17.6	69.7±22.8	0.72
Transitional B (% of CD19⁺ T cells)	0-1	4.3±3.6	4.2±4.0	4.4±3.9	0.48
Natural memory B (% of CD19⁺ T cells)	6-29	21.12±17.4	18.9±11.7	22.0±19.0	0.28
Switched memory B (% of CD19⁺ T cells)	9-29	6.4±6.0	8.0±7.8	5.9±5.0	0.41
Plasmablast (% of CD19⁺ T cells)	0-3.2	0.06±0.01	0.1±0.02	0.04±0.03	0.007
CD21^{low} B (% of CD19⁺ T cells)	0.7-10	15.0±10.9	8.5±4.6	17.2±11.6	0.041

707 * *p* values <0.05 were regarded significant and are **bolded**

Figure legends

Figure 1- Specific antibody response against A(H1N1) pandemic influenza vaccination after 1 and 3 months in 48 patients with CVID.

Figure 2- Clinical phenotyping of 48 patients with CVID including 8 responders and 40 non-responders to A(H1N1) pandemic influenza vaccine. During the course of the disease, all patients presented with the phenotype infections. Thus, the patients had at least the phenotype infections (here presented as “No disease-related complications”) or the phenotype infections together with one or more of the phenotypes in the figure (for detail of clinical phenotyping see **Table S2**). PLI: polyclonal lymphocytic infiltration.

Figure 3- Immunological phenotyping in percentage of 48 patients with CVID including 8 responders and 40 non-responders to A(H1N1) influenza pandemic vaccine and classification according to the **A.** Paris classification (0: low switch memory[SM] and low total memory [M], 1: low SM and normal M, 2: normal SM and normal M), **B.** Freiburg classification (1a: low SM and increased CD21^{low}, 1b: low SM and normal CD21^{low}, 2: normal SM and normal M), **C.** Euroclass classification (according to SM, CD21^{low} and Transitional [Tr] B cells) and **D.** B cell pattern classification (P1:low Tr and low M, P2: low naive and low M, P3: low marginal zone and lowM, P4: lowM and P5: normal).

Figure 4- Cytokine production of interferon gamma (IFN- γ), interleukin 2 (IL-2), IL-12, IL-5 and IL-10 in 48 patients with CVID, including 8 responders (R) and 40 non-responders (NR) to A(H1N1) pandemic influenza vaccine after phytohemagglutinin stimulation.

Fig 1.




Fig. 2




Fig 3





Fig 4



1 **Supplementary Material**

2

3 **Predictive markers for humoral vaccine response in patients with**
4 **common variable immunodeficiency (CVID)**

5

6 Gardulf et al.

7 **Table S1-** Specific antibody response to protein and polysaccharides antigens ^a

Parameters	Total CVID	Responders to Pandemrix [®]	Non-responders to Pandemrix [®]	<i>p</i> -value
	Numbers (%)	Numbers (%)	Numbers (%)	
Diphtheria Ab (normal/total tested)	11/27(40.7)	2/5(40)	9/22(40.9)	1.0
Tetanus Ab (normal/total tested)	13/34(38.2)	3/5(60)	10/29(34.4)	0.34
CMV Ab (normal/total tested)	6/17(35.2)	1/4(25)	5/13(38.4)	1.0
MCV Ab (normal/total tested)	2/10(20)	0/1(0)	2/9(22.2)	1.0
Teichoic acid Ab (normal/total tested)	9/16(56.2)	1/3(33.3)	8/13(61.5)	0.55
Hib Ab (normal/total tested)	6/11(54.5)	1/1(100)	5/10(50)	1.0
Pneumococcus Ab (normal/total tested)	1/9(11.1)	1/1(100)	0/8(0)	0.11
PPS6 Ab (normal/total tested)	7/16(43.7)	1/3(33.3)	6/13(23)	1.0
PPSV23 Ab (normal/total tested)	4 ^b /9(44.4)	1/3(33.3)	3/6(50)	1.0

8 ^a CMV: Cytomegalovirus, Hib: *Haemophilus influenzae* type b, MCV: Meningococcal conjugate
9 vaccine, PPS6: Phospho-p70 S6 Kinase, PPSV23: nonconjugated polysaccharide, Pneumovax 23-
10 valent vaccines.

11 ^b These four patients had defective isohemagglutinins tests and therefore, according to the criteria of
12 the European Society for Immunodeficiencies (ESID), they have a diagnosis of CVID.

13 **Table S2-** Medical manifestations and clinical phenotypes of 48 patients with CVID at diagnosis
 14 including 8 responders and 40 non-responders to A(H1N1) pandemic vaccine.

Parameters	Total CVID Numbers (%)	Responders to Pandemrix® Numbers (%)	Non-responders to Pandemrix® Numbers (%)	<i>p</i> -value*
Medical manifestations				
Recurrent upper respiratory tract infections	48(100)	8(100)	40(100)	1
Lower respiratory tract infections	44	5(62.5)	33(82.5)	0.20
Bronchiectasis	7(14.5)	0	7(17.5)	0.10
Autoimmune cytopenia	5(10.4)	0	5(12.5)	0.29
Inflammatory bowel diseases	6(12.5)	1(12.5)	5(12.5)	1
Celiac disease	4(8.3)	1(12.5)	3(7.5)	0.64
Psoriasis	3(6.2)	0	3(7.5)	0.42
Vitiligo	2(4.1)	0	2(5)	0.51
Diabetes mellitus type 1	2(4.1)	0	2(5)	0.51
Systemic lupus erythematosus	1(2.1)	1(12.5)	1(2.5)	0.19
Waldeyer's lymphadenopathy	5(10.4)	1(12.5)	4(10)	0.83
Generalized lymphadenopathy	4(8.3)	0	4(10)	0.35
Splenomegaly	2(4.1)	0	2(5)	0.51
Granulomatous disease	1(2.1)	0	1(2.5)	0.65
Malignancies	14(29.1)	2(25)	12(30)	0.38
Breast cancer	2(4.1)	1(12.5)	1(2.5)	0.19
Colon cancer	2(4.1)	1(12.5)	1(2.5)	0.19
Thymoma/lymphoma	9(12.5)	0	9(22.5)	0.13
Prostate cancer	1(2.1)	0	1(2.5)	0.65
IgE mediated atopic disorders	13(27.1)	4(50)	9(22.5)	0.12
Asthma	4(8.3)	0	4(10)	0.35
Clinical phenotypes				
Infections without any other disease-related complications	14(29.1)	1(12.5)	13(32.5)	0.12
Autoimmunity phenotype	14(29.1)	2(25)	12(30)	0.38
Enteropathy phenotype	4(8.3)	4(50)	0	<0.001
Polyclonal lymphocytic infiltration phenotype	3(6.2)	0	3(7.5)	0.42
Overlapping phenotypes	13(27.0)	1(12.5)	12(30)	0.30

15 * *p* values <0.05 were regarded significant and are **bolded**