



City Research Online

City St George's, University of London

Citation: Forini, V., Puletti, V. G. M., Griguolo, L., Seminara, D. & Vescovi, E. (2015). Remarks on the geometrical properties of semiclassically quantized strings. *Journal of Physics A: Mathematical and Theoretical*, 48(47), 475401. doi: 10.1088/1751-8113/48/47/475401

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/19733/>

Link to published version: <https://doi.org/10.1088/1751-8113/48/47/475401>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Remarks on the geometrical properties of semiclassically quantized strings

V. Forini^{a,1}, V. Giangreco M. Puletti^{b,2}, L. Griguolo^{c,3}, D. Seminara^{d,4}, E. Vescovi^{a,e,1}

^a*Institut für Physik, Humboldt-Universität zu Berlin, IRIS Adlershof,
Zum Großen Windkanal 6, 12489 Berlin, Germany*

^b*University of Iceland, Science Institute, Dunhaga 3, 107 Reykjavik, Iceland*

^c*Dipartimento di Fisica e Scienze della Terra, Università di Parma and INFN Gruppo Collegato di
Parma, Viale G.P. Usberti 7/A, 43100 Parma, Italy*

^d*Dipartimento di Fisica, Università di Firenze and INFN Sezione di Firenze, Via G. Sansone 1,
50019 Sesto Fiorentino, Italy*

^e*Perimeter Institute for Theoretical Physics, Waterloo, Ontario N2L 2Y5, Canada*

Abstract

We discuss some geometrical aspects of the semiclassical quantization of string solutions in Type IIB Green-Schwarz action on $\text{AdS}_5 \times \text{S}^5$. We concentrate on quadratic fluctuations around classical configurations, expressing the relevant differential operators in terms of (intrinsic and extrinsic) invariants of the background geometry. The aim of our exercises is to present some compact expressions encoding the spectral properties of bosonic and fermionic fluctuations. The appearing of non-trivial structures on the relevant bundles and their role in concrete computations are also considered. We corroborate the presentation of general formulas by working out explicitly a couple of relevant examples, namely the spinning string and the latitude BPS Wilson loop.

¹ {valentina.forini,edoardo.vescovi}@physik.hu-berlin.de

² vgmp@hi.is

³ luca.griguolo@fis.unipr.it

⁴ seminara@fi.infn.it

Contents

1	Introduction	2
2	The background equation	5
3	Bosonic Fluctuations	6
3.1	The bosonic Lagangian	6
3.2	The normal bundle	12
3.3	Mass matrix and sum rules	13
4	Fermionic fluctuations	16
4.1	The fermionic kinetic term	17
4.2	Fermionic “mass” matrix: the flux term	19
4.3	Comparison with bosonic masses and quantum divergences	22
5	Applications	24
5.1	Spinning strings	24
5.2	String dual to latitude Wilson loops	27

1 Introduction

The geometric properties of string worldsheets embedded in a D -dimensional space-time, and of linearized perturbations above them, have been object of various studies since the seminal observation on the relevance of quantizing string models [1]. In the framework of the AdS/CFT correspondence, the semiclassical study of strings in non-trivial backgrounds [2] has played a crucial role, especially in connection with the detection of the underlying integrable structure [3]. At the same time it provides a powerful tool to check, at strong coupling, exact QFT results obtained through localization procedure [4–7] for BPS [8–13] and non-BPS observables [14–22]¹. As a matter of fact a large variety of classical string solutions have been proposed to correspond to CFT gauge-invariant operators, Wilson loops or dimensionally reduced amplitudes, the original suggestions being supported beyond the leading classical order by sometimes non-trivial calculations at one-loop order [9, 18, 20, 22, 23, 25–35] (see also [36, 37] for higher order computations in special cases, the so-called homogenous solutions, for which derivatives of the background fields are constant).

The natural setting in which these analyses have been performed is the Green-Schwarz σ -model on $\text{AdS}_5 \times \text{S}^5$ [38], the relevant string background for $\mathcal{N} = 4$ Super Yang-Mills gauge theory. The first step in order to compute one-loop quantum corrections to classical solutions of the string σ -model is of course to derive the quadratic action for the small fluctuations. Then,

¹The precise match between results obtained via semiclassical quantization and the exact prediction obtained via supersymmetric localization does not go beyond leading order in σ -model perturbation theory, see [23, 24].

after appropriate gauge-fixings conveniently chosen according to the original form of the action (Polyakov or Nambu-Goto) and a careful definition of the path-integral measure, the problem is reduced to the evaluation of a bunch of bosonic and fermionic functional determinants. The geometry of the classical string background is encoded into the structure of the differential operators entering the computations and in the possible appearance of zero-modes, affecting the integration measure. Finally a regularization procedure, compatible with the symmetry of the specific problem, should be exploited to derive sensible results from the formal expression of the one-loop effective action. This project was first addressed in [9], where a systematic treatment of the Green-Schwarz (GS) string in curved $\text{AdS}_5 \times \text{S}^5$ space was initiated and the quadratic fluctuation operators in conformal and static gauges (for Polyakov and Nambu-Goto actions) were found. A careful treatment of the measure factors and ghost determinants was also presented. The aim of that analysis was mainly focussed on the study of particular minimal surfaces, namely the ones associated to the straight, the circular and the antiparallel lines Wilson loops. Some relevant formulas were somehow tailored on the specific examples and on the use of the conformal gauge for the Polyakov path-integral. This pioneering paper was followed along the years by many investigations, both on the open string side (mainly minimal surfaces associated to BPS and non-BPS Wilson loops) and on the closed string side (for example different classes of string solutions related to CFT operators). Although a large number of results were obtained from one-loop quantum corrections, sometimes brilliantly confirming the expectations from integrability and localization, all this analysis relied somehow on the particular form of the classical string configuration. To further increase the effective power of this approach, it would be desirable to have a general and manifestly *covariant* formalism to describe fluctuations, which should be also independent of the particular string solution or of the background in which the string is embedded. Here we attempt some modest steps in this direction, collecting and generalizing some useful formulas previously appeared in the literature and presenting the results of a series of exercises that we hope interesting for people working in the field.

The central point of our analysis is the application of some elementary concepts of intrinsic and extrinsic geometry to the properties of string worldsheet embedded in a D -dimensional curved space-time. We take full advantage of the equations of Gauss, Codazzi, and Ricci for surfaces embedded in a general background to obtain simple and general expressions for perturbations over them. We follow and enlarge earlier investigations [9,39]², starting from the Polyakov formulation and trying to present a systematic and self-consistent perspective to the study of fluctuations in the $\text{AdS}_5 \times \text{S}^5$. The main result consists in general formulas for bosonic and fermionic fluctuation operators above a classical string solution: expressions as (3.56)-(3.59)-(3.60) and (4.22)-(4.28)-(4.31)-(4.46)-(4.47) only require as an input generic properties of the classical configuration and basic information about the space-time background. The inclusion of a suitable choice of orthonormal vectors which are orthogonal to the surface spanned by the string solution will also play a major role. In particular, after fixing the conformal gauge, it allows to decouple explicitly the longitudinal modes arriving to a final

²See also [40,41] and references therein, where this analysis has been exploited for the description of QCD strings or stability effects for membrane solutions, and the more recent [42].

expression, similar to ones that would be obtained in the Nambu-Goto formalism in the static gauge for bosons. In the fermionic sector the reduction of Green-Schwarz fermions to a set of two-dimensional Dirac spinor is equivalently accomplished. We provide explicit expressions in terms of geometric invariants for bosonic and fermionic “masses”, noticing that in all the cases previously analyzed simplifications occur which are associated to the flatness of the normal bundle. While in the bosonic case similar formulas appeared before scattered in the literature, our treatment of the fermionic case, due to the complications related to the flux term, is somehow novel in its generality.

To proceed in the one-loop analysis, one has then to compute the functional determinants associated to the fluctuations operators, which can be done with standard methods for functional determinants (see for example [18, 23, 33–35]) and could involve many regularization subtleties [43]. Here we do not address the problem of regularization procedure and other important issues, as the appropriate definition of integration measure, κ -symmetry ghosts, Jacobians due to change of fluctuation basis are also left to future investigations. These topics should deserve a careful study, especially when BPS configurations are considered and the quantum fluctuations must preserve this property.

A natural generalization of our investigations concerns Type IIA and IIB string backgrounds relevant for the $\text{AdS}_4/\text{CFT}_3$ and $\text{AdS}_3/\text{CFT}_2$ correspondences respectively: we expect it should be possible simply exploiting some general features of their geometry. For example, backgrounds like $\text{AdS}_4 \times \mathbb{C}P^3$, $\text{AdS}_3 \times S^3 \times M^4$, $\text{AdS}_3 \times S^2 \times M^5$, $\text{AdS}_2 \times S^2 \times M^6$ (where $M^4 = T^4, S^3 \times S^1$, $M^5 = S^3 \times T^2$) are direct products of symmetric spaces, which results in a structure of the Riemann tensor resembling the “separability” of (3.40) and allowing the writing of formulas similar to (3.55)-(3.57).

In the perspective adopted in this paper, there is also no explicit reference to the classical integrability of the Green-Schwarz superstring on $\text{AdS}_5 \times S^5$ [3]. In a number of semiclassical studies [18, 33, 35] the underlying integrable structure of the $\text{AdS}_5 \times S^5$ background emerges, for example, in the appearance of certain special, integrable, differential operators [33, 35], whose determinants can be calculated explicitly and result in closed (albeit in integral form) expressions for the one loop partition functions³. The question of a deeper relation between such geometric approach to fluctuations and the integrability of the σ -model of interest should become more manifest within the algebraic curve approach to semiclassical quantization [52–54], likely on the lines of [55] and is an interesting issue to be addressed in the future.

The paper proceeds as follows. The geometrical formulation of classical string solutions as minimal surfaces is briefly recalled in Section 2. In Section 3 we discuss the bosonic sector. After reviewing previous analysis based on background field method for nonlinear σ -models and the expansion in normal coordinates, we write the relevant contributions in terms of intrinsic and extrinsic geometric invariants of the classical solution. We discuss the gauge-fixing and

³In another kind of perturbative analysis of the worldsheet σ -model, i.e. the perturbative evaluation of the massive S-matrix for the elementary excitations around the BMN vacuum [44–46] (see also [15, 47] for reviews), the one-loop computation for the full $\text{AdS}_5 \times S^5$ case [48–50] (in a certain regularization scheme) reproduces exactly the results [51] predicted by (symmetries and) quantum integrability.

the decoupling of the longitudinal modes, as well as the arising of gauge connections in the covariant derivatives associated to the structure of normal bundle. The spectral properties are also investigated, obtaining the mass matrix and deriving some sum rules. Section 4 is instead devoted to the fermionic sector. We explicitly obtain the fermionic kinetic terms by performing suitable rotations that reduce the GS spinor to two-dimensional Dirac fermions and observe the arising of normal bundle gauge connection as in the bosonic case. Then we discuss the mass matrix that, after a careful treatment of the flux contribution, is expressed in terms of geometrical invariants. We conclude with Section 5, where a couple of relevant situations (the well-known spinning string solution of [2, 26] and fluctuations over the minimal surface associated to the 1/4 BPS Wilson loop operator [11, 56, 57]) are considered, in which the general structures previously derived are exemplified.

2 The background equation

We start by recalling some basic fact about classical string theory. In particular we will review the statement that classical string solutions are minimal surfaces, *i.e.* surfaces of vanishing mean curvature. We shall deal with classical backgrounds which are extrema of the Nambu-Goto action (fermions are of course zero at classical level)

$$S_{N.G.} = \int_{\Sigma} d^2\sigma \sqrt{\gamma}, \quad (2.1)$$

where $\gamma_{\alpha\beta} \equiv G_{mn} \partial_{\alpha} X^m \partial_{\beta} X^n$ ($\alpha, \beta = 1, 2$) is the induced metric, namely the pull-back of the target space metric G_{mn} ($l, m, n, p, \dots = 1, \dots, D$) on the worldsheet Σ . The background X^m solves the Euler-Lagrangian equation

$$\partial_{\alpha} \left(\sqrt{\gamma} \gamma^{\alpha\beta} G_{mn} \partial_{\beta} X^n \right) - \frac{1}{2} \sqrt{\gamma} \gamma^{\alpha\beta} \partial_m G_{np} \partial_{\alpha} X^n \partial_{\beta} X^p = 0, \quad (2.2)$$

which is conveniently rewritten as follows

$$G_{mn} \square X^n = \frac{1}{2} \gamma^{\alpha\beta} \partial_m G_{np} \partial_{\alpha} X^n \partial_{\beta} X^p - \gamma^{\alpha\beta} \partial_{\alpha} X^p \partial_p G_{mn} \partial_{\beta} X^n \quad (2.3)$$

where $\square = \frac{1}{\sqrt{\gamma}} \partial_{\alpha} (\sqrt{\gamma} \gamma^{\alpha\beta} \partial_{\beta})$ is the covariant Laplacian on worldsheet scalars. Introducing Γ_{np}^m , the Christoffel connections for G_{np} , we have

$$\square X^m + \gamma^{\alpha\beta} \Gamma_{np}^m \partial_{\alpha} X^n \partial_{\beta} X^p = 0. \quad (2.4)$$

The covariant Laplacian can be further expanded in terms of the induced metric $\gamma_{\alpha\beta}$ and the related Christoffel connections $\Lambda_{\alpha\beta}^{\rho}$ to find

$$\gamma^{\alpha\beta} (\partial_{\alpha} \partial_{\beta} X^m - \Lambda_{\alpha\beta}^{\rho} \partial_{\rho} X^m + \Gamma_{np}^m \partial_{\alpha} X^n \partial_{\beta} X^p) \equiv \gamma^{\alpha\beta} K_{\alpha\beta}^m = 0, \quad (2.5)$$

where the second fundamental form $K_{\alpha\beta}^m$ of the embedding (or *extrinsic curvature*) has been introduced. Then, the string equation of motion simply states that the mean curvature K^m vanishes

$$K^m \equiv \gamma^{\alpha\beta} K_{\alpha\beta}^m = 0. \quad (2.6)$$

As a matter of fact the extrinsic curvature is automatically orthogonal to the two vectors $t_\alpha^m \equiv \partial_\alpha X^m$ ($\alpha = 1, 2$), tangent to the worldsheet,

$$G_{mn} t_\alpha^m K_{\rho\sigma}^n = 0. \quad (2.7)$$

Physically this means that only $D - 2$ of the D equations in (2.6) are independent and they govern the $D - 2$ transverse degrees of freedom. The longitudinal ones are obviously gauge degrees of freedom.

The same result can be of course recovered from the Polyakov action

$$S_P = \int_\Sigma d^2\sigma \sqrt{h} h^{\alpha\beta} G_{mn} \partial_\alpha X^m \partial_\beta X^n \quad (2.8)$$

where the 2d worldsheet metric $h_{\alpha\beta}$ is now an independent field. In this case the dynamical equations for the embedding coordinates X^m are slightly different and read

$$\square_h X^m + h^{\alpha\beta} \Gamma_{np}^m \partial_\alpha X^n \partial_\beta X^p = h^{\alpha\beta} K_{\alpha\beta}^m + h^{\alpha\beta} (\Lambda_{\alpha\beta}^\rho - \tilde{\Gamma}_{\alpha\beta}^\rho) \partial_\rho X^m = 0, \quad (2.9)$$

where \square_h denotes the covariant Laplacian and $\tilde{\Gamma}$ the Christoffel symbols for the auxiliary metric $h_{\alpha\beta}$, whereas Λ are the ones for the induced metric. But if we use that the algebraic equation for the metric field $h_{\alpha\beta}$ is solved by $h_{\alpha\beta} = e^\varphi \gamma_{\alpha\beta}$, the last term in (2.9) vanishes,

$$h^{\alpha\beta} (\Lambda_{\alpha\beta}^\rho - \tilde{\Gamma}_{\alpha\beta}^\rho) = 0, \quad (2.10)$$

and the string equation of motion again reduces to (2.6).

3 Bosonic Fluctuations

In this section we shall discuss the action for bosonic fluctuations around a classical background. After reviewing previous analysis [9, 39] based on background field method for non-linear σ -models and the virtues of the expansion in “geodesic” normal coordinates [58], we write the relevant contributions in terms of intrinsic and extrinsic geometric invariants of the classical solution.

3.1 The bosonic Lagrangian

We will discuss the bosonic fluctuations starting from the Polyakov action

$$S = \int d^2\sigma \sqrt{h} h^{\alpha\beta} \partial_\alpha \tilde{X}^n \partial_\beta \tilde{X}^m G_{mn}(\tilde{X}). \quad (3.1)$$

A well-known subtlety of the expansion of a non-linear σ -model around a classical background X^m [58] is that writing it as a power series in terms of fluctuations, defined as $\delta X^m = \tilde{X}^m - X^m$, does not lead to a manifestly covariant expression for the series coefficients. As a matter of fact the difference between coordinates values at nearby points of the manifold does not transform simply under reparametrization. The easiest way to obtain a manifestly covariant form for the coefficients is to take advantage of the method of normal (or Riemann) coordinates, expressing

δX^m as a local power series in spacetime vectors, those tangent to the spacetime geodesic connecting X^m with $X^m + \delta X^m$ [58]. More precisely one considers a geodesic $X^m(t)$ with t parametrizing the arc length such that

$$X^m(0) = X^m \quad \text{and} \quad X^m(1) = \tilde{X}^m. \quad (3.2)$$

Solving then the geodesic equation for $X^m(t)$

$$\ddot{X}^m(t) + \Gamma_{np}^m \dot{X}^n(t) \dot{X}^p(t) = 0 \quad (3.3)$$

in terms of the tangent vector to this geodesic in $t = 0$

$$\zeta^m \equiv \dot{X}^m(0) \quad (3.4)$$

one finds

$$X^m(t) = X^m + t\zeta^m - \frac{1}{2}t^2\Gamma_{np}^m\zeta^n\zeta^p + O(t^3). \quad (3.5)$$

For $t = 1$, this means ⁴

$$\tilde{X}^m = X^m + \zeta^m - \frac{1}{2}\Gamma_{np}^m\zeta^n\zeta^p + O(\zeta^3) \quad \Rightarrow \quad \tilde{X}^m - X^m = \zeta^m - \frac{1}{2}\Gamma_{np}^m\zeta^n\zeta^p + O(\zeta^3). \quad (3.6)$$

where $\Gamma_{np}^m \equiv \Gamma_{np}^m(X^m)$. The difference $\delta X^m = \tilde{X}^m - X^m$ is now the desired local power series in the vector ζ^m , which can then be conveniently used as a fundamental variable. Combining the expansions of the derivatives of the embedding coordinates

$$\partial_\alpha \tilde{X}^m = \partial_\alpha X^m + \nabla_\alpha \zeta^m - \partial_\alpha X^n \Gamma_{nr}^m \zeta^r - \frac{1}{2} \partial_\alpha X^r (\partial_r \Gamma_{np}^m - 2\Gamma_{nl}^m \Gamma_{rp}^l) \zeta^n \zeta^p - \Gamma_{np}^m \zeta^n \nabla_\alpha \zeta^p + O(\zeta^3),$$

where $\nabla_\alpha \zeta^m \equiv \partial_\alpha \zeta^m + \Gamma_{np}^m \partial_\alpha X^n \zeta^p$, with the contribution of the target metric

$$G_{mn}(\tilde{X}) = G_{mn}(X) + \left(\zeta^r - \frac{1}{2} \Gamma_{pq}^r \zeta^p \zeta^q \right) \partial_r G_{mn}(X) + \frac{1}{2} \zeta^r \zeta^s \partial_r \partial_s G_{mn}(X) + O(\zeta^3), \quad (3.7)$$

we find the fluctation action in the Polyakov formulation (3.1) (see [39] for example)

$$S = S_B^{(0)}(X) + \int d^2\sigma \sqrt{h} h^{\alpha\beta} [\nabla_\alpha \zeta^m \nabla_\beta \zeta^n G_{mn} - R_{rm,sn} \zeta^r \zeta^s \partial_\alpha X^m \partial_\beta X^n] + O(\zeta^3). \quad (3.8)$$

The term $S_B^{(0)}(X)$ denotes the classical action, while the second one describes the quadratic fluctuations and it will be denoted with $S_B^{(2)}$ in the following. In order to have a canonically normalized kinetic term it is convenient to introduce a set of vielbein E_m^A ($A, B, \dots = 1, \dots, D$) for the target metric

$$G_{mn} = \eta_{AB} E_m^A E_n^B \quad (3.9)$$

⁴ At quadratic level for fluctuations, the term linear in ζ does not play a crucial role. In fact it yields only contributions which are proportional to the equation of motions:

$$S = S_0 + \int d^2\sigma \left. \frac{\delta S}{\delta \tilde{X}^m} \right|_{\tilde{X}=X} \zeta^m - \frac{1}{2} \int d^2\sigma \left(\left. \frac{\delta S}{\delta \tilde{X}^m \delta \tilde{X}^n} \right|_{\tilde{X}=X} \zeta^m \zeta^n - \left. \frac{\delta S}{\delta \tilde{X}^m} \right|_{\tilde{X}=X} \Gamma_{ab}^m \zeta^a \zeta^b \right).$$

However, its introduction allows to simplify the algebra involved in the computation.

and a set of zweibein e_α^a ($a, b, \dots = 1, 2$) for $h_{\alpha\beta}$. In terms of the redefined fluctuations fields

$$\xi^A = E_m^A \zeta^m, \quad (3.10)$$

the quadratic fluctuation action for bosons becomes [9, 39]

$$S_{2B} = \int d^2\sigma \sqrt{h} [h^{\alpha\beta} D_\alpha \xi^A D_\beta \xi_A - M_{A,B} \xi^A \xi^B], \quad (3.11)$$

where the *mass matrix* [9, 39]

$$M_{AB} = R_{AM, BN} t_a^M t_a^N \quad (3.12)$$

is defined through two vectors tangent to the worldsheet

$$t_a^A = E_m^A e_a^\alpha \partial_\alpha X^m, \quad (3.13)$$

and the covariant derivative now reads

$$D_\alpha \xi^A = \partial_\alpha X^A + \Omega^A{}_{Bm} \xi^B \partial_\alpha X^m, \quad (3.14)$$

where the spin-connection $\Omega^A{}_{Bm}$ replaced the usual Christoffel symbol. To better understand the geometrical structure of the Lagrangian (3.11), we introduce $(D-2)$ orthonormal vector fields N_i^A orthogonal to the worldsheet, and decompose the field ζ^A in directions tangent (x^a) and orthogonal (y^i) to it

$$\xi^A = x^a t_a^A + y^i N_i^A \quad a = 1, 2 \quad \text{and} \quad i, j, k, l = 1, \dots, D-2. \quad (3.15)$$

As is well-known from the general theory of submanifolds [59], this decomposition carries over to the covariant derivatives and one finds

$$t_a^A D_\beta \xi^A = \mathcal{D}_\beta x^a - K_{A\beta}^a N_i^A y^i, \quad N_i^B D_\beta \xi^B = \mathcal{D}_\beta y^i + x^a N_B^i K_{a\beta}^B. \quad (3.16)$$

Here \mathcal{D}_α is the covariant derivative on the worldsheet and it acts differently on x^a and y^i

$$\mathcal{D}_\alpha x^a \equiv \partial_\alpha x^a + \omega^a{}_{b\alpha} x^b \quad \text{and} \quad \mathcal{D}_\alpha y^i \equiv \partial_\alpha y^i - A^i{}_{j\alpha} y^j, \quad (3.17)$$

since x^a lives in the tangent bundle of the worldsheet, while y^i is a section of the normal bundle. The connection $A^i{}_{j\alpha}$ on the normal bundle⁵ is given by

$$A^i{}_{j\alpha} \equiv N^{jB} D_\beta N_B^i = N_j^B (\partial_\beta N_B^i - N_C^i \Omega^C{}_{B\beta}). \quad (3.18)$$

As usual the action of \mathcal{D}_α on tensors with indices on both bundles is obtained combining the two actions in (3.17). The tensor $K_{A\beta}^a = E_{Am} e^{a\alpha} K_{\alpha\beta}^m$ in (3.16) is the extrinsic curvature (2.5) of the embedding expressed in a mixed basis. In the following we will make use of the *Gauss-Codazzi* equation

$$R_{ACBD} t_\alpha^A t_\rho^C t_\beta^B t_\sigma^D = {}^{(2)}R_{\alpha\rho\beta\sigma} + \eta_{AB} K_{\rho\beta}^A K_{\sigma\alpha}^B - \eta_{AB} K_{\rho\sigma}^A K_{\beta\alpha}^B, \quad (3.19)$$

⁵The normal bundle is an $SO(D-2)$ bundle and $A^i{}_{j\alpha}$ is a gauge connection induced on this bundle by the classical solution.

an integrability condition relating the curvature ${}^{(2)}R_{\alpha\rho\beta\sigma}$ to the extrinsic and background geometry as characterized by the extrinsic curvature $K_{\alpha\beta}^A$ and the space-time Riemann tensor R^A_{CBD} . Another useful constraint on the covariant derivative of the extrinsic curvature is provided by the *Codazzi-Mainardi* equation

$$\mathcal{D}_\alpha K_{\beta\gamma}^i - \mathcal{D}_\beta K_{\alpha\gamma}^i = R_{MNRSt}^M t_\beta^N t_\gamma^S N^{Ri} \quad [K_{\alpha\beta}^i \equiv K_{\alpha\beta}^A N_A^i]. \quad (3.20)$$

Taking into account (3.19), (3.20) and the equation of motion (2.6) for the background, the quadratic Lagrangian (3.11) finally appears to be

$$\begin{aligned} \mathcal{L} = \sqrt{h} [& (h^{\alpha\beta} \mathcal{D}_\alpha x^a \mathcal{D}_\beta x_a - {}^{(2)}R_{ab} x^a x^b) + h^{\alpha\beta} \mathcal{D}_\alpha y^i \mathcal{D}_\beta y_i - \\ & - 2h^{\alpha\beta} (\mathcal{D}_\alpha x^a K_{i,a\beta} y^i - \mathcal{D}_\alpha y^i x^a K_{i,a\beta}) - 2m_{ai} x^a y^i - m_{ij} y^i y^j]. \end{aligned} \quad (3.21)$$

Above, the matrices appearing in the mass terms are defined as follows

$$m_{ai} = -h^{\alpha\beta} \nabla_\alpha K_{i,a\beta} \quad \text{and} \quad m_{ij} = R_{AM,BN} t_c^M t_c^N N_i^A N_j^B - h^{\alpha\beta} h^{\rho\sigma} K_{i,\alpha\rho} K_{j,\beta\sigma}. \quad (3.22)$$

So far we have treated the independent metric $h_{\mu\nu}$ as a non-dynamical field. We should recall that in Polyakov's formulation also $h_{\mu\nu}$ fluctuates,

$$h_{\mu\nu} = \hat{h}_{\mu\nu} + \mathfrak{h}_{\mu\nu}, \quad (3.23)$$

with respect to a classical background $\hat{h}_{\rho\sigma} = e^\varphi \gamma_{\rho\sigma}$ which solves the equations of motion. In particular this means that all the $h_{\mu\nu}$ appearing in the previous analysis must be replaced with $\hat{h}_{\rho\sigma}$. The quadratic part of the Polyakov action involving the fluctuations $\mathfrak{h}_{\mu\nu}$ pedantically reads

$$\begin{aligned} S_{\mathfrak{h}} = \int d^2\sigma e^\varphi \sqrt{\gamma} [& \frac{1}{4} (\gamma^{\sigma\mu} \gamma^{\rho\nu} + \gamma^{\sigma\nu} \gamma^{\rho\mu} - \gamma^{\mu\nu} \gamma^{\rho\sigma}) \mathfrak{h}_{\mu\nu} \mathfrak{h}_{\rho\sigma} - \\ & - (\mathcal{D}^\mu x^\nu + \mathcal{D}^\nu x^\mu - \gamma^{\mu\nu} \mathcal{D}_\alpha x^\alpha - 2y_i K^{i\mu\nu}) \mathfrak{h}_{\mu\nu}], \end{aligned} \quad (3.24)$$

where $x^\alpha = e_\alpha^a x^a$. Of course (3.24) only depends on the traceless part $\bar{\mathfrak{h}}_{\mu\nu}$ of $\mathfrak{h}_{\mu\nu}$ as required by Weyl invariance. To deal with the quadratic fluctuation (3.21) and (3.24), we have different possibilities. For instance, we can choose the *conformal gauge* for the metric fluctuations

$$\mathfrak{h}_{\mu\nu} = e^\varphi \gamma_{\mu\nu} \delta\varphi. \quad (3.25)$$

The action $S_{\mathfrak{h}}$ then vanishes identically, while the ghost action associated to the choice (3.25) is

$$\mathcal{L}_{\text{ghost}} = \frac{1}{\sqrt{2}} b^{\mu\nu} \delta_c^{\text{diff.}} \left(\mathfrak{h}_{\mu\nu} - \frac{1}{2} \gamma_{\mu\nu} \gamma^{\alpha\beta} \mathfrak{h}_{\alpha\beta} \right) = \frac{1}{\sqrt{2}} b^{\mu\nu} (\mathcal{D}_\mu c_\nu + \mathcal{D}_\nu c_\mu - \gamma_{\mu\nu} \mathcal{D}_\alpha c^\alpha). \quad (3.26)$$

Here $\delta_c^{\text{diff.}}$ is the BRST variation under diffeomorphism of $h_{\mu\nu}$ with parameter c , and the ghost $b^{\mu\nu}$ is a symmetric traceless tensor. The full ghost contribution is therefore encoded into a functional determinant, obtained by integrating over c and $b^{\mu\nu}$. It will correct the one-loop quantum result of the bosonic fluctuations, as we will see in the following. Concretely the

computation of the ghost determinant means here to solve the following eigenvalue problem in the background geometry

$$\begin{cases} \frac{1}{\sqrt{2}}(\mathcal{D}_\mu c_{(n)\nu} + \mathcal{D}_\nu c_{(n)\mu} - \gamma_{\mu\nu} \mathcal{D}_\alpha c_{(n)}^\alpha) = \lambda_n b_{(n)\mu\nu} \\ -\sqrt{2} \mathcal{D}_\mu b_{(n)}^{\mu\nu} = \lambda_n c_{(n)}^\nu. \end{cases} \quad (3.27)$$

From the first equation for $b_{\mu\nu}$ one gets

$$-(\square c_{(n)}^\nu + R^\nu{}_\mu c_{(n)}^\mu) = \lambda_n^2 c_{(n)}^\mu \quad (3.28)$$

and therefore the ghost determinant is

$$\Delta_{gh} = \prod_n \lambda_n = \left(\prod_n \lambda_n^2 \right)^{1/2} = [\det(-\square \delta_\mu^\nu - R^\nu{}_\mu)]^{1/2}. \quad (3.29)$$

We can now decouple the longitudinal fluctuation (x^a) from the transverse ones (y^i). We start from the action (3.21), and derive the equation of motion for the fluctuation parallel to the worldsheet:

$$\square x_\alpha + R_{\alpha\beta} x^\beta = \mathcal{D}^\beta B_{\beta\alpha}, \quad B_{\alpha\beta} = 2y_i K_{\alpha\beta}^i \quad (3.30)$$

where we introduced the traceless tensor $B_{\alpha\beta}$. This equation can be equivalently written as follows

$$P_1(x)_{\alpha\beta} = \mathcal{D}_\beta x_\alpha + \mathcal{D}_\alpha x_\beta - \gamma_{\alpha\beta} \mathcal{D}_\rho x^\rho = B_{\alpha\beta}^\parallel \quad (3.31)$$

where we have introduced a projector P_1 , acting on the space of vectors and producing symmetric traceless tensors. We can conveniently decompose the traceless symmetric tensor $B_{\alpha\beta}$ into $B_{\alpha\beta}^\parallel + B_{\alpha\beta}^\perp$, with $B_{\alpha\beta}^\parallel \in \text{range}(P_1)$ and $B_{\alpha\beta}^\perp \in \text{range}(P_1)^\perp = \text{Ker}(P_1^\dagger)$. We remark that we will be only interested in worldsheet with the topology of the sphere or the disc, where $B_{\alpha\beta}^\perp = 0$ and thus $B_{\alpha\beta} = B_{\alpha\beta}^\parallel$ ⁶. Considering now a solution \bar{x} of (3.31) and performing the shift $x^a \mapsto \bar{x}^a + x^a$ in the path-integral, all mixed terms xy in (3.21) disappear and an additional contribution shows up in the quadratic action

$$\bar{x}^\alpha \mathcal{D}^\beta B_{\alpha\beta} = \mathcal{D}^\beta (\bar{x}^\alpha B_{\alpha\beta}) - \frac{1}{2} B^{\alpha\beta} B_{\alpha\beta} = -2y_i y_j K_{\alpha\beta}^i K^{j\alpha\beta}. \quad (3.32)$$

We are left therefore with the quadratic Lagrangian

$$\mathcal{L} \equiv \mathcal{L}_{\text{long}} + \mathcal{L}_{\text{transv}} \quad (3.33)$$

where

$$\mathcal{L}_{\text{long}} = \sqrt{\gamma} (\gamma^{\alpha\beta} \mathcal{D}_\alpha x^a \mathcal{D}_\beta x_a - {}^{(2)}R_{ab} x^a x^b) \quad (3.34)$$

and

$$\begin{aligned} \mathcal{L}_{\text{transv}} &= \sqrt{\gamma} (\gamma^{\alpha\beta} \mathcal{D}_\alpha y^i \mathcal{D}_\beta y_i - \mathcal{M}_{ij} y^i y^j), \\ \mathcal{M}_{ij} &= R_{AM,BN} t_c^{cM} t_c^{cN} N_i^A N_j^B + K_{i,\alpha\beta} K_j^{\alpha\beta}. \end{aligned} \quad (3.35)$$

⁶This fact corresponds to the the well-known property of the absence of non-trivial Beltrami differentials at genus 0.

After the above redefinition, the operator controlling the fluctuations x^a parallel to the worldsheet in (3.34) coincides with the one appearing in the ghost determinant (3.29), but we remark that this does *not* mean in general that the corresponding determinant will be simply cancelled by the ghost contribution. For instance in the case of open strings different boundary conditions should be imposed for the two determinants. Moreover the treatment of the ghost operator requires additional care since it might contain zero modes associated to the Killing vectors of the worldsheet metric $\gamma_{\alpha\beta}$.

We can reach the same expression for $\mathcal{L}_{\text{transv}}$ following an alternative way. Instead of setting to zero the traceless part $\bar{h}_{\mu\nu}$ of $h_{\mu\nu}$ by choosing the conformal gauge fixing (3.25), we can decouple it from the fluctuations of the embedding coordinates through the following shift

$$\bar{h}_{\mu\nu} \mapsto \bar{h}_{\mu\nu} + \mathcal{D}_\mu x_\nu + \mathcal{D}_\nu x_\mu - \gamma_{\mu\nu} \mathcal{D}_\beta x^\beta - 2y_i K_{\mu\nu}^i. \quad (3.36)$$

We find a quadratic algebraic action for $\bar{h}_{\mu\nu}$ and an additional term leading to the bosonic Lagrangian

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} \sqrt{\gamma} \gamma^{\alpha\beta} \gamma^{\rho\sigma} \bar{h}_{\alpha\rho} \bar{h}_{\beta\sigma} + \sqrt{\gamma} [\gamma^{\alpha\beta} \mathcal{D}_\alpha y^i \mathcal{D}_\beta y_i - (m_{ij} + 2K_{i\mu\nu} K_j^{\mu\nu}) y^i y^j] + \\ & + 2\sqrt{\gamma} \gamma^{\alpha\beta} (\mathcal{D}_\alpha x^a K_{i,a\beta} y^i + \mathcal{D}_\alpha y^i x^a K_{i,a\beta} + \gamma^{\alpha\beta} \mathcal{D}_\alpha K_{i,a\beta} x^a y^i), \end{aligned} \quad (3.37)$$

where the last line is a total divergence and can be neglected. The Lagrangian for the transverse fluctuations is again given by (3.35). We notice that in eq. (3.37) the longitudinal fluctuations have completely disappeared from the Lagrangian. This second way to proceed is similar to a sort of *static gauge fixing* (where longitudinal fluctuations are taken to vanish). More precisely, in the present approach the longitudinal degrees of freedom of the metric decouple, and their role in the Lagrangian is taken by the traceless part of the metric fluctuation, which however possesses an algebraic gaussian action. If we integrate it out, we find an ultra-local functional determinant, whose careful regularisation and evaluation hides some of the subtleties discussed below eq. (3.29).

Another option to get rid of the longitudinal fluctuation is to choose the static gauge fixing $x^a = 0$. Then (3.21) reduces to a Lagrangian for the transverse fluctuation only, with a mass matrix given not by \mathcal{M}_{ij} but by m_{ij} . However we must recall that the transverse fluctuation are still coupled to the metric fluctuations (see eq. (3.24)):

$$S_{\bar{h}} = \int d^2\sigma e^\varphi \sqrt{\gamma} \left[\frac{1}{2} \gamma^{\sigma\mu} \gamma^{\rho\nu} \bar{h}_{\mu\nu} \bar{h}_{\rho\sigma} + 2y_i K^{i\mu\nu} \bar{h}_{\mu\nu} \right]. \quad (3.38)$$

If we again eliminate the metric fluctuation $\bar{h}_{\mu\nu}$ through its equation of motion, we get back to the mass matrix \mathcal{M}_{ij} and to $\mathcal{L}_{\text{transv}}$ in (3.35).

We remark finally that starting from the Nambu-Goto action, where no dynamical worldsheet metric is present, in the static gauge $x^a = 0$ we would directly obtain $\mathcal{L}_{\text{transv}}$.

In view of the analysis above, we will focus our attention on transverse fluctuations and will examine closely the structure of the Lagrangian in (3.35) which governs their dynamics. These modes are in general coupled between themselves and we can distinguish two different sources of coupling: the $\text{SO}(D-2)$ gauge connection $A^i_{j\alpha}$ induced on the normal bundle and the mass matrix \mathcal{M}_{ij} . In the following we shall discuss some general properties of these two objects.

3.2 The normal bundle

Let us begin to discuss the geometric structure of the normal bundle. The curvature $F^i_{j\alpha\beta} \equiv \partial_\alpha A^i_{j\beta} - \partial_\beta A^i_{j\alpha} + A^i_{\ell\alpha} A^\ell_{j\beta} - A^i_{\ell\beta} A^\ell_{j\alpha}$ can be easily evaluated in terms of the Riemann curvature of the target space and of the extrinsic curvature through the *Ricci equation*:

$$F^i_{j\alpha\beta} = -R_{ABMN} t_\alpha^A t_\beta^B N^{iM} N_j^N - h^{\rho\sigma} (K_{\rho\alpha}^i K_{j\sigma\beta} - K_{\rho\beta}^i K_{j\sigma\alpha}). \quad (3.39)$$

Focussing on the case of $\text{AdS}^5 \times \text{S}^5$, we see that the contribution of the Riemann tensor to the curvature of the normal bundle vanishes identically as in flat space. In fact the Riemann tensor for this background is given by

$$R_{AB,CD} = -(\hat{P}_{AC}\hat{P}_{BD} - \hat{P}_{AD}\hat{P}_{BC}) + (\bar{P}_{AC}\bar{P}_{BD} - \bar{P}_{AD}\bar{P}_{BC}), \quad (3.40)$$

where \hat{P}_{AB} is the projector on AdS_5 and \bar{P}_{AB} is the on S^5 . Obviously the sum of the two projectors gives the identity:

$$\hat{P}_{AB} + \bar{P}_{AB} = \eta_{AB}. \quad (3.41)$$

The contractions $R_{ABMN} t_\alpha^A t_\beta^B N^{iM} N_j^N$ can be written as

$$R_{ABMN} t_\alpha^A t_\beta^B N^{iM} N_j^N = [(\bar{t}_\alpha \cdot \bar{N}^i)(\bar{t}_\beta \cdot \bar{N}_j) - (\bar{t}_\beta \cdot \bar{N}^i)(\bar{t}_\alpha \cdot \bar{N}_j)] - [(\hat{t}_\alpha \cdot \hat{N}^i)(\hat{t}_\beta \cdot \hat{N}_j) - (\hat{t}_\beta \cdot \hat{N}^i)(\hat{t}_\alpha \cdot \hat{N}_j)], \quad (3.42)$$

where we introduced the inner product $(a \cdot b) \equiv \eta_{CD} a^C b^D$ as a product over flat spacetime indices. Above, the hats and the bars over the vectors denote the projection of the vectors on AdS_5 and S^5 respectively, namely

$$\hat{V}^A = \hat{P}^A_B V^B \quad \text{and} \quad \bar{V}^A = \bar{P}^A_B V^B. \quad (3.43)$$

We can now use that

$$(\bar{t}_\alpha \cdot \bar{N}^i) = (\bar{t}_\alpha \cdot N^i) = ((t_\alpha - \hat{t}_\alpha) \cdot N^i) = -(\hat{t}_\alpha \cdot N^i) = -(\hat{t}_\alpha \cdot \hat{N}^i), \quad (3.44)$$

and we remain with the following expression for the normal curvature

$$F^i_{j\alpha\beta} = -\gamma^{\rho\sigma} (K_{\rho\alpha}^i K_{j\sigma\beta} - K_{\rho\beta}^i K_{j\sigma\alpha}) = (\gamma^{\rho\sigma} \epsilon^{\mu\nu} K_{\rho\mu}^i K_{j\sigma\nu}) \epsilon_{\alpha\beta} \equiv \sqrt{\gamma} \epsilon_{\alpha\beta} \mathcal{F}^i_j \quad (3.45)$$

which also holds in flat space. The normal bundle is *flat* when $F^i_{j,\alpha\beta} = 0$. In that case, we can always choose the normal vectors N^i so that $A^i_{j\alpha} = 0$ and the covariant derivative acting on the transverse fluctuations reduces to the usual one. This occurs, for instance, when the minimal surface is confined in a three-dimensional subspace of our target space: the extrinsic curvature is in fact not vanishing just in one normal direction [9, 60–65].

For a generic worldsheet, which solves the equation of motion in the $\text{AdS}_5 \times \text{S}^5$ background, the extrinsic curvature defines, at most, two independent vector fields normal to the worldsheet⁷.

⁷This follows from the fact that we have just two independent components of the extrinsic curvature (e.g. K_{11}^ℓ and K_{12}^ℓ). Alternatively, we can argue this result, in a covariant way, from the following matrix relation satisfied by \mathcal{F} :

$$\mathcal{F}^4 = \frac{1}{2} \text{Tr}(\mathcal{F}^2) \mathcal{F}^2.$$

In fact the matrix \mathcal{F}^i_j can be always put in the form

$$\mathcal{F}^i_j = (k^i h_j - k_j h^i), \quad (3.46)$$

where $(h^i \cdot t_\alpha) = (k^i \cdot t_\alpha) = 0$ and we can choose $(k \cdot h) = 0$ without loss of generality.

3.3 Mass matrix and sum rules

The next step is to examine more closely the mass matrix (3.35) for the transverse bosonic degrees of freedom. For a generic classical background it has the form

$$\mathcal{M}_{ij} = R_{AM,BN} t_c^{cM} t_c^N N_i^A N_j^B + K_{i,\alpha\beta} K_j^{\alpha\beta}. \quad (3.47)$$

There are few general properties of \mathcal{M}_{ij} that can be easily read from (3.47), since the embedding equations for a sub-manifold do not provide a direct constraint on the contraction $R_{AM,BN} t_c^{cM} t_c^N N_i^A N_j^B$. However, its trace $\text{Tr}(\mathcal{M})$ admits a quite simple and compact expression in terms of geometric quantities. If we use the completeness relation

$$N_i^A N^{iB} = \eta^{AB} - t_c^A t^{cB}, \quad (3.48)$$

we can rewrite the trace of (3.47) as follows

$$\text{Tr}(\mathcal{M}) = R_{MN} t_c^{cM} t_c^N - R_{AM,BN} t_c^{cM} t_c^N t_d^A t^{dB} + \text{Tr}(K^2), \quad (3.49)$$

where

$$\text{Tr}(K^2) = \gamma^{\alpha\beta} \gamma^{\rho\sigma} \eta_{AB} K_{\rho\beta}^A K_{\sigma\alpha}^B. \quad (3.50)$$

By mean of the Gauss equation (3.19) we can reduce the second term in (3.49) to

$$R_{AM,BN} t_c^{cM} t_c^N t_d^A t^{dB} = {}^{(2)}R + \text{Tr}(K^2), \quad (3.51)$$

where ${}^{(2)}R$ is the two-dimensional (intrinsic) scalar curvature. This leads to the sum rule

$$\text{Tr}(\mathcal{M}) = R_{MN} t_c^{cM} t_c^N - {}^{(2)}R \quad (3.52)$$

in terms of the Ricci tensor R_{MN} and of the two-dimensional curvature. In the case of Einstein spaces (where $R_{MN} = n G_{MN}$) this gives $\text{Tr}(\mathcal{M}) = 2n - {}^{(2)}R$.

We now particularize the analysis of the mass matrix (3.47) for a string moving in $\text{AdS}_5 \times \text{S}^5$ background. The Riemann tensor for the background $\text{AdS}_5 \times \text{S}^5$ is given in term of projectors by (3.40). The first contribution to the mass matrix takes the form

$$\begin{aligned} \gamma^{\alpha\beta} R_{ARBS} t_\alpha^A t_\beta^B N_i^R N_j^S &= -\gamma^{\rho\sigma} (\hat{t}_\rho \cdot \hat{t}_\sigma) (\hat{N}_i \cdot \hat{N}_j) + \gamma^{\alpha\beta} (\hat{N}_i \cdot \hat{t}_\alpha) (\hat{N}_j \cdot \hat{t}_\beta) + \\ &+ \gamma^{\rho\sigma} (\bar{t}_\rho \cdot \bar{t}_\sigma) (\bar{N}_i \cdot \bar{N}_j) - \gamma^{\alpha\beta} (\bar{N}_i \cdot \bar{t}_\alpha) (\bar{N}_j \cdot \bar{t}_\beta), \end{aligned} \quad (3.53)$$

where the hats and the bars over the vectors again denote the projection of the vectors on AdS_5 and S^5 respectively and they are defined in (3.43). The relation (3.44) easily shows that the mixed terms in (3.54) cancel and we obtain

$$\gamma^{\alpha\beta} R_{ARBS} t_\alpha^A t_\beta^B N_i^R N_j^S = -\gamma^{\rho\sigma} (\hat{t}_\rho \cdot \hat{t}_\sigma) (\hat{N}_i \cdot \hat{N}_j) + \gamma^{\rho\sigma} (\bar{t}_\rho \cdot \bar{t}_\sigma) (\bar{N}_i \cdot \bar{N}_j). \quad (3.54)$$

With a small abuse of language we shall introduce the following notations

$$m_{AdS_5}^2 \equiv \gamma^{\rho\sigma}(\hat{t}_\rho \cdot \hat{t}_\sigma) \quad \text{and} \quad m_{S^5}^2 \equiv -\gamma^{\rho\sigma}(\bar{t}_\rho \cdot \bar{t}_\sigma). \quad (3.55)$$

Then the complete mass matrix is given by

$$\mathcal{M}_{ij} = -m_{AdS_5}^2(\hat{N}_i \cdot \hat{N}_j) - m_{S^5}^2(\bar{N}_i \cdot \bar{N}_j) + K_{i\alpha\beta}K_j^{\alpha\beta}. \quad (3.56)$$

It is worth noticing that the two scalar quantities $m_{AdS_5}^2$ and $m_{S^5}^2$ can be computed in terms of the two-dimensional scalar curvature ${}^{(2)}R$ and $\text{Tr}(K^2)$. We first observe that

$$m_{AdS_5}^2 - m_{S^5}^2 = \gamma^{\rho\sigma}(\bar{t}_\rho \cdot \bar{t}_\sigma) + \gamma^{\rho\sigma}(\hat{t}_\rho \cdot \hat{t}_\sigma) = \gamma^{\rho\sigma}(t_\rho \cdot t_\sigma) = 2, \quad (3.57)$$

while another relation between these two quantities can be obtained from Gauss equation (3.19) in its contracted form. After some simple manipulations we end up with

$$\begin{aligned} {}^{(2)}R + \text{Tr}(K^2) &= \gamma^{\alpha\beta}\gamma^{\rho\sigma}R_{ACBD}t_\alpha^A t_\rho^C t_\beta^B t_\sigma^D = m_{S^5}^4 - m_{AdS_5}^4 + (\hat{t}_c \cdot \hat{t}^d)(\hat{t}_d \cdot \hat{t}^c) - (\bar{t}_c \cdot \bar{t}^d)(\bar{t}_d \cdot \bar{t}^c) = \\ &= -2(m_{S^5}^2 + 1). \end{aligned} \quad (3.58)$$

We finally obtain:

$$m_{AdS_5}^2 = -\frac{1}{2}({}^{(2)}R + \text{Tr}(K^2)) + 1 \quad \text{and} \quad m_{S^5}^2 = -\frac{1}{2}({}^{(2)}R + \text{Tr}(K^2)) - 1. \quad (3.59)$$

The results (3.59) simplify the explicit evaluation of the trace of the mass matrix $\text{Tr}(\mathcal{M})$ for the case of $\text{AdS}_5 \times S^5$. Since $R_{MN} = -4\hat{P}_{MN} + 4\bar{P}_{MN}$, we find

$$\begin{aligned} \text{Tr}(\mathcal{M}) &= R_{MN}t^{cM}t_c^N - {}^{(2)}R = -4(m_{S^5}^2 + m_{AdS_5}^2) - {}^{(2)}R = 4({}^{(2)}R + \text{Tr}(K^2)) - {}^{(2)}R = \\ &= 3{}^{(2)}R + 4\text{Tr}(K^2). \end{aligned} \quad (3.60)$$

The structure of the mass matrix (3.56) can be further constrained assuming particular properties for the classical background. The simplest geometrical configuration is when in (3.56) either $\hat{t}_c = 0$ or $\bar{t}_c = 0$ ($c = 1, 2$). In that case the minimal surface Σ is confined in one of the two spaces: AdS_5 if $\bar{t}_c = 0$ or S^5 if $\hat{t}_c = 0$. Let us focus on the first possibility; the second one can be discussed in complete analogy. The mass matrix then reduces to

$$\mathcal{M}_{ij} = -2(\hat{N}_i \cdot \hat{N}_j) + K_{i\alpha\beta}K_j^{\alpha\beta}, \quad (3.61)$$

where we have used that $m_{S^5}^2 = (\bar{t}_c \cdot \bar{t}^c) = 0$ and $m_{AdS_5}^2 = m_{AdS_5}^2 - m_{S^5}^2 = 2$. The extrinsic curvature $K_{i\alpha\beta}$ is different from zero only for orthogonal directions lying in AdS_5 . Therefore we have 5 massless scalar ($\mathbf{m}_i = m_{S^5} = 0$ [$i = 1, \dots, 5$]), one for each direction of S^5 . We can always choose a sixth direction (lying in AdS_5) orthogonal to Σ and to the two normal directions defined by $K_{i\alpha\beta}$. The mass \mathbf{m}_6 of this sixth scalar is

$$\mathbf{m}_6^2 = -2. \quad (3.62)$$

Finally, we have to select the last two orthogonal directions ($i=7,8$) and we choose the only two orthonormal eigenvectors of $K_{i\alpha\beta}K_j^{\alpha\beta}$ with non vanishing eigenvalues. They always exist if the normal bundle is not flat. Then the two masses are given by

$$\mathbf{m}_7^2 = \lambda_1 - 2 \quad \text{and} \quad \mathbf{m}_8^2 = \lambda_2 - 2. \quad (3.63)$$

Here λ_1 and λ_2 are the two non-vanishing eigenvalues of $K_{i\alpha\beta}K_j^{\alpha\beta}$ and they are determined in terms of the geometric quantity of the surface through the relations:

$$\lambda_1 + \lambda_2 = \text{Tr}(K^2) = -{}^{(2)}R - 2 \quad \text{and} \quad \lambda_1\lambda_2 = \frac{1}{2}[(\text{Tr}(K^2))^2 - \text{Tr}(K^4)] = \frac{1}{2}\text{Tr}(\mathcal{F}^2), \quad (3.64)$$

where $\text{Tr}(K^4) = K_{\alpha\beta}^i K_j^{\alpha\beta} K_{\rho\sigma}^j K_i^{\rho\sigma}$. If the normal bundle is flat, $\mathcal{F} = 0$ and one of the two eigenvalues vanishes, *e.g.* $\lambda_1 = 0$. Then the two masses collapse to the known result [9, 18]

$$\mathbf{m}_7^2 = -2 \quad \text{and} \quad \mathbf{m}_8^2 = -{}^{(2)}R - 4. \quad (3.65)$$

Let us turn our attention to the general case where the worldsheet extends both in AdS_5 and S^5 spaces and the mass matrix has the general form (3.56). The first step is to choose two of the fluctuations ($i = 7, 8$) along the two orthogonal directions (h and k) with non-vanishing extrinsic curvature. These two directions are defined up to a rotation in (h, k) -plane. We fix this freedom by choosing h and k to be the only two orthonormal eigenvectors of $K_{i\alpha\beta}K_j^{\alpha\beta}$ with non vanishing eigenvalues. Then the only non vanishing component of the field strength in the normal bundle is \mathcal{F}^7_8 as discussed in subsec. 3.2.

The bosonic masses can be analysed in details if the field strength \mathcal{F}^7_8 is essentially *abelian*, namely if the only component of the connection different from zero is given by A^7_8 . In this case the Codazzi-Mainardi equation (3.20) implies for the normal directions $i = 1, \dots, 6$

$$\gamma_{\beta\gamma}(\bar{t}_\alpha \cdot N_i) - \gamma_{\alpha\gamma}(\bar{t}_\beta \cdot N_i) = \mathcal{D}_\alpha K_{\beta\gamma}^i - \mathcal{D}_\beta K_{\alpha\gamma}^i = 0, \quad (3.66)$$

which immediately translates in

$$(\bar{t}_\beta \cdot N_i) = (\hat{t}_\beta \cdot N_i) = 0 \quad \text{for } i \neq 7, 8. \quad (3.67)$$

We find that the remaining six normal directions are orthogonal both to \bar{t}_α and to \hat{t}_α , implying that some of these vectors completely lie in AdS_5 , while the others in S^5 . Generically we expect to find three of them in AdS_5 and three in S^5 (a different partition of the six vectors between the two subspaces may occur when some of \hat{t}_α or of \bar{t}_α vanishes). Because of the orthogonality relations (3.67), the 8×8 mass matrix \mathcal{M} takes the form

$$\mathcal{M} = \begin{pmatrix} -m_{\text{AdS}_5}^2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -m_{\text{AdS}_5}^2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -m_{\text{AdS}_5}^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -m_{S^5}^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -m_{S^5}^2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -m_{S^5}^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & m_{77} & m_{78} \\ 0 & 0 & 0 & 0 & 0 & 0 & m_{87} & m_{88} \end{pmatrix} \quad (3.68)$$

where $m_{\text{AdS}_5}^2$ and $m_{S^5}^2$ are given in (3.59). The trace condition (3.52) still constrains m_{77} and m_{88} and it yields that

$$m_{77} + m_{88} = \text{Tr}(K^2). \quad (3.69)$$

With the help of (3.69), the reduced mass matrix in the directions 7 and 8 can be cast into to the form

$$\mathcal{M}_{\text{red.}} \equiv \begin{pmatrix} m_{77} & m_{78} \\ m_{87} & m_{88} \end{pmatrix} = \begin{pmatrix} -m_{AdS_5}^2 + 2(\bar{N}_7 \cdot \bar{N}_7) + \lambda_1 & 2(\bar{N}_7 \cdot \bar{N}_8) \\ 2(\bar{N}_7 \cdot \bar{N}_8) & m_{AdS_5}^2 - 2(\bar{N}_7 \cdot \bar{N}_7) + \lambda_2 \end{pmatrix} \quad (3.70)$$

where λ_1 and λ_2 again obey (3.64).

In the general case, when \mathcal{F}^7_8 is not generated by only taking A^7_8 different from zero, the structure of the mass matrix may become more intricate. However we can always choose, at least, two orthogonal directions, one in S^5 and one in AdS_5 , which are orthogonal to the minimal surface Σ and to the extrinsic curvature. The masses of these two fluctuations are then given by (3.59).

4 Fermionic fluctuations

The full covariant GS string action in $AdS_5 \times S^5$ has a complicated non-linear structure [38, 66], but to analyze the relevant fermionic contributions it is sufficient to consider only its quadratic part here

$$L_{2F} = i \left(\sqrt{\gamma} \gamma^{\alpha\beta} \delta^{IJ} - \varepsilon^{\alpha\beta} s^{IJ} \right) \bar{\theta}^I \rho_\alpha D_\beta^{JK} \theta^K. \quad (4.1)$$

Above, θ^I ($I = 1, 2$) are two ten-dimensional Majorana-Weyl spinors with the same chirality, $s^{IJ} = \text{diag}(1, -1)$, ρ_α are the worldsheet projections of the ten-dimensional Dirac matrices

$$\rho_\alpha = E_{Am} \partial_\alpha X^m \Gamma^A, \quad (4.2)$$

and D_α^{JK} is the two-dimensional pullback $\partial_\alpha X^m D_m^{JK}$ of the ten-dimensional covariant derivative (here, flat and curved indices span the range from 1 to $D = 10$), sum of an ordinary spinor covariant derivative and an additional ‘‘Pauli-like’’ coupling to the RR flux background, $D_m^{JK} = \mathfrak{D}_m \delta^{JK} - \frac{1}{8 \cdot 5!} F_{m_1 \dots m_5} \Gamma^{m_1 \dots m_5} \Gamma_m \epsilon^{JK}$.

In the $AdS_5 \times S^5$ case, it can be written as follows

$$D_\beta^{JK} \theta^K = \mathfrak{D}_\beta^{JK} \theta^K + \mathcal{F}_\beta^{JK} \theta^K, \quad \mathfrak{D}_\beta^{JK} = \delta^{JK} \left(\partial_\beta + \frac{1}{4} \partial_\beta X^m \Omega_m^{AB} \Gamma_{AB} \right) \theta^K, \quad (4.3)$$

where the flux term \mathcal{F}_β^{JK} , responsible for the fermionic ‘‘masses’’, is ⁸

$$\mathcal{F}_\beta^{JK} = -\frac{i}{2} \epsilon^{JK} \Gamma_\star \rho_\beta, \quad \Gamma_\star = i \Gamma_{01234}. \quad (4.6)$$

⁸ An alternative form for the flux [38] is

$$\mathcal{F}_\beta^{JK} = -\frac{i}{2} \epsilon^{JK} \tilde{\rho}_\beta, \quad \tilde{\rho}_\beta = \Gamma_A \hat{t}_\beta^A + i \Gamma_A \bar{t}_\beta^A \quad (4.4)$$

with which the corresponding part of the gauge-fixed Lagrangian reads

$$L_F^{flux} = -\epsilon^{\alpha\beta} \bar{\theta} \rho_\alpha \tilde{\rho}_\beta \theta. \quad (4.5)$$

Its equivalence with (4.6) and (4.24) below is manifest in the 5 + 5 basis of [38], see also discussion in [9].

Looking for a general formalism for fluctuations, there is a natural choice for the κ -symmetry gauge-fixing that is viable in type IIB string action, where both Majorana-Weyl fermions in the GS action have the same chirality, namely ⁹

$$\theta^1 = \theta^2 \equiv \theta . \quad (4.7)$$

Since $s^{11} = -s^{22} = 1$, in the kinetic part of the gauge-fixed fermionic action only the term proportional to $\gamma^{\alpha\beta}$ will survive after the κ -symmetry gauge-fixing, while in the flux part only the term proportional to $\varepsilon^{\alpha\beta}$.

4.1 The fermionic kinetic term

Let us first focus on the reduction of the ordinary spinor covariant derivative \mathfrak{D}_β in (4.3) and write the relevant part of the κ -symmetry gauge-fixed Lagrangian as

$$L_F^{\text{kin}} = 2i \sqrt{\gamma} \gamma^{\alpha\beta} \bar{\theta} \rho_\alpha (\partial_\beta + \Omega_\beta) \theta, \quad \Omega_\beta = \frac{1}{4} \Omega_\beta^{AB} \Gamma_{AB} . \quad (4.8)$$

The geometrical interpretation of the reduction procedure, already discussed for the bosonic fluctuations, allows us to guess the final result of this section. In fact a straightforward, but tedious computation must yield, at the end, the following form for the kinetic term:

$$\tilde{L}_F^{\text{kin}} = 2i \sqrt{\gamma} \gamma^{\alpha\beta} \bar{\Theta} \Gamma^a e_{a\alpha} \mathcal{D}_\beta \Theta \equiv 2i \sqrt{\gamma} \gamma^{\alpha\beta} \bar{\Theta} \Gamma^a e_{a\alpha} \left(\partial_\beta + \frac{1}{4} \Gamma_{bc} \omega_\beta^{bc} - \frac{1}{4} A_\beta^{ij} \Gamma_{ij} \right) \Theta . \quad (4.9)$$

Above, \mathcal{D}_β is the two-dimensional covariant derivative acting on spinors, and it takes into account that Θ is now a two-dimensional spinor, which also transforms in the spinor representation of the $\text{SO}(8)$ normal bundle. We remind that e_α^a denotes the worldsheet zweibein. The connection A_β^{ij} is defined in (3.18). This pattern is completely analogous to the one obeyed by the normal bosonic fluctuations, which are scalars for the worldsheet, but vectors for the normal bundle.

We shall now see how the result (4.8) emerges from the explicit analysis. Although we will work explicitly in $D = 10$, we remark that the reduction of the canonical covariant derivative (passing from (4.8) to (4.9)) is independent on the dimensionality of the spacetime, exactly as in the bosonic case ¹⁰. The Dirac algebra is naturally decomposed in two subsets: the components along the worldsheet and those orthogonal to the worldsheet, which in the ten-dimensional case means

$$\rho_\alpha = t_{A\alpha} \Gamma^A, \quad \alpha = 1, 2, \quad (4.10)$$

$$\rho_{\underline{i}} = N_{\underline{i}}^A \Gamma^A, \quad \underline{i} = 1, \dots, 8. \quad (4.11)$$

⁹A widely used alternative to (4.7) – especially in the context of AdS/CFT – is the light-cone gauge-fixing $\Gamma^+ \theta^I = 0$, where the light-cone might lie entirely in S^5 [67, 68] or being shared between AdS_5 and S^5 [69] (see [47] and further references therein). One of the obvious advantages of the “covariant” gauge-fixing (4.7) is preservation of global bosonic symmetries of the action. A more general choice is $\theta_1 = k \theta_2$, $\theta_2 \equiv \theta$ where k is a real parameter whose dependence is expected to cancel in the effective action, see discussion in [70]. Yet another κ -symmetry gauge-fixing, albeit equivalent to (4.7) [9], has been used for studying stringy fluctuations in $\text{AdS}_5 \times S^5$ in [25].

¹⁰The dimensionality of space-time will clearly influence the kind of spinors involved.

As used earlier in [60–64] and made explicit in this context in [9], since a two-dimensional Clifford algebra holds by construction for the $\{\rho_\alpha, \rho_\beta\} = 2\gamma_{\alpha\beta}$, it is always possible to find a *local* $\text{SO}(1,9)$ rotation S that transforms ρ_i into two-dimensional Dirac matrices contracted with zweibein

$$\begin{aligned}\rho_\alpha &= S\Gamma^a S^{-1} e_{a\alpha} & a = 1, 2, \\ \rho_{\underline{i}} &= S\Gamma^{\underline{i}+2} S^{-1} & \underline{i} = 1, \dots, 8,\end{aligned}\tag{4.12}$$

where

$$\Gamma^1 = i\tau_2 \otimes \mathbf{1}_{16}, \quad \Gamma^2 = \tau_1 \otimes \mathbf{1}_{16}, \quad \Gamma^{\underline{i}} = \tau_3 \otimes \Sigma^{\underline{i}},\tag{4.13}$$

τ_a are Pauli matrices and $\Sigma^{\underline{i}}$ are (16-dimensional) Dirac matrices in 8 Euclidean dimensions.

Defining now

$$\Theta = S^{-1}\theta \quad \text{and} \quad \hat{\Omega}_\alpha = S^{-1}\Omega_\alpha S + S^{-1}\partial_\alpha S,\tag{4.14}$$

one ends with the following rotated expression

$$\tilde{L}_F^{kin} = 2i\sqrt{\gamma}\gamma^{\alpha\beta} \bar{\Theta}\Gamma^a e_{a\alpha} (\partial_\beta + \hat{\Omega}_\beta) \Theta.\tag{4.15}$$

We remark here that the present analysis is only valid at classical level: as we will see later the local rotation S produces quantum mechanically a non-trivial Jacobian in the path-integral measure, whose contribution is crucial to recover the correct structure of the divergent terms. To evaluate $\hat{\Omega}_\alpha$, we begin expanding

$$\Omega_\beta = S\hat{\Omega}_\beta S^{-1} - \partial_\beta S S^{-1}\tag{4.16}$$

in the basis provided by the Dirac algebra (4.12)

$$\frac{1}{4}\Omega_\beta^{AB}\Gamma_{AB} = \frac{1}{4}(\hat{\Omega}_\beta^{ab} e_a^\mu e_b^\nu \rho_{\mu\nu} + 2\hat{\Omega}_\beta^{a\underline{i}} e_a^\mu \rho_{\mu\underline{i}} + \hat{\Omega}_\beta^{\underline{i}\underline{j}} \rho_{\underline{i}\underline{j}}) + \frac{1}{4}(\rho^\alpha \partial_\beta \rho_\alpha + \rho^{\underline{i}} \partial_\beta \rho_{\underline{i}} - \rho^\alpha \rho^\mu e_\mu^a \partial_\beta e_{a\alpha}).\tag{4.17}$$

The expression for $\partial_\alpha S S^{-1}$ in the second parenthesis above has been derived considering the derivatives of (4.12) with respect to worldsheet coordinates which in turn implies

$$\rho^\alpha \partial_\beta \rho_\alpha + \rho^{\underline{i}} \partial_\beta \rho_{\underline{i}} = -4\partial_\beta S S^{-1} + \rho^\alpha \rho^\mu e_\mu^a \partial_\beta e_{a\alpha}.\tag{4.18}$$

Let us now formally expand all the contributions in the same basis (4.10)

$$\Omega_\beta^{AB}\Gamma_{AB} = \Omega_\beta^{AB} t_A^\mu t_B^\nu \rho_{\mu\nu} + 2\Omega_\beta^{AB} t_A^\mu N_B^{\underline{i}} \rho_{\mu\underline{i}} + \Omega_\beta^{AB} N_A^{\underline{i}} N_B^{\underline{j}} \rho_{\underline{i}\underline{j}}\tag{4.19}$$

in which

$$\begin{aligned}\rho^\alpha \partial_\beta \rho_\alpha - \rho^{\alpha\mu} e_\mu^a \partial_\beta e_{a\alpha} &= \rho^{\alpha\underline{i}} N_{\underline{i}}^A K_{A\alpha\beta} - \rho^{\alpha\mu} e_{a\alpha} e_{b\mu} \omega_\beta^{ab} + \rho^{\alpha\mu} \Omega_\beta^{AB} t_{A\alpha} t_{B\mu} + \rho^{\alpha\underline{i}} \Omega_\beta^{AB} t_{A\alpha} N_{B\underline{i}} \\ \rho^{\underline{i}} \partial_\beta \rho_{\underline{i}} &= \rho^{\alpha\underline{i}} (N_{\underline{i}A} K_{\alpha\beta}^A + \Omega_\beta^{BA} t_{B\alpha} N_{\underline{i}A}) + \rho^{\underline{i}\underline{j}} (N_{\underline{i}}^A D_\beta N_{\underline{j}A} + \Omega_\beta^{BA} N_{\underline{i}B} N_{A\underline{j}}).\end{aligned}\tag{4.20}$$

Collecting the different results we obtain

$$\hat{\Omega}_\alpha^{ab} e_a^\rho e_b^\sigma = e_a^\rho e_b^\sigma \omega_\alpha^{ab}, \quad \hat{\Omega}_\alpha^{a\underline{i}} e_a^\rho = -N_{\underline{i}A}^a K_\alpha^{A\rho}, \quad \hat{\Omega}_\alpha^{\underline{i}\underline{j}} = -A_{\underline{i}\underline{j}}^\alpha,\tag{4.21}$$

and the fermion kinetic part of the rotated Lagrangian takes the form

$$\begin{aligned}\tilde{L}_F^{kin} &= 2i\sqrt{\gamma}\gamma^{\alpha\beta}\bar{\Theta}\Gamma^a e_{a\alpha}(\partial_\beta + \frac{1}{4}\Gamma_{bc}\omega_\beta^{bc} - \frac{1}{4}N_A^i K_\beta^{Ab}\Gamma_{b\bar{i}} - \frac{1}{4}A_\beta^{ij}\Gamma_{i\bar{j}})\Theta \\ &\equiv 2i\sqrt{\gamma}\gamma^{\alpha\beta}\bar{\Theta}\Gamma^a e_{a\alpha}(\partial_\beta + \frac{1}{4}\Gamma_{bc}\omega_\beta^{bc} - \frac{1}{4}A_\beta^{ij}\Gamma_{i\bar{j}})\Theta.\end{aligned}\quad (4.22)$$

Namely, in the rotated basis (4.12)-(4.13) the GS kinetic operator (4.8) results in a standard two-dimensional Dirac fermion action on a curved two-dimensional background with geometry defined by the induced metric [9]. The spinor covariant derivative can be written as a two-dimensional, ordinary, spinor covariant derivative plus two additional terms as in the first line above. The first extra-term is proportional to the extrinsic curvature, it mixes tangential and normal components and drops out naturally in the fermionic action – hence the second line in (4.22) – once contracted with $\gamma^{\alpha\beta}\Gamma^a e_{a\beta}$ and $\varepsilon^{\alpha\beta}\Gamma^a e_{a\beta}$. In fact, using the equations of motion (2.6) and because of the symmetry of $K_{\alpha\beta}^m$ in α, β , it holds

$$\begin{aligned}K^{Aab}\Gamma_a\Gamma_{b\bar{i}} &= K^{Aab}(\Gamma_{ab\bar{i}} + \delta_{ab}\Gamma_{\bar{i}}) = 0, \\ \varepsilon^{\alpha\beta}e^a{}_\alpha K_\beta^{Ab}\Gamma_a\Gamma_{b\bar{i}} &= \varepsilon^{\alpha\beta}e^a{}_\alpha K_\beta^{Ab}(i\varepsilon_{ab}\tau_3 + \delta_{ab})\Gamma_{\bar{i}} = (i\tau_3 \det(e)e_b^\beta K_\beta^{Ab} + \varepsilon^{\alpha\beta}K_{\alpha\beta}^A)\Gamma_{\bar{i}} = 0.\end{aligned}\quad (4.23)$$

The second additional term consists of an extra, normal bundle two-dimensional gauge connection A_α^{ij} , with respect to which the 16 two-dimensional spinors making up the 32-component MW spinor Θ transform in the spinorial representation of $SO(8)$. This interacting term vanishes (*i.e.* a choice of the N^i exists such that $A_\beta^{ij} = 0$) when the field-strength associated to the normal connection vanishes, see discussion around (3.46). As mentioned there, this is always the case, for example, for embeddings of the string worldsheet in AdS_3 , where indeed the normal direction is just one and the normal bundle is then trivial. For a more general embedding extending in both $AdS_5 \times S^5$ subspaces, the presence or not of this interaction term has to be checked case by case. This is what we do in Section 5, where in considering several examples also comment on this aspect.

4.2 Fermionic “mass” matrix: the flux term

We now analyze the flux term (4.6) in the fermionic Lagrangian, (4.1)-(4.3) which after the κ -symmetry gauge-fixing $\theta^1 = \theta^2 \equiv \theta$ reads

$$L_F^{flux} = -\varepsilon^{\alpha\beta}\bar{\theta}\rho_\alpha\Gamma_*\rho_\beta\theta, \quad \Gamma_* = i\Gamma_{01234}.\quad (4.24)$$

In order to understand the geometrical meaning of the terms in (4.24), we again decompose the Gamma matrices in the orthonormal basis formed by the tangent and transverse vectors. Remembering that Γ_* contains only ten-dimensional flat indices belonging to AdS_5 , we have

$$\begin{aligned}\varepsilon^{\alpha\beta}\rho_\alpha\Gamma_*\rho_\beta &= \varepsilon^{\alpha\beta}\Gamma_*(\Gamma_A\hat{t}_\alpha^A - \Gamma_A\bar{t}_\alpha^A)\rho_\beta \\ &= \varepsilon^{\alpha\beta}\Gamma_*\left(\Gamma_B(\gamma^{\lambda\delta}t_\lambda^A t_\delta^B + N_{\bar{i}}^A N_{\bar{i}}^B)\hat{t}_{\alpha A} - \Gamma_B(\gamma^{\lambda\delta}t_\lambda^A t_\delta^B + N_{\bar{i}}^A N_{\bar{i}}^B)\bar{t}_{\alpha A}\right)\rho_\beta \\ &= \varepsilon^{\alpha\beta}\Gamma_*\left(\gamma^{\lambda\delta}\rho_\lambda\rho_\beta[(\hat{t}_\delta \cdot \hat{t}_\alpha) - (\bar{t}_\delta \cdot \bar{t}_\alpha)] - 2\rho_r\rho_\beta(\bar{N}_r \cdot \bar{t}_\alpha)\right), \quad r = 7, 8\end{aligned}\quad (4.25)$$

where we used the completeness relation $\eta^{AB} = \gamma^{\lambda\delta} t_\lambda^A t_\delta^B + N_{\underline{i}}^A N_{\underline{i}}^B$, the fact that $N_{\underline{i}}^A \bar{t}_\alpha^A$, $N_{\underline{i}}^A \hat{t}_\alpha^A$ are non vanishing only for $\underline{i} = r \equiv 7, 8$ and that $\hat{N}_r^A \hat{t}_\alpha^A = -\bar{N}_r^A \bar{t}_\alpha^A$, $r = 7, 8$ ¹¹. Defining the antisymmetric product of the Gamma matrices projected onto the worldsheet as

$$\rho_3 \equiv \frac{1}{2\sqrt{\gamma}} \varepsilon^{\alpha\beta} \rho_{\alpha\beta}, \quad \rho^{\alpha\beta} = -\frac{1}{\sqrt{\gamma}} \varepsilon^{\alpha\beta} \rho_3, \quad (4.26)$$

one can rearrange the flux term in the following way

$$\begin{aligned} \varepsilon^{\alpha\beta} \rho_\alpha \Gamma_* \rho_\beta &= \varepsilon^{\alpha\beta} \Gamma_* \left[\gamma^{\lambda\delta} (\sqrt{\gamma} \varepsilon_{\lambda\beta} \rho_3 + \gamma_{\lambda\beta}) [(\hat{t}_\delta \cdot \hat{t}_\alpha) - (\bar{t}_\delta \cdot \bar{t}_\alpha)] - 2\rho_r \rho_\beta (\bar{N}^r \cdot \bar{t}_\alpha) \right] \\ &= \Gamma_* \left[\sqrt{\gamma} \rho_3 \gamma^{\alpha\delta} [(\hat{t}_\delta \cdot \hat{t}_\alpha) - (\bar{t}_\delta \cdot \bar{t}_\alpha)] - 2\varepsilon^{\alpha\beta} \rho_{r\beta} (\bar{N}^r \cdot \bar{t}_\alpha) \right] \\ &= \Gamma_* \left[\sqrt{\gamma} \rho_3 (m_{AdS_5}^2 + m_{S^5}^2) + 2\varepsilon^{\alpha\beta} \rho_{r\beta} \nabla_\lambda K_\alpha^{r\lambda} \right], \end{aligned} \quad (4.27)$$

where we used that $\varepsilon^{\alpha\beta} (t_\alpha \cdot t_\beta) = 0$, the definition of masses (3.55), and the Gauss-Codazzi equation (3.20)¹². Hence, after the rotation (4.12) and in terms of the spinors $\Theta = S^{-1}\theta$, the flux part of the fermionic Lagrangian takes the form

$$\tilde{L}_F^{flux} = \bar{\Theta} \tilde{\Gamma}_* \left[\sqrt{\gamma} \tau_3 (m_{AdS_5}^2 + m_{S^5}^2) + 2\varepsilon^{\alpha\beta} e_\beta^a \Gamma_{ar} \nabla_\gamma K_\alpha^{r\gamma} \right] \Theta, \quad r = 7, 8, \quad (4.28)$$

where $\tau_3 = S^{-1} \rho_3 S$, and $\tilde{\Gamma}_* = S^{-1} \Gamma_* S$. In general, the rotated Γ_* is written as

$$\begin{aligned} \tilde{\Gamma}_* &= S^{-1} \Gamma_* S = iS^{-1} (\hat{\varepsilon}^{ABCDE} \Gamma_A \dots \Gamma_E) S \\ &= i\hat{\varepsilon}^{ABCDE} (\hat{t}_A^a \Gamma_a + \hat{N}_A^i \Gamma_i + N_A^r \Gamma_r) \dots (\hat{t}_E^e \Gamma_e + \hat{N}_E^j \Gamma_j + N_E^s \Gamma_s), \end{aligned} \quad (4.29)$$

(where the hat in $\hat{\varepsilon}^{ABCDE}$ is to signal that the A, B, C, D, E take values $0, 1, 2, 3, 4$, as clear from (4.24), $i, j = 1, \dots, 6$, $r, s = 7, 8$) and it can be expanded in the same basis. A clever choice among the basis vectors made possible by the string motion, drastically simplifies the above expression. Due to the self-duality of RR flux, we have written the flux contribution only in terms of Γ_* , and thus we can restrict our discussion and attention to the AdS projection. This is the basis we have in mind here with the aim to simplify the product (4.29).

In the more general case discussed in Section 3.2, in order to compute (4.29), we can use the basis build up by the two tangent vectors projected onto AdS₅ \hat{t}_α ($\alpha = 1, 2$) and the three

¹¹Here we choose a basis for the normal directions $\underline{i} = 1, \dots, 8$ which matches the one used in the bosonic analysis. Namely, fluctuations $\underline{i}=7,8$ are along two orthogonal directions with non-vanishing extrinsic curvature, and for the remaining $i = 1, \dots, 6$ it holds (3.67), with three of them in AdS₅ and three in S⁵.

¹²More precisely, from (3.20), it immediately follows

$$\mathcal{D}_\alpha K_{\beta\gamma}^i - \mathcal{D}_\beta K_{\alpha\gamma}^i = -(\bar{t}_\alpha \cdot t_\gamma)(\bar{t}_\beta \cdot N^i) + (\bar{t}_\alpha \cdot N^i)(\bar{t}_\beta \cdot t_\gamma) + (\hat{t}_\alpha \cdot t_\gamma)(\hat{t}_\beta \cdot N^i) - (\hat{t}_\alpha \cdot N^i)(\hat{t}_\beta \cdot t_\gamma).$$

Since $(\bar{t}_\beta \cdot N_i) = -(\hat{t}_\beta \cdot N_i)$, we find

$$\mathcal{D}_\alpha K_{\beta\gamma}^i - \mathcal{D}_\beta K_{\alpha\gamma}^i = [(\bar{t}_\beta \cdot t_\gamma) + (\hat{t}_\beta \cdot t_\gamma)](\bar{t}_\alpha \cdot N^i) - [(\bar{t}_\alpha \cdot t_\gamma) + (\hat{t}_\alpha \cdot t_\gamma)](\bar{t}_\beta \cdot N^i) = \gamma_{\beta\gamma}(\bar{t}_\alpha \cdot N^i) - \gamma_{\alpha\gamma}(\bar{t}_\beta \cdot N^i).$$

Tracing on α and γ and using the equations of motion (2.6), one obtains

$$\mathcal{D}_\alpha K_\beta^i{}^\alpha = -(\bar{t}_\beta \cdot N^i) = (\hat{t}_\beta \cdot N^i).$$

transverse vectors \hat{N}_i ($i = 1, 2, 3$) which entirely lie in AdS. This set forms an orthonormal basis spanning AdS. However, there might be further contributions also from the remaining transverse directions N_r ($r = 7, 8$) and for this we need to project them in the above basis, that is

$$\hat{N}_A^r = \xi_a^r \hat{t}_A^a, \quad \xi_a^r = (\hat{N}^r \cdot \hat{t}_\alpha) \hat{g}^{\alpha\beta} e_{\beta a} \quad r = 7, 8, \quad (4.30)$$

where $\hat{g}^{\alpha\beta}$ is the inverse matrix obtained from worldsheet metric projected in the sub-space AdS, *i.e.* $\hat{g}_{\alpha\beta} \equiv (\hat{t}_\alpha \cdot \hat{t}_\beta)$, $\hat{g}^{\alpha\beta} \hat{g}_{\beta\delta} = \delta_\delta^\alpha$. Clearly, an analogous expression to (4.30) can be written for the projected vectors onto the compact sub-space, with the obvious replacement from hatted to bar vectors. By means of the relation (4.30) we can massage the product (4.29) to the final expression

$$\tilde{\Gamma}_* = i\hat{\epsilon}^{ABCDE} \hat{N}_C^i \hat{N}_D^j \hat{N}_E^k \Gamma_{ijk} \left[\hat{t}_A^a \hat{t}_B^b \left(\Gamma_{ab} + \xi_a^{[r} \xi_b^{s]} \Gamma_{rs} \right) + \frac{1}{2} \xi_d^r \left(\hat{t}_A^c \hat{t}_B^d - \hat{t}_B^c \hat{t}_A^d \right) \Gamma_{cr} \right]. \quad (4.31)$$

In writing the above expression we are assuming that the inner product (4.30) is not degenerate (as well as the projected worldsheet metric $\hat{g}_{\alpha\beta}$). However, whenever (at least) one of the tangent vectors has a vanishing projection in S^5 , the number of transverse directions completely lying in the compact sub-space increases (of at least one), since now we need (at least) another transverse direction to span S^5 . By a simple counting of degrees of freedom, this decreases (at least) by one the remaining \hat{N} . This implies that one of the transverse direction N^r ($r = 7, 8$) can become linearly independent and lie completely in AdS_5 , that is the inner product (4.30) in one of the directions $r = 7, 8$ could be degenerate. This happens for example in the case of the spinning string motion [26] discussed in Section 5.1. In the limiting case when the motion is completely embedded in $AdS_3 \subset AdS_5$, the tangent vectors belong only to AdS, and both the directions N^r can be chosen such that one will completely lie in AdS and the other one in S^5 . Then, we can pick a basis given by t_α , $\alpha = 1, 2$ (since now $\hat{t}_\alpha = t_\alpha$) and three transverse directions \hat{N}_i ($i=1,2$), N_r with r either 7 or 8. The remaining vectors belong only to the sphere. In this case there is only one term contributing to the expression (4.29), namely the first one in (4.31). We illustrate two examples where this takes place: the folded string [26] at the end of Section 5.1, and the string configuration dual to the circular Wilson loop operator [9] at the end of Section 5.2. If the minimal surface is completely embedded in S^5 then this means that only the transverse directions will contribute to the product (4.29), that is

$$\tilde{\Gamma}_* = S^{-1} \Gamma_* S = \hat{\epsilon}^{ABCDE} \hat{N}_A^r \hat{N}_B^s \hat{N}_C^i \hat{N}_D^j \hat{N}_E^k \Gamma_{rsijk}. \quad (4.32)$$

Finally, we want to stress that the arguments explained above work in a completely symmetric manner by swapping the two sub-spaces AdS and S. Due to the self-duality of the flux term, we could have chosen $\Gamma_* = i\Gamma_{56789}$ and then write $\tilde{\Gamma}_* = i\bar{\epsilon}_{ABCDE} \bar{N}_A^r \bar{N}_B^s \bar{N}_C^i \bar{N}_D^j \bar{N}_E^k \Gamma_{rsijk}$ for an AdS_3 motion, keeping also in mind an overall sign originated in the manipulations in equation (4.27). However, we prefer to keep on working with the flux defined in (4.29) for reader's convenience.

4.3 Comparison with bosonic masses and quantum divergences

In this section we work out a sum rule for the fermion “mass matrix” in analogy to the bosonic one (3.60). In this last case the trace of the mass matrix \mathcal{M} appears naturally in the Seeley-De Witt coefficient $a_B^{(2)}$, controlling the logarithmic divergence of the bosonic fluctuations [71]

$$a_B^{(2)} = \text{Tr}\left(\frac{1}{6} R^{(2)} + \mathcal{M}\right). \quad (4.33)$$

A natural candidate for the traced fermionic mass matrix is therefore the companion term appearing in the fermionic Seeley-De Witt coefficient $a_F^{(2)}$, driving the logarithmic fermionic divergences as well. In the most general case, for a self-adjoint Dirac operator $i\rho^\alpha \mathcal{D}_\alpha - \mathcal{M}_F$ this is given by [71]

$$a_F^{(2)} = \text{Tr}(\mathbb{I})\frac{R^{(2)}}{12} - \frac{1}{2}\text{Tr}(\rho^\alpha \mathcal{M}_F \rho_\alpha \mathcal{M}_F), \quad (4.34)$$

and we are at the moment interested in the second term above. In our case, the fermionic mass matrix is

$$\mathcal{M}_F = \frac{1}{2\sqrt{\gamma}} \varepsilon^{\alpha\beta} \rho_\alpha \Gamma_* \rho_\beta, \quad (4.35)$$

where we re-wrote the fermionic Lagrangian as

$$\mathcal{L}_{2F} = 2i\sqrt{\gamma} \bar{\theta} \left(\gamma^{\alpha\beta} \rho_\alpha \mathcal{D}_\beta + \frac{i}{2\sqrt{\gamma}} \varepsilon^{\alpha\beta} \rho_\alpha \Gamma_* \rho_\beta \right) \theta := 2i\sqrt{\gamma} \bar{\theta} D_F \theta. \quad (4.36)$$

We recall that our fermions are worldsheet scalars and the square root of the determinant of the induced metric is the correct normalization in the fermionic norms [9]. It is important to stress that here γ is the absolute value of the determinant. When ρ_α commutes with \mathcal{M}_F the invariant (4.34) reduces to the more familiar $\text{Tr}(\mathcal{M}_F^2)$, analysed in [9] or in [18] for example. This is the case for the folded spinning solution reviewed in Section 5.1. However, for a general string solution, like the latitude configuration, the two matrices do not commute and (4.34) is an invariant, which leads to the sum rule as we explain below.

Notice that due to the cyclicity of the trace we can compute (4.34) before performing any local $\text{SO}(1,9)$ rotations, and actually in this case it turns out a convenient strategy. Let us start by rewriting (4.34) as

$$\begin{aligned} \text{Tr}(\rho^\alpha \mathcal{M}_F \rho_\alpha \mathcal{M}_F) &= \text{Tr} \left(\frac{\gamma^{\alpha\beta}}{4\gamma} \varepsilon^{\mu\nu} \varepsilon^{\lambda\tau} \rho_\alpha \rho_\mu \Gamma_* \rho_\nu \rho_\beta \rho_\lambda \Gamma_* \rho_\tau \right) \\ &= \text{Tr} \left(\frac{\gamma^{\alpha\beta}}{4\gamma} \varepsilon^{\mu\nu} \varepsilon^{\lambda\tau} \rho_\alpha \rho_\mu \Gamma_* (\gamma_{\nu\beta} \rho_\lambda + \gamma_{\beta\lambda} \rho_\nu - \gamma_{\nu\lambda} \rho_\beta) \Gamma_* \rho_\tau \right) \\ &\equiv I_1 + I_2 + I_3. \end{aligned} \quad (4.37)$$

and compute the three terms above separately. The first contribution I_1 gives

$$\begin{aligned} I_1 &= \frac{1}{4\gamma} \text{Tr} \left(\varepsilon^{\mu\nu} \varepsilon^{\lambda\tau} \rho_\nu \rho_\mu \Gamma_* \rho_\lambda \Gamma_* \rho_\tau \right) \\ &= -\frac{1}{4\gamma} \text{Tr} \left(\varepsilon^{\mu\nu} \rho_\mu \rho_\nu \Gamma_* \Gamma_* \left(\sqrt{\gamma} (m_{AdS_5}^2 + m_{S^5}^2) \rho_3 + 2\varepsilon^{\alpha\beta} (\hat{t}_\alpha \cdot N^r) \rho_r \rho_\beta \right) \right), \end{aligned} \quad (4.38)$$

where we have used that

$$\varepsilon^{\alpha\beta} \rho_\alpha \Gamma_* \rho_\beta = \Gamma_* \left\{ \sqrt{\gamma} (m_{AdS_5}^2 + m_{S^5}^2) \rho_3 + 2 \varepsilon^{\alpha\beta} (\hat{t}_\alpha \cdot N^r) \rho_r \rho_\beta \right\}. \quad (4.39)$$

computed in Section 4.2. By recalling that $\rho_3 = \frac{1}{2\sqrt{\gamma}} \varepsilon^{\alpha\beta} \rho_{\alpha\beta}$, and $\rho^{\alpha\beta} = -\frac{\varepsilon^{\alpha\beta}}{\sqrt{\gamma}} \rho_3$, we can compute the following commutation relation

$$\rho_3 \rho_\beta = \frac{\varepsilon^{\mu\nu}}{\sqrt{\gamma}} \gamma_{\nu\beta} \rho_\mu, \quad (4.40)$$

which in I_1 leads to the result

$$I_1 = -\frac{1}{2} (m_{AdS_5}^2 + m_{S^5}^2) \text{Tr}(\rho_3^2). \quad (4.41)$$

The second term I_2 can be manipulated in a similar manner and we obtain

$$I_2 = \frac{1}{4\gamma} \text{Tr} \left(\gamma^{\alpha\beta} \varepsilon^{\mu\nu} \varepsilon^{\lambda\tau} \rho_\alpha \rho_\mu \Gamma_* \gamma_{\beta\lambda} \rho_\nu \Gamma_* \rho_\tau \right) = -\frac{1}{2} (m_{AdS_5}^2 + m_{S^5}^2) \text{Tr}(\mathbb{I}). \quad (4.42)$$

Finally let us look at the last term I_3

$$\begin{aligned} I_3 &= -\frac{1}{4\gamma} \text{Tr} \left(\gamma^{\alpha\beta} \varepsilon^{\mu\nu} \varepsilon^{\lambda\tau} \gamma_{\nu\lambda} \rho_\alpha \rho_\mu \Gamma_* \rho_\beta \Gamma_* \rho_\tau \right) \\ &= -\frac{1}{4} \text{Tr} \left(\gamma^{\alpha\beta} \Gamma_* \rho_\alpha \Gamma_* \rho_\beta \right) + \frac{1}{4} \text{Tr} \left(\gamma^{\alpha\beta} \gamma^{\mu\nu} \rho_{\mu\alpha} \Gamma_* \rho_\beta \Gamma_* \rho_\nu \right). \end{aligned} \quad (4.43)$$

In order to evaluate I_3 we need also the expression for the symmetrized product of Gamma matrices, that is

$$\gamma^{\alpha\beta} \rho_\alpha \Gamma_* \rho_\beta = \Gamma_* \left[(m_{AdS_5}^2 + m_{S^5}^2) + 2 (\hat{t}_\alpha \cdot N_r) \rho^r \rho^\alpha \right], \quad (4.44)$$

which together with (4.39) leads to

$$I_3 = -\frac{1}{4} (m_{AdS_5}^2 + m_{S^5}^2) \text{Tr}(\mathbb{I} - \rho_3^2). \quad (4.45)$$

Hence, collecting the three contributions (4.41), (4.42), and (4.45), and using $\rho_3^2 = \mathbb{I}$, we obtain

$$\text{Tr}(\rho^\alpha \mathcal{M}_F \rho_\alpha \mathcal{M}_F) = - (m_{AdS_5}^2 + m_{S^5}^2) \text{Tr}(\mathbb{I}) = \left({}^{(2)}R + \text{Tr}(K^2) \right) \text{Tr}(\mathbb{I}) \quad (4.46)$$

by means of the expressions (3.59). Armed with (4.46) we can compare the bosonic and fermionic contribution to the total logarithmic divergences, by computing explicitly $a_B^{(2)} + a_F^{(2)}$. The part depending on the traced mass matrices is

$$-\frac{1}{2} \text{Tr}(\rho^\alpha \mathcal{M}_F \rho_\alpha \mathcal{M}_F) + \text{Tr}(\mathcal{M}) = -4 \left({}^{(2)}R + \text{Tr}(K^2) \right) + 3 {}^{(2)}R + 4 \text{Tr}(K^2) = - {}^{(2)}R, \quad (4.47)$$

We have used $\text{Tr}(\mathbb{I}) = 8$ since these are eight $2d$ fermionic modes now, as reviewed at the beginning of Section 4.1. The contribution of the pure intrinsic curvature terms is instead more subtle [9]. The use of the naive Seeley-De Witt fermionic coefficient would lead to a wrong result

$$8 \frac{1}{6} R^{(2)} + 8 \frac{1}{12} R^{(2)} = 2R^{(2)}, \quad (4.48)$$

that combined with (4.47) would not produce the correct coefficient $3R^{(2)}$ [9]. The reason of the apparent disagreement is well known [9, 62–64]: the local $\text{SO}(1,9)$ rotation S that transforms ρ_i into two-dimensional Dirac matrices contracted with zweibein gives rise to a non-trivial Jacobian in the path-integral measure, that contributes additionally to the logarithmic divergence. The net effect of this “anomalous rotation” is to change the coefficient of the conformal anomaly of the relevant two-dimensional Dirac fermions by a factor 4. In our case it amounts to modify the fermionic contribution to (4.48) by a factor 4 and therefore

$$8\frac{1}{6}R^{(2)} + 8\frac{1}{3}R^{(2)} = 4R^{(2)}, \quad (4.49)$$

recovering, in combination with (4.47), the result of [9].

5 Applications

In this Section we work out a few relevant situations in which the general structures previously derived are exemplified. First we discuss the spinning string [2, 26], recovering the known spectrum of the fluctuations. Then we apply our analysis to the minimal surface associated to the 1/4 BPS Wilson loop operator [11, 56, 57], obtaining the bosonic and fermionic sectors and the related mass matrices.

5.1 Spinning strings

The first test of our analysis is the spinning string solution, generalization of [2] and [72], whose semiclassical quantization was worked out in [26]¹³. We choose the following coordinates for the target space metric

$$\begin{aligned} ds^2|_{AdS^5} &= -\cosh^2 \rho dt^2 + d\rho^2 + \sinh^2 \rho (d\beta_1^2 + \cos^2 \beta_1 d\beta_2^2 + \cos^2 \beta_1 \cos^2 \beta_2 d\phi^2), \\ ds^2|_{S^5} &= d\psi_1^2 + \cos^2 \psi_1 [d\psi_2^2 + \cos^2 \psi_2 (d\psi_3^2 + \cos^2 \psi_3 d\psi_4^2 + \cos^2 \psi_3 \cos^2 \psi_4 d\varphi^2)]. \end{aligned} \quad (5.1)$$

The motion of the spinning string in $\text{AdS}_5 \times S^5$ is described by the following ansatz for the embedding coordinates [26]

$$t = \kappa\tau, \quad \rho = \rho(\sigma), \quad \beta_i = 0 \quad (i = 1, 2), \quad \phi = \omega\tau, \quad \psi_i = 0 \quad (i = 1, \dots, 4), \quad \varphi = \nu\tau, \quad (5.2)$$

where (τ, σ) are the worldsheet coordinates, and $\rho(\sigma)$ has to satisfy the equation of motion and Virasoro constraints

$$\rho'' = (\kappa^2 - \omega^2) \cosh \rho \sinh \rho, \quad \rho'^2 = \kappa^2 \cosh^2 \rho - \omega^2 \sinh^2 \rho - \nu^2. \quad (5.3)$$

The solution $\rho(\sigma)$ can be found explicitly and it reads in terms of Jacobi elliptic functions, since we will not need it here we refer the reader to [26] for further details. The parameters κ, ω , and ν are chosen to be positive, and moreover the finite energy condition as well as the request of having a real solution impose certain relations on the parameters, that is

$$\kappa > \nu, \quad \omega > \nu, \quad \coth^2 \rho \geq \frac{\kappa^2 - \nu^2}{\omega^2 - \nu^2}. \quad (5.4)$$

¹³See [20] for the fermionic mass matrix of the string rotating in $\text{AdS}_5 \times S^1$.

In order to compute the quadratic fluctuation Lagrangian, we need to construct an orthonormal basis as in Section 3.3. The tangent vectors are given by

$$t_\tau^A = (\kappa \cos \rho, 0, 0, 0, \omega \sinh \rho, 0, 0, 0, 0, \nu), \quad t_\sigma^A = (0, \rho', 0, 0, 0, 0, 0, 0, 0, 0). \quad (5.5)$$

Then, we choose the 8 orthogonal vectors as follows

$$\begin{aligned} \hat{N}_1^A &= (0, 0, 1, 0, 0, 0, 0, 0, 0, 0), & \hat{N}_2^A &= (0, 0, 0, 1, 0, 0, 0, 0, 0, 0), \\ \bar{N}_1^A &= (0, 0, 0, 0, 0, 1, 0, 0, 0, 0), & \bar{N}_2^A &= (0, 0, 0, 0, 0, 0, 1, 0, 0, 0), \\ \bar{N}_3^A &= (0, 0, 0, 0, 0, 0, 0, 1, 0, 0), & \bar{N}_4^A &= (0, 0, 0, 0, 0, 0, 0, 0, 1, 0) \\ N_7^A &= \frac{1}{\sqrt{\rho'^2 + \nu^2}} (\omega \sinh \rho, 0, 0, 0, \kappa \cosh \rho, 0, 0, 0, 0, 0), \\ N_8^A &= \frac{1}{\rho' \sqrt{\nu^2 + \rho'^2}} (\kappa \nu \cosh \rho, 0, 0, 0, \nu \omega \sinh \rho, 0, 0, 0, 0, \nu^2 + \rho'^2). \end{aligned} \quad (5.6)$$

Finally, the induced worldsheet metric is given by

$$\gamma_{\alpha\beta} = (t_\alpha \cdot t_\beta) = \rho'^2 \text{diag}(-1, 1). \quad (5.7)$$

Once the basis vectors have been chosen, then the computation of the quadratic fluctuation Lagrangian is simply reduced to a mere application of expressions (3.35), (4.22), (4.28).

Bosonic Lagrangian Let us proceed to the construction of the bosonic transverse Lagrangian (3.35). The worldsheet curvature ${}^{(2)}R$ and the trace of the extrinsic curvature $\text{Tr}(K^2)$ turn out to be

$${}^{(2)}R = \frac{2(\kappa^2 - \nu^2)(\omega^2 - \nu^2)}{\rho'(\sigma)^4} - 2, \quad \text{Tr}(K^2) = -\frac{2(\kappa^2 - \nu^2)(\omega^2 - \nu^2)}{\rho'(\sigma)^4} - \frac{2\nu^2}{\rho'(\sigma)^2}. \quad (5.8)$$

The only non-zero components of the extrinsic curvature projected on the transverse directions, *i.e.* $K_{\alpha\beta}^i \equiv N_A^i K_{\alpha\beta}^A$ are

$$K_{\tau\sigma}^7 = K_{\sigma\tau}^7 = \frac{\kappa \omega \rho'}{\sqrt{\nu^2 + \rho'^2}}, \quad K_{\tau\sigma}^8 = K_{\sigma\tau}^8 = -\frac{\nu \rho''}{\sqrt{\nu^2 + \rho'^2}}. \quad (5.9)$$

This implies that the mass matrix \mathcal{M}_{ij} (3.35) for $i, j = 1, \dots, 6$ is diagonal and given by

$$\begin{aligned} \mathcal{M}_{11} = \mathcal{M}_{22} &= -m_{\text{AdS}_5}^2 = -\gamma^{\alpha\beta} (\hat{t}_\alpha \cdot \hat{t}_\beta) = -2 - \frac{\nu^2}{\rho'^2}, \\ \mathcal{M}_{33} = \mathcal{M}_{44} = \mathcal{M}_{55} = \mathcal{M}_{66} &= -m_{\text{S}^5}^2 = \gamma^{\alpha\beta} (\bar{t}_\alpha \cdot \bar{t}_\beta) = -\frac{\nu^2}{\rho'^2}. \end{aligned} \quad (5.10)$$

Along the directions 7, 8 the extrinsic curvature contributes to the mass (in particular, the contribution to the off-diagonal matrix is entirely due to the extrinsic curvature), and we obtain

$$\begin{aligned} \mathcal{M}_{77} &= -2 - \frac{\nu^2}{\rho'^2} - \frac{2\kappa^2 \omega^2}{\rho'^2(\nu^2 + \rho'^2)}, \\ \mathcal{M}_{88} &= -\frac{\nu^2}{\rho'^2} + \frac{2\kappa^2 \omega^2}{\rho'^2(\nu^2 + \rho'^2)} + \frac{2(\kappa^2 - \nu^2)(\nu^2 - \omega^2)}{\rho'^4}, \\ \mathcal{M}_{78} &= \frac{2\kappa \nu \omega \rho''}{\rho'^3(\nu^2 + \rho'^2)}. \end{aligned} \quad (5.11)$$

In order to complete the Lagrangian we construct the covariant derivative, for which we need the connection on the normal bundle A_α^{ij} (3.18). The only non-trivial directions are along the transverse modes 7, 8

$$A_\sigma^{78} = -A_\sigma^{87} = -\frac{\kappa\omega\nu}{\nu^2 + \rho'^2}, \quad A_\tau^{ij} = 0, \quad i, j = 1, \dots, 8. \quad (5.12)$$

Hence, the bosonic Lagrangian agrees with [26].

Fermionic Lagrangian The construction of the fermionic Lagrangian proceeds in two steps: the kinetic part (4.22) and the flux term (4.28). For the kinetic term the only new ingredient we need is the worldsheet spin connection $\omega_{\alpha ab}$,

$$\omega_{\tau 01} = -\omega_{\tau 10} = -\frac{\rho''}{\rho'}, \quad \omega_{\sigma ab} = 0. \quad (5.13)$$

Hence, the kinetic term (4.22) is simply given by [20]

$$\mathcal{L}_{kin} = 2i\bar{\Theta} \left(\sqrt{\gamma}\gamma^{\alpha\beta}\Gamma_a e_\alpha^a \partial_\beta + \frac{1}{2}\rho''\Gamma_1 + \frac{1}{2}\frac{\kappa\nu\omega\rho'}{(\nu^2 + \rho'^2)}\Gamma_{189} \right) \Theta. \quad (5.14)$$

where we have used (5.12) for the third term above, which is the connection-related part. Notice that, as normal bundle connection (5.12) is *flat*, *i.e.* $\partial_\tau A_\sigma^{78} - \partial_\sigma A_\tau^{78} + [A_\tau A_\sigma]^{78} = 0$, we can introduce a rotation which cancels such term - this is what done in [20].

In order to compute the flux term (4.28) we cannot use the final expression (4.31) of the expansion (4.29). This is due to the fact that for the spinning string one of the projected transverse vector, *i.e.* \hat{N}_7 in our convention, is orthogonal to \hat{t}_α . A better choice to simplify the term $S^{-1}\Gamma_\star S$ is then the basis formed by the two tangent vectors \hat{t}_α , the two transverse vectors \hat{N}_1, \hat{N}_2 and finally \hat{N}_7 . The expansion (4.30) is still valid for \hat{N}_8 . Thus, the only non zero terms which can contribute to $S^{-1}\Gamma_\star S$ are

$$\begin{aligned} S^{-1}\Gamma_\star S &= i\hat{e}_{ABCDE} \left(\hat{t}_\alpha^A \hat{t}_\beta^B e_\alpha^a e_b^\beta \hat{N}_i^C \hat{N}_j^D \hat{N}_r^E \Gamma_{ab} \Gamma_{ij} \Gamma_r + \hat{t}_\alpha^A e_\alpha^a \hat{N}_i^B \hat{N}_j^C \hat{N}_r^D \hat{N}_s^E \Gamma_a \Gamma_{ij} \Gamma_{rs} \right) = \\ &= i\frac{\sqrt{\nu^2 + \rho'^2}}{\rho'} \Gamma_{01238} - i\frac{\nu}{\rho'} \Gamma_{12389}. \end{aligned} \quad (5.15)$$

The remaining terms are straightforward to compute, simply from the inner products of the basis vectors, and we obtain

$$\begin{aligned} \mathcal{L}_{flux} &= \frac{i}{2}\bar{\Theta} \left(2(\nu^2 + \rho'^2)\Gamma_{01} + 2\nu\sqrt{\nu^2 + \rho'^2}\Gamma_{19} \right) \left(\frac{\sqrt{\nu^2 + \rho'^2}}{\rho'} \Gamma_{01238} - \frac{\nu}{\rho'} \Gamma_{12389} \right) \Theta \\ &= i\rho'\sqrt{\nu^2 + \rho'^2} \bar{\Theta} \Gamma_{238} \Theta. \end{aligned} \quad (5.16)$$

The sum of the two contributions (5.14) and (5.16) is exactly the fermionic Lagrangian computed in Appendix D of [20].

The special case of the folded string. The folded string of [2, 26] is obtained switching off the angle φ in S^5 (setting the parameter ν to zero in the classical ansatz (5.2)). Then the string motion is confined in AdS_3 , *i.e.* $\bar{t}_\alpha = 0$ for $\alpha = 1, 2$. As explained in the main body, this implies that we can choose 5 orthonormal vectors completely embedded in the compact sub-space S^5 , (in our convention (5.6) they will be $\bar{N}_1, \bar{N}_2, \bar{N}_3, \bar{N}_4, \bar{N}_8 = N_8$ with $\nu = 0$). The remaining vectors (\hat{t}_α with $\alpha = 1, 2$, \hat{N}_1, \hat{N}_2 and $\hat{N}^7 = N^7$) live only in AdS . Namely, the basis vectors is entirely split in the two sub-spaces. It is then immediate to see that the bosonic masses reduce to

$$\begin{aligned} \mathcal{M}_{11} = \mathcal{M}_{22} &= -m_{\text{AdS}_5}^2 = -2, & \mathcal{M}_{77} &= -2 - \frac{2\kappa^2\omega^2}{\rho'^4}, \\ \mathcal{M}_{33} = \mathcal{M}_{44} = \mathcal{M}_{55} = \mathcal{M}_{66} = \mathcal{M}_{88} &= -m_{S^5}^2 = 0, \end{aligned} \quad (5.17)$$

which is the well-known results of 5 massless scalars in S^5 and the 3 massive scalar modes living in $\text{AdS}_3 \subset \text{AdS}_5$ [26]. Now, the cross terms are zero $\mathcal{M}_{78} = 0$ as well as the connection $A_\sigma^{78} = -A_\sigma^{87} = 0$.

The fermionic Lagrangian (5.14) and (5.16) simplifies as well. Being the normal bundle connection zero, the kinetic term reduces to

$$\mathcal{L}_{kin} = 2i \bar{\Theta} \left(\sqrt{\gamma} \gamma^{\alpha\beta} \Gamma_a e_\alpha^a \partial_\beta + \frac{1}{2} \rho'' \Gamma_1 \right) \Theta. \quad (5.18)$$

The only contribution to the flux comes from the AdS_5 mass in (4.28) and the first product in (5.15) (with $\nu = 0$), thus

$$\mathcal{L}_{flux} = i \rho'^2 \bar{\Theta} \Gamma_{238} \Theta. \quad (5.19)$$

5.2 String dual to latitude Wilson loops

In this Section we apply our analysis to the $\text{AdS}_5 \times S^5$ string minimal surface corresponding to the latitude Wilson loop operators of [11, 56, 57]. We choose the following patches for the target space metric:

$$ds^2|_{\text{AdS}_5} = -\cosh^2 \rho dt^2 + d\rho^2 + \frac{\sinh^2 \rho}{(1 + \frac{1}{4} \vec{y} \cdot \vec{y})^2} d\vec{y} \cdot d\vec{y}, \quad ds^2|_{S^5} = \frac{1}{(1 + \frac{1}{4} \vec{z} \cdot \vec{z})^2} d\vec{z} \cdot d\vec{z}, \quad (5.20)$$

where $\vec{y} = (y_1, y_2, y_3)$ with $\vec{y} \cdot \vec{y} = y_1^2 + y_2^2 + y_3^2$, and $\vec{z} = (z_1, \dots, z_5)$, with $\vec{z} \cdot \vec{z} = z_1^2 + \dots + z_5^2$. The classical string solution [57] is described by the ansatz

$$\begin{aligned} t = 0, \quad \rho = \rho(\sigma), \quad y^1 = 2 \sin \varphi, \quad y^2 = 2 \cos \varphi, \quad y^3 = 0 \\ z^1 = 0, \quad z^2 = 0, \quad z^3 = 2 \cos \theta(\sigma), \quad z^4 = 2 \sin \theta(\sigma) \sin \varphi, \quad z^5 = 2 \sin \theta(\sigma) \cos \varphi. \end{aligned} \quad (5.21)$$

where the worldsheet coordinates are $\varphi \in [0, 2\pi)$ and $\sigma \in [0, \infty)$. The classical equations of motion read

$$\begin{aligned} \rho'(\sigma) &= -\sinh \rho(\sigma), & \rho''(\sigma) &= \sinh \rho(\sigma) \cosh \rho(\sigma), \\ \theta'(\sigma) &= \mp \frac{1}{\cosh(\sigma_0 \pm \sigma)} = \mp \sin \theta(\sigma), & \theta''(\sigma) &= \sin \theta(\sigma) \cos \theta(\sigma), \end{aligned} \quad (5.22)$$

with solutions given in terms of simple hyperbolic functions

$$\sinh \rho(\sigma) = \frac{1}{\sinh \sigma}, \quad \sin \theta(\sigma) = \frac{1}{\cosh(\sigma_0 \pm \sigma)}. \quad (5.23)$$

The parameter σ_0 is a constant of integration defined by the internal latitude angle θ_0

$$\cos \theta_0 = \tanh \sigma_0. \quad (5.24)$$

To compute the quadratic fluctuation Lagrangian, we build an orthonormal basis choosing the tangent vectors as (from now on we only consider the solution in (5.22)-(5.23) with the upper sign, as it is this one that corresponds to the minimum of the action)

$$\begin{aligned} t_\varphi^A &= (0, 0, \cos \varphi \sinh \rho, -\sin \varphi \sinh \rho, 0, 0, 0, 0, \cos \varphi \sin \theta, -\sin \varphi \sin \theta), \\ t_\sigma^A &= (0, -\sinh \rho, 0, 0, 0, 0, 0, \sin^2 \theta, -\cos \theta \sin \theta \sin \varphi, -\cos \varphi \sin \theta \cos \theta), \end{aligned} \quad (5.25)$$

where we omit the dependence of ρ and θ on σ . The vectors orthogonal to the worldsheet can be chosen to be

$$\begin{aligned} \bar{N}_1^A &= (0, 0, 0, 0, 0, 1, 0, 0, 0, 0), & \bar{N}_2^A &= (0, 0, 0, 0, 0, 0, 0, 0, \cos \theta, \sin \varphi \sin \theta, \cos \varphi \sin \theta), \\ \bar{N}_3^A &= (0, 0, 0, 0, 0, 0, 1, 0, 0, 0), \\ \hat{N}_1^A &= (-1, 0, 0, 0, 0, 0, 0, 0, 0, 0), & \hat{N}_2^A &= (0, 0, 0, 0, 1, 0, 0, 0, 0, 0), \\ \hat{N}_3^A &= (0, 0, \sin \varphi, \cos \varphi, 0, 0, 0, 0, 0, 0). \end{aligned} \quad (5.26)$$

The basis is completed constructing the normal vectors as

$$\begin{aligned} N_7^A &= \frac{1}{\sqrt{\sin^2 \theta + \sinh^2 \rho}} (0, 0, \cos \varphi \sin \theta, -\sin \varphi \sin \theta, 0, 0, 0, 0, -\cos \varphi \sinh \rho, \sin \varphi \sinh \rho), \\ N_8^A &= \frac{1}{\sqrt{\sin^2 \theta + \sinh^2 \rho}} (0, \sin \theta, 0, 0, 0, 0, 0, \sin \theta \sinh \rho, -\cos \theta \sinh \rho \sin \varphi, -\cos \theta \sinh \rho \cos \varphi). \end{aligned} \quad (5.27)$$

For this string configuration the worldsheet is space-like. The induced worldsheet metric $\gamma_{\alpha\beta}$ has then Euclidean signature and is given by

$$\gamma_{\alpha\beta} = \Omega^2(\sigma) \text{diag}(1, 1), \quad \Omega^2(\sigma) = \sinh^2 \rho + \sin^2 \theta = \rho'^2 + \theta'^2 = \frac{1}{\cosh^2(\sigma + \sigma_0)} + \frac{1}{\sinh^2 \sigma}. \quad (5.28)$$

Instead, the orthonormal vectors on AdS have a time-like component, so that in this Section we will use that

$$\left(\hat{N}_i \cdot \hat{N}_j \right) = \eta_{ij}, \quad \eta_{ij} = \text{diag}(-1, 1, 1), \quad i, j = 1, 2, 3, \quad (5.29)$$

while the remaining scalar products are unchanged with respect to Section 3.3. We report here for completeness the expressions of the worldsheet curvature ${}^{(2)}R$ and of the trace of the squared extrinsic curvature $\text{Tr}(K^2)$:

$${}^{(2)}R = -\frac{2 \partial_\sigma^2 \log \Omega(\sigma)}{\Omega^2(\sigma)}, \quad \text{Tr}(K^2) = 4 \frac{\sinh^2(\sigma) \cosh^2(\sigma + \sigma_0)}{\cosh(\sigma_0) \cosh(2\sigma + \sigma_0)^3}. \quad (5.30)$$

Bosonic Lagrangian We want to compute the transverse Lagrangian (3.35). We proceed first evaluating the mass matrix (3.68). In the case of a space-like worldsheet the matrix is slightly modified: the first element in the diagonal acquires a relative sign. From the definition (3.55) we obtain

$$\begin{aligned} m_{\text{AdS}_5}^2 &= \gamma^{\alpha\beta} (\hat{t}_\alpha \cdot \hat{t}_\beta) = \frac{2 \cosh^2(\sigma + \sigma_0)}{\cosh \sigma_0 \cosh(2\sigma + \sigma_0)}, \\ m_{S^5}^2 &= -\gamma^{\alpha\beta} (\bar{t}_\alpha \cdot \bar{t}_\beta) = -\frac{2 \sinh^2 \sigma}{\cosh \sigma_0 \cosh(2\sigma + \sigma_0)} \end{aligned} \quad (5.31)$$

while the reduced matrix (3.70) vanishes identically. Thus, the matrix (3.68) reads

$$\gamma^{\alpha\beta} R_{AMBNT}{}^A t_\alpha^B N^i M N^j N = - \begin{pmatrix} -m_{\text{AdS}_5}^2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & m_{\text{AdS}_5}^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & m_{\text{AdS}_5}^2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & m_{S^5}^2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & m_{S^5}^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & m_{S^5}^2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad (5.32)$$

with $m_{\text{AdS}_5}^2$ and $m_{S^5}^2$ given in (5.31).

Let us now construct the projection of the extrinsic curvature along the transverse direction, *i.e.* $K_{\alpha\beta}^i \equiv N_A^i K_{\alpha\beta}^A$, (2.5). The only non-zero components are

$$\begin{aligned} K_{\varphi\sigma}^7 &= K_{\sigma\varphi}^7 = -\frac{\sqrt{\cosh \sigma_0}}{\sinh \sigma \cosh(\sigma + \sigma_0) \sqrt{\cosh(2\sigma + \sigma_0)}}, \\ K_{\varphi\varphi}^8 &= K_{\sigma\sigma}^8 = -\frac{\sqrt{\cosh \sigma_0}}{\sinh \sigma \cosh(\sigma + \sigma_0) \sqrt{\cosh(2\sigma + \sigma_0)}}. \end{aligned} \quad (5.33)$$

It is then straightforward to construct the whole mass matrix \mathcal{M}_{ij} in (3.35)

$$\begin{aligned} \mathcal{M}_{11} &= -\mathcal{M}_{22} = -\mathcal{M}_{33} = -m_{\text{AdS}_5}^2 = \frac{2 \cosh^2(\sigma + \sigma_0)}{\cosh \sigma_0 \cosh(2\sigma + \sigma_0)}, \\ \mathcal{M}_{44} &= \mathcal{M}_{55} = \mathcal{M}_{66} = m_{S^5}^2 = -\frac{2 \sinh^2 \sigma}{\cosh \sigma_0 \cosh(2\sigma + \sigma_0)}, \\ \mathcal{M}_{77} &= \mathcal{M}_{88} = \frac{2 \cosh^2(\sigma + \sigma_0) \sinh^2 \sigma}{\cosh \sigma_0 \cosh^3(2\sigma + \sigma_0)}, \quad \mathcal{M}_{78} = \mathcal{M}_{87} = 0. \end{aligned} \quad (5.34)$$

Finally, we calculate the normal connection A_α^{ij} (3.18), whose non-vanishing components read

$$A_\varphi^{78} = -A_\varphi^{87} = -\tanh(2\sigma + \sigma_0). \quad (5.35)$$

The covariant derivatives on the transverse fields (3.17) appearing in (3.35) will be non-trivial only along the directions given by (5.35). Notice that the field-strength $\partial_\sigma A_\varphi^{78} - \partial_\varphi A_\sigma^{78} + [A_\sigma, A_\varphi]^{78} = \partial_\sigma A_\varphi^{78}$ does *not* vanish.

Fermionic Lagrangian The construction of the fermionic Lagrangian proceeds in two steps: the kinetic part (4.22) and the flux term (4.28). We perform the computations in a Lorentzian signature

for the induced worldsheet metric, and only at the end Wick-rotate back. For the kinetic term the only new ingredient we need is the worldsheet spin connection $\omega_{\alpha ab}$,

$$\omega_{\varphi 12} = -\omega_{\varphi 21} = -\frac{1}{\Omega^2(\sigma)} \left(\frac{\coth \sigma}{\sinh^2 \sigma} + \frac{\tanh(\sigma + \sigma_0)}{\cosh^2(\sigma + \sigma_0)} \right), \quad (5.36)$$

then, the final expression of the kinetic term turns out to be

$$\tilde{L}_F^{kin} = 2i \bar{\Theta} \left(\sqrt{\gamma} \gamma^{\alpha\beta} \Gamma^a e_{a\alpha} \partial_\beta - \frac{1}{2\Omega(\sigma)} \left(\frac{\coth \sigma}{\sinh^2 \sigma} + \frac{\tanh(\sigma + \sigma_0)}{\cosh^2(\sigma + \sigma_0)} \right) \Gamma_4 + \frac{\Omega(\sigma)}{2} \tanh(2\sigma + \sigma_0) \Gamma_{389} \right) \Theta. \quad (5.37)$$

For the flux term (4.28), one obtains

$$\tilde{L}_F^{flux} = 2i \bar{\Theta} \left(-\frac{1}{\sinh^2 \sigma} \Gamma_\star \tau_3 - \frac{1}{\cosh^2(\sigma + \sigma_0)} \Gamma_\star \Gamma_{89} \right) \Theta. \quad (5.38)$$

The evaluation of the determinants associated to the fluctuations above is in [43].

String dual to circular Wilson loop. In the limit when $\theta_0 \rightarrow 0$, the latitude on S^5 shrinks to a point, and the Wilson loop reduces to a circular one, whose string dual was studied in [9, 23]. From equations (5.22) and (5.24) the limit is equivalent to $\sigma_0 \rightarrow \infty$ and $\theta \rightarrow \theta_0 \rightarrow 0$, that is the string motion is turned off on the compact space. Taking these limits in our formulas we can reconstruct the bosonic and fermionic Lagrangian of [9, 23]. We will only change the last two normal vectors N , since in this way the normal connection A vanishes by construction (otherwise one can always perform a rotation after to eliminate such a connection since now the corresponding field strength is flat):

$$N_7 = (0, 0, 0, 0, 0, 0, 0, 0, 1, 0), \quad N_8 = (0, 0, 0, 0, 0, 0, 0, 0, 0, 1). \quad (5.39)$$

As for the degenerate case of the folded string, here five normal vectors completely lie in S^5 while AdS_5 is spanned by the two transverse vectors $\hat{t}_\alpha = t_\alpha$ ($\alpha = 1, 2$) and three normal vectors \hat{N}_i ($i = 1, 2, 3$). The rest proceeds as before, and we have

$$\begin{aligned} {}^{(2)}R &= -2, & \text{Tr}(K^2) &= 0, & \Omega^2(\sigma) &= \frac{1}{\sinh^2 \sigma} \\ \mathcal{M}_{11} &= -\mathcal{M}_{22} = -\mathcal{M}_{3t3} = m_{AdS_5}^2 = 2, & \mathcal{M}_{ii} &= -m_{S^5}^2 = 0, & i &= 4, \dots, 8. \end{aligned} \quad (5.40)$$

As mentioned, the choice of vectors (5.39) allows us to immediately eliminate the connection (5.35) without resorting to any further rotation, hence we can write

$$\begin{aligned} \tilde{L}_F^{kin} &= 2i \bar{\Theta} \left(\sqrt{\gamma} \gamma^{\alpha\beta} \Gamma^a e_{a\alpha} \partial_\beta - \frac{1}{2} \coth \sigma \Gamma_4 \right) \Theta, \\ \tilde{L}_F^{flux} &= -2i \frac{1}{\sinh^2 \sigma} \bar{\Theta} \Gamma_\star \tau_3 \Theta. \end{aligned} \quad (5.41)$$

We conclude this section by noticing that we have also carried on the computations for the minimal surface dual to generalized cusped Wilson loops [18]. The quadratic Lagrangian for the string fluctuations was computed in [18], and we have checked that also in this case we reproduces all the bosonic masses of [18]. Notice that the only non-vanishing component is the σ -component

$$A_\sigma^{37} = -\frac{\sqrt{b^4 p^2} \sqrt{(b^2 + 1) p^2 - b^4}}{\sinh^2 \rho(\sigma) \sqrt{b^4 + p^2} \left(\frac{b^4 - p^2}{\sinh^2 \rho(\sigma)} + b^2 p^2 \right)}, \quad (5.42)$$

which makes this case to fall – the solution lies both in AdS_5 and S^5 – in the class of solutions discussed around eqs. (3.66)-(3.70). As $\partial_\tau A_\sigma^{37} - \partial_\sigma A_\tau^{37} = 0$, and so the field-strength vanishes, bosonic masses are described by (3.70) with one of the eigenvalues λ_i vanishing. Also, as mentioned in Section 4.1 a local target space rotation can be found which eliminates the normal bundle contribution to the kinetic part of the fermionic action (see formulas C.16-C.17 in [18]).

Acknowledgments

The work of VF and EV is funded by DFG via the Emmy Noether Programme “*Gauge Field from Strings*”. EV acknowledges support from Research Training Group GK 1504 “*Mass, Spectrum, Symmetry*” and from the Seventh Framework Programme [FP7-People-2010-IRSES] under grant agreement n. 269217 (UNIFY), and would like to thank the Perimeter Institute for Theoretical Physics for hospitality during the completion of this work. All authors would like to thank the Galileo Galilei Institute for Theoretical Physics for hospitality during the completion of this work.

References

- [1] A. M. Polyakov, “*Quantum Geometry of Bosonic Strings*”, *Phys.Lett.* **B103**, 207 (1981).
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “*A semi-classical limit of the gauge/string correspondence*”, *Nucl. Phys.* **B636**, 99 (2002), [hep-th/0204051](#).
- [3] I. Bena, J. Polchinski and R. Roiban, “*Hidden symmetries of the $AdS_5 \times S^5$ superstring*”, *Phys.Rev.* **D69**, 046002 (2004), [hep-th/0305116](#).
- [4] V. Pestun, “*Localization of gauge theory on a four-sphere and supersymmetric Wilson loops*”, *Commun.Math.Phys.* **313**, 71 (2012), [arXiv:0712.2824](#).
- [5] A. Kapustin, B. Willett and I. Yaakov, “*Exact Results for Wilson Loops in Superconformal Chern-Simons Theories with Matter*”, *JHEP* **1003**, 089 (2010), [arXiv:0909.4559](#).
- [6] M. Marino and P. Putrov, “*Exact Results in ABJM Theory from Topological Strings*”, *JHEP* **1006**, 011 (2010), [arXiv:0912.3074](#).
- [7] N. Drukker, M. Marino and P. Putrov, “*From weak to strong coupling in ABJM theory*”, *Commun.Math.Phys.* **306**, 511 (2011), [arXiv:1007.3837](#).
- [8] J. Erickson, G. Semenoff and K. Zarembo, “*Wilson loops in $\mathcal{N} = 4$ supersymmetric Yang-Mills theory*”, *Nucl.Phys.* **B582**, 155 (2000), [hep-th/0003055](#).
- [9] N. Drukker, D. J. Gross and A. A. Tseytlin, “*Green-Schwarz string in $AdS_5 \times S^5$: Semiclassical partition function*”, *JHEP* **0004**, 021 (2000), [hep-th/0001204](#).
- [10] K. Zarembo, “*Supersymmetric Wilson loops*”, *Nucl.Phys.* **B643**, 157 (2002), [hep-th/0205160](#).
- [11] N. Drukker, S. Giombi, R. Ricci and D. Trancanelli, “*Supersymmetric Wilson loops on S^3* ”, *JHEP* **0805**, 017 (2008), [arXiv:0711.3226](#).
- [12] A. Buchel, J. G. Russo and K. Zarembo, “*Rigorous Test of Non-conformal Holography: Wilson Loops in $N=2^*$ Theory*”, *JHEP* **1303**, 062 (2013), [arXiv:1301.1597](#).
- [13] F. Bigazzi, A. L. Cotrone, L. Griguolo and D. Seminara, “*A novel cross-check of localization and non conformal holography*”, *JHEP* **1403**, 072 (2014), [arXiv:1312.4561](#).
- [14] A. A. Tseytlin, “*Review of AdS/CFT Integrability, Chapter II.1: Classical $AdS_5 \times S^5$ string solutions*”, [arXiv:1012.3986](#).
- [15] T. McLoughlin, “*Review of AdS/CFT Integrability, Chapter II.2: Quantum Strings in $AdS_5 \times S^5$* ”, [arXiv:1012.3987](#).
- [16] L. F. Alday, “*Review of AdS/CFT Integrability, Chapter V.3: Scattering Amplitudes at Strong Coupling*”, *Lett.Math.Phys.* **99**, 507 (2012), [arXiv:1012.4003](#).
- [17] T. Klose, “*Review of AdS/CFT Integrability, Chapter IV.3: $N=6$ Chern-Simons and Strings on $AdS_4 \times CP^3$* ”, *Lett.Math.Phys.* **99**, 401 (2012), [arXiv:1012.3999](#).

- [18] N. Drukker and V. Forini, “Generalized quark-antiquark potential at weak and strong coupling”, *JHEP* **1106**, 131 (2011), [arXiv:1105.5144](#).
- [19] D. Correa, J. Henn, J. Maldacena and A. Sever, “An exact formula for the radiation of a moving quark in $\mathcal{N} = 4$ super Yang Mills”, *JHEP* **1206**, 048 (2012), [arXiv:1202.4455](#).
- [20] V. Forini, V. G. M. Puletti and O. Ohlsson Sax, “The generalized cusp in $AdS_4 \times CP^3$ and more one-loop results from semiclassical strings”, *J.Phys. A* **46**, 115402 (2013), [arXiv:1204.3302](#).
- [21] E. I. Buchbinder and A. A. Tseytlin, “One-loop correction to the energy of a wavy line string in AdS_5 ”, *J. Phys. A* **46**, 505401 (2013), [arXiv:1309.1581](#).
- [22] J. Aguilera-Damia, D. H. Correa and G. A. Silva, “Semiclassical partition function for strings dual to Wilson loops with small cusps in ABJM”, *JHEP* **1503**, 002 (2015), [arXiv:1412.4084](#).
- [23] M. Kruczenski and A. Tirziu, “Matching the circular Wilson loop with dual open string solution at 1-loop in strong coupling”, *JHEP* **0805**, 064 (2008), [arXiv:0803.0315](#).
- [24] C. Kristjansen and Y. Makeenko, “More about One-Loop Effective Action of Open Superstring in $AdS_5 \times S^5$ ”, *JHEP* **1209**, 053 (2012), [arXiv:1206.5660](#).
- [25] S. Forste, D. Ghoshal and S. Theisen, “Stringy corrections to the Wilson loop in $N=4$ super Yang-Mills theory”, *JHEP* **9908**, 013 (1999), [hep-th/9903042](#).
- [26] S. Frolov and A. A. Tseytlin, “Semiclassical quantization of rotating superstring in $AdS_5 \times S^5$ ”, *JHEP* **0206**, 007 (2002), [hep-th/0204226](#).
- [27] S. Frolov and A. A. Tseytlin, “Multispin string solutions in $AdS_5 \times S^5$ ”, *Nucl.Phys. B* **668**, 77 (2003), [hep-th/0304255](#).
- [28] S. Frolov and A. A. Tseytlin, “Quantizing three spin string solution in $AdS_5 \times S^5$ ”, *JHEP* **0307**, 016 (2003), [hep-th/0306130](#).
- [29] S. Frolov, I. Park and A. A. Tseytlin, “On one-loop correction to energy of spinning strings in S^5 ”, *Phys.Rev. D* **71**, 026006 (2005), [hep-th/0408187](#).
- [30] I. Park, A. Tirziu and A. A. Tseytlin, “Spinning strings in $AdS_5 \times S^5$: One-loop correction to energy in $SL(2)$ sector”, *JHEP* **0503**, 013 (2005), [hep-th/0501203](#).
- [31] N. Beisert, A. A. Tseytlin and K. Zarembo, “Matching quantum strings to quantum spins: One-loop versus finite-size corrections”, *Nucl.Phys. B* **715**, 190 (2005), [hep-th/0502173](#).
- [32] M. Beccaria, V. Forini, A. Tirziu and A. A. Tseytlin, “Structure of large spin expansion of anomalous dimensions at strong coupling”, *Nucl.Phys. B* **812**, 144 (2009), [arXiv:0809.5234](#).
- [33] M. Beccaria, G. V. Dunne, V. Forini, M. Pawellek and A. A. Tseytlin, “Exact computation of one-loop correction to energy of spinning folded string in $AdS_5 \times S^5$ ”, *J.Phys. A* **43**, 165402 (2010), [arXiv:1001.4018](#).
- [34] V. Forini, “Quark-antiquark potential in AdS at one loop”, *JHEP* **1011**, 079 (2010), [arXiv:1009.3939](#).
- [35] V. Forini, V. G. M. Puletti, M. Pawellek and E. Vescovi, “One-loop spectroscopy of semiclassically quantized strings: bosonic sector”, *J.Phys. A* **48**, 085401 (2015), [arXiv:1409.8674](#).
- [36] S. Giombi, R. Ricci, R. Roiban, A. A. Tseytlin and C. Vergu, “Generalized scaling function from light-cone gauge $AdS_5 \times S^5$ superstring”, *JHEP* **1006**, 060 (2010), [arXiv:1002.0018](#).
- [37] L. Bianchi, M. S. Bianchi, A. Bres, V. Forini and E. Vescovi, “Two-loop cusp anomaly in ABJM at strong coupling”, *JHEP* **1410**, 13 (2014), [arXiv:1407.4788](#).
- [38] R. Metsaev and A. A. Tseytlin, “Type IIB superstring action in $AdS_5 \times S^5$ background”, *Nucl.Phys. B* **533**, 109 (1998), [hep-th/9805028](#).

- [39] C. G. Callan, Jr. and L. Thorlacius, “SIGMA MODELS AND STRING THEORY”, in: “*In *Providence 1988, Proceedings, Particles, strings and supernovae, vol. 2* 795-878.*”.
- [40] R. Capovilla and J. Guven, “*Geometry of deformations of relativistic membranes*”, *Phys.Rev. D51*, 6736 (1995), [gr-qc/9411060](#).
- [41] K. Viswanathan and R. Parthasarathy, “*String theory in curved space-time*”, *Phys.Rev. D55*, 3800 (1997), [hep-th/9605007](#).
- [42] A. Faraggi, W. Mueck and L. A. Pando Zayas, “*One-loop Effective Action of the Holographic Antisymmetric Wilson Loop*”, *Phys. Rev. D85*, 106015 (2012), [arXiv:1112.5028](#).
- [43] V. Forini, L. Griguolo, V. G. M. Puletti, D. Seminara and E. Vescovi, to appear.
- [44] T. Klose, T. McLoughlin, R. Roiban and K. Zarembo, “*Worldsheet scattering in $AdS(5) \times S^{*5}$* ”, *JHEP 0703*, 094 (2007), [hep-th/0611169](#).
- [45] T. Klose, T. McLoughlin, J. A. Minahan and K. Zarembo, “*World-sheet scattering in $AdS(5) \times S^{*5}$ at two loops*”, *JHEP 0708*, 051 (2007), [arXiv:0704.3891](#).
- [46] V. Giangreco Marotta Puletti, T. Klose and O. Ohlsson Sax, “*Factorized world-sheet scattering in near-flat $AdS(5) \times S^{*5}$* ”, *Nucl. Phys. B792*, 228 (2008), [arXiv:0707.2082](#).
- [47] G. Arutyunov and S. Frolov, “*Foundations of the $AdS_5 \times S^5$ Superstring. Part I*”, *J.Phys. A42*, 254003 (2009), [arXiv:0901.4937](#).
- [48] L. Bianchi, V. Forini and B. Hoare, “*Two-dimensional S-matrices from unitarity cuts*”, *JHEP 1307*, 088 (2013), [arXiv:1304.1798](#).
- [49] O. T. Engelund, R. W. McKeown and R. Roiban, “*Generalized unitarity and the worldsheet S matrix in $AdS_n \times S^n \times M^{10-2n}$* ”, *JHEP 1308*, 023 (2013), [arXiv:1304.4281](#).
- [50] R. Roiban, P. Sundin, A. Tseytlin and L. Wulff, “*The one-loop worldsheet S-matrix for the $AdS_n \times S^n \times T^{10-2n}$ superstring*”, *JHEP 1408*, 160 (2014), [arXiv:1407.7883](#).
- [51] N. Beisert, “*The $SU(2|2)$ dynamic S-matrix*”, *Adv. Theor. Math. Phys. 12*, 945 (2008), [hep-th/0511082](#).
- [52] N. Beisert and L. Freyhult, “*Fluctuations and energy shifts in the Bethe ansatz*”, *Phys.Lett. B622*, 343 (2005), [hep-th/0506243](#).
- [53] N. Gromov, “*Integrability in AdS/CFT correspondence: Quasi-classical analysis*”, *J.Phys. A42*, 254004 (2009).
- [54] B. Vicedo, “*Semiclassical Quantisation of Finite-Gap Strings*”, *JHEP 0806*, 086 (2008), [arXiv:0803.1605](#).
- [55] R. Ishizeki, M. Kruczenski and S. Ziam, “*Notes on Euclidean Wilson loops and Riemann Theta functions*”, *Phys. Rev. D85*, 106004 (2012), [arXiv:1104.3567](#).
- [56] N. Drukker and B. Fiol, “*On the integrability of Wilson loops in $AdS_5 \times S^5$: Some periodic ansatze*”, *JHEP 0601*, 056 (2006), [hep-th/0506058](#).
- [57] N. Drukker, “ *$1/4$ BPS circular loops, unstable world-sheet instantons and the matrix model*”, *JHEP 0609*, 004 (2006), [hep-th/0605151](#).
- [58] L. Alvarez-Gaume, D. Z. Freedman and S. Mukhi, “*The Background Field Method and the Ultraviolet Structure of the Supersymmetric Nonlinear Sigma Model*”, *Annals Phys. 134*, 85 (1981).
- [59] S. Kobayashi and K. Nomizu, “*Foundations of differential geometry. Volume II*”, Interscience (1963), New-York.
- [60] A. Kavalov, I. Kostov and A. Sedrakian, “*Dirac and Weyl Fermion Dynamics on Two-dimensional Surface*”, *Phys.Lett. B175*, 331 (1986).

- [61] A. Sedrakian and R. Stora, “Dirac and Weyl Fermions Coupled to Two-dimensional Surfaces: Determinants”, [Phys.Lett. B188, 442 \(1987\)](#).
- [62] F. Langouche and H. Leutwyler, “Two-dimensional fermion determinants as Wess-Zumino actions”, [Phys.Lett. B195, 56 \(1987\)](#).
- [63] F. Langouche and H. Leutwyler, “Weyl fermions on strings embedded in three-dimensions”, [Z.Phys. C36, 473 \(1987\)](#).
- [64] F. Langouche and H. Leutwyler, “Anomalies generated by extrinsic curvature”, [Z.Phys. C36, 479 \(1987\)](#).
- [65] D. R. Karakhanian, “Induced Dirac operator and smooth manifold geometry”.
- [66] R. Kallosh, J. Rahmfeld and A. Rajaraman, “Near horizon superspace”, [JHEP 9809, 002 \(1998\)](#), [hep-th/9805217](#).
- [67] R. Metsaev and A. A. Tseytlin, “Superstring action in $AdS_5 \times S^5$. Kappa symmetry light cone gauge”, [Phys.Rev. D63, 046002 \(2001\)](#), [hep-th/0007036](#).
- [68] R. Metsaev, C. B. Thorn and A. A. Tseytlin, “Light cone superstring in AdS space-time”, [Nucl.Phys. B596, 151 \(2001\)](#), [hep-th/0009171](#).
- [69] J. Callan, Curtis G., H. K. Lee, T. McLoughlin, J. H. Schwarz, I. Swanson et al., “Quantizing string theory in $AdS_5 \times S^5$: Beyond the pp wave”, [Nucl.Phys. B673, 3 \(2003\)](#), [hep-th/0307032](#).
- [70] R. Roiban, A. Tirziu and A. A. Tseytlin, “Two-loop world-sheet corrections in $AdS_5 \times S^5$ superstring”, [JHEP 0707, 056 \(2007\)](#), [arXiv:0704.3638](#).
- [71] P. B. Gilkey, “Invariance Theory, the Heat Equation and the Atiyah-Singer Index Theorem”, CRC Press (1995), Boca Raton.
- [72] D. E. Berenstein, J. M. Maldacena and H. S. Nastase, “Strings in flat space and pp waves from $N=4$ super Yang-Mills”, [JHEP 0204, 013 \(2002\)](#), [hep-th/0202021](#).