



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** Ayers, S., Wright, D. B. & Wells, N. (2007). Post-traumatic stress in couples after birth: Association with the couple's relationship and parent-baby bond. *Journal of Reproductive & Infant Psychology*, 25(1), pp. 40-50. doi: 10.1080/02646830601117175

This is the unspecified version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/2006/>

**Link to published version:** <https://doi.org/10.1080/02646830601117175>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

---

---

City Research Online:

<http://openaccess.city.ac.uk/>

[publications@city.ac.uk](mailto:publications@city.ac.uk)

---

**Post-traumatic stress in couples after birth:  
Association with the couple's relationship and parent-baby bond**

Susan Ayers<sup>1</sup>, Daniel B. Wright<sup>2</sup> & Nicola Wells<sup>3</sup>

<sup>1</sup>City University London, UK; <sup>2</sup>Florida International University, USA, <sup>3</sup>Brunel University,

UK

**Please cite this paper as:**

Ayers, S., Wright, D. B., & Wells, N. (2007). Post-traumatic stress in couples after birth: Association with the couple's relationship and parent-baby bond. *Journal of Reproductive & Infant Psychology*, 25(1), 40-50.

**Key words:** birth, post-traumatic stress disorder, relationship, mother-baby bond

## **Post-traumatic stress in couples after birth:**

### **Association with the couple's relationship and parent-baby bond**

#### **ABSTRACT**

Research suggests a proportion of women develop post-traumatic stress disorder (PTSD) after childbirth. However, the effects of postnatal PTSD on the couple's relationship and parent-baby bond have not been examined. In this study, 64 couples completed questionnaires about birth, symptoms of PTSD, the couple's relationship and parent-baby bond nine weeks after childbirth. Results showed 5% of men and women had severe symptoms of PTSD. Symptoms were significantly associated within couples, and were related to similar birth factors for men and women. PTSD symptoms were not associated with the parent-baby bond or couple's relationship. The mother-baby bond was not associated with any variables measured. The father-baby bond was associated with the couple's relationship.

## **INTRODUCTION**

Research has established that between 1% and 9% of women have PTSD after birth (Ayers, 2004) and a proportion of men may also be affected (Skari et al., 2002). The effect of postnatal PTSD on the mother-infant bond and couple's relationship has been substantiated by case studies and qualitative research (Nicholls & Ayers, in press) but has not been examined quantitatively. This study therefore aimed to examine (i) what proportion of men have severe symptoms of PTSD after birth; (ii) the effect of postnatal PTSD on the parent-baby bond; (iii) the effect of postnatal PTSD on the couple's relationship; and (iv) what birth factors are associated with PTSD in men and women.

## **METHOD**

### **Design**

Postal questionnaire survey of PTSD symptoms, the couple's relationship, and parent-baby bond nine weeks after birth (mean 9.41 weeks, SD 2.06).

### **Sample**

Sixty-four couples were drawn from 207 eligible couples that had a baby at a London hospital during one month. Inclusion criteria were that couples were in a relationship together and the man attended birth. Couples were excluded if their baby was stillborn or transferred to the neonatal unit. Both members of each couple completed the questionnaires.

Participants mean age was 32.4 years (SD = 5.87); 72% were married; 66% were Caucasian. 70% of women had a normal delivery, 14% assisted delivery, 13% emergency caesarean, 3% elective caesarean; 11% had complications of delivery, and 25% had complications with the baby during delivery. Responders did not differ from non-responders on obstetric variables.

### **Measures**

PTSD symptoms (intrusions and avoidance) were measured using the Impact of Events Scale (Horowitz et al., 1979). Scores of >19 are classified as severe symptoms, giving a sensitivity of .94 and a specificity of .33. Experience of birth was measured using the Experience of

Birth Scale (Slade et al., 1993). Parent-infant attachment was measured using the Bethlehem Mother-Infant-Interaction Scale with the wording changed to be non-gender specific. Higher scores indicate a poorer parent-baby bond (Pearce & Ayers, 2005). Couple's relationship was measured using the Dyadic Adjustment Scale (Spanier, 1976).

## **RESULTS**

Five percent of men and women had severe symptoms of PTSD after birth. This comprised two couples where both members had severe symptoms, and two couples where one member had severe symptoms. Overall, men and women did not differ significantly on symptoms of intrusions and avoidance. However, women reported a better parent-baby bond, fewer positive emotions during birth, more support in birth, and more self-blame for things that went wrong during birth. Within couples, symptoms of intrusions and avoidance were significantly associated.

PTSD symptoms were not associated with the couple's relationship or the parent-baby bond. For men, the only median correlation above 0.2 was between the couple's relationship and the father-baby bond (0.26). For women, the only median correlation above 0.2 was between the birth experience and symptoms of PTSD (0.23). Multiple regressions found intrusions and avoidance in men and women were most consistently predicted by reports of something going wrong during birth, delivery problems, and emotions during birth ( $R^2 = .19$  to  $.41$ ,  $p < .01$ ).

- Insert Table 1 about here -

## **DISCUSSION**

Results are inconsistent with previous findings that more women than men have PTSD after birth and other traumatic events. The lack of effect of postnatal PTSD on the parent-baby bond and couple's relationship also contradicts previous research. One explanation is that interference with the couple's relationship and parent-baby bond is not general to all psychopathology but specific to depression. Alternatively, the effect of PTSD on relationships may not be apparent nine weeks after birth. Similarly, effects on the couple's relationship might only be detected when examining *change* in the relationship compared to

before birth. Finally, the measure of parent-baby bond used in this study focused on behavioral aspects of the parent-baby bond, rather than emotional aspects, so could have measured 'care of the baby' rather than emotional bond.

In conclusion, this research suggests postnatal PTSD symptoms are not related to the parent-baby bond or couple's relationship. However, further research should examine the long-term impact of PTSD on change in couples' relationships and emotional attachment to the baby.

## REFERENCES

- Ayers, S. (2004). Delivery as a traumatic event: prevalence, risk factors, and treatment for postnatal posttraumatic stress disorder. *Clinical Obstetrics & Gynecology*, **47**, 552-567.
- Horowitz, M.J., Wilner, N., & Alvarez, W. (1979). Impact of event scale: a measure of subjective stress. *Psychosomatic Medicine*, **41**, 209-218.
- Nicholls, K., & Ayers, S. (in press). Childbirth-related post-traumatic stress disorder in couples: a qualitative study. *British Journal of Health Psychology*.
- Pearce, H., & Ayers, S. (2005) The expected child versus the actual child: Implications for the mother-baby bond. *Journal of Reproductive & Infant Psychology*, **23**, 1-15.
- Skari, H., Skreden, M., Malt, U. F., et al. (2002). Comparative levels of psychological distress, stress symptoms, depression and anxiety after childbirth--a prospective population-based study of mothers and fathers. *British Journal of Obstetrics and Gynaecology*, **109**, 1154-63
- Slade, P., MacPherson, S.A., Hume, A., & Maresh, M. (1993). Expectations, experiences and satisfaction with labour. *British Journal of Clinical Psychology*, **32**, 469-483.
- Spanier, G. (1976). Measuring dyadic adjustment: New scales for assessing the quality of marriage and similar dyads. *Journal of Marriage and the Family*, **38**, 15-28.

**Table 1. Mean scores (columns 1 & 2), correlations between men and women (column 4), and correlations between items for men (upper triangle) and women (lower triangle)**

	Mean scores		3 Wilcoxon z value (p)	4 P	PTSD		Bond		Birth Experience								Relationship					
	1 Women	2 Men			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1. Intrusions	6.58	5.48	1.02 (.31)	<b>.37</b>		<b>.69</b>	.14	.17	<b>.34</b>	.01	.13	.10	.13	<b>.30</b>	-	-	-	-	.19	-	.07	.19
2. Avoidance	3.97	3.41	0.82 (.41)	<b>.50</b>	<b>.52</b>		.23	-	<b>.33</b>	.09	.08	<b>.31</b>	.17	<b>.36</b>	.03	.04	.15	.18	.13	.08	.06	.14
3. Parent-baby bond	9.43	10.83	2.69 (.007)	.19	.00	.09		-	.18	.01	.04	-	-	-	-	.28	.17	.07	.26	.19	<b>.49</b>	.01
4. Positive emotions in birth	28.95	35.44	3.49 (<.001)	<b>.41</b>	.18	-	-	-	-	<b>.53</b>	<b>.48</b>	<b>.56</b>	.22	.14	.24	.11	<b>.40</b>	<b>.43</b>	.24	.13	<b>.30</b>	<b>.31</b>
5. Negative emotions in birth	23.31	21.09	0.78 (.43)	.10	.27	<b>.41</b>	-	-	-	-	.29	<b>.47</b>	<b>.48</b>	<b>.41</b>	.28	<b>.50</b>	-	.16	.08	-	-	-
6. Control in birth	42.03	43.93	0.81 (.42)	<b>.37</b>	.06	-	.02	<b>.45</b>	<b>.37</b>		<b>.54</b>	<b>.43</b>	<b>.32</b>	.04	<b>.46</b>	.15	<b>.52</b>	.22	.20	.16	.12	.06
7. Support in birth	16.32	13.40	3.78 (<.001)	.24	-	-	.00	.27	-	.05		-	-	-	-	-	<b>.40</b>	<b>.35</b>	.04	.17	.18	.08
8. Birth worse than expected	4.87	5.32	1.21 (.23)	<b>.48</b>	.22	<b>.37</b>	.07	-	<b>.52</b>	<b>.36</b>	.11		-	-	-	-	.23	.27	.03	-	.23	.07
9. Life threat	2.64	2.80	0.62 (.54)	<b>.30</b>	<b>.30</b>	<b>.39</b>	.19	.05	<b>.48</b>	.21	.25	.28		<b>.54</b>	<b>.34</b>	.13	.26	.25	.03	.28	.03	.01
10. Something going wrong	2.80	2.72	0.21 (.84)	<b>.69</b>	<b>.37</b>	<b>.51</b>	-	-	<b>.54</b>	.21	<b>.34</b>	<b>.49</b>	<b>.83</b>		.13	<b>.40</b>	.07	.24	.07	.12	.06	.00
11. Self-blame	3.20	0.85	2.88 (.004)	-	<b>.39</b>	<b>.42</b>	.16	.12	<b>.31</b>	.07	.14	.04	.14	<b>.44</b>		.04	-	-	-	-	-	.05
12. Other-blame	5.15	4.35	1.16 (.25)	<b>.62</b>	.07	<b>.35</b>	.31	.04	<b>.33</b>	.16	.03	.28	.18	<b>.39</b>	.28		.06	-	.06	.07	.20	.09
13. Satisfaction	7.77	8.31	1.14 (.25)	-	.08	-	.20	.25	-	.21	.05	.21	.15	.25	.13	.22		<b>.51</b>	.11	.15	.26	.03
14. satisfaction with partner's coping	8.95	9.13	0.48 (.70)	.16	.10	-	.06	.13	-	.11	<b>.37</b>	-	-	-	.10	<b>.52</b>	<b>.47</b>		.16	.28	<b>.41</b>	.07
15. Relationship consensus	51.45	49.19	1.68 (.09)	<b>.56</b>	-	-	-	.13	-	.16	<b>.36</b>	.04	-	-	.02	-	.24	<b>.33</b>		<b>.30</b>	<b>.43</b>	<b>.49</b>
16. Relationship satisfaction	40.11	41.01	1.74 (.08)	<b>.62</b>	.08	.24	.04	.29	.13	.10	<b>.41</b>	.01	.04	.15	<b>.52</b>	.01	.06	<b>.32</b>	<b>.56</b>		<b>.41</b>	.19
17. Relationship cohesion	16.00	16.69	1.81 (.07)	<b>.62</b>	.01	.01	.07	.06	.08	.05	.05	.12	.09	.01	<b>.39</b>	<b>.56</b>	.01	.27	<b>.48</b>	<b>.31</b>		<b>.30</b>
18. Relationship affection	9.19	8.71	1.25 (.21)	<b>.48</b>	-	-	-	.09	-	.12	<b>.39</b>	.05	-	-	.10	.02	.12	<b>.34</b>	<b>.55</b>	<b>.58</b>		.29

N=59 to 64 men, and 60 to 64 women; except q11 and q12 which are based on the 25 women and 24 men who allocated blame for events during birth.

All correlations above .30 (Cohen's medium *r*) are in **bold** and are significant at the 5% level (except for those involving q11 and q12 which need to be above about *r* = .40 to be significant at 5%). All correlations above .50 (Cohen's large *r*) are also **underlined**.