



City Research Online

City, University of London Institutional Repository

Citation: Chinnery, H. & Thompson, S. B. N. (2011). Normative Values for 18-30 Age Group of Benton Visual Retention Test Scores and Pre-morbid Intelligence Quotients: New Data Comparisons for Diagnosing Memory and Visual Spatial Deficits in Alzheimer's Disease and Stroke. WebmedCentral BRAIN, 2(5), WMC001918. doi: 10.9754/journal.wmc.2011.001918

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/21092/>

Link to published version: <https://doi.org/10.9754/journal.wmc.2011.001918>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Normative Values for 18-30 Age Group of Benton Visual Retention Test Scores and Pre-morbid Intelligence Quotients: New Data Comparisons for Diagnosing Memory and Visual Spatial Deficits in Alzheimer's Disease and Stroke

Corresponding Author:

Dr. Simon B Thompson,
Associate Professor, Psychology Research Centre, Bournemouth University, BH12 5BB - United Kingdom

Submitting Author:

Dr. Simon B Thompson,
Associate Professor, Psychology Research Centre, Bournemouth University, BH12 5BB - United Kingdom

Article ID: WMC001918

Article Type: Research articles

Submitted on: 06-May-2011, 01:02:18 PM GMT **Published on:** 07-May-2011, 02:04:28 PM GMT

Article URL: http://www.webmedcentral.com/article_view/1918

Subject Categories: BRAIN

Keywords: Alzheimer's Disease, Benton Visual Retention Test, Dementia, Diagnosis, Intelligence Quotient, Memory; NART, Prognosis, Stroke, Visual Spatial Deficits

How to cite the article: Thompson S B, Chinnery H . Normative Values for 18-30 Age Group of Benton Visual Retention Test Scores and Pre-morbid Intelligence Quotients: New Data Comparisons for Diagnosing Memory and Visual Spatial Deficits in Alzheimer's Disease and Stroke . WebmedCentral BRAIN 2011;2(5):WMC001918

Source(s) of Funding:

None.

Competing Interests:

None.

Normative Values for 18-30 Age Group of Benton Visual Retention Test Scores and Pre-morbid Intelligence Quotients: New Data Comparisons for Diagnosing Memory and Visual Spatial Deficits in Alzheimer's Disease and Stroke

Author(s): Thompson S B, Chinnery H

Abstract

The Benton Visual Retention Test (BVRT) is a well known test used to assess visual memory deficits and visual spatial abilities in patients. Normative data for the 18-30 age group is not presently covered thoroughly; hence, this study continues the work of previous series that examined data comprising BVRT scores, intelligence quotients (IQ), anxiety and depression levels, and gender effects. Correlations between pre-morbid estimates of IQ across different BVRT administrations were examined and discussed in order to compile a database of new data comparisons for this age group.

Introduction

This study was further progression of previous series that examined the use of the Benton Visual Retention Test (BVRT) (Benton, 1974; Benton Sivan, 1992) for the assessment of visual memory functioning and visual spatial ability (see Thompson, Ennis, Coffin & Farman, 2007; Thompson & Gander, 2011). The aim of the study was to create a comprehensive database of normative data collected from using the BVRT in participants aged between 18 and 30 years old. Previously, no normative data existed for BVRT performance for this particular age group apart from that collected from the earlier series (Thompson & Gander, 2011) and the extensive data collected across all other age groups (Benton, 1946; Benton, Eslinger & Damasio, 1981). Data has been combined from this previous study in order to construct a comprehensive data set for reference by examiners of patients with a range of memory deficits.

The normative standard section of the BVRT manual states that performance on the three parallel administrations of the BVRT correlate substantially with intelligence level (Benton Sivan, 1992). Therefore, correlations were carried out across our data to

determine comparisons.

The importance of the BVRT for particular clinical groups has been well established and documented elsewhere (Thompson, Ennis, Coffin & Farman, 2007; Thompson & Gander, 2011). In clinical settings the BVRT is often administered to determine the extent and type of visual spatial disability and visual memory deficit by way of categories of errors according to misplacement, substitution, omission, rotation, etc. This provides valuable information to the examiner especially if the patient has developed a stroke with a certain perceptual difficulty on one side of the body. It has also been a useful instrument in the neuropsychological assessment battery for the diagnosis and prognosis of Alzheimer's disease (Thompson, 2006; 2011a,b).

Methods and Materials

Methods

Rationale

The aim of this study is to gain a set of normative data for performance on the BVRT for normal individuals aged between 18 and 30 years old.

Materials

Benton Visual Retention Test

The BVRT has 3 similar (parallel) forms of task C, D and E, each consisting of 10 designs containing one or more figures. In this study, 3 types of administration were used: A ? showing the images for 10 seconds then requiring immediate reproduction of the images from memory; B ? showing the images for 5 seconds followed by immediate reproduction; and C ? showing the images for 10 seconds, then delaying participants' reproduction of the figures after a further 15 seconds. Only two different versions were used: C and E. Therefore, these conditions were listed as: A(C), A(E), B(C), B(E), C(C) and C(E).

NART

In order to establish an estimate of pre-morbid IQ, the National Adult Reading Test (Nelson, 1992) is often

administered. Benton Sivan (1992) shows BVRT scores together with pre-morbid IQs; hence, this study has collected similar data.

HADS

The Hospital Anxiety and Depression Scale (Snaith & Zigmond, 1994) was used to establish baseline anxiety and depression levels of each participant. It is known the high levels of anxiety and/or depression can affect memory functioning.

Experimental hypotheses

Based on previous experimental series, the following hypotheses were used:

H1 There will be a significant relationship between NART scores and Total Errors score and Total Correct scores on the BVRT.

H2 There will be a significant relationship between NART scores and Total Errors score and Total Correct scores across the three administrations A, B, C.

H3 There will be a significant relationship between Total Errors score and Total Correct scores on the BVRT and Anxiety and Depression scores on the HADS.

study design

In order to determine if age or gender had any significant influence on the data collected, a 3 x 2 unrelated ANOVA was implemented to investigate a significant difference in performance on the BVRT between genders and age groups. The first independent variable was gender, which had two levels, Males and Females. The second independent variable was age group, and was split into 3 levels, 18-20 (mode = 19, median = 19), 21-33 (mode = 21, median = 21), and 24-27 (mode 26, median = 26). The dependent variables were Total Errors score and Total Correct score.

Participants

Fifty undergraduate students attending Bournemouth University were recruited on a volunteer basis. Fourteen were male, 36 were female. Prior to the study, ethics approval was obtained from Bournemouth University Research & Ethics Committee (November 2009) and consent was obtained from each participant according to strict ethical guidelines.

Results

Statistics

Spearman's Rank Order correlation was used to determine the relationship between NART scores with Anxiety and also with Depression, as assessed by the HADS. No statistical evidence was found between

these measures thus refuting H3 (NART plus Anxiety: $r = -.137$, $n = 53$, $p > .05$ two tailed; and NART plus Depression: $r = -.219$, $n = 53$, $p > .05$ two tailed).

No statistical evidence was found between the NART and Total Correct scores on administration A of the BVRT ($r = -.125$, $n = 53$, $p > .05$ two tailed) and also Total Errors scores ($r = .113$, $n = 53$, $p > .05$ two tailed). However, significant evidence of a negative correlation was found between the NART and BVRT Total Correct scores on administration B ($r = -.318$, $n = 53$, p

A significant positive correlation at .05 level was found between NART and BVRT Errors scores on administration B ($r = .318$, $n = 53$, $p < .05$ two tailed). A further positive correlation at .01 level was found between the NART and BVRT Errors scores on administration C ($r = .376$, $n = 53$, $p < .01$ one tailed) (Illustration 1).

A reasonable relationship between NART scores and BVRT Total Correct and Total Error scores on administration B (Illustrations 2 & 3) and administration C (Illustrations 4 & 5) were observed.

A table of statistical comparisons is presented showing support for H1 and H2 (Illustrations 6 & 7).

Discussion & Conclusions

The results obtained from this study support hypotheses H1 and H2 and refute H3. Hence, there are significant correlations between the NART scores and administrations of the BVRT. This allows researchers to refer to the database of correlations for the 18 ? 30 age group. It also allows clinicians to examine patients in these age ranges; previously, this test was constrained due to the lack of comprehensive data in this domain.

Clinical disorders span all age ranges; therefore, it is important to have normative data across the lifespan. By building a picture of abilities across ages, we are in a better position to judge whether or not the patient we are examining is performing below,? above or within their comparable age range. The picture of Alzheimer?s disease and vascular dementia is complicated (Thompson, 2002; Thompson, 2011c), and if we can make sense of people?s abilities across ages then we are better placed to understand the effects of diseases such as the dementias.

References

1. Benton, A.L., 1946. A visual retention test for clinical

- use. New York, The Psychological Corporation.
2. Benton, A.L., 1974. Revised Visual Retention Test: clinical and experimental applications. 4th ed. San Antonio, The Psychological Corporation.
 3. Benton Sivan, A., 1992. Benton Visual Retention Test. 5th ed. San Antonio, The Psychological Corporation.
 4. Benton, A.L., Eslinger, P.J., & Damasio, A.R., 1981. Normative observations on neuropsychological test performances in old age. *Journal of Clinical Neuropsychology*, 3, 33-42.
 5. Nelson, R., 1992. National Adult Reading Test. Revised. Manual. Oxford, NFER-NELSON.
 6. Snaith, R.P., & Zigmond, A.S., 1994. Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, 67, 361-370.
 7. Thompson, S.B.N., 2002. Memory decline, Alzheimer's disease and vascular dementia: the clinical picture. *Journal of Cognitive Rehabilitation*, 20(2), 12-18.
 8. Thompson, S.B.N., 2006. Dementia and memory: a handbook for students and professionals. Aldershot, Ashgate.
 9. Thompson, S.B.N., 2011a. Psychology of trauma: clinical reviews, case histories, research. Portsmouth, Blackwell-Harvard-Academic.
 10. Thompson, S.B.N., 2011b. Advancing knowledge into the clinical assessment of dementia. Extensive review and design of a neuropsychological test battery for assessment. Saarbrücken, Lambert Academic Publishing GmbH & Co. KG.
 11. Thompson, S.B.N., 2011c. Alzheimer's disease: comprehensive aetiology, diagnosis, assessment recommendations and treatment. *WebmedCentral AGING*, 2(3), 1-42.
 12. Thompson, S.B.N., Ennis, E., Coffin, T., & Farman, S., 2007. Design and evaluation of a computerised version of the Benton Visual Retention Test. *Computers in Human Behavior*, 23, 2383-2393.
 13. Thompson, S.B.N., & Gander, J., 2011. Immediate memory functioning and intelligence quotients of 18-30 years age group using new data derived from the Benton Visual Retention Test: applicability to Alzheimer's disease patients. *WebmedCentral GERIATRIC MEDICINE*, 2(3), 1-28.

Illustrations

Illustration 1

Spearman's rank order correlation between NART scores and BVRT Total Correct and Total Error scores on administrations A, B and C

	Rs	P
BVRT – A(C) and NART	-.125	.371
BVRT – A (E) and NART	.113	.420
BVRT – B(C) and NART	-.318	.020*
BVRT – B(E) and NART	.318	.020*
BVRT – C(C) and NART	-.325	.017*
BVRT – C(E) and NART	.376	.005**

*Correlation is significant at the 0.05 level (two-tailed)

**Correlation is significant at the 0.01 level (two tailed)

Illustration 2

BVRT administration B(C) Total Correct scores and NART scores




Illustration 3

BVRT administration B(E) Total Error scores and NART scores




Illustration 4

BVRT administration C(C) Total Correct scores and NART scores




Illustration 5

BVRT administration C(E) Total Error scores and NART scores




Illustration 6

Spearman's rank correlation across all measures

Correlations

			M/F	Anx	Dep	NART	A(C)	A(E)	B(C)	B(E)	C(C)	C(E)
Spearman's rho	M/F	Correlation Coefficient	1.000	.188	.246	-.253	.230	-.231	.187	-.141	.097	-.091
		Sig. (2-tailed)	.	.178	.075	.068	.098	.096	.181	.313	.489	.519
	Anx	N	53	53	53	53	53	53	53	53	53	53
		Correlation Coefficient	.188	1.000	.552	-.137	-.219	.248	-.165	.157	-.169	.196
	Sig. (2-tailed)	.178	.	.000	.327	.116	.073	.239	.261	.226	.160	
	Dep	N	53	53	53	53	53	53	53	53	53	53
		Correlation Coefficient	.246	.552	1.000	-.219	-.154	.171	-.089	.103	.024	.018
	Sig. (2-tailed)	.075	.000	.	.116	.271	.220	.525	.465	.866	.897	
	NART	N	53	53	53	53	53	53	53	53	53	53
		Correlation Coefficient	-.253	-.137	-.219	1.000	-.125	.113	-.318	.318	-.325	.376**
	Sig. (2-tailed)	.068	.327	.116	.	.371	.420	.020	.020	.017	.005	
	A(C)	N	53	53	53	53	53	53	53	53	53	53
		Correlation Coefficient	.230	-.219	-.154	-.125	1.000	-.945	.466	-.465	.511	-.432*
	Sig. (2-tailed)	.098	.116	.271	.371	.	.000	.000	.000	.000	.001	
	A(E)	N	53	53	53	53	53	53	53	53	53	53
		Correlation Coefficient	-.231	.248	.171	.113	-.945	1.000	-.457	.443	-.520	.464**
	Sig. (2-tailed)	.096	.073	.220	.420	.000	.	.001	.001	.000	.000	

Illustration 7

Spearman's rank correlation across all measures (continued)

Correlation Coefficient	-.09	.196	.018	.376**	-.432**	.464**	-.512**	.499**	-.966**	1.00
CE	1									0
Sig. (2-tailed)	.519	.160	.897	.005	.001	.000	.000	.000	.000	.
N	53	53	53	53	53	53	53	53	53	53

** . Correlation is significant at the 0.01 level (2-tailed).

* . Correlation is significant at the 0.05 level (2-tailed).

Descriptive Statistics

	N	Minimum	Maximum	Mean	Std. Deviation
Gender	53	1.0	2.0	1.547	.5025
Anxiety	53	1.0	19.0	7.698	4.2634
Depression	53	.0	18.0	4.113	3.4006
Valid N (listwise)	53				

Disclaimer

This article has been downloaded from WebmedCentral. With our unique author driven post publication peer review, contents posted on this web portal do not undergo any prepublication peer or editorial review. It is completely the responsibility of the authors to ensure not only scientific and ethical standards of the manuscript but also its grammatical accuracy. Authors must ensure that they obtain all the necessary permissions before submitting any information that requires obtaining a consent or approval from a third party. Authors should also ensure not to submit any information which they do not have the copyright of or of which they have transferred the copyrights to a third party.

Contents on WebmedCentral are purely for biomedical researchers and scientists. They are not meant to cater to the needs of an individual patient. The web portal or any content(s) therein is neither designed to support, nor replace, the relationship that exists between a patient/site visitor and his/her physician. Your use of the WebmedCentral site and its contents is entirely at your own risk. We do not take any responsibility for any harm that you may suffer or inflict on a third person by following the contents of this website.