



City Research Online

City, University of London Institutional Repository

Citation: Foye, U., Hazlett, D. E. & Irving, P. (2019). Exploring the role of Emotional Intelligence on disorder eating psychopathology. *Eating and Weight Disorders-Studies on Anorexia Bulimia and Obesity*, 24(2), pp. 299-306. doi: 10.1007/s40519-018-0629-4

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/21151/>

Link to published version: <https://doi.org/10.1007/s40519-018-0629-4>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.



Exploring the role of emotional intelligence on disorder eating psychopathology

Una Foye^{1,2} · D. E. Hazlett³ · Pauline Irving⁴

Received: 25 April 2018 / Accepted: 13 December 2018
© The Author(s) 2018

Abstract

Purpose This study aims to explore the role of Emotional Intelligence (EI) and specific facets that may underpin the aetiology of disordered eating attitudes and behaviours, as a means to understand what aspects of these deficits to target within treatments.

Methods Participants were recruited from the UK and Ireland. Among the sample of 355 participants, 84% were women and 16% were men. Regarding age, 59% were between 18 and 29, 30% were between 30 and 49, and 11% were 50 or older. Using a cross-sectional design, participants completed the Schutte Self-Report Emotional Intelligence Test to measure levels of trait EI and The Eating Attitudes Test (EAT-26) as a measure of eating disorder risk and the presence of disordered eating attitudes.

Results EAT-26 scores were negatively correlated with total EI scores and with the following EI subscales: appraisal of own emotions, regulation of emotions, utilisation of emotions, and optimism. Also, compared to those without an eating disorder history, participants who reported having had an eating disorder had significantly lower total EI scores and lower scores on four EI subscales: appraisal of others emotions, appraisal of own emotions, regulation of emotions, and optimism.

Conclusions Considering these findings, EI (especially appraisal of own emotions, regulation of emotions, and optimism) may need to be addressed by interventions and treatments for eating disorders.

Level of evidence Level V, descriptive cross-sectional study.

Keywords Emotion · Emotional Intelligence · Psychopathology

Introduction

Salovey and Mayer [1] described emotional intelligence (EI) as: "...the ability to perceive emotions, to access and generate emotions so as to assist thought, to understand emotions and emotional knowledge, and to reflectively regulate emotions so as to promote emotional and intellectual growth." (p. 5).

Studies have shown that psychological health is related to an individual's emotional competencies and functioning [2]. Studies show that higher levels of EI are linked to both psychological and physical health, including findings that have found positive effects on psychological well-being and physical health, e.g. immune systems functioning [3]. As individuals with high EI have increased emotional resources, they are better equipped to deal with difficult life events; the emotionally intelligent individual is able to perceive, interpret and understand emotional and situational cues better than those with low EI and, thus, are more prepared

This article is part of the topical collection on Personality and eating and weight disorders.

✉ Una Foye
foye-u@email.ulster.ac.uk

¹ Centre of Psychiatry, Queen Mary University of London, London, UK

² Centre for Mental Health Research, School of Health Sciences, City, University of London, Northampton Square, London EC1V 0HB, UK

³ Centre of Higher Education Research and Practice, Ulster University, Jordanstown Campus, Newtownabbey, Co. Antrim, UK

⁴ Faculty of Social Sciences, Ulster University, Jordanstown Campus, Newtownabbey, Co. Antrim, UK

and able to manage and effectively regulate emotional states and engage with adaptive problem solving [4]. Conversely, those who are low in EI are unable to effectively understand, regulate and control their emotions, therefore, will be less effective in coping suggesting that they are more likely to engage in maladaptive coping or emotional regulation strategies [5]. This has been shown across studies in which participants with low levels of EI have been found to be more likely to engage in maladaptive health behaviours as an attempt to seek alternative methods of mood control and emotional expression such as smoking [6], alcohol use [7] and self-harm behaviours [8].

Emotional intelligence and eating disorders

The role of emotions in eating disorders is not new. Within her seminal works, Hilde Bruch was among the first to address the role of emotional disturbances as smokescreen to the emotional difficulties faced by those suffering from an eating disorder [9]. Explorations of the eating disorder experience have shown the important role of the eating disorder as having a functional purpose related to emotional regulation and coping [10–14]. These findings have been further expanded and focused to show deficits within this population in emotional processing, regulation and awareness [15–23].

In addition, literature has found evidence for deficits within areas of emotional processing including understanding one's own emotions and facial processing in relation to appraising others emotions [24–26]. Somatic awareness and arousal have also been reported as dampened within these groups [27]. Based on this understanding that emotional deficits are strongly associated with disordered eating, a range of emotion-focused treatments have been explored with evidence supporting the acceptability and feasibility this approach for anorexia [28].

These findings correlate with the well-established relationship that exists within the literature between eating disorders and the psychological construct of Alexithymia, defined as a construct meaning literally "...without words for emotions" which described a severe deficit in an individual's representation of emotions symbolically and the understanding, processing and describing of emotions [29, 30]. Empirical findings have formulated this construct to include features such as difficulty in describing and identifying feelings, difficulty in distinguishing between feelings and bodily sensations of emotional arousal, constricted imaginative processes, e.g. paucity of fantasies and externally oriented cognitive style [31]. The highest rates of alexithymia have been found in eating disorder patients with rates of alexithymia within anorexia nervosa patients between 23–77%, and 40–63% in bulimia nervosa patients; compared to rates of 0–28% in non-clinical samples [32–34]. With eating disorders being highly related to alexithymia, this may provide

insight into the low success rates within treatments and psychotherapy as clients with high alexithymia scores have been found to be less able to respond to such interventions due to low insight and inability of emotional expression [35, 36].

While alexithymia research has found significant evidence for its role as a maintenance factor within eating disorder models, these findings express severe dysfunction and clinical levels of eating disorders, thus continuing the trend in eating disorder literature to view these behaviours as polarised. The introduction of a spectrum of emotional functioning, and dysfunction, across the range of emotional aspects described above can be found within the theoretical concept of EI.

The core elements of EI can be closely related to the emotional deficits that have been found within eating disorder populations. This suggests that individuals exhibiting disordered eating attitudes and behaviours, who experience such deficits in their emotional resources, have low levels of EI. Studies have provided evidence towards this application with results showing that those with high disordered eating have significantly lower EI than general populations [37–45]. The findings across these papers show that the relationship between EI and disordered eating spans across adults and adolescents, as well as occurring within sporting populations. These findings support theories linking negative affect and emotion dysregulation with the development of eating disorders, highlighting potential benefits of utilising EI for the identification of at-risk populations or as a potential screening measure for the issue. Furthermore, results highlight the role of emotion in disordered eating attitudes, which is an important finding in terms of the prevention and management of disordered eating.

While these studies provide strong evidence of the relationship between EI and eating disorders, the nature of what elements of EI are salient remains unclear. While there is theoretical recognition that emotional aspects of eating disorders encompass the interpersonal and intrapersonal [16], the majority of the literature places a focus on recognition and coping [12–15, 19, 21, 46] and to a lesser extent aspects of recognition and awareness [20, 23, 24]. Absent from this literature are facets of EI optimism and utilisation of emotion. Utilisation of emotions has been defined as the components of flexible planning, creative thinking, redirected attention and motivation [47] and optimism includes expectations of control over one's own positive results in the future as well as a component of personal effectiveness [48]. To apply recommendations of literature surrounding the entire EI construct as an intervention for screening, treatment and prevention [37–45] requires further investigation of these facets that have been missed within the literature to ensure that programmes developed are able to utilise the correct facets and target the aspects of EI that are relevant.

Aim

The application of emotional dysfunction theories may provide insight into why treatment for eating disorders remains blighted by the high levels of ambivalence, treatment drop-outs and relapse rates that are observed within this patient population [49]. As a result, these disorders observe high rates of chronicity and become incredibly difficult to treat as they are so deeply engrained [50]. The aim of this study is to assess the link between trait EI and disordered eating attitudes to understand the nature of the relationship reported in the literature.

We hypothesise that disordered eating attitudes will be associated with lower EI scores; however, hypothesis that this relationship will be driven by core facets of the EI definition around management and regulation of emotions with facets related to the ability to access and express emotions remains less relevant. We believe understanding the nature of the relationship between EI and eating disorders will allow treatments to focus more effectively on the core facets of the disorder rather than more broadly with the vast area of emotions.

Methodology

The findings reported within this paper are part of a larger UK-based study which explored the variables and barriers impacting on the onset and maintenance of eating disorders. The study used a sequential, mixed methods design made up of two phases. Following this research design, the current paper will explore the data emerging related to emotional functioning.

Participants

Participants were recruited in the UK and Ireland from two local universities and a local further education institution, three community groups focused on the area of eating disorders, and five general health organisations. To ensure anonymity data collected, we did not note where participants were recruited from. Groups were encouraged to share the study link for additional snowball sampling [51]. Exclusion criteria included participants under 16 years of age, individuals in treatment, or those living outside of the UK or Ireland. From these sampling procedures, 435 agree to participate, and of them, 355 completed all survey measures that were included in the data analyses. Among this final sample of 355, 84% were women and 16% were men. Regarding age, 59% were between 18 and 29, 30% were between 30 and 49, and 11% were 50 or older. From this final sample, 23% identified themselves as having had an eating disorder, with 29 reporting more

than one of these disorders at different points in time (41 with anorexia, 21 with bulimia, 28 with binge eating disorder, and 29 with eating disorder not otherwise specified). Of these 83 individuals who reported having had an eating disorder, only 42 reported having received treatment, 20 received treatment for anorexia ($n=20$), five for EDNOS, three for bulimia, three for binge eating disorder and 11 for complex, multiple or atypical presentations. Females accounted for the majority of those having received treatment ($n=41$) with only one male reporting having received treatment for their eating disorder. Due to the unequal distribution between eating disorder experiences, non-parametric testing was used to carry out the data analysis related to this variable.

Questionnaires and procedure

Institutional ethical approval was granted by Ulster University's Research Ethics Committee prior to the study commencing. The questionnaires below, along with a demographic survey, were placed online for completion by participants with a web link to allow the study to be distributed widely. When participants accessed the link to the survey, they were first required to give informed consent.

Schutte Self-Report Emotional Intelligence Test

EI was measured using the Schutte Self Report EI Test [47]. This measure was selected as this measure of EI is one of the most widely used scales and can be used by a wide variety of participant groups. The SSEIT is a 33-item measure that uses a 5-point likert scale to measure levels of trait EI. The measure represents all portions of the Salovey and Mayer model of EI [1], with 13 items representing the appraisal and expression of emotion category, e.g. I know when to speak about my personal problems to others, 10 items representing the regulation of emotion, e.g. I have control over my emotions and 10 items representing the utilisation of emotion, e.g. I motivate myself by imagining a good outcome to tasks I take on. This measure has been widely used within the literature testing Emotional Intelligence levels. Reports show the measure to have an internal consistency of 0.90, with a 2-week test-retest reliability of 0.78 from the same sample [47] with further validation studies carried out by Austin et al. [52]. The Cronbach alpha coefficient was 0.897. Scored ranged between 67 and 162 with a mean of 119.738 (SD 14.251, trimmed mean 120.050).

The EI scale was also recorded using the subscales, according to the analysis by Lane et al. [53] to provide a scoring for six facets of EI; Appraisal of Others Emotions consisting of seven items and an internal consistency of

0.79, Appraisal of Own Emotions consisting of five items and an internal consistency of 0.77, Regulation consisting of five items and an internal consistency of 0.6, Social Skills consisting of five items and an internal consistency of 0.62, Utilisation of Emotions consisting of seven items and an internal consistency of 0.67, and Optimism consisting of four items and an internal consistency of 0.66.

Eating Attitudes Test (EAT-26)

The Eating Attitudes Test (EAT-26) [54] was selected as a measure of eating disorder risk and presence of disordered eating attitudes as this is the most widely utilised means of disordered eating. The EAT-26 is a non-clinical self-report 26 item measure of individuals disordered eating attitudes. This measure has replaced the 40 item Eating Attitudes Test as one of the most widely used measures of disordered eating, including questions related to dieting and pre-occupation with being thinner; “I am terrified about being overweight”, bulimia and food pre-occupation; “I have the impulse to vomit after meals”, and oral control; “I avoid eating when I am hungry”. Questions are scored on a 4 point likert scale ranging from Always scored as 3, Usually scored as 2, Often scored as 1 and Never as 0; one item is reverse scored with Never scoring 3 and Always scoring 0. Scores at or higher than 20 indicate high levels of concern about dieting and possible disordered eating in non-clinical samples. Research has shown that the EAT-26 has an accuracy rate of at least 90% when used to differentially diagnose those with and without eating disorders and that mean EAT-26 scores differed among eating-disordered, symptomatic, and asymptomatic participants [55]. The measure has demonstrated internal consistency of between 0.79 and 0.94 [54, 55].

Data analysis

Data collected from the study were analysed using SPSS 21.0. Preliminary statistical tests to identify linearity, outliers and distribution were carried out to test the validity of

these data. Pearson correlation analyses were conducted to explore the relationships between EAT-26 and all EI measures, first for all participants combined and then separately for participants with and without an eating disorder history. To further understand these relationships, a series of Mann–Whitney *U* tests were conducted to examine differences in the EI measures between those with and without an eating disorder history.

Results

Preliminary analyses revealed that those with a history of having an eating disorder had significantly higher EAT-26 scores ($M=35.46$) than those with no history of an eating disorder ($M=17.13$), $U=3431.00$, $Z=-9.22$, $p<0.001$. There was no significant difference in EAT-26 scores between those who had treatment for their eating disorder and those who had no treatment.

Table 1 shows the results of the Pearson correlation analyses that tested the relationships between EAT-26 scores and both the EI total score and EI subscale scores. For all participants combined, whereas there was a significant negative correlation between EAT-26 scores and total EI score, this relationship remained significant for only four facets of the EI construct: appraisal of own emotions, emotional regulation, emotional utilisation, and optimism. As indicated previously, these analyses were conducted again but separately for the different groups. For those who reported having had an eating disorder, these same relationships remained significant. For those with no history of an eating disorder, the relationships with emotional regulation and emotional utilisation were no longer significant.

Results of the Mann–Whitney *U* analyses that tested whether the EI total and subscale scores differed between groups. Compared to those without an eating disorder history, participants who reported having had an eating disorder had significantly lower total EI scores and lower scores on four facets of EI: appraisal of others emotions, appraisal of own emotions, emotional regulation, and optimism.

Table 1 Correlations between EAT-26 scores and EI subscale scores

Scale	Total sample ($n=355$)	Group 1: history of an ED ($n=83$)	Group 2: no ED ($n=272$)
Overall EI	$r = -0.221, p < 0.001$	$r = -0.371, p < 0.001^*$	$r = -0.118, p = 0.064$
Appraisal of Own Emotions	$r = -0.332, p < 0.001^*$	$r = -0.363, p = 0.09^*$	$r = -0.280, p < 0.001^*$
Emotional Regulation	$r = -0.335, p < 0.001^*$	$r = -0.477, p < 0.001^*$	$r = 0.085, p = 0.2840.05$
Emotional Utilisation	$r = -0.272, p = 0.005^*$	$r = -0.346, p = 0.004^*$	$r = 0.030, p = 0.656$
Optimism	$r = -0.059, p < 0.001^*$	$r = -0.379, p = 0.001^*$	$r = -0.131, p = 0.033^*$
Appraisal of Others Emotions	$r = 0.059, p = 0.291$	$r = 0.158, p = 0.171$	$r = -0.061, p = 0.338$
Social Skills	$r = -0.056, p = 0.313$	$r = -0.092, p = 0.426$	$r = 0.008, p = 0.898$

*Indicate a significant result

Discussion

Currently, there is no clear aetiological model for eating disorders, with a wealth of causal models suggested to explain their onset [56]. With a considerable range of biological, psychological and social theories evidenced within the wide range of literature, the results from this study provide insights into the aspects of emotional dysfunction that may influence the psychopathology of disordered eating.

The results of this study found that individuals with high EI had significantly lower disordered eating attitudes, while those with low EI had significantly higher levels of disordered eating attitudes. This supports previous research in which EI is strongly related to disordered eating attitudes and behaviours with low EI scores predictive of higher levels of disordered eating [37]. These findings suggest that EI skills may provide protection against the development of disordered eating attitudes while their absence create a vulnerability. These findings support previous literature that has found this relationship between disordered eating and EI [44, 45, 57–59]. These results add to our understanding of the

nature of this relationship and leads to questions regarding how low EI may contribute to the development of eating disorders.

Overall, there is a depth of research demonstrating a salient relationship between EI and disordered eating attitudes as well as the emerging evidence of the specific facets of EI that may contribute to this relationship. This may present an opportunity to tackle these illnesses from a secondary intervention perspective. Secondary prevention can produce targeted programs in which EI skills such as problem-solving can be built alongside emotional psychoeducation as a means to build coping mechanisms and resilience. By changing perspective from interventions to remove the triggers and difficulties that create risk for individuals, the evidence that high EI may protect against onset of disordered eating attitudes provides the opportunity to build the individual to cope with and manage the risk.

This modelling may help to explain why some triggers such as low self-esteem are managed effectively by the majority of the general population with no negative impact; however, a small number will go on to develop an eating disorder. It may be inferred that EI can act as a buffer to




Fig. 1 Model exploring the development of disordered eating attitudes and behaviours

moderate against the harmful negative effects of triggering events or low self-esteem, as modelling in Fig. 1. Those with low EI may become vulnerable due to the lack of emotional resources at their disposal, leaving individuals unable to cope, therefore, an alternative is sought [5]. As found in previous qualitative examination of emotional intelligence for eating disordered individuals, it was found that the emotional function that the eating disorder provided acted as a mediator between more normative disordered eating attitudes and the development of more clinical and dangerous symptomology [46]. This supports our understanding of eating disorders as associated with coping and regulation of negative emotion previously explored within the literature [10–12].

Examination of the EI subscales was carried out to provide additional understanding of how EI deficits impact on eating disorder development and maintenance. Appraisal of one's own emotions, emotional regulation and optimism were found to be significantly lower in all measures of disordered eating, with emotional utilisation significantly lower in relation to attitudes and behaviour scores, suggesting that these elements of EI may be more important in understanding eating disorder aetiology. This is supported by previous research within the field which has suggested that eating disorders occur as a result of emotional dysfunction and regulation difficulties [6–8, 13, 25]. This suggests that disordered eating behaviours may occur due to an individual's inability to regulate their emotions in response to stressors or triggering events. For example, while low self-esteem is widely considered a core feature that underpins disordered eating psychopathology [60], the co-occurring low abilities to regulate, express and manage emotions have the role to act as a vulnerability factor for individuals with low self-esteem to engage with disordered eating as a form of coping.

Limitations

It must be acknowledged that the study faced a number of limitations that should be accounted for and addressed within future research. The cross-sectional design of this study limits the conclusions that may be made regarding whether EI deficits contribute to either the onset or the maintenance of eating disorders. As a result, participants included in the analysis are not well distributed or representative in relation to gender, age or history of clinical diagnosis. Future work should consider exploring this relationship in populations that are often underrepresented, e.g. males, as well as replicating this work in clinical samples to further understand the potential unique role EI plays for this group. Furthermore, due to the skewed distribution of EAT-26 scores and population sizes being considerably smaller for the clinical sample, analysis remains simplistic and underpowered

to perform mediation analysis. Further research is necessary to apply the theoretical modelling outcome from this study into mediation analysis to further understand the nature of this pathway. A further limitation to address is the use of self-report measurements. All the measurement tools used in the study used self-report scales in which participants reflected on their own perception of their skills, attitudes and behaviours. While each of the tools used has been previously validated, research has shown that the individual's perception of themselves and their skills are often incompatible to their true abilities.

Conclusion

The role of EI has been well identified in the literature with these findings showing that lower EI correlates with the development of eating disorders. The results from this study uncovered that some core elements of the EI construct such as appraisal of one's own emotions and emotional regulation of emotions may be salient to this relationship than other aspects of EI. By understanding the nature of this relationship between EI and eating disorders, and the core aspects of EI that are related to disordered eating, this will support targeted interventions that may support treatments to focus more effectively on the core facets of the disorder rather than more broadly with the vast area of emotions.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Institutional ethical approval was granted prior to the study commencing.

Informed consent Informed consent was obtained from all individual participants included in the study.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Salovey P, Mayer JD (1990) Emotional intelligence. *Imagin Cogn Personal* 9(3):185–211. <https://doi.org/10.2190/DUGG-P24E-52WK-6CDG>
- Tsaousis I, Nikolaou I (2005) Exploring the relationship of emotional intelligence with physical and psychological health functioning. *Stress Health* 21:77–86. <https://doi.org/10.1002/smi.1042>
- Day AL, Therrien DL, Carroll SA (2005) Predicting psychological health: assessing the incremental validity of emotional intelligence beyond personality, type A behaviour, and daily hassles. *Eur J Personal* 19(6):519–536. <https://doi.org/10.1002/per.552>
- Matthews G, Emo A, Funke G, Zeidner M, Roberts RD (2006) Emotional intelligence, personality, and task-induced stress. *J Exp Psychol* 12(2):96–107. <https://doi.org/10.1037/1076-898X.12.2.96>
- Saklofske DH, Austin EJ, Galloway J, Davidson K (2007) Individual difference correlates of health related behaviours: preliminary evidence for links between emotional intelligence and coping. *Person Individ Differ* 42:491–502. <https://doi.org/10.1016/j.paid.2006.08.006>
- Trinidad DR, Unger JB, Chou CP, Johnson CA (2004) The protective association of emotional intelligence with psychosocial smoking risk factors for adolescents. *Personal Individ Differ* 36:945–954. [https://doi.org/10.1016/S0191-8869\(03\)00163-6](https://doi.org/10.1016/S0191-8869(03)00163-6)
- Peterson K, Malouff J, Thorsteinsson FB (2011) A meta-analytic investigation of emotional intelligence and alcohol involvement. *Subst Use Misuse* 46(14):1726–1733. <https://doi.org/10.3109/10826084.2011.618962>
- Mikolajczak M, Petrides KV, Hurry J (2009) Adolescents choosing self-harm as an emotion regulation strategy: the protective role of trait emotional intelligence. *Br J Clin Psychol* 48:181–193. <https://doi.org/10.1348/014466508X386027>
- Bruch H (1988) *Conversations with anorexics*. Basic Books, New York
- Schmidt U, Treasure J (2006) Anorexia nervosa: valued and visible. A cognitive-interpersonal maintenance model and its implications for research and practice. *Br J Clin Psychol* 45:343–366. <https://doi.org/10.1348/014466505X53902>
- McNamara C, Chur-Hansen A, Hay PJ (2008) Emotional responses to food in adults with an eating disorder: a qualitative investigation. *Eur Eat Disord Rev* 16:115–123
- Reid M, Burr J, Williams S, Hammersley R (2008) Eating disorders patients' views on their disorder and on an outpatient service: a qualitative study. *J Health Psychol* 13(7):956–960
- Bento S, Ferreira C, Mendes AL, Marta-Simões J (2017) Emotion regulation and disordered eating: the distinct effects of body image-related cognitive fusion and body appreciation. *Psychologica* 60:11–25. <https://doi.org/10.14195/1647-8606>
- Southward MW, Christensen KA, Fettich KC, Weissman J, Berona J, Chen EY (2014) Loneliness mediates the relationship between emotion dysregulation and bulimia nervosa/binge eating disorder psychopathology in a clinical sample. *Eat Weight Disord* 19:509–513. <https://doi.org/10.1007/s40519-013-0083-2>
- Sim L, Zeman J (2005) Emotion regulation factors as mediators of bulimic symptoms in early adolescent girls. *J Early Adolesc* 25:478–496
- Treasure J, Tchanturia K, Schmidt U (2005) Developing a model of the treatment for eating disorder: using neuroscience research to examine the how rather than what of change. *Couns Psychother Res* 5(3):1–12
- Holiday J, Uher R, Landau S, Collier D, Treasure J (2006) Personality pathology among individuals with a lifetime history of AN. *J Personal Disord* 20(4):417–430
- Harrison A, Sullivan S, Tchanturia K, Treasure J (2010) Emotional functioning in eating disorders: attentional bias, emotion recognition and emotion regulation. *Psychol Med* 40(11):1887–1897
- Weinbach N, Sher H, Bohon CJ (2018) Differences in emotion regulation difficulties across types of eating disorders during adolescence. *Abnorm Child Psychol* 46:1351. <https://doi.org/10.1007/s10802-017-0365-7>
- Shouse SH, Nilsson J (2011) Self-silencing, emotional awareness, and eating behaviors in college women. *Psychol Women Q* 35(3):451–457
- Lulé D, Schulze UM, Bauer K, Schöll F, Müller S, Fladung AK, Uttner I (2014) Anorexia nervosa and its relation to depression, anxiety, alexithymia and emotional processing deficits. *Eat Weight Disord* 19(2):209–216. <https://doi.org/10.1007/s40519-014-0101-z>
- Mendes AL, Ferreira C, Marta-Simões J (2017) experiential avoidance versus decentering abilities: the role of different emotional processes on disordered eating. *Eat Weight Disord* 22:467–474. <https://doi.org/10.1007/s40519-016-0291-7>
- Kessler H, Schwarze M, Filipic S, Traue HC, von Wietersheim J (2006) Alexithymia and facial emotion recognition in patients with eating disorders. *Int J Eat Disord* 39(3):245–251
- Castro L, Davies H, Hale L, Surguladze S, Tchanturia K (2010) Facial affect recognition in AN: is obsessiveness a missing piece of the puzzle? Australian and New Zealand. *J Psychiatry* 44(12):1118–1125
- Sanchez-Ruiz MJ, El-Jor C, Abi Kharma J, Bassil M, Zeeni N (2018) Personality, emotion-related variables, and media pressure predict eating disorders via disordered eating in Lebanese university students. *Eat Weight Disord*. <https://doi.org/10.1007/s40519-017-0387-8>
- Davies H, Schmidt U, Stahl D, Tchanturia K (2010) Evoked facial emotional expression and emotional experience in people with AN. *Int J Eat Disord* 44:531–539
- Myers TA, Crowther JH (2008) Is self-objectification related to interoceptive awareness? An examination of potential mediating pathways to disordered eating attitudes. *Psychol Women Q* 32:172–180
- Sala M, Heard A, Black EA (2016) Emotion-focused treatments for anorexia nervosa: a systematic review of the literature. *Eat Weight Disord* 21:147–164. <https://doi.org/10.1007/s40519-016-0257-9>
- Sifneos PE (1973) The prevalence of 'alexithymic' characteristics in psychosomatic patients. *Psychother Psychosom* 22(2):255–262
- Apfel RJ, Sifneos PE (1979) Alexithymia: concept and measurement. *Psychother Psychosom* 32:180–190
- Taylor GJ, Bagby BM, Parker JDA (1991) The alexithymia construct: a potential paradigm for psychosomatic medicine. *Psychosomatics* 32:153–164
- Bourke MP, Taylor GJ, Parker JDA, Bagby RM (1992) Alexithymia in women with AN. A preliminary investigation. *Br J Psychiatry* 161:240–243
- Taylor GJ, Parker JD, Bagby RM, Bourke MP (1996) Relationships between alexithymia and psychological characteristics associated with eating disorders. *J Psychosom Res* 41:561–568
- Parling T, Mortazavi M, Ghaderi A (2010) Alexithymia and emotional awareness in AN: time for a shift in the measurement of the concept? *Eat Behav* 11:205–210
- Horney K (1952) The paucity of inner experiences. *Am J Psychoanal* 12:3–9
- Bar-On R (2001) Emotional intelligence and self-actualization. In: Ciarrochi J, Forgas J, Mayer JD (eds) *Emotional intelligence in everyday life: a scientific inquiry*. Psychology Press, New York, pp 82–97
- Markey MA, Vander Wal JS (2007) The role of emotional intelligence and negative affect in bulimic symptomatology.

- Comp Psychiatry 48:458–464. <https://doi.org/10.1016/j.comppsy.2007.05.006>
38. Costarelli V, Stamou D (2009) Emotional intelligence, body image and disordered eating attitudes in combat sports athlete. *J Exerc Sci Fit* 7:104–111
 39. Swami V, Begum S, Petrides KV (2010) Associations between trait emotional intelligence, actual-ideal weight discrepancy, and positive body image. *Personal Individ Differ* 49:485–489. <https://doi.org/10.1016/j.paid.2010.05.009>
 40. Zysberg L, Rubanov A (2010) Emotional intelligence and emotional eating patterns: a new insight into the antecedents of eating disorders? *J Nutr Educ Behav* 42(5):345–348
 41. Filaire E, Larue J, Rouveix M (2010) Eating behaviours in relation to emotional intelligence. *Int J Sports Med* 32:309–315
 42. Hambrook D, Brown G, Tchanturia K (2012) Emotional intelligence in AN: is anxiety a missing piece of the puzzle? *Psychiatry Res* 200:12–19. <https://doi.org/10.1016/j.psychres.2012.05.017>
 43. Zysberg L (2014) Emotional intelligence, personality, and gender as factors in disordered eating patterns. *J Health Psychol* 19:1035–1042. <https://doi.org/10.1177/1359105313483157>
 44. Cuesta-Zamora C, González-Martí I, García-López LM (2018) The role of trait emotional intelligence in body dissatisfaction and eating disorder symptoms in preadolescents and adolescents. *Personal Individ Differ* 126:1–6. <https://doi.org/10.1016/j.paid.2017.12.021>
 45. Peres V, Corcos M, Robin M, Pham-Scottez A (2018) Emotional intelligence, empathy and alexithymia in anorexia nervosa during adolescence. *Eat Weight Disord*. <https://doi.org/10.1007/s40519-018-0482-5>
 46. Foye U, Hazlett HE, Irving P (2018) ‘The body is a battleground for unwanted and unexpressed emotions’: exploring eating disorders and the role of emotional intelligence. *Eat Disord*. <https://doi.org/10.1080/10640266.2018.1517520>
 47. Schutte NS, Malouff JM, Hall LE, Haggerty D, Cooper JT, Golden C, Dornheim L (1998) Development and validation of a measure of emotional intelligence. *Personal Individ Differ* 25:167–177. [https://doi.org/10.1016/S0191-8869\(98\)00001-4](https://doi.org/10.1016/S0191-8869(98)00001-4)
 48. Gillham JE, Shatte AJ, Reivich KJ, Seligman MEP (2001) Optimism, pessimism and explanatory style. In: Chang EC (ed) *Optimism and pessimisms. Implications for theory, research and practice*. American Psychological Association, Washington, DC, pp 53–75
 49. Williams S, Reid M (2010) Understanding the experience of ambivalence in AN: the maintainer’s perspective. *Psychol Health* 25(5):551–567. <https://doi.org/10.1080/08870440802617629>
 50. Geller J, Williams K, Srikaneswaran S (2001) Clinician stance in the treatment of chronic eating disorders. *Eur Eat Disord Rev* 9:1–9
 51. Hendricks VM, Blanken P, Adriaans N (1992) Snowball sampling: a pilot study on cocaine use. IVO, Rotterdam
 52. Austin EJ, Saklofske DH, Huang SHS, McKenney D (2004) Measurement of emotional intelligence: testing and cross-validating a modified version of Schutte et al’s (1998) measure. *Personal Individ Differ* 36:555–562. [https://doi.org/10.1016/S0191-8869\(03\)00114-4](https://doi.org/10.1016/S0191-8869(03)00114-4)
 53. Lane AM, Meyer BB, Devonport TJ, Davies KA, Thelwell R, Gill GS, Diehl CDP, Wilson M, Weston N (2009) Validity of the emotional intelligence scale for use in sport. *J Sport Sci Med* 8:289–295 (PMID: 24149540, PMCID: PMC3761491)
 54. Garner DM, Olmsted MP, Bohr Y, Garfinkel PE (1982) The Eating Attitudes Test: psychometric features and clinical correlates. *Psychol Med* 12:871–878. <https://doi.org/10.1017/S003329170049163>
 55. Mintz LB, Halloran MS (2000) The Eating Attitudes Test: validation with DSM-IV eating disorder criteria. *J Personal Assess* 74(3):489–503
 56. Schmidt U (2003) Aetiology of eating disorders in the 21st century. New answers to old questions. *Eur Child Adolesc Psychiatry* 12:30–37. <https://doi.org/10.1007/s00787-003-1105-9>
 57. Costarelli V, Demerzi M, Stamou D (2009) Disordered eating attitudes in relation to body image and emotional intelligence in young women. *J Hum Nutr Diet* 22:239–245. <https://doi.org/10.1111/j.1365-277X.2009.00949.x>
 58. Pettit ML, Jacobs SC, Page KS, Porras CV (2010) An assessment of perceived emotional intelligence and eating attitudes among college students. *Am J Health Educ* 41(1):46–52. <https://doi.org/10.1080/19325037.2010.10599126>
 59. Zysberg L, Tell E (2013) Emotional intelligence, perceived control, and eating disorders. *SAGE Open* 3(3):1–7. <https://doi.org/10.1177/2158244013500285>
 60. Fairburn CG (2008) *Cognitive behavioural therapy and eating disorders*. The Guilford Press, New York

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.