



City Research Online

City, University of London Institutional Repository

Citation: Egan, C., Zhu, H., Lee, A., Sim, D., Mitry, D., Bailey, C., Johnston, R., Chakravarthy, U., Denniston, A., Tufail, A., et al (2016). The United Kingdom Diabetic Retinopathy Electronic Medical Record Users Group, Report 1: baseline characteristics and visual acuity outcomes in eyes treated with intravitreal injections of ranibizumab for diabetic macular oedema. *British Journal of Ophthalmology*, 101(1), pp. 75-80. doi: 10.1136/bjophthalmol-2016-309313

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/22083/>

Link to published version: <https://doi.org/10.1136/bjophthalmol-2016-309313>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk



OPEN ACCESS

The United Kingdom Diabetic Retinopathy Electronic Medical Record Users Group, Report 1: baseline characteristics and visual acuity outcomes in eyes treated with intravitreal injections of ranibizumab for diabetic macular oedema

Catherine Egan,^{1,2} Haogang Zhu,³ Aaron Lee,⁴ Dawn Sim,^{5,6} Danny Mity,¹ Clare Bailey,⁷ Robert Johnston,⁸ Usha Chakravarthy,⁹ Alastair Denniston,¹⁰ Adnan Tufail,¹¹ Rehna Khan,¹² Sajjad Mahmood,¹³ Geeta Menon,¹⁴ Toks Akerele,¹⁵ Louise Downey,¹⁶ Martin McKibbin,¹⁷ Atul Varma,¹⁸ Aires Lobo,¹⁹ Elizabeth Wilkinson,²⁰ Alan Fitt,²¹ Christopher Brand,²² Marie Tsaloumas,²³ Kaveri Mandal,²⁴ Vineeth Kumar,²⁵ Salim Natha,²⁶ David Crabb,²⁷
on behalf of the UK AMD and DR EMR Users Group

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjophthalmol-2016-309313>).

For numbered affiliations see end of article.

Correspondence to

Catherine Egan, Department of Medical Retina, Moorfields Eye Hospital NHS Trust, 162 City Road, London EC1V 2PD, UK; catherine.egan@moorfields.nhs.uk

Received 11 August 2016
Revised 12 October 2016
Accepted 15 November 2016

ABSTRACT

Aims To describe baseline characteristics and visual outcome for eyes treated with ranibizumab for diabetic macular oedema (DMO) from a multicentre database.

Methods Structured clinical data were anonymised and extracted from an electronic medical record from 19 participating UK centres: age at first injection, ETDRS visual acuity (VA), number of injections, ETDRS diabetic retinopathy (DR) and maculopathy grade at baseline and visits. The main outcomes were change in mean VA from baseline, number of injections and clinic visits and characteristics affecting VA change and DR grade.

Results Data from 12 989 clinic visits was collated from baseline and follow-up for 3103 eyes. Mean age at first treatment was 66 years. Mean VA (letters) for eyes followed at least 2 years was 51.1 (SD=19.3) at baseline, 54.2 (SD: 18.6) and 52.5 (SD: 19.4) at 1 and 2 years, respectively. Mean visual gain was five letters. The proportion of eyes with VA of 72 letters or better was 25% (baseline) and 33% (1 year) for treatment naïve eyes. Eyes followed for at least 6 months received a mean of 3.3 injections over a mean of 6.9 outpatient visits in 1 year.

Conclusions In a large cohort of eyes with DMO treated with ranibizumab injections in the UK, 33% of patients achieved better than or equal to 6/12 in the treated eye at 12 months compared with 25% at baseline. The mean visual gain was five letters. Eyes with excellent VA at baseline maintain good vision at 18 months.

ranibizumab prevents central vision loss and improves mean visual acuity (VA), when given at monthly or according to pro re nata intervals in eyes with DMO.

In the UK, the National Institute for Health and Care Excellence approved the use of ranibizumab for DMO in February 2013, leading to universal availability of this drug in the NHS, but only if the central retinal thickness was 400 microns or more. Delivering the therapy with a recommendation of monthly assessment and then retreatment for 'active' disease makes substantial demands on healthcare services, funding authorities and patients. A recent review of the use of ranibizumab for any indication shows a more than twofold geographic variation in usage across England.⁸ The intervention is costly and intensive, so it is important to understand what visual outcomes are achieved and how clinical trial results 'translate' into clinical practice in the 'real world'.

Clinical trials are limited by entry criteria, for example, excluding patients with uncontrolled medical conditions or severe DR, and have a limited number of trial subjects. In particular for DMO, the haemoglobin A1c and blood pressure in pivotal trials can be substantially different from 'real-world' patients,^{9 10} although a recent study suggests this may not affect outcomes of treatment.¹¹ Electronic medical record (EMR) systems, if well-designed, offer more complete, prospective, real-time data collection as a by-product of routine care. They can be designed to mandate capture of a defined minimum dataset and allow the collection of more enriched datasets. The NHS Connecting for Health's 'Do Once and Share' programme has defined EMR datasets in ophthalmology including: diabetic eye disease, the cataract national dataset and the glaucoma dataset.¹²

This study aims to report the visual outcome and define benchmark standards of care for patients treated with ranibizumab for DMO at a large number of the UK centres.

INTRODUCTION

Intravitreal injections of ranibizumab are an established therapy to treat a sight-threatening complication of diabetic retinopathy (DR), diabetic macular oedema (DMO). In the UK, the second highest annual National Health Service (NHS) expenditure for any single drug is for ranibizumab (£244 million).¹ Clinical practice was initially informed by pivotal clinical trials,²⁻⁷ which demonstrated that



CrossMark

To cite: Egan C, Zhu H, Lee A, et al. *Br J Ophthalmol* Published Online First: [please include Day Month Year] doi:10.1136/bjophthalmol-2016-309313

METHODS

Ethical approval

The lead clinician and Caldicott Guardian at each centre gave written approval for extraction of anonymised data. The study protocol was approved by the head of research governance at the lead clinical centre. This study was conducted in accordance with the Declaration of Helsinki and the UK Data Protection Act.

Data collection

Anonymised data were remotely extracted from 19 centres using the same EMR system (Medisoft Ophthalmology, Medisoft, Leeds, UK) in November 2014. Each site is the only NHS provider of DMO care to their local population and very few patients switch between providers or access care privately. Data were extracted through the EMR compulsory DR structured assessment module (see online supplementary file for detailed explanation). Demographic data were extracted from the hospital's patient administration system to the EMR.

All patients had data extracted from the time of their first DR structured assessment entry onto the EMR, including the data from the time of their first injection of ranibizumab up to the date of their last clinical entry before the data extraction on 26 November 2014.

Missing data

The EMR data extraction does not record values on visits where the EMR was not used, so no missing value substitutions were performed. The only exception to this rule was baseline VA. Some treatment centres operate '2 stop' or 'injection only' clinics, where treatment is given without measuring vision. Six centres have over 10% of injections without recording a VA measurement on the same day. For these services, the baseline VA was taken from the prior assessment visit if within 4 weeks of the injection date. This was therefore not missing data per se but reflects variation in treatment delivery. Therefore, data on number of visits represent a normalised value to allow standardised comparison between centres, rather than the precise number of attendances by the patient. Analyses of all patients initiated into the study were compared with the cohort of patients that completed follow-up.

Analysis

The primary analysis was restricted to treatment-naïve eyes undergoing ranibizumab therapy as the only treatment for DMO during the follow-up period. Secondary analyses of eyes with other treatments prior to undergoing intravitreal ranibizumab therapy, or who were treated with combination therapy, were also undertaken. Eyes that had cataract surgery within 3 months of their first ranibizumab injection or during the period of follow-up were excluded.

Eyes were assigned to two groups according to their treatment history and were analysed separately. Group 1 eyes were treatment naïve at baseline for any treatment for DMO (including intravitreal injections of any drug, macular laser treatments and vitrectomy), but could have had previous peripheral scatter retinal laser at the time of their first injection of ranibizumab and were managed solely with ranibizumab during the course of follow-up until the end of the data extraction period. DMO group 2 patients had received other treatments for diabetic macular disease prior to their first injection of ranibizumab.

RESULTS

Participants

Data were extracted on 123 968 eyes of 61 984 patients with DR. There were 33 967 male patients and 28 002 female patients (in 15 cases, gender was not recorded).

The 19 UK hospitals treated a total of 3103 eyes from 2416 patients, who received 15 537 ranibizumab injections for CIDMO during 12 989 clinic visits. No patients in this dataset received ranibizumab for non-CIDMO. Of these, 28% (n=687) patients received bilateral treatments.

The 19 sites entered their first DMO ranibizumab treatment episodes into the EMR systems during the following years: 2008 (n=4 sites), 2009 (n=1), 2010 (n=5), 2011 (n=5), 2012 (n=3), 2014 (n=1). The first recorded ranibizumab injection for DMO was dated 10 June 2008.

The mean age (at the time of the first DMO injection) was 66 years (SD 13 years). The female to male ratio was 1.45:1 (1430/986 patients). There were 268 (16% of the patients with a known diabetic status) patients with type 1 diabetes mellitus. There were 1380 patients (84%) recorded with type 2 diabetes mellitus. The rate of endophthalmitis was 0.015%.

Treatment history

There were 1584 treatment-naïve eyes in group 1. Group 2 had 1471 eyes (1112 macular laser treatments, 77 vitrectomies, 332 bevacizumab, 212 with triamcinolone). Of the patients in group 2 who had received previous treatment, the median (IQR) time between the first ranibizumab injection and the previous diabetic macular treatment was 0.6 (0.2, 1.2) years.

Visual acuity

About 34.1% of BCVA tests were converted by the EMR software at source from Snellen to letter score equivalent. The rest were recorded as ETDRS letters. Baseline VA is shown in online supplementary figure S1 for all treatment groups. There was a mean gain of five letters at 12 months when both groups were analysed.

Treatment-naïve eyes

The proportion of eyes with VA of 20/40 or better (72 or more letters) in the better-seeing eye was 25% at baseline, 33% at year 1, 24% at year 2. A more complete understanding of the change in vision is obtained by plots of VA at different time points (figure 1) demonstrating mean, and SD of the data stratified by starting VA.

Figure 2 shows the association of number of injections and VA difference between the last and first injection to the baseline VA for treatment-naïve eyes. There was no significant association between the number of injections and baseline VA, but the VA difference is negatively associated with the baseline VA.

Table 1 shows the visual change and number of injections from baseline; 17.3% of eyes gained at least 15 letters. Sixty per cent of eyes were in the 0–15 letter change from baseline.

The baseline DR grade within 4 months of the first injection did not seem to influence the average VA outcomes (see online supplementary figure S2) in the first year of treatment.

Online supplementary figure S3 highlights the change in mean VA between patients with up to 1 year of follow-up (n=1136) and those with >1 year of follow-up (n=363).

Effect of baseline characteristics on VA change

The treatment effect, quantified as the VA difference between the last and first injection, is negatively associated with baseline acuity (figure 2) for treatment-naïve eyes, there was a 'ceiling effect' for those with better vision, who showed a reduced visual gain compared with baseline. Of note, the VA change from the baseline for eyes receiving at least three ranibizumab treatments was stratified by the baseline VA of 70–100 letters (548 eyes), 55–69 letters (813 eyes), 40–54 letters (422 eyes) and 0–39 letters (403 eyes) and is illustrated in figure 1. Eyes in




Figure 1 Visual acuity (VA) change in letters from baseline stratified by starting vision in ETDRS letter for treatment-naïve eyes (group 1).




Figure 2 The association of the number of injections and visual acuity difference between the last and first injection to the baseline visual acuity for the treatment-naïve eyes.

the 55–70 letter group gained a mean of 5 letters at month 12 but not at month 18, and in the 40–55 letter group gained a mean of 10 letters, which was maintained at month 12 and at month 18. The poor baseline VA group (0–40 letters) gained a mean of 12 letters at month 12, but the VA results from 12 to 18 months, while showing mean gains, were highly variable. Figure 3 shows mean VA versus months since the first injection.

Number of injections and visits

The frequency of ranibizumab injections at various treatment lengths is shown in online supplementary figure S4. Eyes

followed-up for at least 6 months received a mean of 3.3 injections. The mean number of outpatient visits (normalised to allow comparison of one stop and two stop services as discussed in the Methods section) in the first year of follow-up was 6.9 visits.

DISCUSSION

This study confirms that clinics using EMRs that mandate collection of nationally agreed datasets can prospectively collect large volumes of ‘real-life’ outcomes data on patient characteristics, DMO treatments and visual outcomes that can be rapidly

257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320


321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384

Table 1 Visual change and number of injections from baseline for eyes receiving at least three injections and with a baseline visual acuity recorded within 4 weeks of the first injection, vision change (letters) from baseline and the number of injections (SD) during the whole period of follow-up.

Visual acuity letters from baseline	Loss 30 or more	Loss 15–29	Loss 1–14	Stable/gain 0–14	Gain 15–29	Gain 30 or more
Number of eyes (total=2292) (%)	30 (1.3)	0	490 (21.4)	1375 (60.0)	318 (13.9)	79 (3.4)
Number of injections (SD)	4.6 (2.6)	0	5.4 (2.8)	5.3 (2.9)	6.1 (3.6)	5.8 (3.3)

Visual change and number of injections from baseline for eyes receiving at least three injections and with a baseline visual acuity recorded within 4 weeks of the first injection, vision change (letters) from baseline and the number of injections (SD) during the whole period of follow-up.

Figure 3 Upper: visual acuity (mean and SD) versus months since the first injection for all treatment groups (upper graph) and the number of eyes contributing to each data point (lower). Lower: number of eyes treated plotted against months from first injection of ranibizumab for eyes with a baseline visual acuity recorded within 1-month of the first injection.



extracted for analysis. Data were collected as part of routine clinical care and form part of the patient's contemporaneous clinical care record. One potential weakness of this study is that some centres used the EMR only for patients receiving injections and that patients apparently lost to follow-up were in fact still under review in outpatient clinics. Patients with either very good or very poor responses to therapy may have been lost to this dataset in those centres. Key missing data points that represent areas of influence in designing care pathways (eg, ethnicity data, or validated type of diabetes mellitus, VA data on non-treatment clinic visits) will be important in the future designs of ophthalmic EMRs.

In this study, there was a mean gain of five letters at 12 months after initiation of treatment when both treatment-naïve and previous treatment groups were analysed together. This is quite disappointing compared with the pivotal trials,^{2 3 5–7 9 13–15} which observed a mean gain of 10.6–11.1 letters at 12 months. DMO observations from smaller open-label prospective, phase IIIb studies such as PRIDE (n=515), RELIGHT (n=108) and RETAIN (n=332) observed slightly greater VA acuity gains at >12 months ranging from four to eight letters.^{16 17} However, the gain at 12 months is similar to real-world studies^{16 18} and within the range from an analysis of several trials.¹⁹

Our findings from the UK real-world practice may reflect the more chronic nature of the disease in some patients, who had waited for access to ranibizumab or received other treatments for DMO prior to national approval; the presence of other ocular or systemic conditions that may limit the efficacy of treatment that were excluded from clinical trials or undertreatment

and insufficient follow-up visits. It is also worth noting that patients were only eligible for initiating ranibizumab therapy when their macular thickness was >400 microns. This imposed threshold may have resulted in the increased chronicity of disease in our UK-based cohort. The impact of a longer duration of DMO and previous macular laser on reducing VA gains has been suggested in the outcomes from clinical trials.⁵ Furthermore, our findings may also reflect service delivery issues—fewer injections per eye, longer intervals between injections than clinically indicated, longer delays before initiation of treatment or recommencement of treatment after a recurrence, variability in interpretation of retreatment criteria or stopping and starting criteria.

Nonetheless, analysis of this dataset does highlight some encouraging findings. Figure 1 shows that eyes with excellent starting VA (70–100 letters) at baseline maintained good mean VA throughout the 18 months follow-up, although showed a slight negative gain (–2 letters at 18 months). This emphasises the ceiling effect of visual gain as an outcome measure and indicates that gain is not a good measure of the quality of care. In contrast, eyes with worse starting acuity achieved a greater gain in acuity. However, the final VA achieved in these patients, which is often the most important outcome for the patient, was not as good as those with a good baseline VA. This has been similarly observed in other diseases, such as age-related macular degeneration treated with ranibizumab²⁰ and in another study of DMO outcomes.¹⁹

There were 373 eyes in the worst VA group and the reasons for the variability in vision in the second year of follow-up warrant further study. It is possible that the first-year VA results

reflect resolution of macular oedema and the second-year results reflect other ocular pathologies, such as diabetic macular ischaemia or poor glycaemic control. A recent study suggested that the level of diabetic control may not influence VA outcomes.⁹

Role for benchmarking standards

‘Translating’ the results of clinical trials into real-world clinical settings is a well-recognised problem that well-designed EMR systems can inform. In order to understand how well we are delivering care, we need to understand what can be achieved in unselected patient populations and how they differ from clinical trials.

This study represents the visual outcomes achieved in routine clinical practice in the UK prior to 2014 and provides a ‘real-world’ benchmark to compare local outcomes. It is important to note, however, that these results were obtained with fewer injections and fewer clinical visits than the pivotal studies.

An important benchmark for visual function is the proportion of patients who achieve 20/40 or better, which approximates the UK driving standard threshold. In this study, 33% of patients achieved better than or equal to 20/40 in the treated eye at 12 months. In the RISE and RIDE trials at baseline, this proportion of eyes was 19.7% and 19.2% in the control arms and 19.2% and 19.7% in the 0.5 mg ranibizumab treatment arms. At 12 months, this proportion of eyes was 34.6% and 37.8% in the control/crossover arms and 62.2% and 63.2%²¹ in the 0.5 mg ranibizumab arms.

Another important benchmark is the stability of vision after the maximum VA gained. Vision stability is thought to be an important outcome in other studies^{7 21} and is clearly of primary importance to patients. We suggest that VA stability is an important measure of the quality of service delivery. It is independent of baseline acuity and may therefore be used as a metric for comparisons between different population groups. In our cohort, VA gains at 6 months were stable at 18 months, except the group with the worst starting VA.

Author affiliations

¹Department of Medical Retina, Moorfields Eye Hospital NHS Foundation Trust, London, UK

²Institute of Ophthalmology, London, UK

³City University, London, UK

⁴Department of Ophthalmology, University of Washington, Seattle, USA

⁵Department of Medical Retina, Moorfields Eye Hospital, London, UK

⁶Department of Cell Biology, UCL Institute of Ophthalmology, London, UK

⁷Bristol Eye Hospital, Bristol, UK

⁸Gloucestershire Hospitals NHS Foundation Trust, Cheltenham, UK

⁹Queens University, Belfast, UK

¹⁰Birmingham and Midland Eye Centre, Birmingham, UK

¹¹Moorfields Eye Hospital, London, UK

¹²Calderdale and Huddersfield NHS Foundation Trust, Huddersfield, UK

¹³Manchester Royal Eye Hospital, Manchester, UK

¹⁴Eye Department, Frimley Park Hospital, Frimley, UK

¹⁵Hinchingbrooke Health Care NHS Trust, Huntingdon, Cambridgeshire, UK

¹⁶Department of Ophthalmology, Hull Royal Infirmary, Hull, UK

¹⁷Leeds, UK

¹⁸Mid Yorkshire Hospitals NHS Trust, Yorkshire, UK

¹⁹Moorfields Eye Hospital NHS Foundation Trust, London, London, UK

²⁰Northern Devon Healthcare NHS Trust, Barnstaple, Devon, UK

²¹Peterborough and Stamford Hospitals NHS Foundation Trust, Peterborough, Peterborough, UK

²²Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK

²³Department of Ophthalmology, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK

²⁴Department of Ophthalmology, Warrington and Halton Hospitals NHS Foundation Trust, Warrington, UK

²⁵Wirral University Teaching Hospital NHS Foundation Trust, Wirral, UK

²⁶Wrightington, Wigan and Leigh NHS Foundation Trust, Wigan, UK

²⁷Division of Optometry & Visual Science, City University, London, UK

Twitter Follow David Crabb @crabblab

Collaborators UK AMD and DR EMR Users Group: *Contributing Centres and Lead Clinician at each centre*: Belfast Health and Social Care Trust, Professor Usha Chakravarthy; Calderdale and Huddersfield NHS Foundation Trust, Mrs Rehna Khan; Cambridge University Hospitals NHS Foundation Trust, Mr Jong Min Ong; Central Manchester University Hospitals NHS Foundation Trust, Mr Sajjad Mahmood; Frimley Park Hospital NHS Foundation Trust, Mrs Geeta Menon; Gloucestershire Hospitals NHS Foundation Trust, Mr Robert Johnston; Heart of England NHS Foundation Trust, Miss Saher Al-Husainy; Hinchingbrooke Health Care NHS Trust, Ms Toks Akerele; Hull and East Yorkshire Hospitals NHS Foundation Trust, Miss Louise Downey; Leeds Teaching Hospitals NHS Trust, Mr Martin McKibbin; Mid Yorkshire Hospitals NHS Trust, Mr Atul Varma; Moorfields Eye Hospital at Bedford Hospital, Mr Aires Lobo; Northern Devon Healthcare NHS Trust, Dr Elizabeth Wilkinson; Peterborough and Stamford Hospitals NHS Foundation Trust, Mr Alan Fitt; Sheffield Teaching Hospitals NHS Foundation Trust, Mr Christopher Brand; University Hospitals Birmingham NHS Foundation Trust, Miss Marie Tsaloumas; University Hospitals Bristol NHS Foundation Trust, Miss Clare Bailey; Warrington and Halton Hospitals NHS Foundation Trust, Miss Kaveri Mandal; Wirral University Teaching Hospital NHS Foundation Trust, Mr Vineeth Kumar; Wrightington, Wigan and Leigh NHS Foundation Trust, Mr Salim Natha.

Contributors All the authors have contributed to the planning, conduct and reporting of the work described in the article.

Funding This work was supported in part by an unrestricted research award by Novartis Pharmaceuticals. This research has received a proportion of its funding from the Department of Health’s NHR Biomedical Research Centre for Ophthalmology at Moorfields Eye Hospital and UCL Institute of Ophthalmology.

Competing interests RJ is the Medical Director of Medisoft, the Electronic Medical Record software provider from which data were extracted.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Hawkes N. Cost of hospital prescribing rose 15% in a year owing to new rheumatoid arthritis and cancer drugs. *BMJ* 2014;349:g6867.
- Elman MJ, Aiello LP, Beck RW, *et al.*, Diabetic Retinopathy Clinical Research Network. Randomized trial evaluating ranibizumab plus prompt or deferred laser or triamcinolone plus prompt laser for diabetic macular edema. *Ophthalmology* 2010;117:1064–77.e1035.
- Elman MJ, Qin H, Aiello LP, *et al.* Diabetic Retinopathy Clinical Research Network. Intravitreal ranibizumab for diabetic macular edema with prompt versus deferred laser treatment: three-year randomized trial results. *Ophthalmology* 2012;119:2312–18.
- Domalpally A, Ip MS, Ehrlich JS. Effects of intravitreal ranibizumab on retinal hard exudate in diabetic macular edema: findings from the RIDE and RISE Phase III clinical trials. *Ophthalmology* 2015;122:779–86.
- Elman MJ, Ayala A, Bressler NM, *et al.* Intravitreal ranibizumab for diabetic macular edema with prompt versus deferred laser treatment: 5-year randomized trial results. *Ophthalmology* 2015;122:375–81.
- Elman MJ, Bressler NM, Qin H, *et al.* Expanded 2-year follow-up of ranibizumab plus prompt or deferred laser or triamcinolone plus prompt laser for diabetic macular edema. *Ophthalmology* 2011;118:609–14.
- Brown DM, Nguyen QD, Marcus DM, *et al.* Long-term outcomes of ranibizumab therapy for diabetic macular edema: the 36-month results from two phase III trials: RISE and RIDE. *Ophthalmology* 2013;120:2013–22.
- ETDRS report 19.pdf.
- Bansal AS, Khurana RN, Wieland MR, *et al.* Influence of glycosylated hemoglobin on the efficacy of ranibizumab for diabetic macular edema: a post hoc analysis of the RIDE/RISE trials. *Ophthalmology* 2015;122:1573–9.
- Lane M, Mathewson PA, Sharma HE, *et al.* Social deprivation as a risk factor for late presentation of proliferative diabetic retinopathy. *Clin Ophthalmol* 2015;9:347–52.
- Singh RP, Habbu K, Ehlers JP, *et al.* The impact of systemic factors on clinical response to ranibizumab for diabetic macular edema. *Ophthalmology* 2016;123:1581–7.
- Do once and share web address. http://rcophth-website.www.premierhosting.com/standards/do_once_and_share
- Wells JA, Glassman AR, Ayala AR, *et al.*, Diabetic Retinopathy Clinical Research Network. Aflibercept, bevacizumab, or ranibizumab for diabetic macular edema. *N Engl J Med* 2015;372:1193–203.

641	14	Bressler NM, Varma R, Suñer JJ, <i>et al.</i> , RIDE and RISE Research Groups. Vision-related function after ranibizumab treatment for diabetic macular edema: results from RIDE and RISE. <i>Ophthalmology</i> 2014;121:2461–72.	705
642			706
643	15	Nguyen QD, Brown DM, Marcus DM, <i>et al.</i> Ranibizumab for diabetic macular edema: results from 2 phase III randomized trials: RISE and RIDE. <i>Ophthalmology</i> . 2012;119:789–801.	707
644			708
645	16	Menchini U, Bandello F, De Angelis V, <i>et al.</i> Ranibizumab for visual impairment due to diabetic macular edema: real-world evidence in the Italian population (PRIDE Study). <i>J Ophthalmol</i> 2015;2015:324841.	709
646			710
647	17	Pearce I, Banerjee S, Burton BJ, <i>et al.</i> Ranibizumab 0.5 mg for diabetic macular edema with bimonthly monitoring after a phase of initial treatment: 18-month, multicenter, phase IIIB RELIGHT study. <i>Ophthalmology</i> 2015;122:1811–19.	711
648			712
649	18	Patrao NV, Antao S, Egan C, <i>et al.</i> Real-world outcomes of ranibizumab treatment for diabetic macular edema in a United Kingdom National Health Service setting. <i>Am J Ophthalmol</i> 2016.	713
650			714
651	19	Dugel PU, Hillenkamp J, Sivaprasad S, <i>et al.</i> Baseline visual acuity strongly predicts visual acuity gain in patients with diabetic macular edema following anti-vascular endothelial growth factor treatment across trials. <i>Clin Ophthalmol</i> 2016;10:1103–10.	715
652			716
653	20	Zarranz-Ventura J, Liew G, Johnston RL, <i>et al.</i> The neovascular age-related macular degeneration database: report 2: incidence, management, and visual outcomes of second treated eyes. <i>Ophthalmology</i> 2014;121:1966–75.	717
654			718
655	21	Boyer DS, Nguyen QD, Brown DM, <i>et al.</i> Outcomes with as-needed ranibizumab after initial monthly therapy: long-term outcomes of the phase III RIDE and RISE trials. <i>Ophthalmology</i> 2015;122:2504–13.e2501.	719
656			720
657			721
658			722
659			723
660			724
661			725
662			726
663			727
664			728
665			729
666			730
667			731
668			732
669			733
670			734
671			735
672			736
673			737
674			738
675			739
676			740
677			741
678			742
679			743
680			744
681			745
682			746
683			747
684			748
685			749
686			750
687			751
688			752
689			753
690			754
691			755
692			756
693			757
694			758
695			759
696			760
697			761
698			762
699			763
700			764
701			765
702			766
703			767
704			768