



City Research Online

City, University of London Institutional Repository

Citation: Marshall, C. R., Rowley, K., Mason, K., Herman, R. & Morgan, G. (2013). Lexical organization in deaf children who use British Sign Language: Evidence from a semantic fluency task. *Journal of Child Language*, 40(1), pp. 193-220. doi: 10.1017/s0305000912000116

This is the unspecified version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/2223/>

Link to published version: <https://doi.org/10.1017/s0305000912000116>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Lexical organisation in deaf children who use British Sign Language: Evidence from a semantic fluency task

Marshall, C. R., Rowley, K., Mason, K., Herman, R., & Morgan, G.

Journal of Child Language, 2012 doi:10.1017/S0305000912000116

Acknowledgements

We thank the children who participated in this study, and their teachers and parents.

This work was supported by the Economic and Social Research Council of Great Britain (Grant RES-620-28-6001; Deafness, Cognition and Language Research Centre (DCAL)), and by a Leverhulme Early Career Fellowship awarded to the first author. We thank Joanna Atkinson and Nicola Botting for discussions about data coding.

Abstract

We adapted the semantic fluency task into British Sign Language (BSL). In Study 1, we present data from 22 deaf signers aged 4-15. We show that the same 'cognitive signatures' that characterise this task in spoken languages are also present in deaf children, for example, the semantic clustering of responses. In Study 2, we present data from 13 deaf children with Specific Language Impairment (SLI) in BSL, in comparison to a subset of children from Study 1 matched for age and BSL exposure. The two groups' results were comparable in most respects. However, the group with SLI made occasional word-finding errors and gave fewer responses in the first 15 seconds. We conclude that deaf children with SLI do not differ from their controls in terms of the semantic organisation of the BSL lexicon, but that they access signs less efficiently.

Introduction

Sign languages are independent, fully fledged languages created by deaf people in different countries (for a review, see Brentari, 2010). Lexical items, be they signed or spoken, are mappings between a phonological form and a meaning or set of meanings. As children's vocabulary grows, items become organised into a semantic network, with strong links between items that are closely related, weaker links between items that are less closely related, and a hierarchical organisation that reflects taxonomic relationships (for a review of lexical acquisition, see Clark, 1993). The learning of lexical items, and their organisation within a semantic network, is just as central to the acquisition of a signed language as it is to spoken language acquisition.

This paper investigates lexical organisation in two groups of deaf children who are acquiring British Sign Language (BSL): those who are learning BSL without any difficulty, and those who have Specific Language Impairment (SLI) in BSL. We investigate these children's lexical organisation using a semantic fluency task adapted for BSL. This is the first investigation of semantic fluency with deaf children in any signed language.

The introduction is structured as follows. After a general introduction to lexical acquisition in deaf signing children, we discuss the main features of (hearing) children's performance on the semantic fluency task and discuss what the task measures. We also discuss the only previous study of semantic fluency in signers, which tested deaf adults who use BSL. We then turn our attention to the characteristics of SLI in signed languages, and to previous results of semantic fluency

in hearing children with SLI. We end by setting out our predictions on the semantic fluency task for two groups of deaf signing children: those whose language is developing appropriately, and those who have SLI in BSL.

Lexical acquisition in deaf children is interesting for several reasons that can be linked to the nature of language exposure in this group. Research on language development in deaf native signers (i.e. those who acquire a natural sign language from birth, and from their parents) has shown that early exposure to sign enables children to reach developmental milestones at the same pace as their hearing peers acquiring spoken languages (Anderson & Reilly, 2002; Lillo-Martin, 1994; Woolfe et al, 2010). However, only 5-10% of deaf children receive sign language input from their deaf parents (Mitchell & Karchmer, 2004), which leaves the remaining 90 - 95% of children who are born to hearing parents with little or no early experience of sign language. This latter group of children grow up with widely differing language-learning backgrounds. Research on deaf children growing up with hearing parents suggests a slower pace of sign lexical acquisition and a smaller lexicon size (Anderson, 2006; Blamey, 2003; Lederberg & Spencer, 2009; Prezbindowski & Lederberg, 2003). This may be largely due to reduced incidental exposure to sign language: hearing parents tend to use sign only when directly addressing their deaf child and not with other hearing family members. This means that the child has few opportunities for picking up vocabulary through observing the interactions of others (Marschark, 1997).

What are the consequences of these differences in language exposure for lexical acquisition and the ensuing organisation of the lexicon? There are very few studies on this topic. In some ways, sign vocabulary acquisition in *native* signers appears to be very similar to that of hearing children in spoken languages. For example, Anderson and Reilly (2002) report that native deaf signers' acquisition of American Sign Language (ASL) vocabulary within particular semantic domains, such as question words, emotion words and cognitive verbs, is comparable to that found in hearing peers. On the other hand, a recent study on early British Sign Language (BSL) development in deaf children of hearing parents suggests a higher frequency of certain verbs or signs based on actions (e.g., CATCH, PLAY, SWIM, CROCODILE¹), compared to native signing children and hearing children who are acquiring a spoken language (Marschark & Woll, in preparation). This action bias is also seen in deaf children's homesigns (conventionalised gestures created between children and their hearing parents; Goldin-Meadow, Butcher, Mylander & Dodge, 1994).

In the present study we investigated the lexical organisation of nouns, within two particular semantic domains: food and animals. These domains have been widely studied in spoken language (Crowe & Prescott, 2003; Lucariello, Kyatzis & Nelson, 1992; Nelson, 1974, *inter alia*). The task we use – semantic fluency – is straightforward to administer: participants name as many exemplars as they can from a particular semantic category within a limited period of time (usually one minute). Semantic fluency has been used in many spoken languages with a range of age groups and with children who have various developmental disorders, including Down Syndrome (Nash & Snowling, 2008), High Functioning Autism (Boucher, 1988), and

¹ The sign CROCODILE is made with two hands making repeated contact at the palms, representing the opening and closing of the crocodile's jaws. Note that here and throughout the paper we use capital letters to indicate the English gloss for BSL signs.

Attention Deficit/ Hyperactivity Disorder and Tourette's Syndrome (Mahone, Koth, Cutting, Singer & Denckla, 2001). It also has the advantage that many different aspects of performance can be analysed beyond just the number of items produced. We therefore considered it an appropriate tool for adaptation into BSL, for testing deaf children with SLI, and for investigating potential group differences in lexical organisation between deaf children with SLI and those with typically developing signing skills.

When a word is spoken (or a sign produced) it is assumed that this will in turn activate other words or concepts that are semantically similar or associatively related to it (Crowe & Prescott, 2003). Hence it is also assumed that the order in which words are produced during the semantic fluency task will indicate, indirectly, their proximity to each other in the lexicon. Given the limited amount of time that participants are given to respond, the task does not provide an exhaustive list of the words that they know, but it does reveal those items that come most readily to mind.

Performance in this task shows a number of consistent characteristics, termed 'cognitive signatures' (Koren, Kofman & Berger, 2005; Riva, Nicelli & Devoti, 2000; Sauzéon, Lestage, Raboutet, N'Kaoua & Claverie, 2004; Troyer, Moscovitch & Wincour, 1997). There is a hyperbolic decline in the rate of production of new items over the duration of the task, and items are produced in bursts of semantically-related words. More prototypical category exemplars are produced with higher frequency (i.e. by more participants) than less typical ones. The task is generally considered to provide a measure of both semantic organisation and executive function. If participants can generate exemplars in response to a superordinate label, e.g. food,

then this suggests that semantic knowledge is organised taxonomically. Furthermore, there is internal clustering, whereby words that are even more closely related are produced together (for example, a cluster of farm animals, or a cluster of fruits). Good performance on the task requires good semantic memory, i.e. the component of long-term memory that contains the permanent representation of our knowledge of objects, facts, and concepts as well as words and their meaning. The task also requires the use of word-retrieval strategies, which in turn rely on executive functions, namely switching (i.e. set-shifting between different clusters), working memory (to keep track of items that have already been produced), and inhibition (so as to avoid repeating previous responses, and irrelevant responses). These skills enable the participant to retrieve lexical items more efficiently.

An obvious question is whether semantic fluency in a signed language shows the same cognitive signatures as those reported for spoken languages. Despite its widespread use as a tool in spoken language, there is only one other study of semantic fluency in a signed language (Marshall, Rowley & Atkinson, submitted). Marshall et al (submitted) tested 30 native or near-native users of BSL aged between 18 and 60 years old, with the same categories used in the present study, namely 'food' and 'animals'. They discovered the same cognitive signatures as reported for spoken language fluency, i.e. a hyperbolic decline in the rate of production of new items over the duration of the task, clusters of semantically-related words, and more prototypical category exemplars being produced by more participants than less typical ones. Importantly, the number of items produced in each category was comparable to reports of adults in spoken languages (e.g. English: Harrison, Buxton, Husain & Wise, 2000; Greek: Kosmidis, Vlahou, Panagiotaki, & Kiosseoglou, 2004; Hebrew: Kavé,

2005; Spanish: Buriel, Gramunt, Bohm, Rodes & Peña-Casanova, 2004), despite the smaller established lexicon of BSL compared to spoken languages (Sutton-Spence & Woll, 1999).

The existence of fingerspelling (i.e. a manual alphabet) is another difference between spoken and signed languages. Fingerspellings of a few highly frequent and short items (e.g. HAM, h-a-m; EGG, e-g-g) have become lexicalised. However, fingerspelling is also used for low frequency items for which there is no established sign. In addition, In Marshall et al's (submitted) study, fewer than 2% of items were fingerspellings, indicating that during the semantic fluency task signers were retrieving signs from the established BSL lexicon, rather than retrieving items from their English lexicon and spelling them out.

The present study is the first documented investigation of semantic fluency in signing children. In addition to reporting data from deaf children who are acquiring BSL without any evidence of difficulty, we investigate semantic fluency in deaf children with SLI in their acquisition of BSL.

SLI is a significant impairment in acquiring language despite normal nonverbal IQ and no gross level of impairment in neurological function, motor development, or social interaction, alongside normal hearing (Leonard, 1998). The requirement for normal hearing means that profoundly deaf children are excluded from a diagnosis of SLI by default. Yet given that 7% of the general hearing child population have SLI (Tomblin, Records, Buckwater, Zhang, Smith, & O'Brien, 1997), this would also be expected to

be the case for deaf children, including those whose primary mode of communication is a signed language.

The characterisation of SLI in signed languages is just beginning, and has so far been reported in only two signed languages: BSL (Mason, Rowley, Marshall, Atkinson, Herman, Woll & Morgan, 2010; Morgan, Herman & Woll, 2007), and ASL (Quinto-Pozos, Forber-Pratt & Singleton, 2011). A major difficulty in identifying SLI with confidence in children acquiring a signed language is the aforementioned confound with delayed language exposure – over 90% of deaf children are born to hearing parents, who are not able to provide fluent sign language input. Deaf children may be exposed to fluent models of sign language outside the family, for example if they attend preschool settings with deaf signing staff, but for most their first contact with sign language will be when they start school. Their language development will hence be delayed, although many will go on to be proficient signers. Yet experienced teachers of the deaf and speech and language therapists do report working with children who are not acquiring sign language as well as would be expected in comparison to peers who have had the same (delayed) language experience (Edwards, 2010; Mason et al, 2010; Quinto-Pozos et al, 2011).

Alongside the issue of late language exposure, another important factor that complicates the diagnosis of SLI in signed languages (in common, indeed, with many lesser-studied spoken languages) is the lack of standardised language assessments. For many signed languages, even those in the developed world, reliable language assessments are not available (Haug, 2005). Professionals may, of course, be able draw upon their own knowledge of sign language acquisition to determine when a

child seems learning language more slowly than expected. Yet in these cases identification relies on many years of experience on the part of the professional.

One of the few signed languages for which standardised measures of receptive and expressive language are available and suitable for children is BSL, the assessment instruments being the BSL Receptive Skills Test (Herman, Holmes & Woll, 1999) and the BSL Production Test (Herman, Grove, Holmes, Morgan, Sutherland, & Woll, 2004). These two tasks have been used as the basis for identifying SLI in deaf children who use BSL in a couple of studies to date (Mason et al, 2010; Morgan et al, 2007). In Mason et al's group study of SLI in signers, children were considered to have SLI when a teacher of the deaf or a specialist speech and language therapist reported language concerns after comparing their development to other deaf children in the same classes with comparable exposure to BSL. Children with additional diagnoses of special educational need, e.g. autism, were excluded, but those with reading difficulties were not, given the close relationship between language and literacy development (Cain, 2010) and the difficulties that many deaf children face in learning to read (Allen, 1986; Conrad, 1979). In addition the children were required to display non-verbal abilities in the normal range but impaired performance on one or both of the BSL standardised assessments.

Mason et al's (2010) study of 13 deaf children with SLI, aged 5;10-14;08 showed that 7/13 children scored -1.3 SD or worse on the BSL Receptive Skills Test and that all scored at or below the 10th percentile on one or more subtest of the BSL Production Test. Children's narratives almost invariably showed minimal use of grammatical morphology, unclear signing, and no introduction of characters or setting. A more in-

depth characterisation of the linguistic features of SLI in sign language users is required, and the present study contributes to this endeavour.

Only a small number of studies have used the semantic fluency task with hearing children who have SLI in spoken language. Recently, Henry, Messer & Nash (2012) reported that a group of English-speaking children with SLI performed below their chronological age-matched controls on both verbal and non-verbal fluency tasks. However, their difficulties were particularly marked for phonological fluency, a task where participants are asked to generate as many words as they can that begin with a particular sound or letter. For phonological fluency children performed more poorly than they did for semantic and non-verbal fluency (Nash, Henry & Messer, 2010).

In a different test probing lexical organisation, an association task whereby children were asked to generate three words associated to each of a list of 48 words, a subgroup of children with SLI performed more poorly than typically developing peers matched for expressive vocabulary ability (Sheng & McGregor, 2010). Such children generated fewer semantically related responses and more unrelated responses than expected. However, the SLI group as a whole was characterised by variable performance, and some children performed age-appropriately.

The findings from these studies confirm the general view in the field of SLI research that although these children vary greatly in their lexical abilities, lexical deficits do not characterise the disorder in the way that morphological and syntactic impairments do (Leonard, 1998). Moreover, in their case study of a deaf native signer with SLI, Morgan and colleagues report that the boy had a good sign vocabulary. Even though

he was 5 years old he mostly communicated with single signs, and his deficits were argued to lie principally in the morphology and syntax of BSL (Morgan et al, 2007). That preliminary study raises the possibility that lexical deficits may not characterise SLI in signed languages either. However, this has not yet been tested in a fluency task, which measures lexical organisation and speed of access to lexical representations.

Predictions for our study

We set out to investigate semantic fluency performance in deaf children who are acquiring BSL typically, and in deaf children who have SLI in their signing. Semantic fluency offers a rich dataset over which several different analyses can be undertaken. In particular, we calculated various measures and made the following predictions:

- 1) **Total number of responses and number of correct and incorrect responses.** We predicted that the task would be sensitive to development, and therefore that the total and correct number of responses given by both groups would increase with age. We compared the total number of correct responses with figures available from the spoken language literature. Given that both of our experimental groups contain children with late exposure to BSL, productivity (i.e. total number of responses) was predicted to be lower than for children of the same age who are acquiring spoken language.
- 2) **Rate of decline of responses.** We calculated the number of responses during each quadrant of the time available for the task, i.e. at 1-15 seconds, 16-30s, 31-45s and 46-60s. For both groups we predicted a decline in response rate over the course of the minute, as has been reported for spoken language and in deaf adults doing the task in BSL.

- 3) **Semantic clusters.** We calculated the number of semantic clusters, where clusters were defined as two or more successive words belonging to a conventional subcategory. We predicted that, as has been reported in previous studies of deaf adults and of hearing children and adults, such clusters would be identifiable. We also calculated how many items there were in each cluster, and how many switches there were between clusters and/or non-clustered items. This allowed us to investigate whether increased productivity was related to an increase in the number of clusters generated and the number of times children switched between clusters, or alternatively to an increase in the size of the clusters.
- 4) **Item analysis.** We investigated which items emerged as most 'typical', and how these compared to studies of hearing children from the USA and UK doing the task in English. Given that hearing English-speaking children and children who use BSL are growing up in the same westernised society, we did not expect differences here.

Study 1: Typically developing deaf children

Introduction

We first investigated semantic fluency in typically developing deaf signers, with the aim of comparing performance to that of hearing children doing the task in spoken languages, and to provide a comparison group to the children with SLI who participated in Study 2.

Methods

Participants

22 deaf signing children, aged 4;00 to 15;2, participated in this study. None had any identified educational need (e.g. Autism, Attention Deficit/Hyperactivity Disorder, intellectual disability) other than deafness. All were acquiring BSL without any difficulties, as reported by their teachers and parents. Table 1 shows the range of backgrounds with respect to whether there is a Deaf family member and the type of school attended. This range is representative of the variable language background of deaf children. Current scores on one or both of the standardised tasks of BSL (BSL Receptive Skills Test, Herman et al, 1999; BSL Production Test, Herman et al, 2004) and/or the Nonsense Sign Repetition Test (Mann, Marshall, Mason & Morgan, 2010) were available for only 12 of these children, and were made available to us by their schools. All 12 had scores within the normal range, and these are reported in Table 1. As we already had an experimental battery lasting between one and two hours, we were not able to ask for an additional hour to administer the tests ourselves to the other ten children. However, each child was seen for the experimental battery and a warm-up conversation by the second author, who is a Deaf native signer experienced at working with deaf children. In no case did she suspect any difficulty in BSL acquisition – in her judgement, all children had BSL skills at the expected level for their age and exposure.

TABLE 1 ABOUT HERE

Procedure

We used two semantic categories: 'food' and 'animals', which are the most widely-used categories in the spoken language literature (Koren et al, 2005; Kosmidis et al,

2004; Nash & Snowling, 2008; Riva et al, 2000; *inter alia*). Instructions were delivered in BSL by the experimenter (second author, a Deaf native signer, or third author, hearing signer with advanced BSL skills). The instructions were straightforward: “Please tell me the names of as many animals/food items as you can. Be as quick as possible. You have one minute. Ready? Go”. No examples were given, but ‘colours’ was used as a practice category. Responses were filmed and subsequently glossed into English.

Results

Data loss

Two children (N004 and N010; the youngest children at 5;6 and 4;0 respectively) appeared not to understand the task and were unable to respond without prompting. Another child (N016; aged 9;9) responded to just one category (animals), and so her partial data were excluded from the analysis. We therefore present data from only 19 of the 22 children who participated.

Coding of responses

The signs were glossed into English semantic equivalents, timed (i.e. it was noted how many seconds into the minute they were produced) and coded as correct/incorrect by the second and third authors working together. Each incorrect response was coded as one of three types, and these categories captured all the errors:

- Repetition of an item
- Intrusion (i.e. an item that was from a category other than food/animals)
- Uninterpretable

The coding was checked by the first author (hearing linguist with advanced BSL skills), who discussed the very few discrepancies (all involving uninterpretable items) with the second and third authors until a consensus was reached. The first author then further coded the correct and repeated responses according to semantic clusters. A cluster was defined as two or more adjacent responses that were semantically related in some way. We allowed categories to emerge from the data, rather than imposing them. For example, in one recent study the taxonomic categories ‘mammal’, ‘bird’, ‘reptile’, ‘amphibian’ ‘fish’ and ‘insect’ were used to code data from hearing children (Nash & Snowling, 2008). However, in our view this coding scheme does not reflect how our participants were grouping their responses. We therefore followed an emergent approach to coding clusters (e.g. Kosmidis et al, 2004). Animal categories were: ‘zoo’, ‘pet’, ‘farm’ ‘water’, ‘invertebrate’, ‘bird’, and ‘British wild’. The number of food categories was much greater, and included: ‘fruit’, ‘vegetables’, ‘meat’, ‘carbohydrates’, ‘desserts’, ‘snacks’, ‘meals with chips’, ‘takeaway meals’, ‘breakfast foods’, ‘Italian foods’, and ‘roast dinner foods’.

This emergent approach is supported by evidence from Crowe & Prescott (2003) that children cluster animals around their environmental context (e.g. home, farm, zoo). It meant, however, that certain responses could potentially fall into more than one category. For example, the animal FISH could fall into the categories ‘pet’ or ‘water’, and DUCK into ‘farm’, ‘bird’ or ‘water’. In each case the category was chosen based on the answers before and after. For example, CROCODILE was coded as ‘reptile’ when it occurred in the sequence ‘SWAN-SNAKE-CROCODILE-SHARK’ but in the category ‘zoo’ when it occurred in the sequence ‘LION-CROCODILE-ELEPHANT’. In assigning categories we endeavoured to be as inclusive as possible, meaning that we

tried to ensure that as many responses as possible fell within clusters. An example of the coding for one child's responses is presented in the Appendix.

Responses for eight children (four typical, four SLI (i.e. for study 2)) were then independently coded by the fourth author (hearing Speech and Language Therapist with advanced BSL skills) with respect to semantic clusters using the coding instructions exactly as they appear below. Despite allowing categories to emerge from the data rather than imposing them, and despite more categories emerging for food than for animals, inter-coder agreement was high and equivalent across food and animals: 88.71% for animals and 88.53% for food.

Table 2 presents the total number of items within each category, and the mean score across both categories. There was no significant difference between the number of responses to food and animals, $t(18)=0.594$, $p = 0.560$. Nor did the two categories differ in the number of correct items, repeated items, irrelevant items or uninterpretable items, all t s < 0.6 . Error rates across all three error types were low.

TABLE 2 ABOUT HERE

A correlational analysis with age revealed that both the number of total responses and the number of correct responses (averaged across 'food' and 'animals') increased with age, as shown in Figure 1: $r(19) = 0.601$, $p = 0.007$ and $r(19) = 0.648$, $p = 0.003$ respectively. The correlations between total responses and years of BSL exposure, and between correct responses and BSL exposure, were not significant however, $r(19) = 0.144$, $p = 0.556$ and $r(19) = 0.192$, $p = 0.430$.

FIGURE 1 ABOUT HERE

The total number of responses in each quadrant of the minute did not differ significantly between 'food' and 'animals'; for quadrant 1 (i.e. 1-15s), $t(18) = -1.764$, $p = 0.095$; for quadrant 2 (16-30s), $t(18) = 0.601$, $p = 0.555$; for quadrant 3 (31-45s), $t(18) = 1.326$, $p = 0.201$ and for quadrant 4 (46-60s), $t(18) = 1.312$, $p = 0.206$. The decline in responses over the course of the minute is shown for each category in Figure 2. Bars indicate 1SD above and below the mean.

FIGURE 2 ABOUT HERE

No significant differences were found between food and animals in terms of the number of clusters produced or average cluster size, $t(18) = 0.515$, $p = 0.613$ and $t(18) = 1.249$, $p = 0.228$, respectively. Nor was the number of switches significantly different, $t(18) = 1.312$, $p = 0.206$. Responses to food and animals were therefore collapsed in the analysis that follows.

In order to investigate how cluster size, cluster number and the number of switches relate to age and productivity (i.e. do children who produce more correct responses do so because they produce bigger clusters, or because they produce more clusters, and switch more often between clusters and/or individual items?), correlations between number of correct responses and those three measures were carried out. The number of correct responses correlated significantly with the number of clusters, $r(19) = 0.888$, $p < 0.001$ and number of switches, $r(19) = 0.771$, $p < 0.001$, but not with cluster size,

$r(19) = 0.272$, $p = 0.260$. The full correlation matrix between number of correct responses, total number of responses, age, number of clusters, cluster size and number of switches is shown in Table 3.

Finally, all responses given by more than 33% of the children (a cut-off selected arbitrarily) are shown in Table 4.

TABLE 4 ABOUT HERE

Discussion

The aim of Study 1 was to investigate the cognitive signatures of semantic fluency in typically developing deaf children who use BSL. We expected to find an increase in fluency with age. We also expected to find the same cognitive signatures as have been found in children and adults who use spoken language and in deaf adults who use BSL, namely the production of items in bursts of semantically-related words, similar category exemplars produced with high frequency (i.e. by a large proportion of participants), and a decline in rate of production of new items over the course of the minute. All of these signatures were indeed found to characterise semantic fluency in children acquiring BSL.

The mean of 14.82 correct responses (SD 4.28) is difficult to compare directly to that reported in the literature for hearing children, as there are few studies encompassing the wide age range of the present study. Nash and Snowling (2008) found a mean fluency of 13.24 (averaged across ‘food’ and ‘animals’) for English-speaking children

aged 5;06-9;05. In Italian, Riva et al (2000), found that for children aged 5;11-11;4, productivity increased from a mean of approximately 10 items in the youngest children to 17 items in the oldest group (again, averaged across 'food' and 'animals'). Koren et al (2005) reported for Hebrew-speaking children aged 9-11, a mean production of 15 animals and 10 food items. Therefore the children in the present study were performing at an approximately similar level to the reported literature, despite many of the group not having exposure to BSL from birth (only 5/22 were native signers). Therefore, it does not appear that deaf children, providing they are able to understand the task requirements, find the task in BSL more difficult than hearing children doing the task in a spoken language.

Nevertheless, despite the small sample size, and the variability in BSL exposure across the group, we found an increase in productivity with age, as has been reported for spoken languages (Koren et al, 2005; Riva et al, 2000; Sauz  on et al, 2004). There is still the potential for a developmental increase in productivity, given that native adult signers averaged 23 items in BSL (Marshall et al, submitted). We found that increased productivity was related to an increase in cluster number and the number of switches, rather than to cluster size. Again, this mirrors the results for spoken language (Koren et al, 2005). In other words, the most fluent children produce more responses because they retrieve a greater number of subcategories within 'food' and 'animals', and not because they produce more items in each subcategory. The standard interpretation in the literature is that it is an increase in cognitive flexibility that drives the switch to a new semantic subcategory once lexical retrieval within a particular subcategory slows down (Koren et al, 2005; Troyer et al, 1997). Older children do of course also tend to have larger vocabularies (although we were unable

to measure this directly in our study because there was no standardised BSL vocabulary test available), but with respect to the increase in fluency, it appears that executive functions are the main driver.

The items produced by the deaf children in BSL are very similar to those reported in studies of English. For example, Nelson (1974) reports amongst 5 and 8 year-olds in the USA that the most common animal responses are 'giraffe', 'lion', 'elephant', 'tiger', 'horse', 'cat' and 'dog'. Crowe and Prescott (2003) also report a high frequency of these items in the responses of 5-10 year-old children from England. These were also the most common responses in our study. Nelson (1974) additionally tested the category 'fruit' and found the most common fruits were 'orange', 'apple' and 'banana', also the three most common fruit responses in our food category. This similarity in responses is not surprising given the similar experiences that children in westernised cultures are likely to have, regardless of their hearing status.

Finally, the characteristic decline in the number of items produced during the course of the minute was also observed in our data, with most items produced in the first 15 seconds and fewest items in last 15 seconds.

Study 2: Children with SLI

Introduction

We next tested semantic fluency in a group of deaf signing children who have SLI in their signing. We compared their performance to that of subset of children from study 1 matched for age and years of exposure, in order to investigate any differences in semantic fluency between typically developing signers and signers with SLI.

Methods

Participants

13 deaf signers (10 male), identified as having SLI in their acquisition of BSL by teacher report and follow-up testing with standardised tests of BSL, were recruited to the study. All had non-verbal abilities in the normal range as measured by the matrices, recall of designs and pattern construction subtests of the British Ability Scales 2nd edition (Elliott, Smith & McCullouch, 1996), yet scored at or below -1.3SD on the BSL Receptive Skills Test (Herman et al, 1999) and/or below the 10th percentile on one or more of the BSL Production Test subtests (Herman et al, 2004). Aside from deafness and SLI, they had no additional recognised special needs other than teacher-reported difficulties with reading (N=12), which is not unusual for deaf children (Conrad, 1979; Kyle & Harris, 2006). They ranged in age from 7;5-14;10, mean 10;9, SD = 2;2. Background details for each of the SLI participants are shown in Table 5. 10 of these 13 were participants in Mason et al's (2010) study, and the additional three were selected according to the same criteria as those described in that study.

TABLE 5 ABOUT HERE

13 control children (9 male) were selected from study 1 and individually matched with SLI children to within + or - 6 months of age. The age range of the control group was 7;6-14;10, mean = 10;10, SD = 2;2. The groups had similar experience of BSL: for the SLI group, years of exposure to BSL ranged from 3;0-10;4, mean 6;8, SD = 2;1; for the control group, years of exposure ranged from 1;6-11;9, mean 7;5, SD = 3;7. Two independent samples t-tests revealed no significant differences between the groups with respect to either age, $t(24) = 0.106$, $p = 0.917$, or years of BSL exposure, $t(19.30) = 0.640$, $p = 0.528^2$. Note that the control children were selected before the data were coded, in order to avoid the risk of selection bias. Note also that this group contained N016, one of the children whose data could not be analysed for study 1 as she responded only to the 'animals' category.

Procedure

The procedure for the deaf children with SLI was identical as for the children in study 1.

Results

Two children with SLI did not understand the task and did not provide responses. One of these was the youngest, at 7;05, but the other was older, at 10;09. A third child did respond but refused to be filmed. As filming was essential for accurate glossing of the responses and for timing how many seconds into the minute they were produced, this child's data could not be used. We therefore present data from 10 children with SLI,

² For age of exposure to BSL, the variances of the two groups were significantly different according to Levene's Test for the Equality of Variances ($F(24) = 7.390$, $p = 0.012$). The SLI group has less variance than the control group. Therefore we have not assumed equal variances, and have reduced the degrees of freedom as appropriate.

compared to the 12 remaining controls. Rerunning the t-tests to compare age and years of BSL exposure in these smaller groups revealed that the groups were still well-matched for both measures, both $t_s < 0.4$. The data are averaged across both categories (i.e. 'food' and 'animals') and presented in Table 6.

TABLE 6 ABOUT HERE

A set of t-tests was carried out to compare the two groups on the following measures: total number of responses, number of correct responses, number of incorrect responses (repetitions, irrelevant and uninterpretable responses), number of clusters, average cluster size, and the number of switches. None of these comparisons was significant (see Table 6).

We also compared the two groups' number of responses per quadrant of the minute, using a 4(quadrant) x 2(group) ANOVA. We found a significant interaction between group and quadrant, $F(3,60) = 4.35$, $p = 0.008$, partial $\eta^2 = 0.179$. There was no main effect of group, $F(1,20) = 0.88$, $p = 0.360$, partial $\eta^2 = 0.042$. The main effect of quadrant was strongly significant, $F(3,60) = 84.02$, $p < 0.001$, partial $\eta^2 = 0.808$, reflecting a sharp decline in responses over the course of the minute.

To investigate the interaction, we conducted four independent samples t-tests comparing the two groups' performance in each quadrant, with the alpha level reduced to $p = 0.013$ in order to compensate for multiple comparisons ($N=4$). As shown in Table 6, there is a significant difference between groups only for the first quadrant, $t(20) = 2.698$, $p = 0.013$. This difference is accounted for by the control group

producing significantly more items in the first 15 seconds of the minute compared to the SLI group.

The interaction was further investigated with a set of paired samples t-tests for each group comparing items produced in successive quadrants, again with the alpha level reduced to $p = 0.013$. For the control group, there were significantly more responses for the first versus the second quadrant, $t(11) = 8.742$, $p < 0.001$, but the difference between the second and third quadrant did not reach significance, $t(11) = 1.541$, $p = 0.152$, and nor did the difference between the third and fourth quadrants, $t(11) = 2.191$, $p = 0.051$. For the SLI group showed the same pattern as the controls over the course of the minute, with significantly more responses for the first versus the second quadrants, $t(9) = 8.728$, $p < 0.001$, and no significant difference between the second and third, $t(10) = 2.795$, $p = 0.021$, and third and fourth, $t(9) = -0.307$, $p = 0.766$.

In an attempt to understand what might be driving fluency, we ran correlations to investigate whether the total number of responses and the number of responses in each of the four quadrants were related to performance on the only standardised test of BSL for which there was sufficient variance in the scores: the BSL Receptive Skills Test (Herman et al, 1999). The correlation with BSL Receptive Skills score was significant for the first quadrant, $r(10) = 0.674$, $p = 0.033$, but not (at the 2-tailed level) for overall number of items produced, $r(10) = 0.578$, $p = 0.080$, nor for the remaining 3 quadrants, $r(10) = 0.456$, $p = 0.185$, $r(10) = 0.285$, $p = 0.425$, and $r(10) = 0.353$, $p = 0.318$ respectively. Because we had BSL Receptive Skills scores for six of the controls, we added them to the sample, and reran the correlations. While the relationship between Receptive Skills performance and fluency in quadrants two to

four remained insignificant, for the first quadrant it remained significant, $r(16) = 0.6662$, $p = 0.005$, and was now also significant for the total number of items produced, $r(16) = 0.645$, $p = 0.007$. Correlations with such small group sizes have to be treated with caution, but they are consistent with the interpretation that children who are more fluent, particularly in the first fifteen seconds of the task, also have better BSL skills as measured by a sentence comprehension task.

Given the small numbers in the SLI group, it would be misleading to produce a list of the items produced by 33% or more of participants as we did for the children in Study 1. However, the five most common food responses by children with SLI, APPLE, CHIPS, ORANGE, BANANA and CHICKEN, were all produced by more than 33% of the typically developing deaf children in Study 1, as were the top eight animals, CAT, DOG, ELEPHANT, RABBIT, COW, LION, MONKEY, and TIGER.

Finally, it was observed that five children in the SLI group made types of errors that weren't found in the control group. One child, SLI019, fingerspelt EGG incorrectly as g-g-e-e, which could reflect uncertainty with the phonology of the fingerspelt form and/or the orthography of the English word. Four children evidenced word-finding difficulties, and made the following errors. Child SLI004 signed MOUSE IN WHEEL – YOU KNOW – (7 seconds later) HAMSTER! Child SLI027 signed ORANGE BUT NOT HORSE, and never found the correct sign for the animal she was searching for. Child SLI002 signed the letter S, and then the signs for DOG and WHISTLE. He was given credit for DOG, but presumably he was searching for SHEEPDOG. SLI003 created many compound signs which in some instances were acceptable (DOGFISH, CATFISH, GOLDFISH), but in other instances were not (REDBERRY, SEABIRD (not

specific enough – SEAGULL would have been acceptable), SILVERFISH (as a fish, not an insect). There were no examples of any such word-finding behaviours in the control group.

General discussion

We carried out two studies of semantic fluency in children with typical and atypical sign language development. The task probes both the semantic organisation of the lexicon and executive functions related to lexical retrieval. The aim of Study 1 was to investigate semantic fluency in typically developing deaf children, aged 4 to 15 years. The aim of Study 2 was to compare the performance of children with SLI in BSL to a subset of the children in Study 1, matched for chronological age and years of exposure to BSL. Both groups of children produced the same characteristic ‘cognitive signatures’ as are reported for studies of semantic fluency in hearing children and adults, and in signing adults. These were: (i) a decline in the rate of production of new items over the course of the task, (ii) the production of items in semantically-related bursts (‘clusters’), and (iii) production of more prototypical category members by a greater number of participants. It appears that, despite the difference in modality between signed and spoken languages, their lexicons are semantically organised in similar ways.

Although the task can be successfully completed by deaf children who are acquiring a signed language, it proved harder for certain participants: 2/22 children in Study 1, and 2/13 children with SLI in Study 2, were unable to understand the demands of the task, at ages (4 to 10 years) and a further child in Study 1, aged 9, could only do the task

for “animals” and not for “food”. These are ages where no difficulties, as far as we are aware, have been reported for hearing children. For example, in Nelson’s (1974) study, all 63 children aged 4;6-5;7 were able to attempt ‘animals’, and in Nash and Snowling’s (2008) study all 17 children aged 5;6-9;5 were able to respond to ‘animals’ and ‘food’. It is possible that the semantic fluency task is more demanding in BSL, perhaps linked to deaf children having smaller vocabularies. We also speculate that the metalinguistic nature of the task might be challenging for some deaf children, but that with some training they would be able to do it.

Nevertheless, for those participants (the majority) who did complete the task, the number of responses is within the range that has been reported for hearing children in a variety of spoken languages. This is despite our expectations of lower productivity given delayed BSL exposure for many of our participants. Presumably ‘foods’ and ‘animals’ are categories that contain enough early acquired items for deaf children of the age range tested here to be able to produce a similar number of items to hearing children. Very little age of acquisition data is available for foods and animals in BSL, so this is speculation, but it seems plausible. There is only one norming study of BSL with just 20 signers (Vinson, Cormier, Denmark, Schembri & Vigliocco, 2008), and it contains only nine food items (of which ICE CREAM is the earliest acquired, at 3.6 years), and eleven animals (of which DUCK and RABBIT are the earliest acquired at 4.5 years). The semantic task is therefore an appropriate one for use with deaf children who are learning a signed language.

There is nevertheless still room for development beyond the ages that we tested here; the two groups averaged around 15 or 16 items, but adults (Marshall et al, submitted)

averaged 23 or 24. Adults not only produce more clusters (an average of 6, compared to 3.9 and 3.7 for the control and SLI groups respectively in study 2), but their clusters are a little larger, with a mean number of 3.8 per cluster (compared to 3.4 and 3.3 for the control and SLI groups). This indicates that there is development between childhood and adulthood in both the number of lexical items that signers are able to retrieve these categories (as indexed by larger clusters), which is presumably linked to their larger vocabulary size, and in their ability to switch to new clusters in order to continue to retrieve items fluently (as indexed by the number of clusters produced). Given that in Study 1 productivity was very strongly related to the number of clusters rather than to cluster size, it would appear that the development of executive functions is the principal driver of improved performance on this task. Here, as throughout our analysis, we are struck by the comparability of our results compared to those reported for spoken languages: for example, Koren et al (2005) also found that cluster number rather than cluster size drives productivity in Hebrew. We further found that fluency, particularly in the first fifteen seconds, is related to BSL skills as indexed by accuracy on the BSL Receptive Skills test (Herman et al, 1999). Unfortunately there does not exist a standardised vocabulary test for BSL, but it seems likely that fluency is also related to vocabulary skills more generally.

The group of children with SLI in BSL did not differ from the control group on any measure related to the number of responses produced (whether correct or incorrect), types of responses or to anything related to semantic clusters. We therefore conclude that there are no significant differences between the two groups in terms of the types of words that they know, the semantic organisation of their lexicon, or executive functions related to word retrieval. We do of course recognise that this only one

particular semantic task, and other tasks (e.g. the word association task used by Sheng & McGregor, 2010), might probe the organisation of the lexicon in a different and perhaps more sensitive way. We also recognise that significant differences might come to light with a larger sample size, but the population of deaf children with SLI in a signed language is, by its very nature, small. Furthermore, the diagnosis of SLI in a signed language is tentative, as so far we are the only research team to investigate a group of deaf children with SLI: our results need to be replicated by other teams, and in signed languages other than BSL.

Nevertheless, there are two ways in which the SLI group differed from their controls on the semantic fluency task: They produced significantly fewer responses in the first 15 seconds, and there were some examples of word-finding behaviours (although these were not frequent and not demonstrated by every child). We interpret both these differences as resulting from the same underlying cause, namely access to signs being slower in the SLI group. This could be due to slower access to the semantic component of the sign, or to less efficient mapping from the semantic to the phonological form, meaning that the phonological form of the sign is retrieved more slowly or not at all. Slow picture naming, even for successfully retrieved high frequency words, has been reported in hearing children with SLI (Leonard, Nippold, Kail & Hale, 1983), Kail has since taken this work further, and hypothesised that children with SLI have generalised slow processing across a range of linguistic and non-linguistic tasks (Kail, 1994). Similarly, word-finding difficulties in hearing children with SLI were reported in some very early studies of the disorder (Menyuk, 1975; Wiig, Semel & Nystrom, 1982). However, word-finding difficulties are not found in all children with SLI and there is debate over whether these reflect semantic or

phonological impairments (Messer & Dockrell, 2006; Sheng & McGregor, 2010; *inter alia*).

Despite the subtle difficulties of the group of deaf signers with SLI on the semantic fluency task, their overall success on this particular word-level task contrasts with their very poor performance on sentence level tasks (Mason et al, 2010; Morgan et al, 2007) and narrative tasks (Mason et al, 2010; and data for ASL reported in Quinto-Pozos et al, 2011). What emerges from these studies is that for children with SLI in a signed language, it may not be the acquisition of vocabulary that is challenging, but the acquisition of morphology, syntax and discourse-level language. Of course, it is also possible that the potentially slower lexical access we have identified in this study does affect morphosyntactic processing in deaf signers with SLI, but this is a question for future research. Research into SLI in signed languages is only just beginning, but we see that, at least at a broad level, it is remarkably similar to SLI in spoken languages.

References

- Allen, T. E. (1986). Patterns of academic achievement among hearing impaired students: 1974 and 1983. In A. N. Schildroth & M. A. Karchmer (Eds.), *Deaf children in America* (pp. 161–206). San Diego, CA: College-Hill Press.
- Anderson, D. (2006). Lexical development of deaf children acquiring signed languages. In B. Schick, M. Marschark & P. E. Spencer (Eds.), *Advances in the sign language development of deaf children*. Oxford University Press.
- Anderson, D., & Reilly, J. (2002). The MacArthur Communicative Development Inventory: Normative data for American Sign Language. *Journal of Deaf Studies and Deaf Education*, 7, 83-119.
- Blamey, P. J., (2003). Development of spoken language by deaf children. In M. Marschark & P. Spencer (Eds), *Oxford Handbook of Deaf Studies, Language, and Education* (pp. 232-246). Oxford University Press.
- Boucher, J. (1988). Word fluency in high-functioning autistic children. *Journal of Autism and Developmental Disorders*, 18, 637-645.
- Brentari, D., Ed. (2010). *Sign Languages*. Cambridge, UK: Cambridge University Press.
- Brien, D., Ed. (1992). *The Dictionary of British Sign Language/ English*. London, UK: Faber & Faber.

Buriel, Y., Gramunt, N., Bohm, P., Rodes, E., & Peña-Casanova, J. (2004). Verbal fluency: Preliminary normative data in a Spanish sample of young adults (20–49 years of age). [Fluencia verbal. Estudio normativo piloto en una muestra española de adultos jóvenes (20 a 49 años)]. *Neurología*, 19, 153–159.

Cain, K. (2010). *Reading development and difficulties*. Chichester, W. Sussex: BPS Blackwell.

Clark, E. (1993). *The Lexicon in Acquisition*. Cambridge, UK: Cambridge University Press.

Conrad, R. (1979). *The deaf school child: Language and cognitive function*. London: Harper & Row.

Crowe, S. J. & Prescott, T. J. (2003). Continuity and change in the development of category structure: Insights from the semantic fluency task. *International Journal of Behavioural Development*, 27, 467-479.

Edwards, L. (2010). Learning disabilities in deaf and hard-of-hearing children. In Marschark, M., & Spencer, P. E. (Eds) *The Oxford Handbook of Deaf Studies, Language and Education*. Oxford, UK: Oxford University Press. (pp. 425-438).

Elliot, C. D., Smith, P. & McCullough, K. (1996). *British Ability Scales II*. Windsor: NFER-Nelson.

Goldin-Meadow, S., Butcher, C., Mylander, C., & Dodge, M. (1994). Nouns and verbs in a self-styled gesture system: What's in a name? *Cognitive Psychology*, 27, 259-319.

Harrison J. E., Buxton P., Husain M., & Wise R. (2000). Short test of semantic and phonological fluency: Normal performance, validity and test-retest reliability. *British Journal of Clinical Psychology*, 39, 181-191.

Haug, T. (2005) Review of sign language assessment instruments. *Sign Language & Linguistics*, 8, 61–98.

Henry, L. A., Messer, D. J., & Nash, G. (2012). Executive functioning in children with specific language impairment. *Journal of Child Psychiatry and Psychology*, 53, 37-45.

Herman, R., Holmes, S., & Woll, B. (1999). *Assessing BSL Development: Receptive Skills Test*. Forest Bookshop. Gloucestershire, UK.

Herman, R., Grove, N., Holmes, S., Morgan, G., Sutherland, H., & Woll, B. (2004). *Assessing BSL Development: Production Test (Narrative Skills)*. City University Publication.

Kail, R. (1994). A method for studying the generalised slowing hypothesis in children with specific language impairment. *Journal of Speech and Hearing Research*, 37, 418-421.

Kavé, G. (2005). Phonemic fluency, semantic fluency, and difference scores:

Normative data for adult Hebrew speakers. *Journal of Clinical and Experimental Neuropsychology*, 27, 690-699.

Koren, R., Kofman, O., & Berger, A. (2005). Analysis of word clustering in verbal

fluency of school-aged children. *Archives of Clinical Neuropsychology*, 20, 1087-1104.

Kosmidis, M. H., Vlahou, C. H., Panagiotaki, P., & Kiosseoglou, P. (2004). The verbal

fluency task in the Greek population: Normative data, and clustering and switching strategies. *Journal of the International Neuropsychological Society*, 10, 164–172.

Kyle, F..E., & Harris, M. (2006). Concurrent correlates and predictors of reading and

spelling in deaf and hearing school children *Journal of Deaf Studies and Deaf Education*, 11, 273-288.

Lederberg, A. R., & Spencer, P. E. (2009). Word-learning abilities in deaf and hard-of-

hearing preschoolers: effect of lexicon size and language modality, *Journal of Deaf Studies and Deaf Education*, 14, 44-62.

Leonard, L. B. (1998). *Children with Specific Language Impairment*. Cambridge, MA:

MIT Press.

Leonard, L. B., Nippold, M. A., Kail, R., & Hale, C. A. (1983). Picture naming in

language-impaired children. *Journal of Speech and Hearing Research*, 26, 609-615.

Lillo-Martin, D. (1994). Setting the null argument parameters: Evidence from American Sign Language and other languages. In B. Lust, M. Suñer & G. Hermon (Eds.), *Syntactic Theory and First Language Acquisition: Cross Linguistic Perspectives, Volume 2: Anaphora* (pp. 301-318). Hillsdale, NJ: Lawrence Erlbaum Associates.

Lucariello, J., Kyratzis, A., & Nelson, K. (1992). Taxonomic knowledge: What kind and when. *Child Development*, 63, 978-998.

Mahone, C. M., Koth, C. W., Cutting, L., Singer, H. S., & Denckla, M. B. (2001). Executive function in fluency and recall measures among children with Tourette syndrome or ADHD. *Journal of the International Neuropsychological Society*, 7, 102-111.

Marschark, M. (1997). *Raising and educating a deaf child*. New York: Oxford University Press.

Marschark, M., & Woll, B. (in preparation). Deaf children and the vocabulary spurt.

Marshall, C. R., Rowley, K., & Atkinson, J. R. (submitted). Modality-dependent and -independent factors in the organisation of the signed language lexicon: Insights from semantic and phonological fluency tasks in BSL.

Mason, K., Rowley, K., Marshall, C. R., Atkinson, J. R., Herman, R., Woll, B., & Morgan, G. (2010). Identifying specific language impairment in deaf children acquiring

British Sign Language: Implications for theories and practice. *British Journal of Developmental Psychology*, 28, 33-49.

Menyuk, P. (1975). The language-impaired child: Linguistic or cognitive impairment? *Annals of the New York Academy of Sciences*, 263, 59–69.

Messer, D., & Dockrell, J. E. (2006). Children's naming and word-finding difficulties: Descriptions and explanations. *Journal of Speech, Hearing and Language Research*, 49, 309–324.

Mitchell, R., & Karchmer, M. (2004). Chasing the mythical ten percent: Parental hearing status of deaf and hard of hearing students in the United States, *Sign Language Studies*, 4, 138-163.

Morgan, G., Herman, R., & Woll, B. (2007). Language impairments in sign language: Breakthroughs and puzzles. *International Journal of Language and Communication Disorders*, 42, 97-105.

Nash, G., Henry, L. A., & Messer, D. J. (2010). Verbal and non-verbal fluency in children with Specific Language Impairment. Paper presented at the Child Language Seminar, City University, London, June 2010.

Nash, H. M., & Snowling, M. J. (2008). Semantic and phonological fluency in children with Down syndrome: Atypical organization of language or less efficient retrieval strategies? *Cognitive Neuropsychology*, 25, 690-703.

Nelson, K. (1974). Variations in children's concepts by age and category. *Child Development*, 45, 577-584.

Prezbindowski, A. K., & Lederberg, A. R. (2003). Vocabulary assessment of deaf and hard-of-hearing children from infancy through the preschool years. *Journal of Deaf Studies and Deaf Education*, 8, 383- 400.

Quinto-Pozos, D., Forber-Pratt, A., & Singleton, J. (2011). Do developmental communication disorders exist in the signed modality? Reporting on the experience of language professionals and educators from Schools for the Deaf. *Language, Speech and Hearing Services in Schools*, 42, 423-443.

Riva, D., Nichelli, F., & Devoti, M. (2000). Developmental aspects of verbal fluency and confrontational naming in children. *Brain and Language*, 71, 267-284.

Sauzéon, H., Lestage, P., Raboutet, C., N'Kaoua, B., & Claverie, B. (2004). Verbal fluency output in children aged 7–16 as a function of the production criterion: Qualitative analysis of clustering, switching processes, and semantic network exploitation. *Brain and Language*, 89, 192–202.

Sheng, L., & McGregor, K. K. (2010). Lexical-semantic organisation in children with Specific Language Impairment. *Journal of Speech, Language and Hearing Research*, 53, 146-159.

Sutton-Spence, R., & Woll, B. (1998). *The linguistics of British Sign Language: An Introduction*. Cambridge: Cambridge University Press.

Tomblin, B., Records, N., Buckwater, P. Zhang, X., Smith, E., & O'Brien, M., (1997). Prevalence of Specific Language Impairment in kindergarten children. *Journal of Speech, Language and Hearing Research*, 40, 1245-1260.

Troyer, A. K., Moscovitch, M., & Winocur, G. (1997). Clustering and switching as two components of verbal fluency: Evidence from younger and older healthy adults. *Neuropsychology*, 11, 138-146.

Vinson, D., Cormier, K., Denmark, T., Schembri, A., & Vigliocco, G. (2008). The British Sign Language (BSL) norms for age of acquisition, familiarity and iconicity. *Behavior Research Methods*, 40, 1079-1087.

Wiig, E., Semel, E., & Nystrom, L. (1982). Comparison of rapid naming abilities in language-learning-disabled and academically achieving eight-year-olds. *Language, Speech, and Hearing Services in Schools*, 13, 11–23.

Woolfe, T., Herman, R., Roy, P., & Woll, B. (2010). Early vocabulary development in deaf native signers: a British Sign Language adaptation of the communicative development inventories. *Journal of Child Psychology and Psychiatry*, 51, 322 – 331.

Appendix

An example of the coding: Participant N021, category 'Animals'

Seconds after start	Quadrant	English gloss	Response	Correct	Repeat	Irrelevant	Uninterpretable	Switches	Clusters	Items in cluster
1	1	cat	1	1					pet	1
2		dog	1	1						1
3		fish	1	1						1
4		lion	1	1				1	zoo	1
5		tiger	1	1						1
6		monkey	1	1						1
7		bird	1	1				1	bird	1
8		swan	1	1						1
12		snake	1	1				1	reptile	1
13		crocodile	1	1						1
14		shark	1	1				1		
22	2	gorilla	1	1				1		
26		spider	1	1				1		
36	3	giraffe	1	1				1	zoo	1
38		elephant	1	1						1
41		kangaroo	1	1						1
51	4	koala bear	1	1						1
53		monkey	1		1					1
TOTALS			18	17	1	0	0	7	5	15

Table 1. Background information for typically developing children in Study 1.

Participant code	Age (years; months)	Male or female	Years of BSL exposure	Deaf family members?	Type of school	BSL Receptive Skills Test	BSL Narrative Skills Test			Non-sign Repetition Test
							Narrative content	Narrative structure	Grammar	
N001	13;5	M	9;5	No	Mainstream with specialist unit					
N002	6;10	M	2;4	No	Mainstream with specialist unit					
N003	6;4	F	1;10	No	Mainstream with specialist unit					
N004	5;6	F	1;0	No	Mainstream with specialist unit					
N005	13;11	M	13;11	Yes – sibling	Deaf school	112				125
N008	15;2	M	10;8	No	Deaf school	118	25	25	25	116
N009*	14;4	M	11;4	Yes – sibling	Deaf school		50	50	75	125
N010	4;0	F	4;0	Yes – parents	Not yet in school					
N011*	10;5	M	10;5	Yes – parents	Mainstream with specialist unit					
N012*	8;5	M	6;5	No	Mainstream with specialist unit					
N013*	10;11	M	10;11	Yes – parents	Mainstream with specialist unit					
N014*	9;1	M	4;7	No	Mainstream with specialist unit					
N015*	7;6	M	7;6	Yes – sibling	Deaf school					
N016*	9;9	F	4;9	No	Deaf school	95				
N017*	10;0	F	4;0	No	Deaf school	90	25	50	90	
N018*	9;9	F	8;9	Yes – sibling	Deaf school	92	25	75	50	

N019	8;0	M	8;0	Yes – parents	Deaf school	129	75	75	50	
N020*	11;9	M	11;9	Yes – parents	Deaf school	101	25	50	50	
N021*	11;4	M	11;4	No	Deaf school	95	25	90	50	109
N024*	14;10	M	1;6	No	Deaf school	112	50	75	50	
N025	11;5	M	1;0	No	Deaf school	118	50	75	25	
N026*	13;0	F	3;0	No	Deaf school	116	75	50	50	116

Note: The asterisks indicate the children who took part in Study 2, as age-matched controls to the children with SLI.

Table 2: Results for 'food' and 'animals' in Study 1

	Food		Animals		Average across both categories	
	M	(SD)	M	(SD)	M	(SD)
Total number of responses	16.31	(5.18)	15.68	(5.22)	16.00	(4.65)
Correct responses	15.05	(4.47)	14.58	(4.87)	14.82	(4.28)
Repeated responses	0.68	(0.82)	0.58	(1.07)	0.63	(0.76)
Irrelevant responses	0.21	(0.71)	0.16	(0.37)	0.18	(0.54)
Uninterpretable responses	0.37	(0.68)	0.39	(1.01)	0.38	(0.55)
Number of responses in 1 st quadrant (i.e. 1-15s)	7.16	(2.01)	7.89	(2.49)	7.53	(2.07)
Number of responses in 2 nd quadrant (i.e. 16-30s)	3.89	(1.66)	3.53	(1.84)	3.71	(1.13)
Number of responses in 3 rd quadrant (i.e. 31-45s)	3.11	(2.33)	2.32	(1.86)	2.71	(1.66)
Number of responses in 4 th quadrant (i.e. 46-60s)	2.16	(1.46)	1.95	(1.47)	2.05	(1.21)
Number of clusters	4.00	(1.70)	3.79	(1.61)	3.89	(1.40)
Average cluster size	3.16	(0.82)	3.58	(1.20)	3.37	(1.01)
Number of switches	6.58	(2.65)	5.42	(2.67)	6.00	(1.84)

Table 3. Correlation matrix for number of age, total number of responses, correct responses, cluster size, number of clusters and number of switches in Study 1

		Total number of items	Number of correct items	Cluster size	Number of clusters	Number of switches
Age	Correlation	.601**	.648**	.373	.525*	.414
	Sig.	.007	.003	.116	.021	.078
Total number of items	Correlation		.977**	.321	.864**	.774**
	Sig.		<.001	.180	<.001	<.001
Number of correct items	Correlation			.272	.888**	.771**
	Sig.			.260	<.001	<.001
Cluster size	Correlation				-.068	-.192
	Sig.				.784	.431
Number of clusters	Correlation					.738**
	Sig.					<.001
Number of switches	Correlation					
	Sig.					

Table 4. Responses from 33% or more of children in Study 1

Food		Animals	
Response	% children	Response	% children
Chips	58	Lion	84
Chocolate	58	Cat	79
Chicken	53	Dog	68
Meat	53	Giraffe	58
Orange	53	Elephant	53
Sausages	53	Tiger	53
Apple	42	Horse	47
Bread	42	Bird	47
Banana	37	Monkey	47
Burger	37	Cow	42
Crisps	37	Fish	42
Fish	37	Pig	37
Pizza	37	Mouse	37
Potatoes	37	Rabbit	37
		Snake	37
		Zebra	37

Table 5. Background information for participants with SLI in Study 2.

Participant code	Age(years; months)	Male or female	Years of BSL exposure	Deaf family members?	Type of school	BSL Receptive Skills Test	Narrative Skills Test			Non-sign Repetition Test
							Narrative content	Narrative structure	Grammar	
S002	9;3	M	4;9	No	Mainstream with specialist unit	57	<10	<10	<10	80
S003	14;5	M	9;11	No	Mainstream with specialist unit	116	10	10	25	107
S004	14;10	F	10;4	No	Mainstream with specialist unit	78	10	10	10	98
S005	7;5	M	3;0	No	Mainstream with specialist unit	69	<10	<10	<10	84
S006	11;0	M	6;6	No	Mainstream with specialist unit	101	25	10	50	74
S009	9;1	F	4;7	Yes - sibling	Mainstream with specialist unit	66	<25	10	25	113
S010	10;7	M	6;1	Yes - sibling	Mainstream with specialist unit	78	10	10	10	103
S011	10;9	M	6;3	No	Mainstream with specialist unit	56	<10	<10	<10	79
S016	12;8	M	8;2	No	Mainstream with specialist unit	95	<25	<25	<25	85
S019	9;8	M	5;2	No	Deaf school	116	<10	10	<25	93
S027	9;11	F	7;0	No	Deaf school	88	10	25	25	87
S031	9;1	M	7;0	Yes - sibling	Mainstream with specialist unit	85	10	10	10	79
S032	11;3	M	8;0	No	Mainstream with specialist unit	90	10	50	10	96

Table 6. Data for group of children with SLI and their age-matched controls.

Mean across both categories	Control group			SLI group			Independent samples t-test	
	M	(SD)	Range	M	(SD)	Range	t	p
Total number of responses	16.42	(3.47)	11-21	14.80	(4.61)	9.5-24	0.936	0.360
Correct responses	15.13	(3.26)	10.5-20.5	13.10	(4.45)	8-22	1.230	0.233
Repeated responses	0.79	(0.86)	0-2.5	0.50	(0.62)	0-1.5	0.890	0.384
Irrelevant responses	0.13	(0.23)	0-0.5	0.55	(1.40)	0-4.5	-1.038	0.312
Uninterpretable responses	0.38	(0.43)	0-1	0.65	(0.63)	0-2	-1.215	0.238
Number of responses in 1 st quadrant (i.e. 1-15s)	7.88	(1.46)	6-11	6.15	(1.53)	3-8.5	2.698	0.013
Number of responses in 2 nd quadrant (i.e. 16-30s)	3.50	(1.04)	2.5-6	3.80	(1.40)	1-5.5	-0.572	0.571
Number of responses in 3 rd quadrant (i.e. 31-45s)	2.92	(1.46)	1-6	2.35	(1.08)	1-5	1.016	0.322
Number of responses in 4 th quadrant (i.e. 46-60s)	2.13	(0.98)	0.5-3.5	2.50	(1.96)	0-6	-0.583	0.566
Number of clusters	3.88	(1.30)	2-6.5	3.65	(1.73)	2-7.5	0.348	0.746
Average cluster size	3.42	(0.57)	2.9-4.4	3.32	(0.56)	2.8-4.8	0.421	0.530
Number of switches	6.00	(2.03)	3-10	5.00	(2.20)	2.5-8.5	1.107	0.281

Figure 1. Mean total and mean number of correct responses for each participant in Study 1, plotted against age




Figure 2. Rate of decline of responses over the four quadrants of the minute for participants in Study 1

