



City Research Online

City, University of London Institutional Repository

Citation: Fonagy, P., Lemma, A., Target, M., O'Keeffe, S., Constantinou, M. P., Ventura Wurman, T., Luyten, P., Allison, E., Roth, A., Cape, J. & et al (2019). Dynamic interpersonal therapy for moderate to severe depression: A pilot randomized controlled and feasibility trial. *Psychological Medicine*, 50(6), pp. 1010-1019. doi: 10.1017/s0033291719000928

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/22433/>

Link to published version: <https://doi.org/10.1017/s0033291719000928>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Dynamic Interpersonal Therapy for moderate to severe depression: a pilot randomized controlled and feasibility trial

Peter Fonagy^{1,2}, Alessandra Lemma¹, Mary Target¹, Sally O’Keeffe^{1,2}, Matthew P.

Constantinou¹, Tamara Ventura Wurman¹, Patrick Luyten^{1,3}, Elizabeth Allison¹, Anthony Roth¹, John Cape¹ and Stephen Pilling¹

¹ *Research Department of Clinical, Educational and Health Psychology, University College London, London, UK*

² *Anna Freud National Centre for Children and Families (AFNCCF), London, UK*

³ *Faculty of Psychology and Educational Sciences, University of Leuven, Leuven, Belgium*

* Address for correspondence: Peter Fonagy, Research Department of Clinical, Educational and Health Psychology, University College London, 1–19 Torrington Place, London WC1E 7HB, UK. (E-mail: p.fonagy@ucl.ac.uk)

Word count of main text =5000 words

Abstract

Background: Improving Access to Psychological Therapy (IAPT) services treat most patients in England who present to primary care with major depression. Psychodynamic psychotherapy is one of the psychotherapies offered. Dynamic Interpersonal Therapy (DIT), is a psychodynamic and mentalization-based treatment for depression. 16 sessions are delivered over approximately 5 months. Neither DIT's effectiveness relative to low-intensity treatment (LIT), nor the feasibility of randomizing patients to psychodynamic or cognitive-behavioural treatments (CBT) in an IAPT setting, has been demonstrated.

Methods: 147 patients were randomized in a 3:2:1 ratio to DIT (n = 73), LIT (control intervention; n = 54) or CBT (n = 20) in four IAPT treatment services in a combined superiority and feasibility design. Patients meeting criteria for major depressive disorder were assessed at baseline, mid-treatment (3 months) and post-treatment (6 months) using the Hamilton Rating Scale for Depression (HRSD-17), Beck Depression Inventory-II (BDI-II) and other self-rated questionnaire measures. Patients receiving DIT were also followed up 6 months post-completion.

Results: The DIT arm showed significantly lower HRSD-17 scores at the 6-month primary end-point compared with LIT ($d = 0.70$). Significantly more DIT patients (51%) showed clinically significant change on the HRSD-17 compared with LIT (9%). The DIT and CBT arms showed equivalence on most outcomes. Results were similar with the BDI-II. DIT showed benefit across a range of secondary outcomes.

Conclusions: DIT delivered in a primary care setting is superior to LIT and can be appropriately compared with CBT in future RCTs.

Introduction

Depressive disorders constitute a major global health problem (World Health Organization 2001; Lopez *et al.* 2006). Psychosocial treatments are as effective as medication (National Institute for Health and Clinical Excellence 2009; Spielmans *et al.* 2011; Cuijpers *et al.* 2013) and are often preferred by clients (Zimmerman *et al.* 2008; McHugh *et al.* 2013; Hanson *et al.* 2016). Cognitive-behavioural therapy (CBT) has a large evidence base, including data on cost-effectiveness (Cuijpers *et al.* 2013; The Community Mental Health Team 2016).

Interpersonal psychotherapy (IPT), a brief psychotherapy addressing interpersonal problems of individuals with depression, has a smaller evidence base (Lemmens *et al.* 2015). Longer-term recovery is not guaranteed even by CBT or IPT (the ‘best’ evidence-based treatments) (Hollon *et al.* 2006; Karyotaki *et al.* 2016; The Community Mental Health Team 2016).

Choice of therapy is important because no single modality has been firmly established as superior (Cuijpers *et al.* 2008; Cuijpers *et al.* 2011; Cuijpers *et al.* 2013; Cuijpers *et al.* 2014).

Several randomized controlled trials (RCTs) have examined the effectiveness of psychodynamic psychotherapy for depression. Evidence for its efficacy is accumulating (Fonagy 2015; Leichsenring *et al.* 2015). Meta-analytic findings reveal large pre–post treatment effects, with results maintained at long-term follow-up (Driessen *et al.* 2010; Abbass *et al.* 2014; Driessen *et al.* 2015). PDT has been shown to be non-inferior to other active treatments, including cognitive therapy (Driessen *et al.* 2015; Connolly Gibbons *et al.* 2016; Steinert *et al.* 2017). The National Institute for Health and Care Excellence (NICE) (2017) identifies short-term psychodynamic psychotherapy (STPP) as an option for people who have declined CBT and IPT, but there are several implementations.

Some UK providers have adopted Dynamic Interpersonal Therapy (DIT; Lemma *et al.* 2010, 2011) as a ‘prototype’ of STPP. DIT’s structured framework enables therapists without extensive training in psychodynamic psychotherapy to practise. DIT is a time-limited

16-session treatment that formulates the presenting symptoms of depression as responses to interpersonal difficulties or perceived threats to attachments. It is therefore particularly well suited to implementation in a primary-care, community-based context.

The Improving Access to Psychological Therapies (IAPT) programme in England is an internationally unique implementation of evidence-based psychological therapies for common mental disorders in the community (Clark *et al.* 2017). It provides a number of evidence-based treatments for depressive disorders (the most common problem for which psychological therapy is offered (Perfect *et al.* 2016)). In line with guidelines from NICE (2017), CBT is currently offered to most patients presenting with depression in IAPT services; DIT is offered in relatively few services. IAPT is committed to improving access to and widening the choice of psychological therapies (Guy *et al.* 2012). Therefore, evidence for the effectiveness of alternative treatments is crucial. DIT's effectiveness as a STPP for patients with depression within community services has not yet been established. The REDIT (Randomized Evaluation of Dynamic Interpersonal Therapy) study aimed to (i) evaluate DIT's effectiveness for moderate to severe depression in adults within IAPT, comparing the outcomes associated with DIT with a low-intensity treatment (LIT) condition, and (ii) evaluate the feasibility of comparing the treatment of adult depression with DIT and CBT within IAPT services.

Method

Trial design

The trial was conducted within IAPT services, which are based on a stepped-care model, where patients recognized as depressed in primary care or hospital settings (Step 1) are offered triage or low-intensity (Step 2) or high-intensity (Step 3) treatment, depending on clinical need. Triage followed the IAPT protocol and focused on evaluating severity and

suicide/self-harm risk. Low-intensity treatment (LIT), delivered by a Psychological Wellbeing Practitioner (PWP), involves guided self-help. High-intensity treatment typically involves face-to-face treatment, such as CBT, with an accredited therapist. Participants were recruited from four sites: two for the DIT *v.* LIT comparison, and two separate sites for the DIT *v.* CBT comparison. Participants in the low-intensity condition who had not recovered were offered high-intensity therapy after completing 4–6 months of guided self-help treatment. In parallel, at two IAPT sites, patients were randomized to either DIT or CBT. The trial was granted ethical approval by NHS Research Ethics Committees and was registered with the ISRCTRN Registry (ISRCTRN38209986; ISRCTRN06629587).

Sample size calculation

Recruitment was powered for a superiority trial of DIT *v.* LIT. An *a priori* power analysis suggested that a sample size of 54 (27 in each group) was required to detect a mean difference of 5 ($SD = 5.62$) on the Hamilton Rating Scale for Depression (HRSD) at 90% power and a 5% type I error rate. The study was thus well powered to detect superiority in the DIT *v.* LIT comparison ($n_{DIT} = 48$ and $n_{LIT} = 31$ at 6 months). There was no intention to incorporate a CBT *v.* LIT comparison ($n_{CBT} = 15$ at 6 months) because the CBT arm was included only to assess the feasibility of comparing DIT and CBT¹. We report on results obtained from CBT without comment other than in relation to the acceptability, attrition, and timing of assessments for the sake of comprehensiveness.

Participants

Recruitment took place from November 2012 to January 2015. Patients consecutively

¹ A formal test of equivalence between DIT and CBT groups will require 205 patients in each group, based on a Type I error rate of 5%, 90% power, an equivalence limit of 2 on the HRSD, and a population standard deviation of 5.62 (the mean of 52 standard deviations of post-CBT HRSD scores; Cuijpers *et al.* 2010).

referred to four metropolitan IAPT sites, aged 18 and above, who met DSM-IV criteria for a major depressive episode (American Psychiatric Association 1994) established by research staff using the MINI Plus 6.0 interview (Sheehan *et al.* 1998), scored >14 on the HRSD and >10 on the Patient Health Questionnaire (PHQ-9) (Kroenke *et al.* 2001), and were identified as likely to require high-intensity treatment after triage, were approached to participate. As in most instances there would be significant delay in implementing high-intensity treatments, the referral to a low-intensity treatment with the offer of high-intensity treatment to follow if required was considered ethically acceptable. Exclusion criteria were current psychotic symptoms or bipolar disorder, clinical contraindications for short-term psychotherapy including current use of antipsychotic medication, historical clinical diagnosis of Axis II disorder, historical or current repeated non-suicidal self-injury, disclosure of current suicidal intent and plans, historical or current eating disorder, and historical or current substance abuse. Furthermore, non-English speakers, those who had participated in another clinical trial within the past year in which they had received CBT or STPP for depression, those who had had previous unsuccessful CBT, and those with highly unstable or insecure life arrangements were excluded from the trial. Figure 1 shows recruitment to the trial.

Assessments

Clinicians at recruiting IAPT sites referred suitable patients to the trial team. Referred patients were given a baseline assessment by a research assistant to assess eligibility. This included structured clinical interviews and self-report measures. Participants were followed up mid-treatment (3 months; approximately 90 days) and after the end of treatment (6 months; approximately 180 days). Those in the DIT arm were followed up on average 12 months (approximately 360 days) post-randomization to establish whether treatment gains were maintained.

Randomization

Participant randomization was undertaken by an administrator independent of the trial, based at a different location from the research team and blind to the hypotheses or trial conditions. Following the completion of a baseline assessment by the research team, minimization criteria (including sex, severity of depression and age) were e-mailed with the request for randomization. Depending on the site at which they were recruited, participants were randomized to either the DIT or LIT condition, or the DIT or CBT condition, using a minimization algorithm with an 80% bias to minimize imbalance in a ratio of 3:2:1 for the three treatment groups.

Interventions

Dynamic Interpersonal Therapy. DIT is a brief dynamically oriented intervention informed by attachment and mentalization theory (Lemma *et al.* 2010, 2011; Luyten and Blatt 2012) which owes much to conceptualizations of dynamic therapy advanced by Luborsky and Crits-Christoph (1998) and Kernberg (1988). It views symptoms of depression and anxiety as responses to interpersonal difficulties or perceived threats to attachments (loss/separation) and hence also as threats to the self. DIT aims to help patients improve their ability to cope with current attachment-related interpersonal challenges through better understanding their subjective reactions to them as threats, making implicit anxieties and concerns explicit, and improving their ability to reflect on their own and others' thoughts and feelings. DIT helps patients understand the connection between their presenting symptoms and their relationships by identifying a core, non-conscious, repetitive pattern of relating. This pattern, termed the interpersonal affective focus, becomes the therapy's central focus, and the therapist works on the conscious cost of such repetitive patterns. DIT primarily targets the assumed dynamic

origins of depressive symptoms and, secondarily, enhanced interpersonal functioning through improved capacity to reflect on the individual's own states of mind as well as those of others.

DIT was delivered within IAPT services by trained DIT practitioners with an approved DIT supervisor. Participants were offered 16 weekly sessions, each lasting approximately 1 hour.

Low-intensity treatment. LIT had a comparable time span to DIT. It consisted of fortnightly individual sessions with a PWP using a self-guided manual-based programme commonly used in IAPT services (Coull and Morris 2011). The LIT condition was designed to reflect interventions usually offered at Step 2 within IAPT services, including guided self-help, psychoeducation, and face-to-face and telephone support using NICE-recommended interventions. The maximum number of sessions offered was nine. After the 16–24-week LIT period, all participants who had not recovered were offered high-intensity therapy (CBT, IPT, counselling or DIT).

Cognitive-behavioural therapy. CBT was delivered within IAPT services by trained CBT practitioners under supervision following the standard IAPT high-intensity protocol (Roth and Pilling 2015). Therapists followed a cognitive therapy model (Beck and Haigh 2014). Clients were offered 14–18 sessions of CBT over a 16–24-week period, each session lasting approximately 1 hour.

Trial therapists

Seventeen DIT therapists saw an average of 4.2 patients (SD = 3.0, range: 1–12). All sessions were recorded and up to three sessions per case (M = 2.5, SD = 0.9, range: 1–3), sampled from the beginning, middle and end of each treatment, were assessed using the DIT

adherence/competence measure (Lemma *et al.* 2012), rated by the developers (M.T. and A.L.) and an expert rater independent of the study. Agreement between the developers and the independent rater was high (intraclass correlation coefficient = 0.82).

Ten PWPs offered LIT, each seeing 5.3 patients on average (SD = 3.0, range: 3–10). Their sessions were recorded, but not coded for fidelity, although PWPs received routine supervision.

Thirteen CBT therapists saw an average of 1.54 patients (SD = 0.66, range: 1–3); their sessions were recorded and they were routinely supervised.

Outcomes

The prespecified primary outcome measure was change in mean scores on the 17-item version of the HRSD at 6 months indicating full or partial remission (Hamilton 1960), rated by research assistants who were blind to treatment allocation. Secondary outcome measures included the Beck Depression Inventory-II (BDI; Beck *et al.* 1996), Brief Symptom Inventory (BSI; de Beurs and Zitman 2006), revised Social Adjustment Scale (SAS; Weissman *et al.* 1978), EQ-5D (The EuroQol Group 1990), Experiences in Close Relationships Questionnaire-Revised (ECR-R; Fraley *et al.* 2000) and Inventory of Interpersonal Problems-64 (IIP-64; Horowitz *et al.* 1988).

Statistical analyses

Therapeutic benefit was assessed by end-point scores and the rate of change on the HRSD-17 (primary outcome) and BDI-II (secondary outcome) using linear mixed-effects models and marginal means. Models included the fixed effects of treatment group (with DIT as the

reference group², e.g., DIT v. LIT and DIT v. CBT), linear time (baseline = 0, 3 months = 1, 6 months = 2), quadratic time (baseline = 0, 3 months = 1, 6 months = 4), age, sex, ethnicity, marital status, higher education status, income bracket, concurrent medical problems, and assessment variability (days until assessment relative to the respective assessment period), as well as the random effects of patient and time. We tested a random intercept for randomization site but this explained almost no variance when covariates were added to the model (see Supplement 3 online). Treatment differences between the DIT and LIT/CBT groups at the primary end-point (6 months) were evaluated with marginal means and Wald chi-square tests.

All available data post-randomization was used (intention-to-treat analysis). There were no missing observations on the HRSD-17 or BDI-II at baseline. However, 24% of observations on the HRSD-17 and BDI-II were missing at mid-treatment (3 months), and 35% at post-treatment (6-months; 57% of missing observations occurred at both mid- and post-treatment). Sixty percent of DIT cases were lost to follow-up. Covariates also showed small amounts of missing data: higher education status (7%), marital status (5%), ethnicity (5%), and income (24%). There were no signs of major bias in the cases showing missing values on the outcome measures or covariates, but for completeness we replicated the analysis while handling missing data with multiple imputation by chained equations (see Supplement 2 online).

Reliable and clinically significant change on the HRSD-17 and BDI-II was calculated using Jacobson and Truax's (1991) criteria. Specifically, reliable change indices were calculated using the sample's baseline standard deviation on each measure, and reliability estimates (Cronbach's α) from meta-analyses [e.g., HRSD-17 = .79 (Trajkovic *et al.* 2011);

² We collapsed the DIT groups over all randomization sites. Sensitivity analyses showed that outcomes did not vary by randomization site or study (e.g., pilot trial or feasibility study; see Supplement 3 online).

BDI-II = .89 (Erford *et al.* 2017)]. Reliable improvement (or deterioration) was defined as pre–post difference scores >4.81 (or <-4.81) on the HRSD-17, and >7.60 (or <-7.60) on the BDI-II. No reliable change was defined as pre–post difference scores between -4.81 and 4.81 on the HSRD, and between -7.60 and 7.60 on the BDI-II. Validated clinical cut-offs defined clinically significant improvement. For instance, baseline HRSD-17 scores >7 or BDI-II scores >12 (i.e. mild to severe depression) and 6-month HRSD-17 scores <8 or BDI-II scores <13 (i.e. no depression) defined clinically significant improvement (Dozois *et al.* 1998; Zimmerman *et al.* 2013). The proportion of patients in each treatment group showing reliable improvement, deterioration, no reliable change, and clinically significant change was compared using logistic regressions while controlling for covariates (age, sex, ethnicity, marital status, higher education status, income bracket, concurrent medical problems, and end-point assessment variability). Results were also replicated while handling missing data with multiple imputation (see Supplement 2 online).

Additional secondary measures included the ECR-R, BSI, EQ-5D health status measure, IIP-64, and SAS. Missing data rates for these measures were non-ignorable (range: 5–40% at baseline and 34–53% at end of treatment). As the missing values appeared Missing At Random (see Supplement 2 online), we applied multiple imputation to handle missingness. We imputed and analysed all subscales apart from the child, family, and marital subscales of the SAS due to disproportionate amounts of missing data ($>70\%$). Treatment differences on each secondary outcome at the primary end-point were evaluated using linear regressions on the imputed datasets, with treatment contrast (DIT *v.* LIT, DIT *v.* CBT), HRSD-17 and BDI-II scores at baseline, age, sex, and end-point assessment variability. To ensure that no further bias was introduced by the imputation model, we replicated the analyses with the observed data (see Supplement 2 online).

Results

Sample characteristics

The final sample included 147 patients (39% mild, 47% moderate, and 14% severe on the HRSD-17, and 6% mild, 23% moderate, and 71% severe on the BDI-II). Of these, 73 were randomized to DIT, 20 to CBT and 54 to LIT, until the end of DIT treatment (minimal treatment control). Three percent of observations were discarded because the discrepancy between the actual and planned assessment period was too large (greater than a month) to assign the observation to a specific time-point.

Table 1 summarizes demographics for the three groups. They did not differ significantly on any demographic variable, but there was a trend towards a difference on the percentage of patients with long-term health conditions (e.g. diabetes); therefore, a binary variable coding for medical problems was used as covariate in all subsequent analyses. The treatment groups showed similar levels of moderate to severe depression at baseline.

Primary Outcome

Analyses were based on an intent-to-treat sample of 147 patients using all available data. A mixed-effects linear regression that included linear and quadratic slopes for time, treatment contrast (DIT *v.* LIT, DIT *v.* CBT), covariates (age, sex, higher education status, marital status, ethnicity, income bracket, co-occurring medical problems, and assessment variability; see Supplementary Table 1 for regression coefficients), a random intercept for patient, and random linear slope for time, fitted the data well (fixed portion: $\chi^2(17) = 137.71, p < .001$; random portion: $\chi^2(2) = 81.05, p < .001$). Patients varied in their baseline scores (random intercept = 7.96, 95% CI [4.85, 12.07]) and amount of change over time (random slope = 5.49, 95% CI [3.15, 9.57]).

HRSD-17 scores showed a linear decline over time ($b = -3.81, z = -3.11, p = .002$,

95% CI [-6.21, -1.41]). The rate of change also differed between treatment groups: the LIT group showed a stronger quadratic pattern of change compared with the DIT group ($b = 2.75$, $z = 2.89$, $p = .004$, 95% CI [0.89, 4.62]). That is, the DIT group showed a steady rate of decline over the treatment period, while the LIT group plateaued between the middle and end of treatment (Fig. 2). A separate model that included all four assessment points for the DIT group only (e.g., baseline, mid-treatment, post-treatment, and follow-up) showed that there was no significant difference between marginal HRSD-17 scores at post-treatment and follow-up (post-treatment = 10.0 *v.* follow-up = 10.7; Wald $\chi^2(1) = 0.28$, $p = .596$, 95% CI [-2.28, 3.97]). Therefore, the DIT group maintained their gains at follow-up.

At post-treatment, the DIT group scored significantly lower on the HRSD-17 than the LIT group (marginal means: 10.0 *v.* 14.8; Wald $\chi^2(1) = 8.47$, $p = .004$, 95% CI [1.59, 8.16]; marginal $d = .70$, 95% CI [0.24, 1.16]). There were no significant differences between the DIT and CBT groups in end-point marginal means (DIT = 10.0, CBT = 13.2, $\chi^2(1) = 2.80$, $p = .094$, 95% CI [-0.56, 7.07]; $d = .50$, 95% CI [-0.07, 1.07]).

A logistic regression model with treatment contrast (DIT *v.* LIT, DIT *v.* CBT), baseline HRSD-17 scores, and covariates (age, sex, higher education status, marital status, ethnicity, income bracket, co-occurring medical problems, and end-point assessment variability) showed that more DIT patients achieved clinically significant change than LIT patients (marginal percentages: 51% *v.* 9%; $\chi^2(1) = 14.70$, $p < .001$, 95% CI [-.64, -.21]; RR = 5.17, 95% CI [1.75, 15.25]) and CBT patients (51% *v.* 20%; $\chi^2(1) = 5.94$, $p = .01$, 95% CI [-.56, -.06]; RR = 2.55, 95% CI [1.00, 6.54]). All treatment groups showed moderate-to-high levels of reliable improvement, moderate levels of no reliable change, and low levels of deterioration (see Table 2). Sensitivity analyses demonstrated that the results were maintained when (i) multiple imputation was used to handle missing data on the HRSD-17 (see Supplement 2 online), (ii) the BDI-II was used as the outcome measure (see Supplement 1

online and Supplementary Table 4), and (iii) mild cases were excluded (see Supplement 3 online).

Secondary outcomes

We used linear regression models with multiple imputation to examine treatment differences on each secondary outcome subscale at post-treatment. Models included treatment group (DIT v. LIT, DIT v. CBT), baseline scores on the subscale in question, and the covariates listed above. Marginal means at post-treatment are described below and reported in Table 2. Marginal means at baseline are presented in Supplementary Table 7 (available online).

The DIT group reported lower end-point scores on the BSI compared with the LIT group, specifically in depression (1.03 v. 1.87; $t(141) = 3.58, p = 0.001, 95\% \text{ CI } [0.37, 1.32], d = 0.67, 95\% \text{ CI } [0.30, 1.02]$), anxiety (1.00 v. 1.52; $t(141) = 2.29, p < .026, 95\% \text{ CI } [0.06, 0.98], d = .43, 95\% \text{ CI } [0.08, 0.79]$), obsessive-compulsiveness (1.26 v. 2.07; $t(141) = 3.28, p = .002, 95\% \text{ CI } [.31, 1.29], d = .59, 95\% \text{ CI } [.23, .95]$), psychoticism (0.77 v. 1.17; $t(141) = 2.42, p = .01, 95\% \text{ CI } [.07, .75], d = .44, 95\% \text{ CI } [.08, .79]$), general severity (0.90 v. 1.35; $t(141) = 2.63, p = .01, 95\% \text{ CI } [0.11, 0.81], d = .49, 95\% \text{ CI } [.13, .84]$), and symptom distress (1.61 v. 2.13, $t(141) = 2.98, p = .005, 95\% \text{ CI } [0.17, 0.88], d = .57, 95\% \text{ CI } [.21, .93]$). There were no significant differences between the DIT and CBT groups.

On the SAS, the DIT group reported fewer social problems overall compared with the LIT group (2.24 v. 2.50; $t(141) = 2.47, p = 0.016, 95\% \text{ CI } [.05, .57], d = 0.45, 95\% \text{ CI } [.09, .81]$). Specific improvements were found at work (1.99 v. 2.36; $t(141) = 2.66, p = 0.01, 95\% \text{ CI } [.09, .65], d = 0.45, 95\% \text{ CI } [.09, .80]$) and with friends (2.39 v. 2.72; $t(141) = 2.48, p = 0.016, 95\% \text{ CI } [.07, .60], d = 0.44, 95\% \text{ CI } [.08, .80]$) compared with the LIT group.

Finally, DIT patients reported higher health-related quality of life scores on the EQ-5D than LIT patients (69.66 v. 60.14; $t(141) = -3.02, p = 0.003, 95\% \text{ CI } [-15.81, -3.24], d =$

0.55, 95% CI [0.19, 0.90]). Results were replicated using the observed rather than imputed data (see Supplementary Table 2).

Therapist fidelity and competence

Competency ratings were high for all DIT therapists ($M = 53.3$, $SD = 10.6$, range: 19–65). All therapists were coded as adherent on 80% of recordings (see Supplementary Table 2).

We explored whether competency ratings predicted treatment outcomes. A mixed-effects model was used to predict HRSD or BDI-II scores at 6 months from the fixed effects of therapist competence (i.e., competence ratings averaged across sessions for the 52 participants who completed DIT) and covariates (age, sex, education, medical problems, and assessment variability at 6 months), and random effects of baseline HRSD (or BDI-II) scores. Random effects for therapist and site were not included as they did not significantly improve model fit ($\chi^2(2) = 0.013$, $p > 0.05$). Therapist competence did not significantly predict 6-month HRSD ($t(34.81) = 0.483$, $p > 0.05$) or BDI-II ($t(23.43) = 0.459$, $p > 0.05$) scores. Results were no different when using competence ratings at the initial, mid, or late phases of treatment, or when using a log-normal distribution to account for the mild skewness in residuals.

Discussion

This is the first RCT investigating the effect of DIT on moderate to severe depression and the first trial of STPP in an IAPT context. Participants randomized to DIT reported significantly lower levels of depression following treatment than those who received LIT on both clinician-rated (HRSD-17) and self-report (BDI-II) measures. Over half of DIT patients made clinically significant improvements, i.e. full recovery, while only 9% of LIT patients achieved this. The benefits of DIT were maintained at 12-month follow-up. At post-

treatment, the DIT group showed reduced overall symptom severity and distress, better social adjustment, and fewer health-related problems than the LIT group. Benefits on measures of attachment (ECR-R; Fraley *et al.* 2000) and interpersonal problems (IIP-64) were less marked. The findings are nevertheless encouraging as, unlike many other trials of psychodynamic psychotherapy (Gerber *et al.* 2011), this study made use of an active control group (LIT) that represents a bona fide form of lower-intensity treatment actually regularly provided within the IAPT system by providers who believe in the method.

These results are consistent with meta-analytic findings that reveal medium or large pre–post treatment effects of psychodynamic therapy, with results maintained at longer-term follow-up (Abbass *et al.* 2014; Driessen *et al.* 2015; Connolly Gibbons *et al.* 2016). This study offers further evidence for STPP in the treatment of depression with moderate severity, and confirms that these effects are maintained at least in the short term. The relatively large number of trial therapists involved increases confidence in the generalizability of the findings. The current trial was not designed to investigate equivalence between STPP and CBT, but results suggest that an equivalence trial could be performed.

The NICE depression guidelines recommend CBT as a first-line psychological intervention because of its relatively strong evidence base. This study adds to the evidence base for STPP. Recently published data for 2015–2016 show that the vast majority of patients were offered CBT and counselling, with DIT offered to only 1% of patients in England (The Community Mental Health Team 2016). This may reflect the lack of RCTs supporting the specific implementation of dynamic therapy or, more likely, the lack of trained staff able to deliver STPP. DIT is unique among the psychodynamic therapies in offering a treatment manual and curriculum to enable those without backgrounds in psychodynamic therapies to deliver it (Lemma *et al.* 2017). It is part of a family of dynamic treatments designed to be delivered by a broad range of healthcare professionals to optimize accessibility to these

approaches (Bateman and Fonagy 2009; Fonagy *et al.* 2014; Bateman *et al.* 2016). DIT may have a significant role in increasing the general availability of psychodynamic approaches by increasing the accessibility of training in this modality. There have been a number of different trials of psychodynamic for therapy for depression based on different manuals, but DIT attempts to be unique by providing an introduction to psychoanalytic ideas and introducing a clinical approach to those with very limited experience of working with these models in the hope of making a dynamic approach available at scale. The evaluation of DIT training protocols in relation to clinical outcomes is the next hurdle in the dissemination of this approach within IAPT.

This paper also reports a smaller-scale study, which aimed to assess the feasibility of an RCT comparing DIT and CBT for depression. Participants readily accepted randomization to DIT and CBT, per-protocol treatments and assessments were delivered to almost all participants, and 6-month assessments indicated the potential for meaningful comparisons between DIT and CBT. No patient, when offered randomization to DIT or CBT refused being randomized and once assigned refused to take up the treatment. Although the proportion of early terminations was significant (approximately 25% in each group), it was no higher than may be expected in routine care at these IAPT sites. Patients found DIT acceptable. 85% of the CBT group and 80% of the DIT group were available for assessments at 6 months, indicating comparable fidelity to the research protocol. The findings support the feasibility of randomizing individuals to either of these high-intensity therapies in the context of routine service provision and thus provide the basis for implementing a protocol to test the hypothesis that certain patients with major depressive disorder and co-occurring disorders may specifically benefit from STPP or CBT. On the basis of this preliminary investigation, given the large number of patients with depression currently seen in IAPT, we plan to examine the hypothesis that DIT may be considered as a potential alternative to CBT in an

adequately powered RCT comparing the two treatments.

The study has a number of important limitations. The LIT group, while realistic as a comparison given the stepped-care protocol implemented in IAPT settings, is not an adequate control, as participants may not have had comparable expectations of benefit in the two arms of the trial. Participants agreed to randomization to LIT (guided self-help) in the knowledge that they would be offered a high-intensity psychological therapy at the end of that treatment if they felt they needed it. Many participants did not take up this offer, but the LIT condition has features in common with placebo designs, which may have slightly worse outcomes for depressed patients than no-treatment control conditions (Furukawa *et al.* 2014). Given that participants may have perceived the offer of further treatment as contingent on their non-responsiveness to the LIT intervention, it is possible that the observed impact of DIT was exaggerated in this comparison. However, although comparison with the CBT arm was not part of the trial design, the lack of a marked difference in the outcomes of the two high-intensity arms suggests that the superiority of DIT to LIT may be genuine. While adherence to the assessment and treatment protocol was relatively good, a substantial proportion of the sample failed to complete the additional outcome measures, and multiple imputations were needed to provide adequate analysis of the data. Furthermore, 3% of the assessments fell outside the pre-specified window and had to be discarded, further reducing the power of available contrasts. While the size of the sample for the key comparison of DIT *v.* LIT was adequate, the study is relatively small.

This study also investigated whether therapist competence was associated with outcomes. No evidence of this was found, perhaps because the DIT therapists working in the study were all functioning significantly above a minimal level of competence and were supervised by the developers. The fact that neither adherence nor competence were examined in the CBT pilot group is a further limitation of the study highlighting the need for caution in

relation to these findings.

Conclusion

As IAPT services offer psychodynamic therapy, its efficacy in real-world settings is highly relevant to patients, therapists, and the healthcare system. Evidence for the efficacy of psychodynamic therapy for common mental disorders is available (Leichsenring *et al.* 2015), but this study is the first to demonstrate DIT's effectiveness in primary care. Encouraging remission rates were obtained, particularly given the severity of the sample and the low recovery rate in the LIT condition. Further work is warranted in the IAPT setting to identify patient groups for whom DIT may be indicated. We noted informally that referrals to DIT were often individuals with adverse childhood experiences, possibly because of the implicit belief that DIT may be better at addressing persistent distortions of relationship representations, perhaps associated with a history of trauma. This, in turn, in the context of large-scale effectiveness studies will require the implementation of DIT at scale in the IAPT setting.

Financial support

This work was supported by the National Institute for Health Research (NIHR) under its Research for Patient Benefit Programme (grant number PB-PG-0610-22287). P.F. is in receipt of a NIHR Senior Investigator Award (grant number NF-SI-0514-10157) and was in part supported by the NIHR Collaboration for Leadership in Applied Health Research and Care (CLAHRC) North Thames at Barts Health NHS Trust. S.P. is in receipt of funding from the National Institute for Health and Care Excellence (grant number 541506) and the NIHR (grant number 541918), and receives support from the UCLH Biomedical Research Centre. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR

or the Department of Health.

Declaration of interest

P.F. delivers training on DIT organized by the AFNCCF, of which he is Chief Executive Officer. He also receives royalties from books related to DIT. A.L. and M.T. deliver training on DIT organized by the AFNCCF, for which they are paid, and receive royalties from books related to DIT. P.L. has been involved in the development, training and evaluation of psychodynamic treatments. S.O’K., M.C., T.V.W., E.A., A.R., J.C., and S.P. have no interests to declare.

References

- Abbass, A. A., Kisely, S. R., Town, J. M., Leichsenring, F., Driessen, E., De Maat, S., Gerber, A., Dekker, J., Rabung, S., Rusalovska, S. & Crowe, E. (2014). Short-term psychodynamic psychotherapies for common mental disorders. *Cochrane Database of Systematic Reviews*, CD004687.
- American Psychiatric Association** (1994). *Diagnostic and statistical manual of mental disorders: DSM-IV*. American Psychiatric Association: Washington, DC.
- Bateman, A. & Fonagy, P.** (2009). Randomized controlled trial of outpatient mentalization-based treatment versus structured clinical management for borderline personality disorder. *American Journal of Psychiatry* **166**, 1355-1364.
- Bateman, A., O'Connell, J., Lorenzini, N., Gardner, T. & Fonagy, P.** (2016). A randomised controlled trial of mentalization-based treatment versus structured clinical management for patients with comorbid borderline personality disorder and antisocial personality disorder. *BMC Psychiatry* **16**, 304.
- Beck, A., Steer, R. & Brown, G.** (1996). *Manual for the Beck Depression Inventory-II (BDI-II)*. Psychological Corporation: San Antonio.
- Beck, A. T. & Haigh, E. A.** (2014). Advances in cognitive theory and therapy: The generic cognitive model. *Annual Review of Clinical Psychology* **10**, 1-24.
- Clark, D. M., Canvin, L., Green, J., Layard, R., Pilling, S. & Janecka, M.** (2017). Transparency about the outcomes of mental health services (IAPT approach): An analysis of public data. *Lancet*.
- Connolly Gibbons, M. B., Gallop, R., Thompson, D., Luther, D., Crits-Christoph, K., Jacobs, J., Yin, S. & Crits-Christoph, P.** (2016). Comparative effectiveness of cognitive therapy and dynamic psychotherapy for major depressive disorder in a community mental health setting: A randomized clinical noninferiority trial. *JAMA Psychiatry* **73**, 904-911.
- Coull, G. & Morris, P. G.** (2011). The clinical effectiveness of CBT-based guided self-help interventions for anxiety and depressive disorders: a systematic review. *Psychological Medicine* **41**, 2239-2252.
- Cuijpers, P., Berking, M., Andersson, G., Quigley, L., Kleiboer, A. & Dobson, K. S.** (2013). A meta-analysis of cognitive-behavioural therapy for adult depression, alone and in comparison with other treatments. *Canadian Journal of Psychiatry* **58**, 376-385.
- Cuijpers, P., Geraedts, A. S., van Oppen, P., Andersson, G., Markowitz, J. C. & van Straten, A.** (2011). Interpersonal psychotherapy for depression: A meta-analysis. *American Journal of Psychiatry* **168**, 581-592.
- Cuijpers, P., Koole, S. L., van Dijke, A., Roca, M., Li, J. & Reynolds, C. F., 3rd** (2014). Psychotherapy for subclinical depression: Meta-analysis. *British Journal of Psychiatry* **205**, 268-274.
- Cuijpers, P., van Straten, A., Andersson, G. & van Oppen, P.** (2008). Psychotherapy for depression in adults: A meta-analysis of comparative outcome studies. *Journal of Consulting and Clinical Psychology* **76**, 909-922.
- de Beurs, E. & Zitman, F. G.** (2006). De Brief Symptom Inventory (BSI). De betrouwbaarheid en validiteit van een handzaam alternatief voor de SCL-90. *Maandblad Geestelijke Volksgezondheid* **61**, 120-141.
- Dozois, D. J. A., Dobson, K. S. & Ahnberg, J. L.** (1998). A psychometric evaluation of the Beck Depression Inventory-II. *Psychological Assessment* **10**, 83-89.
- Driessen, E., Cuijpers, P., de Maat, S. C. M., Abbass, A. A., de Jonghe, F. & Dekker, J. J. M.** (2010). The efficacy of short-term psychodynamic psychotherapy for depression: a meta-analysis. *Clinical Psychology Review* **30**, 25-36.
- Driessen, E., Hegelmaier, L. M., Abbass, A. A., Barber, J. P., Dekker, J. J., Van, H. L.,**

- Jansma, E. P. & Cuijpers, P.** (2015). The efficacy of short-term psychodynamic psychotherapy for depression: A meta-analysis update. *Clinical Psychology Review* **42**, 1-15.
- Erford, B. T., Johnson, E. & Bardoshi, G.** (2017). Meta-analysis of the English version of the Beck Depression Inventory–Second Edition. *Measurement and Evaluation in Counseling and Development* **49**, 3-33.
- Fonagy, P.** (2015). The effectiveness of psychodynamic psychotherapies: an update. *World Psychiatry* **14**, 137-150.
- Fonagy, P., Rossouw, T., Sharp, C., Bateman, A., Allison, L. & Farrar, C.** (2014). Mentalization-based treatment for adolescents with borderline traits. In *Handbook of borderline personality disorder in children and adolescents* (ed. C. Sharp and J. L. Tackett), pp. 313-332. Springer: New York.
- Fraley, R. C., Waller, N. G. & Brennan, K. A.** (2000). An item response theory analysis of self-report measures of adult attachment. *Journal of Personality and Social Psychology* **78**, 350-365.
- Furukawa, T. A., Noma, H., Caldwell, D. M., Honyashiki, M., Shinohara, K., Imai, H., Chen, P., Hunot, V. & Churchill, R.** (2014). Waiting list may be a nocebo condition in psychotherapy trials: a contribution from network meta-analysis. *Acta Psychiatrica Scandinavica* **130**, 181-192.
- Gerber, A. J., Kocsis, J. H., Milrod, B. L., Roose, S. P., Barber, J. P., Thase, M. E., Perkins, P. & Leon, A. C.** (2011). A quality-based review of randomized controlled trials of psychodynamic psychotherapy. *American Journal of Psychiatry* **168**, 19-28.
- Guy, A., Loewenthal, D., Thomas, R. & Stephenson, S.** (2012). Scrutinising NICE: The impact of the National Institute for Health and Clinical Excellence Guidelines on the provision of counselling and psychotherapy in primary care in the UK. *Psychodynamic Practice* **18**, 25-50.
- Hamilton, M.** (1960). A rating scale for depression. *Journal of Neurology, Neurosurgery & Psychiatry* **23**, 56-62.
- Hanson, K., Webb, T. L., Sheeran, P. & Turpin, G.** (2016). Attitudes and preferences towards self-help treatments for depression in comparison to psychotherapy and antidepressant medication. *Behavioural and Cognitive Psychotherapy* **44**, 129-139.
- Hollon, S. D., Stewart, M. O. & Strunk, D.** (2006). Enduring effects for cognitive behavior therapy in the treatment of depression and anxiety. *Annual Review of Psychology* **57**, 285-315.
- Horowitz, L. M., Rosenberg, S. E., Baer, B. A., Ureno, G. & Villasenor, V. S.** (1988). Inventory of Interpersonal Problems: psychometric properties and clinical applications. *Journal of Consulting and Clinical Psychology* **56**, 885-892.
- Jacobson, N. S. & Truax, P.** (1991). Clinical significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology* **59**, 12-19.
- Karyotaki, E., Smit, Y., de Beurs, D. P., Henningsen, K. H., Robays, J., Huibers, M. J., Weitz, E. & Cuijpers, P.** (2016). The long-term efficacy of acute-phase psychotherapy for depression: A meta-analysis of randomized trials. *Depression and Anxiety* **33**, 370-383.
- Kernberg, O. F.** (1988). Object relations theory in clinical practice. *Psychoanalytic Quarterly* **57**, 481-504.
- Kroenke, K., Spitzer, R. L. & Williams, J. B.** (2001). The PHQ-9: validity of a brief depression severity measure. *Journal of General Internal Medicine* **16**, 606-613.
- Leichsenring, F., Luyten, P., Hilsenroth, M. J., Abbass, A., Barber, J. P., Keefe, J. R., Leweke, F., Rabung, S. & Steinert, C.** (2015). Psychodynamic therapy meets evidence-based medicine: a systematic review using updated criteria. *Lancet Psychiatry* **2**, 648-660.
- Lemma, A., Target, M. & Fonagy, P.** (2010). The development of a brief psychodynamic

protocol for depression: Dynamic Interpersonal Therapy (DIT). *Psychoanalytic Psychotherapy* **24**, 329-346.

Lemma, A., Target, M. & Fonagy, P. (2011). The development of a brief psychodynamic intervention (dynamic interpersonal therapy) and its application to depression: a pilot study. *Psychiatry* **74**, 41-48.

Lemma, A., Target, M. & Fonagy, P. (2012). *Adherence rating scale for Dynamic Interpersonal Therapy*. Unpublished manuscript. University College London: London.

Lemma, A., Target, M. & Fonagy, P. (2017). *Manual for Dynamic Interpersonal Therapy (DIT), 2nd edn*. Anna Freud National Centre for Children and Families/Tavistock Clinic: London.

Lemmens, L. H. J. M., Arntz, A., Peeters, F., Hollon, S. D., Roefs, A. & Huibers, M. J. H. (2015). Clinical effectiveness of cognitive therapy v. interpersonal psychotherapy for depression: results of a randomized controlled trial. *Psychological Medicine* **45**, 2095-2110.

Lopez, A. D., Mathers, C. D., Ezzati, M., Jamison, D. T. & Murray, C. J. L. (2006). *Global burden of disease and risk factors*. Oxford University Press: New York.

Luborsky, L. & Crits-Christoph, P. (1998). *Understanding transference: The Core Conflictual Relationship Theme method*. American Psychological Association: Washington, DC.

Luyten, P. & Blatt, S. J. (2012). Psychodynamic treatment of depression. *Psychiatric Clinics of North America* **35**, 111-129.

McHugh, R. K., Whitton, S. W., Peckham, A. D., Welge, J. A. & Otto, M. W. (2013). Patient preference for psychological vs pharmacologic treatment of psychiatric disorders: a meta-analytic review. *Journal of Clinical Psychiatry* **74**, 595-602.

National Institute for Health and Care Excellence (2017). *Depression: The treatment and management of depression in adults (guidelines under consultation)*. National Institute for Health and Care Excellence: London.

National Institute for Health and Clinical Excellence (2009). *Depression: The treatment and management of depression in adults. Clinical Guideline 90* National Institute for Health and Clinical Excellence: London.

Perfect, D., Jackson, C., Pybis, J. & Hill, A. (2016). *Choice of therapies in IAPT: An overview of the availability and client profile of step 3 therapies*. British Association for Counselling & Psychotherapy: Lutterworth.

Roth, A. D. & Pilling, S. (2015). A competency framework for the practice of psychological interventions: Cognitive and behavioural therapy.

<https://www.ucl.ac.uk/pals/research/clinical-educational-and-health-psychology/research-groups/core/competence-frameworks-2>

Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T., Baker, R. & Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry* **59 Suppl 20**, 22-33.

Spielmanns, G. I., Berman, M. I. & Usitalo, A. N. (2011). Psychotherapy versus second-generation antidepressants in the treatment of depression: a meta-analysis. *Journal of Nervous and Mental Disease* **199**, 142-149.

Steinert, C., Munder, T., Rabung, S., Hoyer, J. & Leichenring, F. (2017). Psychodynamic therapy: As efficacious as other empirically supported treatments? A meta-analysis testing equivalence of outcomes. *American Journal of Psychiatry* **174**, 943-953.

The Community Mental Health Team (2016). *Psychological Therapies: Annual report on the use of IAPT services. England, 2015-16*. NHS Digital: Leeds.

The EuroQol Group (1990). EuroQol - a new facility for the measurement of health-related quality of life. *Health Policy* **16**, 199-208.

- Trajkovic, G., Starcevic, V., Latas, M., Lestarevic, M., Ille, T., Bukumiric, Z. & Marinkovic, J.** (2011). Reliability of the Hamilton Rating Scale for Depression: A meta-analysis over a period of 49 years. *Psychiatry Research* **189**, 1-9.
- Weissman, M. M., Prusoff, B. A., Thompson, W. D., Harding, P. S. & Myers, J. K.** (1978). Social adjustment by self-report in a community sample and in psychiatric outpatients. *Journal of Nervous and Mental Disease* **166**, 317-326.
- World Health Organization** (2001). *The World Health Report 2001: Mental health: new understanding, new hope*. World Health Organization: Geneva.
- Zimmerman, M., Martinez, J. H., Young, D., Chelminski, I. & Dalrymple, K.** (2013). Severity classification on the Hamilton Depression Rating Scale. *Journal of Affective Disorders* **150**, 384-388.
- Zimmerman, M., McGlinchey, J. B., Posternak, M. A., Friedman, M., Boerescu, D. & Attiullah, N.** (2008). Remission in depressed outpatients: more than just symptom resolution? *Journal of Psychiatric Research* **42**, 797-801.

Table 1. Participant demographics in each treatment group

Demographic	Treatment Group			Test values		
	LIT (n = 54)	DIT (n = 73)	CBT (n = 20)	F	df	p
Age (years)	37.2 (10.0)	38.1 (13.0)	38.8 (11.9)	0.160	2, 144	0.853
Assessment time (weeks)						
3 months	11.8 (2.1)	13.4 (3.4)	12.7 (1.9)	3.8	2, 115	0.025
6 months	22.0 (2.0)	26.1 (4.4)	26.3 (3.4)	16.8	2, 109	0.001
12 months		62.8 (9.5)				
				χ^2	df	p
n (%) Female	34 (63%)	49 (67%)	14 (70%)	0.406	2	0.816
n (%) Caucasian	31 (61%)	54 (79%)	15 (75%)	5.12	2	0.077
n (%) Married/cohabiting	15 (30%)	22 (32%)	7 (35%)	0.22	2	0.895
Income bracket				0.794	4	0.939
<£10,000-20,000	19 (50%)	13 (34%)	6 (16%)			
£20,000-50,000	24 (41%)	22 (38%)	12 (21%)			
£50,000-100,000+	8 (47%)	41 (36%)	3 (18%)			
n (%) Completed higher education	27 (55%)	44 (65%)	16 (80%)	3.88	2	0.143
n (%) With medical problems	29 (58%)	37 (67%)	14 (74%)	5.569	2	0.062
HRSD-17 severity				2.814	4	0.589
Mild	33%	44%	35%			
Moderate	48%	45%	55%			
Severe	19%	11%	10%			
BDI-II severity				5.205	4	0.267
Mild	6%	3%	15%			
Moderate	22%	26%	15%			
Severe	72%	71%	70%			

Note. CBT, cognitive-behavioural therapy; DIT, Dynamic Interpersonal Therapy; LIT, low-intensity therapy; HRSD, Hamilton Rating Scale for Depression-17; BDI-II, Beck Depression Inventory-II. Count variables differ in the number of respondents. For continuous measures, values in brackets represent standard deviations. Significant results are in bold.

Table 2. Marginal means and percentages on the HRSD-17 for each treatment group at each assessment point

Outcome	Marginal Estimates (SE)			Contrast [95% CI]	
	LIT	DIT	CBT	DIT v. LIT	DIT v. CBT
Mean Scores					
Baseline	18.92 (0.70)	18.11 (0.57)	19.14 (1.03)	0.82[-0.94, 2.58]	1.03 [-1.26, 3.32]
3 months	14.26 (0.89)	14.17 (0.71)	16.35 (1.24)	0.10 [-2.15, 2.34]	2.18 [-0.62, 4.98]
6 months	14.84 (1.39)	9.96 (0.93)	13.22 (1.70)	4.88 [1.60, 8.16]**	3.26 [-0.56, 7.07]
12 months		10.70 (1.55)			
RCI					
CSC	9% (0.08)	51% (0.07)	20% (0.11)	-42% [-0.64, -0.21]***	-31% [-0.56, -0.06]*
Improvement	42% (0.12)	64% (0.07)	50% (0.15)	-22% [-0.51, 0.07]	-14% [-0.48, 0.19]
No Change	56% (0.12)	35% (0.07)	49% (0.15)	20% [-0.08, 0.49]	14% [-0.18, 0.46]
Deterioration ^a	6% (0.04)	0% (0)	0% (0)	N/A	N/A

Note. SE, Standard error; CI, confidence interval; LIT, low-intensity therapy; DIT, Dynamic Interpersonal Therapy; CBT, cognitive-behavioural therapy; RCI, Reliable Change Indices; CSC, clinically significant change. Outcomes at 12 months were not collected for the LIT and CBT groups.

^aChi-square difference testing could not be performed due to lack of variation in at least one of the groups estimates.

Significant results are in bold: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Table 3. Pro-rated and imputed marginal means and contrasts for the secondary measures at post-treatment

Outcome measure	Marginal Mean (SE)			Contrast [95% CI]	
	LIT	DIT	CBT	DIT v. LIT	DIT v. CBT
ECR					
Avoidance	4.07 (0.17)	4.22 (0.13)	3.80 (0.25)	-0.15 [-0.58, 0.28]	-0.42 [-0.98, 0.13]
Anxiety	4.36 (0.16)	4.47 (0.14)	4.21 (0.26)	-0.12 [-0.56, 0.33]	-0.26 [-0.90, 0.37]
EQ-5D					
Index	0.79 (0.02)	0.83 (0.01)	0.83 (0.03)	-0.03 [-0.08, 0.01]	0 [-0.06, 0.06]
Continuous	60.14 (2.44)	69.66 (2.02)	64.92 (4.05)	-9.52 [-15.81, -3.24]**	-4.75 [-13.83, 4.34]
BSI					
Somatic	0.98 (0.16)	0.66 (0.11)	0.77 (0.20)	0.31 [-0.08, 0.70]	0.10 [-0.35, 0.56]
OC	2.07 (0.20)	1.26 (0.15)	1.57 (0.27)	0.80 [0.31, 1.29]**	0.31 [-0.32, 0.93]
IS	1.68 (0.19)	1.22 (0.16)	1.42 (0.25)	0.46 [-0.03, 0.95]	0.20 [-0.40, 0.80]
Depression	1.87 (0.18)	1.03 (0.15)	1.20 (0.25)	0.85 [0.37, 1.32]***	0.18 [-0.40, 0.75]
Anxiety	1.52 (0.19)	1.00 (0.13)	0.98 (0.23)	0.52 [0.06, 0.98]*	-0.02 [-0.55, 0.51]
Hostility	0.70 (0.14)	0.61 (0.10)	0.70 (0.18)	0.09 [-0.26, 0.44]	0.09 [-0.32, 0.51]
Phobic	1.01 (0.19)	0.76 (0.13)	0.59 (0.24)	0.25 [-0.22, 0.72]	-0.17 [-0.71, 0.38]
PI	1.10 (0.16)	0.89 (0.12)	1.08 (0.20)	0.22 [-0.19, 0.62]	0.19 [-0.28, 0.66]
Psychoticism	1.17 (0.13)	0.77 (0.11)	0.81 (0.17)	0.41 [0.07, 0.75]*	0.05 [-0.36, 0.46]
GSI	1.35 (0.14)	0.90 (0.10)	1.01 (0.18)	0.46 [0.11, 0.81]**	0.11 [-0.30, 0.52]
PSDI	2.13 (0.14)	1.61 (0.10)	1.73 (0.17)	0.52 [0.17, 0.88]**	0.12 [-0.27, 0.51]
IIP-64					
Total distress	1.60 (0.10)	1.46 (0.08)	1.56 (0.14)	0.14 [-0.12, 0.40]	0.10 [-0.22, 0.43]
Domineering	-0.60 (0.08)	-0.58 (0.07)	-0.60 (0.12)	-0.01 [-0.20, 0.17]	0.00 [-0.27, 0.26]
Vindictive	-0.40 (0.07)	-0.47 (0.06)	-0.27 (0.10)	0.07 [-0.10, 0.24]	0.19 [-0.04, 0.42]
Cold	0.00 (0.09)	-0.14 (0.07)	0.03 (0.12)	-0.04 [-0.07, 0.35]	-0.08 [-0.11, 0.44]
Socially Inhibited	0.40 (0.08)	0.32 (0.07)	0.38 (0.12)	0.08 [-0.13, 0.29]	0.06 [0.23, 0.34]
Non-assertive	0.53 (0.09)	0.57 (0.07)	0.53 (0.12)	-0.04 [-0.24, 0.16]	-0.04 [-0.33, 0.24]
OA	0.23 (0.08)	0.34 (0.06)	0.27 (0.11)	-0.11 [-0.30, 0.08]	-0.07 [-0.32, 0.18]
Self-sacrificing	0.24 (0.08)	0.29 (0.07)	0.14 (0.12)	-0.04 [-0.25, 0.16]	-0.15 [-0.42, 0.13]
Intrusive	-0.40 (0.08)	-0.34 (0.07)	-0.49 (0.11)	-0.06 [-0.26, 0.13]	-0.15 [-0.41, 0.11]
SAS					
Total	2.50 (0.08)	2.24 (0.07)	2.50 (0.12)	0.26 [0.05, 0.47]*	0.26 [-0.01, 0.53]
Work	2.36 (0.11)	1.99 (0.09)	2.33 (0.17)	0.37 [0.09, 0.65]**	0.35 [-0.04, 0.73]
Social (friends)	2.72 (0.11)	2.39 (0.09)	2.50 (0.15)	0.33 [0.07, 0.60]**	0.11 [-0.23, 0.46]
Extended family	2.41 (0.10)	2.27 (0.08)	2.58 (0.14)	0.14 [-0.11, 0.39]	0.31 [-0.01, 0.63]
Household	2.45 (0.14)	2.26 (0.13)	2.40 (0.21)	0.19 [-0.19, 0.58]	0.14 [-0.37, 0.65]

Note. SE, Standard error; CI, confidence interval; LIT, low-intensity therapy; DIT, Dynamic Interpersonal Therapy; CBT, cognitive-behavioural therapy; BSI, Brief Symptom Inventory; OC, Obsessive-compulsive; IS, Interpersonal sensitivity; PI, Paranoid ideation; GSI, Global Severity Index; PSDI, Positive Symptom Distress Index; IIP-64, Inventory of Interpersonal Problems-64; OA; Overly Accommodating; SAS, Social Adjustment Scale. Significant results are in bold: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Fig. 1. CONSORT diagram.

Fig. 2. Predicted and observed mean scores on the HRSD-17 at baseline, mid-treatment, and post-treatment for (a) the DIT and LIT groups, and (b) the DIT and CBT groups.