



City Research Online

City St George's, University of London

Citation: Ballotta, L., Eberlein, E., Schmidt, T. & Zeineddine, R. (2020). Variable annuities in a Lévy-based hybrid model with surrender risk. *Quantitative Finance*, 20(5), pp. 867-886. doi: 10.1080/14697688.2019.1687929

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/23052/>

Link to published version: <https://doi.org/10.1080/14697688.2019.1687929>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

VARIABLE ANNUITIES IN A LÉVY-BASED HYBRID MODEL WITH SURRENDER RISK

LAURA BALLOTTA^(a), ERNST EBERLEIN^{(b),(c),(d)}, THORSTEN SCHMIDT^{(b),(c),(d),*},
RAGHID ZEINEDDINE^(d)

ABSTRACT. This paper proposes a market consistent valuation framework for variable annuities with guaranteed minimum accumulation benefit, death benefit and surrender benefit features. The setup is based on a hybrid model for the financial market and uses time-inhomogeneous Lévy processes as risk drivers. Further, we allow for dependence between financial and surrender risks. Our model leads to explicit analytical formulas for the quantities of interest, and practical and efficient numerical procedures for the evaluation of these formulas. We illustrate the tractability of this approach by means of a detailed sensitivity analysis of the fair value of the variable annuity and its components with respect to the model parameters. The results highlight the role played by the surrender behaviour and the importance of its appropriate modelling.

Keywords: Finance; variable annuities; hybrid models; Lévy processes; surrender risk

JEL Classification: G13, G12, G22, C63

1. INTRODUCTION

Variable Annuities (VAs) are unit-linked investment policies providing a post-retirement income, which is generated by the returns on a suitably managed financial portfolio. Various guarantees are applied with the aim of providing protection of the policyholders' saving accounts. VAs are popular insurance products in the US, Japan, the UK, and are increasingly present in the other European markets as well. According to the Life Insurance and Market Research Association (LIMRA) Secure Retirement Institute and the Insured Retirement Institute (IRI), VAs sales for 2018 in the US were more than \$100 billion - a 2% increase compared to 2017.

Common types of guarantees offered by variable annuity contracts are the so-called Guaranteed Minimum Accumulation Benefit (GMAB), the Guaranteed Minimum Death Benefit (henceforth DB) - which applies in case of early death - the Guaranteed Minimum Income Benefit and the Guaranteed Minimum Withdrawal Benefit. The former two offer protection during the accumulation period, i.e. up to the expiration of the contract, whilst the latter two provide payouts after expiration, during the so-called 'annuitization' period. For an extensive overview and classification of these products we refer to [Bacinello et al. \(2011\)](#) and references therein.

Date: August 29, 2019.

^(a) Faculty of Finance, Cass Business School, City, University of London, UK.

^(b) Freiburg Institute for Advanced Studies (FRIAS), Germany.

^(c) Department of Mathematical Stochastics, University of Freiburg, Germany.

^(d) University of Strasbourg Institute for Advanced Study (USIAS), France.

*Corresponding Author: Thorsten Schmidt; email: thorsten.schmidt@stochastik.uni-freiburg.de.

Due to the construction of these contracts, the underwriting insurance companies are exposed to financial and mortality risk, as well as surrender risk originated by the policyholder behaviour. Indeed, the option to leave the contract prior to maturity is a common additional feature of insurance contracts, which might cause significant cash outflows for the insurer, and negatively impact on the market growth of VAs (LIMRA Secure Retirement Institute).

The study of the pricing of life insurance contracts in presence of financial risk has been pioneered by [Brennan and Schwartz \(1976\)](#) and [Boyle and Schwartz \(1977\)](#); an extensive literature has developed since, from the seminal contributions of [Albizzati and Geman \(1994\)](#), [Bacinello and Ortu \(1996\)](#) and [Grosen and Jørgensen \(2002\)](#), to the more recent ones of [Bacinello et al. \(2011\)](#), [Deelstra and Rayée \(2013\)](#), [Giacinto et al. \(2014\)](#) and [Gudkov et al. \(2018\)](#), to mention a few. These contributions distinguish themselves in terms of the specific product under consideration, although they are all based on diffusion-driven market models. Extensions to financial dynamics driven by Lévy processes are considered in [Ballotta \(2005, 2009\)](#), [Bacinello et al. \(2016\)](#), [Kélani and Quittard-Pinon \(2017\)](#) and [Alonso-Garcia et al. \(2018\)](#) to mention a few. In these papers, though, the focus is primarily on equity markets as the rate of interest is assumed constant and the possibility of surrender is not considered.

In light of the above, the aim of this paper is to provide a realistic framework for the modelling of these risks and the market consistent valuation of variable annuity contracts. We focus specifically on the case of the GMAB, the DB and the quantification of the surrender risk by means of the so-called Surrender Benefit (SB), contributing to the current state of the literature in a number of ways. Firstly, contrary to the literature mentioned above, we propose a general joint model for financial and insurance risks, which is driven by time-inhomogeneous Lévy processes. Our choice is motivated by the increased distributional flexibility offered by these stochastic processes in portraying observed market trends (see, for example, [Eberlein and Keller, 1995](#), for an extensive empirical analysis of equity markets).

In more details, for the financial market we adopt the hybrid construction of [Eberlein and Rudmann \(2018\)](#) in which the stochastic dependence between interest rate markets and stock markets is taken into account explicitly. This feature is an important aspect in this context given the typically long maturity of insurance contracts.

Further, we subdivide insurance risk into surrender risk and mortality risk. For the surrender risk, we follow market practice by considering two components: one capturing the baseline surrender behaviour due to non-economic factors and personal contingencies, and the second one which instead is responsive to changes in the conditions of the financial markets (see [Kolkiewicz and Tan, 2006](#), [Le Courtois and Nakagawa, 2011](#), [Ducuroir et al., 2016](#), for example). This component in particular includes a function of the spread between the rate of the contract and the rate offered on the market for equivalent products, and therefore incorporates the stochastic inputs from the financial market model. The function of choice is designed as to suitably accommodate for surrender triggering arguments based on the money-ness hypothesis (see [Knoller et al., 2016](#), for example), the interest rate hypothesis and the emergency fund hypothesis (see, for example, [Nolte and Schneider, 2017](#), and references therein).

For mortality risk, we adopt an extended Gompertz-Makeham model with a stochastic mortality improvement ratio; furthermore, we make the natural assumption of stochastic independence between the demographic and the financial risks.

Secondly, through the proposed general framework we obtain expressions for the value of the guarantees and the SB, which are analytical up to a multidimensional integral. The dimensionality of these integrals is dictated by the frequency with which the policyholder is allowed to terminate the contract. These formulas are quite general and do not depend on any particular distribution of the relevant random quantities.

Thirdly, we also develop a practical and efficient numerical scheme for the evaluation of these formulas by means of Monte Carlo integration with importance sampling, and we illustrate its applicability by performing a sensitivity analysis of the contract's value with respect to the model parameters. The results from this analysis underline the importance of a correct calibration of the model to the observed surrender rates.

We conclude by noting that the availability of both analytical expressions and a practical numerical scheme for the evaluations of the relevant multidimensional integrals - as opposed to pricing by Monte Carlo simulations of the market dynamics, stochastic mortality and the policyholder behaviour - could provide a significant support to the development of suitable risk management strategies based on, for example, the Greeks of the contract, or the local risk-minimization principle as discussed in [Kélani and Quittard-Pinon \(2017\)](#).

The paper is organized as follows. In section 2 we introduce the model for the financial market; the contract features of the VAs are introduced in section 3 together with the additional modelling assumptions regarding mortality and surrender risk. The closed analytical formulas are derived in section 4; the corresponding numerical scheme is offered in section 5. We present the results of the sensitivity analysis in section 6, whilst section 7 concludes.

Some proofs are offered in the appendix, and additional material is provided in an online companion.

2. THE INTEREST RATE AND EQUITY MARKET

The aim of this section is to introduce a joint model for the interest rate and the equity markets used to develop the valuation framework for variable annuities. Specifically, we follow [Eberlein and Rudmann \(2018\)](#) and build this hybrid model on time-inhomogeneous Lévy processes. Thus we first offer some necessary preliminary results with the pricing application in view.

2.1. Time-inhomogeneous Lévy process and pricing. Given a finite time horizon $T^* > 0$, consider a stochastic basis $(\Omega, \mathcal{F}, \mathbb{F}, Q)$ with a filtration $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T^*]}$ satisfying the usual conditions. Due to our focus on pricing, the probability measure Q represents a risk-neutral martingale measure. Let L^1 and L^2 be two independent time-inhomogeneous Lévy processes, i.e. continuous-time processes with independent increments, and with characteristic function

$$E_Q \left[e^{iuL_t^j} \right] = \exp \left(\int_0^t \left(iub_s^j - \frac{1}{2}c_s^j u^2 + \int_{\mathbb{R}} (e^{iux} - 1 - iux \mathbb{1}_{\{|x| \leq 1\}}) F_s^j(dx) \right) ds \right) \quad (1)$$

for $j = 1, 2$; the local characteristic $(b_s^j, c_s^j, F_s^j)_{s \in [0, T^*]}$ satisfies the integrability condition

$$\int_0^{T^*} \left(|b_s^j| + c_s^j + \int_{\mathbb{R}} (\min\{|x|^2, 1\}) F_s^j(dx) \right) ds < \infty.$$

We note that a financial market built on (exponential) Lévy processes is in general incomplete, and consequently the risk-neutral martingale measure is not unique. Standard practice in this case is to fix the pricing measure via calibration, using market quotes for traded derivatives contracts written on the quantities of interest, i.e. bonds and stocks in the specific case of our application. As model calibration is not in the scope of this paper, we refer to [Eberlein and Rudmann \(2018\)](#) for a detailed illustration of the calibration procedure for the adopted hybrid model. Nevertheless, for any risk-neutral martingale measure to be well defined, we require that the exponential moments of a certain order exist. To this purpose, the following assumption holds throughout the rest of this paper.

Assumption 2.1 (Exponential moments). For $j = 1, 2$, there exist positive constants M_j and ϵ_j such that for each $u \in [-(1 + \epsilon_j)M_j, (1 + \epsilon_j)M_j]$

$$\int_0^{T^*} \int_{\{|x|>1\}} e^{ux} F_s^j(dx) ds < \infty.$$

All standard processes typically used in mathematical finance such as hyperbolic, Normal Inverse Gaussian, Variance Gamma and CGMY processes satisfy the above condition. Assumption 2.1 implies the existence of the first moment of the process; this allows us to rewrite (1) as

$$E_Q \left[e^{iuL_t^j} \right] = \exp \left(\int_0^t \left(iub_s^j - \frac{1}{2} c_s^j u^2 + \int_{\mathbb{R}} (e^{iux} - 1 - iux) F_s^j(dx) \right) ds \right).$$

Note that the term b^j is now different from the one in equation (1). In this setting, we define the cumulant function of L^j

$$\theta_s^j(z) = b_s^j z + \frac{1}{2} c_s^j z^2 + \int_{\mathbb{R}} (e^{zx} - 1 - zx) F_s^j(dx),$$

for any $z \in \mathbb{C}$ such that $\text{Re}(z) \in [-(1 + \epsilon_j)M_j, (1 + \epsilon_j)M_j]$. Then, $E_Q[\exp(zL_t^j)] < \infty$ and

$$E_Q \left[\exp(zL_t^j) \right] = \exp \left(\int_0^t \theta_s^j(z) ds \right).$$

Further, let $f : \mathbb{R}^+ \rightarrow \mathbb{C}$ be a continuous function with $|\text{Re}(f)| \leq M_j$, then

$$E_Q \left[\exp \left(\int_0^t f(s) dL_s^j \right) \right] = \exp \left(\int_0^t \theta_s^j(f(s)) ds \right), \quad (2)$$

where the integrals are defined component-wise for the real and imaginary part (see [Eberlein and Raible, 1999](#), for the derivation of this formula).

2.2. The fixed income market. For the modelling of the fixed income market we follow the approach in [Eberlein et al. \(2005\)](#) (see also [Eberlein and Raible, 1999](#)), so that the starting point is the definition of the dynamics of the instantaneous forward rates $(f(t, T)_{0 \leq t \leq T \leq T^*})$.

Let us assume that

$$f(t, T) = f(0, T) + \int_0^t \alpha(s, T) ds - \int_0^t \sigma_1(s, T) dL_s^1, \quad 0 \leq t \leq T \leq T^*, \quad (3)$$

for a deterministic and bounded function $f(0, T)$. The drift function $\alpha(\cdot)$ and the volatility function $\sigma_1(\cdot)$ are assumed to satisfy the usual conditions of measurability and boundedness (see Eberlein et al., 2005, (2.5)). The price of a zero coupon bond at time t with maturity $T \geq t$ is

$$B(t, T) = \exp\left(-\int_t^T f(t, s) ds\right).$$

Let us denote

$$A(s, T) := \int_s^T \alpha(s, u) du, \quad \Sigma(s, T) := \int_s^T \sigma_1(s, u) du.$$

From Fubini's theorem and equation (3) it follows that the dynamics of the bond price is

$$B(t, T) = B(0, T) \exp\left(\int_0^t (f(s, s) - A(s, T)) ds + \int_0^t \Sigma(s, T) dL_s^1\right),$$

We remind that the short rate $r(t)$ is implicitly given by the forward rate dynamics in equation (3) by setting $T = t$. Finally, we assume that

$$\Sigma(s, T) \leq \frac{M_1}{3}, \quad (4)$$

where M_1 is the constant from Assumption 2.1; this guarantees that the exponential of the stochastic integral has finite expectation.

For the market to be arbitrage free, we require that $(B(t)^{-1}B(t, T))_{0 \leq t \leq T}$, with $B(t) = \exp\left(\int_0^t r(s) ds\right)$, is a martingale; it follows from (2) that Q is a martingale measure if

$$A(s, T) = \theta_s^1(\Sigma(s, T)), \quad 0 \leq s \leq T. \quad (5)$$

In the following we will always assume that the drift condition (5) holds.

2.3. The stock market. For the modelling of the equity market, we consider the case of a single asset, be it a single stock or a stock index. More general settings can be obtained in a straightforward manner.

It is well known from a number of empirical studies that equity and fixed income markets influence each other; this interaction is of particular importance in the context of long-dated insurance contracts. Thus, following Eberlein and Rudmann (2018) we choose an approach which allows for stochastic dependence between the two markets. Consequently, we model the price process of the asset as

$$S_t = S_0 \exp\left(\int_0^t r(s) ds + \int_0^t \sigma_2(s) dL_s^2 + \int_0^t \beta(s) dL_s^1 - \omega(t)\right). \quad (6)$$

In this hybrid approach, the driver of the interest rate dynamics affects explicitly the stock price in reason of $\beta(\cdot)$ through the term $\int_0^t \beta(s) dL_s^1$. Further dependence is originated endogenously by the (integrated) short rate, i.e. the classical risk-neutral term $\int_0^t r(s) ds$. $\sigma_2(\cdot)$ is a positive function and denotes the volatility of the stock price. Both, $\sigma_2(\cdot)$ and $\beta(\cdot)$, could be chosen as random processes, but having numerical aspects in mind, in the following we consider deterministic functions $\sigma_2(\cdot)$ and $\beta(\cdot)$. To ensure the existence of exponential

moments we require

$$\sigma_2(s) \leq \frac{M_2}{2}, \quad |\beta(s)| \leq \frac{M_1}{3}, \quad (7)$$

with M_1, M_2 the constants from Assumption 2.1. The drift term $\omega(t)$ in (6) is chosen such that the discounted stock price $(B(t)^{-1}S_t)_{t \in [0, T^*]}$ is a Q -martingale. Equation (2) and the independence of the two driving processes L^1 and L^2 imply that

$$\omega(t) = \int_0^t [\theta_s^2(\sigma_2(s)) + \theta_s^1(\beta(s))] ds. \quad (8)$$

Hence, under (5) and (8) the joint market model for bonds and security S is free of arbitrage.

3. THE VARIABLE ANNUITY CONTRACT

A variable annuity (VA) is an insurance contract which gives the holder a variety of benefits depending on the notional I in exchange for a single premium paid at inception of the VA contract. We note that in the following we focus on the risk-neutral valuation of the benefits offered by the contract and therefore ignore any fees aimed at covering the cost of the guarantees and other management expenses. The proposed model, though, can be adapted to cater for these fees, for example by means of a fee rate which is continuously deducted from the underlying fund value. The fair rate would then be obtained as to ensure equality between the total value of the VA contract and the initial premium. We refer the interested reader to, for example, [Bacinello et al. \(2011\)](#) and [Kélani and Quittard-Pinon \(2017\)](#) and references therein.

The maturity T of the contract is assumed to satisfy $0 < T \leq T^*$. In the specification considered here, the VA includes three features: a Guaranteed Minimum Accumulation Benefit (GMAB), a Surrender Benefit (SB), and a Death Benefit (DB).

In details, at maturity T the GMAB pays to the policyholder

$$\max(IS_T, G(T)),$$

with $G(T) = I \exp(\delta T)$, $\delta > 0$. In other words, the GMAB offers the best of the investment of an amount I in either the asset S or in a risk-free account with guaranteed rate δ . Note that, in order to simplify the notation, in the following we assume that the price process of the asset is normalized so that $S_0 = 1$. This payoff, however, can only be claimed if the policyholder is still alive at time T and did not exercise the surrender option before.

In case of early surrender, the right of refund is restricted to the current value of the fund reduced by a compulsory surrender penalty. To this purpose, let $P : [0, T] \rightarrow (0, 1]$ be an increasing function with $P(T) = 1$, and define $\mathbf{t} := (t_0, t_1, \dots, t_K)^\top$ with $0 = t_0 < t_1 < \dots < t_K < T$. We assume that premature surrender is possible at any time point $t_i \in \mathbf{t}$ with $i = 1, \dots, K - 1$, in which case the policyholder would receive the amount

$$IS_{t_i} P(t_i),$$

Thus, $P(t)$ represents the proportion of the benefit that the policyholder is entitled to in case of surrender; consequently $1 - P(t)$ amounts to the actual surrender penalty. Small values of P at early dates would allow the insurer to recover any expenses related to the writing of the contract. In the case in which the policyholder leaves the scheme while the equity

market experiences a significant rise, the function $P(t)$ could also be considered as a control (in fact a partial hedge) on the loss incurred by the insurer.

Finally, in case of death before maturity T , the death benefit pays (to the beneficiaries)

$$\max(IS_{\bar{t}_i}, G(\bar{t}_i)),$$

for $\bar{t}_i \in \bar{\mathbf{t}}$, $i = 1, \dots, N$, and $\bar{\mathbf{t}} := (\bar{t}_0, \bar{t}_1, \dots, \bar{t}_N)^\top$ with $0 = \bar{t}_0 < \bar{t}_1 < \dots < \bar{t}_N = T$. The time points \bar{t}_i denote the frequency with which mortality is monitored by the insurer over the lifetime of the contract.

In principle, the time scales \mathbf{t} and $\bar{\mathbf{t}}$ could be arbitrary, but in typical cases they are not: we assume that $\mathbf{t} \subset \bar{\mathbf{t}}$ in the sense that any surrender time t_i is contained in $\{\bar{t}_1, \dots, \bar{t}_N\}$. This is a very natural assumption since death of the policyholder might be monitored by the insurer at the end of every month or every quarter, whereas surrender of the contract might be allowed only at the end of each year during the life of the contract.

The description of the contract benefits given above highlights the need for an accurate financial model, but also an appropriate modelling for mortality risk and surrender risk. This is offered in the rest of this section, which we conclude with the discussion of the market consistent valuation of the VA.

3.1. Mortality model. For the modelling of mortality risk, we follow standard literature in the field and adopt a stochastic intensity-based approach. Pioneered by [Milevsky and Promislow \(2001\)](#), further developed by [Dahl \(2004\)](#), [Dahl and Møller \(2006\)](#), and more recently generalized by [Li and Szimayer \(2011\)](#), this framework builds on a given initial curve for mortality rates by superimposing a stochastic process capturing random improvements and fluctuations.

In details. Let $\tau^m(x)$ be a random time capturing the remaining lifetime of a x years old individual. The corresponding survival probability with respect to the given probability measure is

$$Q(\tau^m(x) > t) = \mathbb{E}_Q \left(e^{-\int_0^t \lambda_u^m(x+u) du} \right),$$

where $\lambda_t^m(x+t)$, $t > 0$, is the stochastic mortality intensity for an individual aged $x+t$ at time t . This intensity process is modelled as

$$\lambda_t^m(x+t) = \lambda^{m,0}(x+t)\xi_t, \tag{9}$$

for an initial curve of the mortality intensity, $\lambda^{m,0}$, and a strictly positive process ξ_t such that $\xi_0 = 1$, capturing the mortality improvements from time 0 to time t for a person aged $x+t$. Finally, the mortality intensity satisfies the property that $\lambda_0^m(x) = \lambda^{m,0}(x)$ (see [Dahl and Møller, 2006](#) and references therein for full details).

A popular choice for the initial mortality curve is represented by the Gompertz-Makeham model ([Dahl, 2004](#), [Dahl and Møller, 2006](#), for example). An alternative specification corresponding to the structure for the UK standard tables for annuitant and pension population is adopted in [Ballotta and Haberman \(2006\)](#). Numerous choices for the process of the mortality improvement ratio, ξ_t , have been put forward as well: [Dahl \(2004\)](#), [Biffis \(2005\)](#), [Dahl and Møller \(2006\)](#) for example focus on affine diffusion specifications with particular emphasis on time-inhomogeneous CIR processes; [Ballotta and Haberman \(2006\)](#) instead extend the

generalized linear model by superimposing a standard mean reverting Ornstein-Uhlenbeck process and suitably accommodating longevity effects.

In light of the above, in the following we choose the standard Gompertz-Makeham model for the initial curve, so that

$$\lambda^{m,0}(x+t) = \frac{1}{b} e^{\left(\frac{x+t-z}{b}\right)},$$

and a generalized Ornstein-Uhlenbeck process with mean reversion level $e^{-\lambda t}$ for the mortality improvement ratio, of the form

$$d\xi_t = \kappa(\exp(-\lambda t) - \xi_t)dt + dL_t^3.$$

We assume that L^3 is a suitably chosen Lévy process independent of (L^1, L^2) , z, κ are non-negative, b is positive and $\lambda \in \mathbb{R}$. This implies that the mortality intensity is independent of the financial market.

The moment generating function of the integrated intensity process, and thus the survival probability, can be recovered by standard argument based on the theory of affine processes (for fuller details, we refer the interested reader to [Duffie et al., 2003](#), or [Eberlein and Kallsen, 2019](#), Chapter 6). The above formulation is the generalization of [Krazzler et al. \(2016\)](#) to the case of a driver modelled via Lévy processes.

Let us denote by $\mathbf{F}^{L^1, L^2} = (\mathcal{F}_t^{L^1, L^2})_{0 \leq t \leq T^*}$ the filtration generated by L^1 and L^2 . Then, for any set $A \in \mathcal{F}_t^{L^1, L^2}$, it follows that

$$E_Q \left[\mathbb{1}_{\{\tau^m(x) > t\}} \mathbb{1}_A \right] = Q(A) E_Q \left[e^{-\int_0^t \lambda_u^m(x+u) du} \right]. \quad (10)$$

The above assumption of independence between the mortality risk driver, L^3 , and the financial risk drivers, L^1 and L^2 , is plausible from the perspective of the insurer: precise modelling would be highly client-specific, and would require information rarely accessible by insurance companies. Moreover, the independence assumption implies a high degree of tractability which is important for complex products such as VAs considered here. Finally, as a consequence of the independence between demographic and financial risks, the computation of the survival probability is carried out under the risk-neutral measure Q .

3.2. Surrender model. Surrender risk is notoriously difficult to assess and model due to the nature of the decisions leading to it. Policyholders can indeed surrender both because of rising alternative financial opportunities, and apparently non rational (in the financial sense) behaviours due to personal considerations and contingencies. Nevertheless, surrender represents one of the main risks faced by life insurance companies due to the liquidity issues it can generate, with potential loss of market share (see, for example, [Loisel and Milhaud, 2011](#), and references therein).

A common market and academic practice to surrender modelling (see [Kolkiewicz and Tan, 2006](#), [Le Courtois and Nakagawa, 2011](#), [Ducuroir et al., 2016](#), for example) is to consider

two components: a deterministic baseline hazard rate function capturing lapses¹ due to non-economic factors, and a stochastic process representing additional shocks to the baseline due to changes in the market.

This random component is usually linked to the spread between the return offered by the contract and the one offered on the market for equivalent products. Indeed, dependence of the surrender on interest rates is relatively intuitive: higher interest rates are strong incentives for the policyholder to switch to higher yield investments, whilst very low interest rates - such as the ones currently observed in all major economies - could represent advantageous opportunities for refinancing. The random component should also be linked to the change in value of the underlying asset, as it directly impacts on the amount received in case of surrender.

Thus, following this line of reasoning, let τ^s denote the random time of the policyholder decision to surrender. As specified above, surrender is allowed at time points t_i , $i = 1, \dots, K-1$; by convention $\{\tau^s = \infty\}$ corresponds to no surrender. Let λ^s denote the corresponding intensity of surrender; then $\lambda^s(t) = 0$ for $t \in [0, t_1) \cup [t_K, T]$. Further, we model the baseline surrender behaviour due to non-economic factors and personal contingencies by a non-negative constant C .

Let $D(t)$ be the process driving the dynamic lapse component; consistently with the considerations offered above, we build this process on the spread between the return offered by the surrender benefit plus the market rates at which this amount can be re-invested, and the total yield of the policy represented by the value at maturity of the guaranteed amount. Thus, let $Y_t = \log S_t$, and $p(t) = -\log P(t)$. Then

$$D(t) = Y_t - p(t) + \int_t^T f(t, s) ds - \delta T, \quad 0 \leq t \leq T. \quad (11)$$

The overall surrender intensity consequently is defined as

$$\lambda^s(t) = \beta D^2(t_i) + C, \quad t_i \leq t < t_{i+1}, \quad (12)$$

so that it is piecewise constant on the interval $[t_i, t_{i+1})$ for $i = 1, \dots, K-1$.

The non-negative constant β captures the dependence between the surrender intensity and the market and is a measure of the investors' rationality (in the pure economic sense), and their response to personal financial motivations. Equation (12) uses the square of the spread function $D(t)$ in order to capture both situations of favourable market conditions offering more remunerative investment opportunities, and financial market turmoils in which policyholders might lack sufficient resources to finance their expenses (emergency fund hypothesis).

Although in spirit similar to others in the literature (see for example [Le Courtois and Nakagawa, 2011](#) and [Escobar et al., 2016](#)), our construction distinguishes itself also for the non-Gaussian dynamics of the underpinning risk drivers L^1 and L^2 .

The resulting probability of no surrender is given by

$$Q(\tau^s \geq t_i | \mathcal{F}_{t_i}^{L^1, L^2}) = \exp\left(-\int_0^{t_i} \lambda^s(u) du\right), \quad (13)$$

¹'Lapse' was originally used to denote termination of an insurance policy and loss of coverage because the policyholder had failed to pay premia, whilst 'surrender' denotes termination accompanied by the payout of a surrender benefit. Nowadays 'lapse' often denotes both situations.

for all $1 \leq i \leq K - 1$, and

$$\begin{aligned} Q(\tau^s \geq t | \mathcal{F}_t^{L^1, L^2}) &= Q(\tau^s = \infty | \mathcal{F}_t^{L^1, L^2}) \\ &= \exp\left(-\int_0^{t_K} \lambda^s(u) du\right), \end{aligned} \quad (14)$$

for $t_{K-1} < t$. We note that the last integral equals $\exp(-\int_0^T \lambda^s(u) du)$. The set-up chosen here can be obtained in a doubly-stochastic model or in a model where immersion holds, see [Aksamit and Jeanblanc \(2017\)](#) for a comprehensive treatment in this regard. An alternative form of the intensity is investigated in the online Appendix [D](#).

Finally, we observe that the intensity functions λ^s and λ^m are independent due to the assumed independence between demographic and financial risks. This is consistent with the literature in the field for which we refer in particular to the studies of [Milhaud et al. \(2011\)](#), [Knoller et al. \(2016\)](#) and [Nolte and Schneider \(2017\)](#) and references therein.

3.3. Valuation of the variable annuity. Using the notation introduced above, we can now formulate the actual cash-flows associated with the considered variable annuity. Firstly, recall that the GMAB provides a payoff only at the maturity time T if the policyholder is alive (i.e., $\{\tau^m(x) > T\}$) and if there was no surrender until this time (i.e., $\{\tau^s > T\}$). Consequently, the associated cash-flow at maturity T is

$$\text{GMAB}(T) = \mathbb{1}_{\{\tau^m(x) > T\}} \mathbb{1}_{\{\tau^s > T\}} \max(IS_T, G(T)). \quad (15)$$

Secondly, the surrender option can be exercised only once if the policyholder is still alive (i.e., $\{\tau^s < \tau^m(x)\}$). Therefore, should surrender occur, the surrender benefit at time t_i pays

$$\text{SB}(t_i) = \mathbb{1}_{\{\tau^s = t_i\}} \mathbb{1}_{\{\tau^s < \tau^m(x)\}} IS_{t_i} P(t_i), \quad (16)$$

where t_i is one of the possible premature surrender dates. Finally, the death benefit provides a payoff only in case of no early surrender, and is quantified as

$$\text{DB}(\bar{t}_i) = \mathbb{1}_{\{\bar{t}_{i-1} \leq \tau^m(x) < \bar{t}_i\}} \mathbb{1}_{\{\tau^m(x) < \tau^s\}} \max(IS_{\bar{t}_i}, G(\bar{t}_i)), \quad (17)$$

where \bar{t}_i is one of the possible payoff dates of the death benefit.

The value P^{VA} of the variable annuity at time $t = 0$ is equal to the sum of the values of its constituents, i.e.

$$P^{\text{VA}} = P^{\text{GMAB}} + P^{\text{SB}} + P^{\text{DB}}, \quad (18)$$

with, by standard risk-neutral valuation argument,

$$\begin{aligned} P^{\text{GMAB}} &= E_Q \left[e^{-\int_0^T r(u) du} \text{GMAB}(T) \right], \\ P^{\text{SB}} &= \sum_{i=1}^{K-1} E_Q \left[e^{-\int_0^{t_i} r(u) du} \text{SB}(t_i) \right], \\ P^{\text{DB}} &= \sum_{i=1}^N E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \text{DB}(\bar{t}_i) \right]. \end{aligned}$$

Tractable formulae for these expressions are provided in the following section.

4. MARKET-CONSISTENT VALUATION

In this section we derive analytical expressions for the components of the variable annuity contract discussed above in the model setup provided in section 2 in presence of mortality and surrender risk. Useful results and representations needed in the following are provided in Appendix A.

4.1. Guaranteed minimum accumulation benefit. The independence between $\tau^m(x)$ and the financial market and result (14) imply that

$$\begin{aligned} P^{\text{GMAB}} &= E_Q \left[e^{-\int_0^T r(u)du} \mathbb{1}_{\{\tau^m(x) > T\}} \mathbb{1}_{\{\tau^s > T\}} \max(IS_T, G(T)) \right] \\ &= Q(\tau^m(x) > T) E_Q \left[e^{-\int_0^T r(u)du} \max(IS_T, G(T)) E_Q \left(\mathbb{1}_{\{\tau^s > T\}} | \mathcal{F}_T^{L^1, L^2} \right) \right] \end{aligned}$$

and therefore

$$P^{\text{GMAB}} = Q(\tau^m(x) > T) E_Q \left[e^{-\int_0^T r(u)du} e^{-\int_0^T \lambda^s(u)du} \max(IS_T, G(T)) \right].$$

We introduce the T -forward measure Q^T defined as

$$\frac{dQ^T}{dQ} = \frac{1}{B(0, T)B(T)}. \quad (19)$$

Denoting the expectation with respect to Q^T by E^T , we obtain

$$P^{\text{GMAB}} = Q(\tau^m(x) > T) B(0, T) E^T \left[e^{-\int_0^T \lambda^s(u)du} \max(IS_T, G(T)) \right].$$

Observe that

$$\max(IS_T, G(T)) = G(T) \left[1 + \left(\frac{IS_T}{G(T)} - 1 \right)^+ \right], \quad (20)$$

consequently

$$\begin{aligned} \frac{P^{\text{GMAB}}}{Q(\tau^m(x) > T) B(0, T) G(T)} &= E^T \left[e^{-\int_0^T \lambda^s(u)du} \right] + E^T \left[e^{-\int_0^T \lambda^s(u)du} \left(\frac{IS_T}{G(T)} - 1 \right)^+ \right] \\ &=: A_1 + A_2, \end{aligned} \quad (21)$$

with obvious definitions in the last line. The term A_1 can be interpreted as a weight factor due to the policyholder not leaving the scheme prior to maturity; the term A_2 , instead, captures the cost of the option embedded in the GMAB, conditional on no early surrender.

Let

$$w_l := \int_0^{t_l} A(s, T) ds + \int_0^T f(0, s) ds - \delta T - \omega(t_l) - p(t_l) \quad (22)$$

for $l = 1, \dots, K - 1$ and

$$w_K := \int_0^T A(s, T) ds + \int_0^T f(0, s) ds - \delta T - \omega(T). \quad (23)$$

Further, define $\Delta t_l := t_l - t_{l-1}$, as well as $R := (0, \dots, 0, r) \in \mathbb{R}^K$, with $1 < r < 2$, and let for all $0 \leq s \leq T$, and $u \in \mathbb{R}^{K-1}$, $v \in \mathbb{C}^K$

$$\begin{aligned} D(u, T) &:= \exp\left(i \sum_{l=1}^{K-1} u_l w_l\right), \\ \tilde{D}(v, T) &:= D(v_1, \dots, v_{K-1}, T) \exp\left(i v_K w_K\right), \\ E(s, u, T) &:= \Sigma(s, T) + i(\beta(s) - \Sigma(s, T)) \sum_{l=1}^{K-1} u_l \mathbb{1}_{\{0 \leq s \leq t_l\}}, \\ \tilde{E}(s, v, T) &:= E(s, v_1, \dots, v_{K-1}, T) + i(\beta(s) - \Sigma(s, T)) v_K, \end{aligned} \quad (24)$$

$$\begin{aligned} F(s, u) &:= i\sigma_2(s) \sum_{l=1}^{K-1} u_l \mathbb{1}_{\{0 \leq s \leq t_l\}}, \\ \tilde{F}(s, v) &:= F(s, v_1, \dots, v_{K-1}) + i\sigma_2(s) v_K, \end{aligned}$$

$$M(u, T) := D(u, T) e^{\int_0^T \theta_s^1(E(s, u, T)) ds + \int_0^T \theta_s^2(F(s, u)) ds} \prod_{l=2}^K \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-u_{l-1}^2 / (4\beta \Delta t_l)},$$

$$\begin{aligned} N(v, T) &:= \tilde{D}(v - iR, T) e^{\int_0^T \theta_s^1(\tilde{E}(s, v - iR, T)) ds + \int_0^T \theta_s^2(\tilde{F}(s, v - iR)) ds} \\ &\quad \times \frac{1}{(i v_K + r - 1)(i v_K + r)} \prod_{l=2}^K \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-v_{l-1}^2 / (4\beta \Delta t_l)}, \text{ for } v \in \mathbb{R}^K. \end{aligned}$$

Theorem 4.1. *The value P^{GMAB} is given by*

$$P^{\text{GMAB}} = Q(\tau^m(x) > T) B(0, T) G(T) (A_1 + A_2)$$

with

$$\begin{aligned} A_1 &= \frac{e^{-C(t_K - t_1)}}{(2\pi)^{K-1}} e^{-\int_0^T A(s, T) ds} \int_{\mathbb{R}^{K-1}} M(u, T) du, \\ A_2 &= \frac{e^{-C(t_K - t_1)}}{(2\pi)^K} e^{-\int_0^T A(s, T) ds} \int_{\mathbb{R}^K} N(u, T) du, \end{aligned}$$

with $A(s, T)$ as in (5).

Proof. By the definition of λ^s in (12), we have

$$\begin{aligned} A_1 &= e^{-C(t_K - t_1)} E^T \left[\prod_{i=2}^K e^{-\beta \Delta t_i D(t_{i-1})^2} \right] \\ &= e^{-C(t_K - t_1)} E^T \left[f(D(t_1), \dots, D(t_{K-1})) \right], \end{aligned}$$

where $f(x_1, \dots, x_{K-1}) := \prod_{l=2}^K (e^{-\beta \Delta t_l x_{l-1}^2}) = \prod_{l=2}^K f_l(x_{l-1})$. For a generic function f we denote by \hat{f} its Fourier transform. Then, for any $y \in \mathbb{C}$,

$$\hat{f}_l(y) = \int_{\mathbb{R}} e^{iyt} e^{-\beta \Delta t_l t^2} dt = \sqrt{\frac{\pi}{\beta \Delta t_l}} \exp\left(-y^2/(4\beta \Delta t_l)\right). \quad (25)$$

This implies that

$$\hat{f}(y_1, \dots, y_{K-1}) = \prod_{l=2}^K \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-y_{l-1}^2/(4\beta \Delta t_l)}, \quad (26)$$

and we observe that $\hat{f} \in L^1(\mathbb{R}^{K-1})$. By Theorem 3.2 in [Eberlein et al. \(2010\)](#)

$$E^T \left[f(D(t_1), \dots, D(t_{K-1})) \right] = \frac{1}{(2\pi)^{K-1}} \int_{\mathbb{R}^{K-1}} \tilde{M}(iu) \hat{f}(-u) du, \quad (27)$$

where for any $u = (u_1, \dots, u_{K-1})$, $\tilde{M}(iu)$ is defined as follows

$$\tilde{M}(iu) := E^T \left[e^{iu_1 D(t_1) + \dots + iu_{K-1} D(t_{K-1})} \right].$$

Using representation (A.1) of D given in the appendix together with equations (3) and (11), we obtain

$$\begin{aligned} \tilde{M}(iu) &= \exp \left[i \sum_{l=1}^{K-1} u_l \left(-p(t_l) - \delta T + \int_0^{t_l} f(0, s) ds + \int_0^{t_l} A(s, T) ds - \omega(t_l) \right) \right] \\ &\quad \times E^T \left[\exp \left(i \sum_{l=1}^{K-1} u_l \left(\int_0^{t_l} \sigma_2(s) dL_s^2 + \int_0^{t_l} (\beta(s) - \Sigma(s, T)) dL_s^1 \right) \right) \right]. \end{aligned}$$

Moreover, in virtue of representation (A.3), the density of Q^T given in (19) can be written as

$$\frac{dQ^T}{dQ} = \exp \left(- \int_0^T A(s, T) ds + \int_0^T \Sigma(s, T) dL_s^1 \right).$$

Consequently

$$\begin{aligned} &E^T \left[\exp \left(i \sum_{l=1}^{K-1} \left(\int_0^{t_l} u_l \sigma_2(s) dL_s^2 + \int_0^{t_l} u_l (\beta(s) - \Sigma(s, T)) dL_s^1 \right) \right) \right] \\ &= E_Q \left[\exp \left(\int_0^T \Sigma(s, T) dL_s^1 - \int_0^T A(s, T) ds \right) \right. \\ &\quad \left. \times \exp \left(i \sum_{l=1}^{K-1} \left(\int_0^{t_l} u_l \sigma_2(s) dL_s^2 + \int_0^{t_l} u_l (\beta(s) - \Sigma(s, T)) dL_s^1 \right) \right) \right] \\ &= e^{-\int_0^T A(s, T) ds} E_Q \left[\exp \left(\int_0^T E(s, u, T) dL_s^1 + \int_0^T F(s, u) dL_s^2 \right) \right] \\ &= \exp \left(- \int_0^T A(s, T) ds + \int_0^T \theta_s^1(E(s, u, T)) ds + \int_0^T \theta_s^2(F(s, u)) ds \right), \end{aligned}$$

with E and F as in (24). The last equality follows from the independence of L^1 and L^2 and equation (2). Therefore, with $D(u, T)$ defined in (24) we have

$$\tilde{M}(iu) = D(u, T) \exp \left(- \int_0^T A(s, T) ds + \int_0^T \theta_s^1(E(s, u, T)) ds + \int_0^T \theta_s^2(F(s, u)) ds \right)$$

and the representation for A_1 follows from (27).

We continue by calculating

$$A_2 = E^T \left[e^{-\int_0^{t_K} \lambda^s(u) du} \left(\frac{IS_T}{G(T)} - 1 \right)^+ \right].$$

With the definitions $S_T = \exp(Y_T)$, $G(T) = I \exp(\delta T)$, and (A.1), we obtain that

$$\frac{IS_T}{G(T)} = \exp(D(T)).$$

Therefore, by the same arguments used in the calculation of A_1 , we can prove that

$$\begin{aligned} A_2 &= e^{-C(t_K - t_1)} E^T \left[f(D(t_1), \dots, D(t_{K-1})) \left(e^{D(T)} - 1 \right)^+ \right] \\ &= e^{-C(t_K - t_1)} E^T \left[F(D(t_1), \dots, D(t_{K-1}), D(T)) \right], \end{aligned}$$

for

$$F(x_1, \dots, x_K) := f(x_1, \dots, x_{K-1}) (e^{x_K} - 1)^+.$$

To ensure integrability, we define $g(x_1, \dots, x_K) := F(x_1, \dots, x_K) e^{-rx_K}$, with $1 < r < 2$.

Further, let

$$g_K(x_K) := (e^{x_K} - 1)^+ e^{-rx_K}.$$

Then, $g_K \in L^1(\mathbb{R})$, such that $g \in L^1(\mathbb{R}^K)$. Moreover, elementary integration shows that for all $y \in \mathbb{R}$

$$\hat{g}_K(y) = \frac{1}{(iy - r + 1)(iy - r)}.$$

Observe that $|\hat{g}_K(y)|_{\mathbb{C}} = (((1 - r)^2 + y^2)(r^2 + y^2))^{-1/2}$, thus, $\hat{g}_K \in L^1(\mathbb{R})$. Therefore, combining the last result with (26), we deduce that $\hat{g} \in L^1(\mathbb{R}^K)$, and

$$\hat{g}(y_1, \dots, y_K) = \frac{1}{(iy_K - r + 1)(iy_K - r)} \prod_{l=2}^K \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-y_{l-1}^2 / (4\beta \Delta t_l)}. \quad (28)$$

Since $g, \hat{g} \in L^1(\mathbb{R}^K)$, we can apply Theorem 3.2 in Eberlein et al. (2010) and obtain

$$E^T \left[F(D(t_1), \dots, D(t_{K-1}), D(T)) \right] = \frac{1}{(2\pi)^K} \int_{\mathbb{R}^K} \tilde{N}(R + iu) \hat{F}(iR - u) du, \quad (29)$$

with $R := (0, \dots, 0, r) \in \mathbb{R}^K$, $1 < r < 2$, and $\tilde{N}(R + iu)$ defined as

$$\tilde{N}(R + iu) := E^T \left[e^{iu_1 D(t_1) + \dots + iu_{K-1} D(t_{K-1}) + (iu_K + r) D(T)} \right].$$

As above, using the notation from (24), we derive

$$\begin{aligned} \tilde{N}(R + iu) &= \tilde{D}(u - iR, T) e^{-\int_0^T A(s, T) ds} \\ &\quad \times E_Q \left[\exp \left(\int_0^T \tilde{E}(s, u - iR, T) dL_s^1 + \int_0^T \tilde{F}(s, u - iR) dL_s^2 \right) \right]. \end{aligned}$$

Observe that as $1 < r < 2$, and in virtue of (4) and (7), $r\sigma_2(s) \leq M_2$ and $|r\beta(s) + (1 - r)\Sigma(s, T)| \leq (2r - 1)\frac{M_1}{3} \leq M_1$. Thus, the above expectation exists. Using independence of

L^1 and L^2 and (2), we obtain that

$$\begin{aligned} \tilde{N}(R + iu) &= \tilde{D}(u - iR, T) e^{-\int_0^T A(s, T) ds} \\ &\quad \times \exp\left(\int_0^T \theta_s^1(\tilde{E}(s, u - iR, T)) ds + \int_0^T \theta_s^2(\tilde{F}(s, u - iR)) ds\right). \end{aligned} \quad (30)$$

On the other hand, observe that for any $u \in \mathbb{R}^K$,

$$\hat{g}(u) = \int_{\mathbb{R}^K} e^{i\langle u, x \rangle} e^{-\langle R, x \rangle} F(x) dx = \hat{F}(u + iR).$$

Thus, we deduce that

$$\hat{F}(iR - u) = \hat{g}(-u) = \frac{1}{(iu_K + r - 1)(iu_K + r)} \prod_{l=2}^K \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-u_{l-1}^2 / (4\beta \Delta t_l)}. \quad (31)$$

Plugging (30) and (31) into (29), the claim follows. \square

4.2. Death benefit. In this section we compute the value of the death benefit previously defined as

$$\text{P}^{\text{DB}} = \sum_{i=1}^N E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \text{DB}(\bar{t}_i) \right].$$

Recall the definition of w_l from Equations (22) and (23). Further, for $i \in \{1, \dots, N\}$, we define

$$w_{\bar{t}_i} := \int_0^{\bar{t}_i} A(s, \bar{t}_i) ds + \int_0^{\bar{t}_i} f(0, s) ds - \delta \bar{t}_i - \omega(\bar{t}_i) - p(\bar{t}_i). \quad (32)$$

We also recall that $\Delta t_l := t_l - t_{l-1}$. For all $j \in \{1, \dots, K-1\}$, let $R := (0, \dots, 0, r) \in \mathbb{R}^{j+1}$ with $1 < r < 2$, $u \in \mathbb{R}^j$, and $v \in \mathbb{C}^{j+1}$. For all $0 \leq s \leq T$, for all $j \in \{1, \dots, K-2\}$ and all $i \in \{1, \dots, N\}$ such that $t_j < \bar{t}_i \leq t_{j+1}$, and for $j = K-1$ and all i such that $t_{K-1} < \bar{t}_i \leq T$, we define

$$\begin{aligned} D^{j,i}(u, T) &:= \exp\left(i \sum_{l=1}^j u_l w_l\right), \\ \tilde{D}^{j,i}(v, T) &:= D^{j,i}(v_1, \dots, v_j, T) \exp\left(i v_{j+1} w_{\bar{t}_i}\right), \\ E_{j,i}(s, u, T) &:= \Sigma(s, \bar{t}_i) + i(\beta(s) - \Sigma(s, T)) \sum_{l=1}^j u_l \mathbb{1}_{\{0 \leq s \leq t_l\}}, \\ \tilde{E}_{j,i}(s, v, T) &:= E_{j,i}(s, v_1, \dots, v_j, T) + i(\beta(s) - \Sigma(s, \bar{t}_i)) v_{j+1}, \\ F_j(s, u) &:= i\sigma_2(s) \sum_{l=1}^j u_l \mathbb{1}_{\{0 \leq s \leq t_l\}}, \\ \tilde{F}_j(s, v) &:= F_j(s, v_1, \dots, v_j) + i\sigma_2(s) v_{j+1}, \end{aligned} \quad (33)$$

$$M^{j,i}(u, T) := D^{j,i}(u, T) \exp \left(\int_0^{\bar{t}_i} \left(\theta_s^1(E_{j,i}(s, u, T)) + \theta_s^2(F_j(s, u)) \right) ds \right) \\ \times \prod_{l=2}^{j+1} \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-u_{l-1}^2 / (4\beta \Delta t_l)},$$

$$N^{j,i}(v, T) := \tilde{D}^{j,i}(v - iR, T) \exp \left(\int_0^{\bar{t}_i} \left(\theta_s^1(\tilde{E}_{j,i}(s, v - iR, T)) + \theta_s^2(\tilde{F}_j(s, v - iR)) \right) ds \right) \\ \times \frac{\exp(p(\bar{t}_i)(iv_{j+1} + r))}{(iv_{j+1} + r - 1)(iv_{j+1} + r)} \prod_{l=2}^{j+1} \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-v_{l-1}^2 / (4\beta \Delta t_l)}, \text{ for } v \in \mathbb{R}^{j+1}.$$

Finally, we define for $0 \leq s \leq T$, $u \in \mathbb{R}$ and $i \in \{1, \dots, N\}$

$$E_1(s, u) := \Sigma(s, \bar{t}_i) + (r + iu)(\beta(s) - \Sigma(s, \bar{t}_i)),$$

$$F_1(s, u) := (r + iu)\sigma_2(s),$$

$$N^i(u) := \exp \left((r + iu)w_{\bar{t}_i} \right) \exp \left(\int_0^{\bar{t}_i} \theta_s^1(E_1(s, u)) ds + \int_0^{\bar{t}_i} \theta_s^2(F_1(s, u)) ds \right) \\ \times \frac{\exp(p(\bar{t}_i)(iu + r))}{(iu + r - 1)(iu + r)}.$$

The following result for the DB value holds.

Theorem 4.2. *The value P^{DB} is given by*

$$P^{DB} = \sum_{i: \bar{t}_i \leq t_1} Q(\tau^m(x) \in [\bar{t}_{i-1}, \bar{t}_i]) (G(\bar{t}_i)B(0, \bar{t}_i) + G(\bar{t}_i)B(0, \bar{t}_i)A_{0,i}) \\ + \sum_{j=1}^{K-2} \sum_{i: \bar{t}_i \in (t_j, t_{j+1}]} Q(\tau^m(x) \in [\bar{t}_{i-1}, \bar{t}_i]) G(\bar{t}_i)B(0, \bar{t}_i) (A_{j,i}^1 + A_{j,i}^2) \\ + \sum_{i: \bar{t}_i \in (t_{K-1}, T]} Q(\tau^m(x) \in [\bar{t}_{i-1}, \bar{t}_i]) G(\bar{t}_i)B(0, \bar{t}_i) (A_{K-1,i}^1 + A_{K-1,i}^2),$$

where, using the notation from (33), for $j \in \{1, \dots, K-1\}$

$$A_{0,i} = \frac{e^{-\int_0^{\bar{t}_i} A(s, \bar{t}_i) ds}}{2\pi} \int_{\mathbb{R}} N^i(u) du, \\ A_{j,i}^1 = \frac{e^{-C(t_{j+1}-t_1)}}{(2\pi)^j} e^{-\int_0^{\bar{t}_i} A(s, \bar{t}_i) ds} \int_{\mathbb{R}^j} M^{j,i}(u, T) du, \\ A_{j,i}^2 = \frac{e^{-C(t_{j+1}-t_1)}}{(2\pi)^{j+1}} e^{-\int_0^{\bar{t}_i} A(s, \bar{t}_i) ds} \int_{\mathbb{R}^{j+1}} N^{j,i}(u, T) du.$$

The proof of Theorem 4.2 is similar to the proof of Theorem 4.1. For this reasons we defer it to Appendix B.

Remark 4.1. We have

$$E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \text{DB}(\bar{t}_i) \right] = Q(\tau^m(x) \in [\bar{t}_{i-1}, \bar{t}_i]) E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \mathbf{1}_{\{\tau^s \geq \bar{t}_i\}} \max(IS_{\bar{t}_i}, G(\bar{t}_i)) \right].$$

As shown in the previous theorem, this last expectation can be computed directly. However, this expression can also be traced back to P^{GMAB} . Note, that by definition,

$$E_Q \left[e^{-\int_0^{\bar{t}_i} r(u)du} \mathbb{1}_{\{\tau^s \geq \bar{t}_i\}} \max(IS_{\bar{t}_i}, G(\bar{t}_i)) \right] = \frac{P^{\text{GMAB}}(\bar{t}_i)}{Q(\tau^m(x) > \bar{t}_i)}.$$

To facilitate the computation of $P^{\text{GMAB}}(\bar{t}_i)$ we suggest the following approximation: first, for j such that $t_j < \bar{t}_i \leq t_{j+1}$, we obtain that

$$P^{\text{GMAB}}(\bar{t}_i) = Q(\tau^m(x) > \bar{t}_i) E_Q \left[e^{-\int_0^{\bar{t}_i} r(u)du} e^{-\int_0^{t_{j+1}} \lambda^s(u)du} \max(IS_{\bar{t}_i}, G(\bar{t}_i)) \right].$$

Now, we define $\hat{D}(\bar{t}_i) = Y_{\bar{t}_i} - \delta \bar{t}_i$, then

$$\max(IS_{\bar{t}_i}, G(\bar{t}_i)) = G(\bar{t}_i) \left[1 + \left(\exp(\hat{D}(\bar{t}_i)) - 1 \right)^+ \right].$$

The suggested approximation is

$$\hat{P}^{\text{GMAB}}(\bar{t}_i) = Q(\tau^m(x) > \bar{t}_i) G(\bar{t}_i) E_Q \left[e^{-\int_0^{\bar{t}_i} r(u)du} e^{-\int_0^{t_{j+1}} \lambda^s(u)du} \left(1 + \left(\exp(\hat{D}(\bar{t}_i)) - 1 \right)^+ \right) \right],$$

which can be evaluated using the results from Theorem 4.1, replacing T by \bar{t}_i and K by the $j+1$ for which $t_j < \bar{t}_i \leq t_{j+1}$.

4.3. Surrender benefit. Finally, we compute the value of the surrender benefit

$$P^{\text{SB}} = \sum_{i=1}^{K-1} E_Q \left[e^{-\int_0^{t_i} r(u)du} \text{SB}(t_i) \right].$$

We use the spot measure, i.e. the measure with the stock price chosen as numéraire. This allows to exploit the dependence between the surrender intensity and the stock price.

Recall the definition of w_l from Equations (22) and (23). Further, for $0 \leq s \leq T$, $i \in \{2, \dots, K-1\}$, $u \in \mathbb{R}^{i-1}$, and $v \in \mathbb{R}^i$, we define

$$\begin{aligned} D^i(u, T) &:= \exp \left(i \sum_{l=1}^{i-1} u_l w_l - \omega(t_i) \right), \\ \tilde{D}^i(v, T) &:= D^i(v_1, \dots, v_{i-1}, T) \exp \left(i v_i w_i \right), \\ E^i(s, u, T) &:= i(\beta(s) - \Sigma(s, T)) \sum_{l=1}^{i-1} u_l \mathbb{1}_{\{0 \leq s \leq t_l\}} + \beta(s), \\ \tilde{E}^i(s, v, T) &:= E^i(s, v_1, \dots, v_{i-1}, T) + i(\beta(s) - \Sigma(s, T)) v_i, \\ F^i(s, u) &:= i\sigma_2(s) \sum_{l=1}^{i-1} u_l \mathbb{1}_{\{0 \leq s \leq t_l\}} + \sigma_2(s), \\ \tilde{F}^i(s, v) &:= F^i(s, v_1, \dots, v_{i-1}) + i\sigma_2(s) v_i, \end{aligned} \tag{34}$$

$$\begin{aligned}
M^i(u, T) &:= D^i(u, T) \exp\left(\int_0^{t_i} \left(\theta_s^1(E^i(s, u, T)) + \theta_s^2(F^i(s, u))\right) ds\right) \\
&\quad \times \prod_{l=2}^i \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-u_{l-1}^2 / (4\beta \Delta t_l)}, \\
N^i(v, T) &:= \tilde{D}^i(v, T) \exp\left(\int_0^{t_i} \left(\theta_s^1(\tilde{E}^i(s, v, T)) + \theta_s^2(\tilde{F}^i(s, v))\right) ds\right) \\
&\quad \times \prod_{l=2}^{i+1} \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-v_{l-1}^2 / (4\beta \Delta t_l)}.
\end{aligned}$$

Theorem 4.3. *The value P^{SB} is given by*

$$P^{\text{SB}} = I \sum_{i=1}^{K-1} P(t_i) Q(\tau^m(x) > t_i) (B_i^1 - B_i^2),$$

where $B_1^1 = 1$ and for $i \in \{2, \dots, K-1\}$,

$$B_i^1 = \frac{e^{-C(t_i - t_1)}}{(2\pi)^{i-1}} \int_{\mathbb{R}^{i-1}} M^i(u, T) du,$$

and, for $i \in \{1, \dots, K-1\}$,

$$B_i^2 = \frac{e^{-C(t_{i+1} - t_1)}}{(2\pi)^i} \int_{\mathbb{R}^i} N^i(u, T) du.$$

Proof. By construction $P^{\text{SB}} = \sum_{i=1}^{K-1} E_Q \left[e^{-\int_0^{t_i} r(u) du} \text{SB}(t_i) \right]$. As

$$\text{SB}(t_i) = \mathbf{1}_{\{\tau^s = t_i\}} \mathbf{1}_{\{\tau^s < \tau^m(x)\}} I S_{t_i} P(t_i),$$

we are interested in computing the expression

$$\begin{aligned}
E_Q \left[e^{-\int_0^{t_i} r(s) ds} \mathbf{1}_{\{\tau^s = t_i\}} \mathbf{1}_{\{\tau^s < \tau^m(x)\}} S_{t_i} \right] &= E_Q \left[e^{-\int_0^{t_i} r(s) ds} \mathbf{1}_{\{\tau^s = t_i\}} \mathbf{1}_{\{t_i < \tau^m(x)\}} S_{t_i} \right] \\
&= Q(\tau^m(x) > t_i) E_Q \left[e^{-\int_0^{t_i} r(s) ds} \mathbf{1}_{\{\tau^s = t_i\}} S_{t_i} \right]. \quad (35)
\end{aligned}$$

It follows from (13) that, for $A \in \mathcal{F}_{t_i}^{L^1, L^2} \subset \mathcal{F}_{t_{i+1}}^{L^1, L^2}$,

$$\begin{aligned}
E_Q \left[\mathbf{1}_A \mathbf{1}_{\{\tau^s = t_i\}} \right] &= E_Q \left[\mathbf{1}_A (\mathbf{1}_{\{t_i \leq \tau^s\}} - \mathbf{1}_{\{t_{i+1} \leq \tau^s\}}) \right] \\
&= E_Q \left[\mathbf{1}_A \left(e^{-\int_0^{t_i} \lambda^s(u) du} - e^{-\int_0^{t_{i+1}} \lambda^s(u) du} \right) \right].
\end{aligned}$$

Consequently, equation (35) can be written as

$$\begin{aligned}
E_Q \left[e^{-\int_0^{t_i} r(s) ds} \mathbf{1}_{\{\tau^s = t_i\}} \mathbf{1}_{\{\tau^s < \tau^m(x)\}} S_{t_i} \right] &= Q(\tau^m(x) > t_i) E_Q \left[e^{-\int_0^{t_i} r(u) du} S_{t_i} e^{-\int_0^{t_i} \lambda^s(u) du} \right] \\
&\quad - Q(\tau^m(x) > t_i) E_Q \left[e^{-\int_0^{t_i} r(u) du} S_{t_i} e^{-\int_0^{t_{i+1}} \lambda^s(u) du} \right].
\end{aligned}$$

Let us write B_i^1 for the first expectation and B_i^2 for the second one. In order to compute these two expectations, we introduce the spot probability measure $Q^{S,i}$, $i = 1, \dots, K-1$

defined by its Radon-Nikodym derivative

$$\frac{dQ^{S,i}}{dQ} = e^{-\int_0^{t_i} r(u)du} S_{t_i}. \quad (36)$$

The above defines indeed a density process as the discounted stock price $e^{-\int_0^t r(u)du} S_t$ is a Q -martingale. We can use the new measure to simplify B_i^1 and B_i^2 as follows

$$\begin{aligned} B_i^1 &= E_Q \left[e^{-\int_0^{t_i} r(u)du} S_{t_i} e^{-\int_0^{t_i} \lambda^s(u)du} \right] \\ &= E_{Q^{S,i}} \left[e^{-\int_0^{t_i} \lambda^s(u)du} \right], \end{aligned} \quad (37)$$

$$\begin{aligned} B_i^2 &= E_Q \left[e^{-\int_0^{t_i} r(u)du} S_{t_i} e^{-\int_0^{t_{i+1}} \lambda^s(u)du} \right] \\ &= E_{Q^{S,i}} \left[e^{-\int_0^{t_{i+1}} \lambda^s(u)du} \right]. \end{aligned} \quad (38)$$

Note that $\lambda^s(u)$ for $t_i \leq u < t_{i+1}$ is defined by $D(t_i)$ and consequently it is $\mathcal{F}_{t_i}^{L^1, L^2}$ -measurable. Focusing first on B_i^1 , by construction $\lambda^s(u) = 0$ for $u \in [0, t_1)$, and therefore $B_1^1 = 1$. For $i \in \{2, \dots, K-1\}$, we follow the same strategy as in the proof of Theorem 4.1 for the computation of A_1 and consider

$$e^{C(t_i-t_1)} E_{Q^{S,i}} \left[e^{-\int_0^{t_i} \lambda^s(u)du} \right] = E_{Q^{S,i}} \left[f(D(t_1), \dots, D(t_{i-1})) \right], \quad (39)$$

where $f(x_1, \dots, x_{i-1}) := \prod_{l=2}^i e^{-\beta \Delta t_l x_{l-1}^2}$ with $\Delta t_l = t_l - t_{l-1}$. As in (27) we obtain that the last expectation is

$$\frac{1}{(2\pi)^{i-1}} \int_{\mathbb{R}^{i-1}} \tilde{M}^{i-1}(iu) \hat{f}(-u) du, \quad (40)$$

with

$$\hat{f}(u_1, \dots, u_{i-1}) = \prod_{l=2}^i \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-u_{l-1}^2 / (4\beta \Delta t_l)},$$

and $\tilde{M}^{i-1}(iu)$ defined as

$$\tilde{M}^{i-1}(iu) = E_{Q^{S,i}} \left[e^{iu_1 D(t_1) + \dots + iu_{i-1} D(t_{i-1})} \right].$$

From (A.1) it follows that

$$\begin{aligned} \tilde{M}^{i-1}(iu) &= E_Q \left[e^{iu_1 D(t_1) + \dots + iu_{i-1} D(t_{i-1}) + \int_0^{t_i} \sigma_2(s) dL_s^2 + \int_0^{t_i} \beta(s) dL_s^1 - \omega(t_i)} \right] \\ &= \exp \left(i \sum_{l=1}^{i-1} u_l (-p(t_l) - \delta T) + \int_0^T f(0, s) ds + \int_0^{t_i} A(s, T) ds - \omega(t_i) \right) \\ &\quad \times E_Q \left[\exp \left(i \sum_{l=1}^{i-1} \left(\int_0^{t_l} u_l \sigma_2(s) dL_s^2 + \int_0^{t_l} u_l (\beta(s) - \Sigma(s, T)) dL_s^1 \right) \right. \right. \\ &\quad \left. \left. + \int_0^{t_i} \sigma_2(s) dL_s^2 + \int_0^{t_i} \beta(s) dL_s^1 - \omega(t_i) \right) \right]. \end{aligned}$$

By equation (2) it follows that

$$\begin{aligned} E_Q \left[\exp \left(\int_0^{t_i} E^i(s, u, T) dL_s^1 + \int_0^{t_i} F^i(s, u) dL_s^2 \right) \right] \\ = \exp \left(\int_0^{t_i} \left(\theta_s^1(E^i(s, u, T)) + \theta_s^2(F^i(s, u)) \right) ds \right) \end{aligned}$$

Therefore, given the definition of $D^i(u, T)$ in (34) we have

$$\tilde{M}^{i-1}(iu) = D^i(u, T) \exp \left(\int_0^{t_i} \left(\theta_s^1(E^i(s, u, T)) + \theta_s^2(F^i(s, u)) \right) ds \right)$$

and the representation of B_i^1 follows from (40).

Finally, we note that

$$B_i^2 = e^{-C(t_{i+1}-t_1)} E_{Q^{S,i}} \left[\prod_{l=2}^{i+1} e^{-\beta \Delta t_l D(t_{l-1})^2} \right]. \quad (41)$$

Repeating mutatis mutandis the above arguments, the expression for B_i^2 follows. \square

5. NUMERICAL IMPLEMENTATION AND TESTING

This section discusses the numerical computation of the equations in Theorems 4.1, 4.2 and 4.3, based on the model features introduced in section 2. For the purpose of the numerical illustration, assumptions are required regarding the processes driving the stock price, the instantaneous forward rate and the intensity of mortality.

Starting from the financial market, we choose as a relevant Lévy process the Normal Inverse Gaussian (NIG) process introduced by Barndorff-Nielsen (1997) with cumulant function

$$\theta(u) = \mu u + \delta \left(\sqrt{\alpha^2 - \beta^2} - \sqrt{\alpha^2 - (\beta + u)^2} \right), \quad -\alpha - \beta < u < \alpha - \beta,$$

for $\mu \in \mathbb{R}$, $\delta > 0$, $0 \leq |\beta| < \alpha$. The parameter α controls the steepness of the density (and therefore its tail behaviour), β primarily controls the skewness of the distribution, whilst δ is the scale parameter; the location parameter μ is instead set to zero, without loss of generality.

Further, we assume a simplified Vasiček structure for the function $\sigma_1(s, T)$ so that for $a > 0$

$$\sigma_1(s, T) = \begin{cases} ae^{-a(T-s)}, & s \leq T \\ 0, & s > T \end{cases}$$

and

$$\Sigma(s, T) = \begin{cases} 1 - e^{-a(T-s)}, & s \leq T \\ 0, & s > T \end{cases}.$$

For the equity part, we assume $\sigma_2(s) = \sigma_2 > 0$ and $\beta(s) = b \in \mathbb{R}$.

Concerning the stochastic mortality model, we follow Krazzler et al. (2016) and choose a Brownian motion W as the driver of the mortality improvement ratio (i.e. $L^3 = \sigma W$) so that

$$d\xi_t = \kappa(\exp(-\lambda t) - \xi_t)dt + \sigma dW_t, \quad \sigma > 0,$$

with κ, b, z, λ as in section 3.1.

TABLE 1. Parameters for the Variable Annuity contract. Reference process: NIG. Financial Market Model parameters - source: [Eberlein and Rudmann \(2018\)](#). Mortality Model parameters - source: [Escobar et al. \(2016\)](#).

Variable Annuity		Financial Market Model			Surrender Model		Mortality Model	
			L_t^1	L_t^2				
I	100	α	4	5.73	β	0.05	b	12.1104
T	3, 4, 10 years	β	-3.8	-2.13	C	0.01	z	76.139
δ	0.01 p.a.	δ	1.34	8.3			κ	0.4806
$P(t_l)$	$0.95 + 0.05t_l/T$	a	0.0020898	-			λ	0.0195
Δt_l	1 year	σ_2	-	0.1818			σ	0.0254
$\bar{t}_i - \bar{t}_{i-1}$	6 months	b	-	0.0065				

A few considerations are in order. Firstly, as ξ is modelled as a Gaussian Ornstein-Uhlenbeck process, the mortality intensity can become negative with positive probability. However, this probability is negligible (less than 10^{-5}) as shown in Appendix A.2 in [Escobar et al. \(2016\)](#). For the occurring difficulties with negative values in the intensity, see for example [Bielecki and Rutkowski \(2002\)](#), Lemma 9.1.4 and the related remarks.

It follows that (see also [Escobar et al., 2016](#)) for any $0 \leq t \leq T$

$$Q(\tau^m(x) > t) = \exp\left(A_x(t) + B_x(t)\lambda_0^m(x)\right), \quad (42)$$

with

$$\begin{aligned} A_x(t) &:= \frac{c_1 \exp(c_2 t)}{c_3(c_2 + c_3)} [1 - \exp(-(c_2 + c_3)t)] \\ &\quad + \frac{1}{4} \left(\frac{c_4}{c_3}\right)^2 \frac{\exp(2c_5 t)}{c_5} [1 - \exp(-2c_5 t)] \\ &\quad - \left(\frac{c_4}{c_3}\right)^2 \frac{\exp(2c_5 t)}{2c_5 + c_3} [1 - \exp(-(2c_5 + c_3)t)] \\ &\quad - \frac{c_1 \exp(c_2 t)}{c_2 c_3} [1 - \exp(-c_2 t)] \\ &\quad + \frac{1}{4} \left(\frac{c_4}{c_3}\right)^2 \frac{\exp(2c_5 t)}{c_3 + c_5} [1 - \exp(-2(c_3 + c_5)t)], \\ B_x(t) &:= \frac{1}{c_3} [\exp(-c_3 t) - 1], \end{aligned}$$

and

$$c_1 := \frac{\kappa}{b} \exp\left(\frac{x-z}{b}\right), c_2 := \frac{1}{b} - \lambda, c_3 := \kappa - \frac{1}{b}, c_4 := \frac{\sigma}{b} \exp\left(\frac{x-z}{b}\right), c_5 := \frac{1}{b}.$$

All numerical experiments refer to contracts with a parameter setting as in Table 1; the parameters of the financial model are taken from the calibration exercise of [Eberlein and Rudmann \(2018\)](#), whilst for the stochastic mortality model we refer to the parameters in [Escobar et al. \(2016\)](#). The short maturity contracts are used for benchmarking purposes, whilst the 10 year maturity contract represents a realistic specification for practical purposes. Concerning the possible termination dates, for illustration purposes we use a 1 year frequency, i.e. $\Delta t_l = 1$, whilst the time grid for mortality, which is assumed to have finer granularity, is built with a frequency of 6 months. We note though that the results obtained in Theorems 4.1, 4.2 and 4.3 hold for any choice of the relevant time steps. The numerical schemes




FIGURE 1. $N(u, T)$, $T = 3$ years ($K = 2$). Left panel: real part of the function $N(u, T)$. Right panel: imaginary part of $N(u, T)$. Parameters: Table 1.

are implemented in MATLAB R2018A and run on a computer with an Intel i5-6500, 3.20 gigahertz CPU, and 8 gigabytes of RAM.

5.1. Implementation. The multidimensional integrals in Theorems 4.1, 4.2 and 4.3 are computed by means of Monte Carlo integration (see for example Pharr and Humphreys, 2010, and references therein), due to lack of efficient deterministic quadrature methods for very high dimensional problems as the ones considered here. Thus, the Monte Carlo estimate of a high-dimensional integral over the domain $\Omega \subseteq \mathbb{R}^d$

$$I = \int_{\Omega} f(x) dx$$

is obtained by evaluating the function $f(x)$ at M points x , drawn randomly in Ω with a given probability density $p(x)$, so that

$$I_{MC} = \frac{1}{M} \sum_{i=1}^M \frac{f(x_i)}{p(x_i)}.$$

The error is measured by means of the (unbiased) sample variance

$$\frac{\sum_{i=1}^M (I_i - I_{MC})^2}{M - 1};$$

this point shows that the rate of convergence is independent of the dimensionality of the integrand, which is the advantage of Monte Carlo integration compared to standard numerical quadrature techniques (if available at all) in computing even relatively low-dimensional integrals.

Classical Monte Carlo integration uses the uniform probability density. However, visual inspection of the integrand functions $M(u, T)$ and $N(u, T)$ of Theorem 4.1 reveals that $M(u, T)$ is strongly peaked around the origin, whilst $N(u, T)$ is characterised by alternations of peaks and troughs of significant magnitude - as shown in Figure 1 for $T = 3$ ($K = 2$). Similar considerations hold for the integrand functions in Theorems 4.2 and 4.3, due to the similarities between the corresponding payoff functions. These features, in general, can induce large variances in the Monte Carlo estimate of the corresponding integral.

TABLE 2. Benchmarking Monte Carlo integration with importance sampling - the case of the GMAB. Parameters: Table 1. ‘Quadrature’: MATLAB built-in functions `integral`, `integral2` and `integral3`. Bias/standard error expressed as percentage of the actual value. 100 batches of size 10^6 . CPU time expressed in seconds and referred to the average time of 1 batch of 10^6 iterations.

T	K		Quadrature	Monte Carlo integration (Imp. Sampling)		
			Value	Value	Bias (%)	Std. Error (%)
3 years	2	A_1	0.9867	0.9867	0.0035	0.0050
		A_2	0.1487	0.1482	0.3584	0.2683
	CPU	6.6323	31.2944			
			(A_1 : 0.1280) (A_2 : 6.5043)			
4 years	3	A_1	0.9703	0.9702	0.0139	0.0076
		A_2	0.1669	0.1669	0.0155	0.0647
	CPU	589.0926	51.6269			
			(A_1 : 1.3042) (A_2 : 587.7884)			

TABLE 3. Variable Annuity by Monte Carlo integration with importance sampling. Parameters: Table 1. Standard error expressed as percentage of the actual value. 100 batches of size 10^6 . CPU time expressed in seconds and referred to the average time of 1 batch of 10^6 iterations.

T	GMAB	DB	SB	VA
4 years	112.5121	4.9280	2.6997	120.1399
(Std. Error %)	0.0001	0.0256	0.1142	
CPU	51.6269	264.3202	31.7341	
10 years	93.0783	14.2344	15.4533	122.7661
(Std. Error %)	0.0003	0.0098	0.0534	
CPU	179.0842	1971.3826	717.3945	

Consequently, for variance reduction purposes and in order to speed up convergence, we implement importance sampling; in details, given that the integrand functions are strongly peaked around the origin, the importance distribution of choice is the multivariate Gaussian distribution with zero mean (i.e. centered around the peak), independent components, and a given variance matrix, which is treated as a parameter. Full deployment of the pricing algorithm in the setting of VEGAS and MISER Monte Carlo integration is left to future research.

5.2. Benchmarking and Testing. Numerical results for the valuation functions considered in this paper are reported in Table 2 and 3. In order to provide a reliable benchmark for the Monte Carlo integration procedure introduced above, we consider first some low dimension examples, for which the relevant multidimensional integrals can be tackled with standard quadrature packages (for example on the MATLAB platform, deterministic quadrature methods are available up to dimension 3). We illustrate the results obtained for the case of the GMAB. Similar performances are obtained for the DB and the SB as well, and we refer the interested reader to the online Appendix C for more details.

In details. Table 2 reports values obtained with deterministic quadrature methods, the corresponding estimate from Monte Carlo integration, and the CPU time. Together with the Monte Carlo estimate, I_{MC} , we also report measures of its accuracy in terms of the absolute value of the bias of the estimator expressed as percentage of the value obtained by quadrature I_Q , i.e.

$$100 \times \frac{|I_{MC} - I_Q|}{I_Q}.$$

In addition, we also report the percentage standard error of the Monte Carlo estimate

$$100 \times \frac{1}{I_{MC}} \sqrt{\frac{\sum_{i=1}^M (I_i - I_{MC})^2}{M(M-1)}}.$$

For increased reliability of the estimate of the CPU time, we base the numerical experiment on 100 repetitions of the Monte Carlo integration algorithm with sample size 10^6 .

We consider two examples for a contract with maturity $T = 3$ years (i.e. $K = 2$) and $T = 4$ years (i.e. $K = 3$) respectively. With annually spaced termination dates t_i , these choices imply that A_1 and A_2 are respectively one- and two-dimensional integrals in the first case, and two- and three-dimensional integrals in the second case, which can also be computed using packages for deterministic quadrature methods.

For importance sampling, numerical experiments show that relatively small biases and standard errors can be obtained by using the same variance fixed at 0.25 for the first $K - 1$ dimensions, and increasing this value to 1 for the last K^{th} dimension, as to cater for the higher variability of the integrand function $N(u, T)$. The results in Table 2 confirm the quality of the estimates as all biases and standard errors are below 0.5%. We also note the required CPU time, which shows the advantage of Monte Carlo integration already at dimension 3.

The value of the VA together with its components are reported in Table 3. We observe that the GMAB value decreases with increasing maturities, due to the larger number of dates at which the contract can be terminated by either death or surrender of the policyholder. Consistently with these findings, we also observe the increase in the value of both the DB and the SB for contracts with longer maturities. The overall value of the VA increases as well with longer maturities, denoting the dominant impact of the DB and the SB components.

Finally, we observe the higher computational cost of the DB and SB parts of the full variable annuity; this is due to the large number of integrals of the cumulant functions of interest (see definitions (33) and (34)) to be computed. This number depends on the granularity of both the \mathbf{t} and $\bar{\mathbf{t}}$ grids: the finer these grids, the more computational demanding these components of the contract become.

6. RESULTS: SENSITIVITY ANALYSIS

The sensitivity analysis is carried out by perturbing the parameters of interest one at a time, ceteris paribus. The reference case is given by a 10 years contract with the parameters set as in Table 1.

In Figure 2 we show the sensitivity of the VA and its components to the parameters β and C , controlling the surrender intensity. Consistently with intuition, the higher the value of β ,




FIGURE 2. Sensitivity Analysis: the surrender parameters (β, C). Top panels: left-hand-side - GMAB; right-hand-side: SB. Bottom panels: left-hand-side - DB; right-hand-side - VA. Maturity: $T = 10$ years. Other parameters: Table 1.




FIGURE 3. Sensitivity Analysis: the dependence parameters b and β . Top panels: left-hand-side - GMAB; right-hand-side: SB. Bottom panels: left-hand-side - DB; right-hand-side - VA. Maturity: $T = 10$ years. Other parameters: Table 1.




FIGURE 4. Sensitivity Analysis: the guarantee δ for different values of β . Top panels: left-hand-side - GMAB; right-hand-side: SB. Bottom panels: left-hand-side - DB; right-hand-side - VA. Maturity: $T = 10$ years. Other parameters: Table 1.

the stronger the influence of the financial market on the policyholder decision to surrender. This is captured in particular by the value of the surrender benefit, SB, which increases with β . Due to the higher surrender probability, the values of the GMAB and the DB reduce, and so does the resulting value of the VA.

The impact of the parameter β though becomes less relevant in correspondence of higher values of the constant C which represents the baseline surrender behaviour. In this case, the non-economic factors leading to the decision of surrender dominate the influence of the financial market to the point that the VA as well as its components are almost insensitive to β .

These results highlight that a correct quantification of the baseline surrender parameter C is of paramount importance given the significant impact on the value of the VA and its components. In this respect, it is crucial for insurance companies to formulate suitable surrender models, and to correctly calibrate them on the basis of the information that they could collect regarding the policyholder behaviour towards lapses.

Figure 3 shows that the impact of the parameter β is also affected by the dependence between the equity and the fixed income markets, here captured by the parameter b . Indeed, in presence of explicit dependence between the financial markets (i.e. $b \neq 0$), the behaviour of the policyholders becomes more responsive to changes in the market conditions. This is reflected in the rate of change of the contract's value.

Finally, in Figure 4 we illustrate the behaviour of the contract with respect to the guaranteed rate δ . A higher value of this rate corresponds to a more valuable VA, as one would

expect; the SB though is relatively insensitive to δ , also due to the low values of the latter used in this analysis.

7. CONCLUSION

We proposed a general framework for the valuation of a number of guarantees commonly included in variable annuity products, such as guaranteed minimum accumulation benefit, death benefit and surrender benefit, in the setting of a hybrid model based on multivariate Lévy processes, with surrender risk captured endogenously.

This framework proves to be tractable, and allows for the deployment of efficient numerical schemes based on Monte Carlo integration. We used this setting to gain insights into the contract sensitivity with respect to the model parameters, with special focus on surrender risk. The emphasis on surrender in particular finds its motivation in the large percentage of early terminations experienced by the insurance companies issuing variable annuities, and the resulting net cash outflow they sustain as a consequence (as also pointed out by LIMRA Secure Retirement Institute and IRI). The results obtained in our analysis confirm the importance of the surrender behaviour and its appropriate modelling.

We envisage the applicability of the model setup offered in this paper, and the proposed numerical scheme, also for other benefits offered by VAs, such as the post-retirement guaranteed minimum income benefit. This would represent a promising route for further research.

ACKNOWLEDGEMENTS

The Authors gratefully acknowledge the financial support from the Freiburg Institute for Advanced Studies (FRIAS) within the FRIAS Research Project ‘Linking Finance and Insurance: Theory and Applications (2017–2019)’.

REFERENCES

- Aksamit, A. and Jeanblanc, M. (2017), *Enlargement of Filtration with Finance in View*, Springer.
- Albizzati, M.-O. and Geman, H. (1994), ‘Interest rate risk management and valuation of the surrender option in life insurance policies’, *The Journal of Risk and Insurance* **61**(4), 616–637.
- Alonso-Garcia, J., Wood, O. and Ziveyi, J. (2018), ‘Pricing and hedging guaranteed minimum withdrawal benefits under a general Lévy framework using the COS method’, *Quantitative Finance* **18**(6), 1049–1075.
- Bacinello, A. R., Millosovich, P. and Montealegre, A. (2016), ‘The valuation of GMWB variable annuities under alternative fund distributions and policyholder behaviours’, *Scandinavian Actuarial Journal* **2016**(5), 446–465.
- Bacinello, A. R., Millosovich, P., Olivieri, A. and Pitacco, E. (2011), ‘Variable annuities: A unifying valuation approach’, *Insurance: Mathematics and Economics* **49**(3), 285–297.
- Bacinello, A. R. and Ortu, F. (1996), ‘Fixed income linked life insurance policies with minimum guarantees: Pricing models and numerical results’, *European Journal of Operational Research* **91**(2), 235–249.

- Ballotta, L. (2005), ‘A Lévy process-based framework for the fair valuation of participating life insurance contracts’, *Insurance: Mathematics and Economics* **37**(2), 173–196.
- Ballotta, L. (2009), ‘Pricing and capital requirements for with profit contracts: modelling considerations’, *Quantitative Finance* **9**(7), 803–817.
- Ballotta, L. and Haberman, S. (2006), ‘The fair valuation problem of guaranteed annuity options: The stochastic mortality environment case’, *Insurance: Mathematics and Economics*. **38**(1), 195 – 214.
- Barndorff-Nielsen, O. E. (1997), ‘Normal inverse Gaussian distributions and stochastic volatility modelling’, *Scandinavian Journal of Statistics* **24**(1), 1–13.
- Bielecki, T. and Rutkowski, M. (2002), *Credit Risk: Modeling, Valuation and Hedging*, Springer Verlag. Berlin Heidelberg New York.
- Biffis, E. (2005), ‘Affine processes for dynamic mortality and actuarial valuations’, *Insurance: Mathematics and Economics* **37**(3), 443 – 468.
- Boyle, P. P. and Schwartz, E. S. (1977), ‘Equilibrium prices of guarantees under equity-linked contracts’, *The Journal of Risk and Insurance* **44**(4), 639.
- Brennan, M. J. and Schwartz, E. S. (1976), ‘The pricing of equity-linked life insurance policies with an asset value guarantee’, *Journal of Financial Economics* **3**(3), 195–213.
- Dahl, M. (2004), ‘Stochastic mortality in life insurance: market reserves and mortality-linked insurance contracts’, *Insurance: Mathematics and Economics* **35**(1), 113–136.
- Dahl, M. and Møller, T. (2006), ‘Valuation and hedging of life insurance liabilities with systematic mortality risk’, *Insurance: Mathematics and Economics* **39**(2), 193–217.
- Deelstra, G. and Rayée, G. (2013), ‘Pricing variable annuity guarantees in a local volatility framework’, *Insurance: Mathematics and Economics* **53**(3), 650–663.
- Ducuroir, F., Miller, A. and Zians, J. (2016), ‘Lapse rate models in life insurance and a practical method to foresee interest rates dependencies’, A Reacfin White Paper.
- Duffie, D., Filipović, D. and Schachermayer, W. (2003), ‘Affine processes and applications in finance’, *Ann. Appl. Probab.* **13**, 984–1053.
- Eberlein, E., Glau, K. and Papapantoleon, A. (2010), ‘Analysis of Fourier transform valuation formulas and applications’, *Applied Mathematical Finance* **17**(3), 211–240.
- Eberlein, E., Jacod, J. and Raible, S. (2005), ‘Lévy term structure models: No-arbitrage and completeness’, *Finance Stochastics* **9**(1), 67–88.
- Eberlein, E. and Kallsen, J. (2019), *Mathematical Finance*, Springer. Forthcoming.
- Eberlein, E. and Keller, U. (1995), ‘Hyperbolic distributions in finance’, *Bernoulli* **1**(3), 281.
- Eberlein, E. and Raible, S. (1999), ‘Term structure models driven by general Lévy processes’, *Mathematical Finance* **9**, 31–53.
- Eberlein, E. and Rudmann, M. (2018), ‘Hybrid Lévy models: Design and computational aspects’, *Applied Mathematical Finance* **25**(5-6), 533–556.
- Escobar, M., Krayzler, M., Ramsauer, F., Saunders, D. and Zagst, R. (2016), ‘Incorporation of stochastic policyholder behavior in analytical pricing of GMABs and GMDBs’, *Risks* **4**(4), 41.
- Giacinto, M. D., Federico, S., Gozzi, F. and Vigna, E. (2014), ‘Income drawdown option with minimum guarantee’, *European Journal of Operational Research* **234**(3), 610–624.

- Grosen, A. and Jørgensen, P. L. (2002), ‘Life insurance liabilities at market value: An analysis of insolvency risk, bonus policy, and regulatory intervention rules in a barrier option framework’, *The Journal of Risk and Insurance* **69**(1), 63–91.
- Gudkov, N., Ignatieva, K. and Ziveyi, J. (2018), ‘Pricing of guaranteed minimum withdrawal benefits in variable annuities under stochastic volatility, stochastic interest rates and stochastic mortality via the componentwise splitting method’, *Quantitative Finance* **19**(3), 501–518.
- Kélani, A. and Quittard-Pinon, F. (2017), ‘Pricing and Hedging Variable Annuities in a Lévy Market: A Risk Management Perspective’, *Journal of Risk and Insurance* **84**(1), 209–238.
- Knoller, C., Kraut, G. and Schoenmaekers, P. (2016), ‘On the propensity to surrender a variable annuity contract: An empirical analysis of dynamic policyholder behavior’, *Journal of Risk and Insurance* **83**(4), 979–1006.
- Kolkiewicz, A. W. and Tan, K. S. (2006), ‘Unit-linked life insurance contracts with lapse rates dependent on economic factors’, *Annals of Actuarial Science* **1**(01), 49–78.
- Krayzler, M., Zagst, R. and Brunner, B. (2016), ‘Closed-form solutions for guaranteed minimum accumulation and death benefits’, *European Actuarial Journal* **6**(1), 197–231.
- Le Courtois, O. and Nakagawa, H. (2011), ‘On surrender and default risks’, *Mathematical Finance* **23**(1), 143–168.
- Li, J. and Szimayer, A. (2011), ‘The uncertain mortality intensity framework: Pricing and hedging unit-linked life insurance contracts’, *Insurance: Mathematics and Economics* **49**(3), 471–486.
- Loisel, S. and Milhaud, X. (2011), ‘From deterministic to stochastic surrender risk models: Impact of correlation crises on economic capital’, *European Journal of Operational Research* **214**(2), 348–357.
- Milevsky, M. A. and Promislow, S. D. (2001), ‘Mortality derivatives and the option to annuitise’, *Insurance: Mathematics and Economics* **29**(3), 299–318.
- Milhaud, X., Loisel, S. and Maume-Deschamps, V. (2011), ‘Surrender triggers in life insurance: what main features affect the surrender behavior in a classical economic context?’, *Bulletin Français d’Actuariat, Institut des Actuaire* **11**(22), 5–48.
- Nolte, S. and Schneider, J. C. (2017), ‘Don’t lapse into temptation: A behavioral explanation for policy surrender’, *Journal of Banking & Finance* **79**, 12 – 27.
- Pharr, M. and Humphreys, G. (2010), Monte Carlo integration I: Basic concepts, in M. Pharr and G. Humphreys, eds, ‘Physically Based Rendering’, second edn, Morgan Kaufmann, Boston, pp. 636 – 676.

APPENDIX A. SOME USEFUL RESULTS AND REPRESENTATIONS

Let us recall that $A(u, T) = \int_u^T \alpha(u, s) ds$ and $\Sigma(u, T) = \int_u^T \sigma_1(u, s) ds$. We derive here a representation result in the Lévy forward rate framework.

Lemma A.1. *For any $0 \leq t \leq T$, we have that*

$$-\int_t^T f(t, s) ds = \int_0^t r(s) ds - \int_0^T f(0, s) ds - \int_0^t A(u, T) du + \int_0^t \Sigma(u, T) dL_u^1,$$

where $f(t, s)$ is defined in (3).

Proof. Using Fubini's theorem for stochastic integrals, we deduce that

$$\begin{aligned} -\int_t^T f(t, s) ds &= \int_0^t r(s) ds - \int_0^t r(s) ds - \int_t^T f(0, s) ds - \int_t^T \int_0^t \alpha(u, s) du ds \\ &\quad + \int_t^T \int_0^t \sigma_1(u, s) dL_u^1 ds \\ &= \int_0^t r(s) ds - \left[\int_0^t f(0, s) ds + \int_0^t \left(\int_0^s \alpha(u, s) du \right) ds - \int_0^t \left(\int_0^s \sigma_1(u, s) dL_u^1 \right) ds \right] \\ &\quad - \int_t^T f(0, s) ds - \int_0^t \int_t^T \alpha(u, s) ds du + \int_0^t \int_t^T \sigma_1(u, s) ds dL_u^1 \\ &= \int_0^t r(s) ds - \left[\int_0^t f(0, s) ds + \int_0^t \int_u^t \alpha(u, s) ds du - \int_0^t \int_u^t \sigma_1(u, s) ds dL_u^1 \right] \\ &\quad - \int_t^T f(0, s) ds - \int_0^t \int_t^T \alpha(u, s) ds du + \int_0^t \int_t^T \sigma_1(u, s) ds dL_u^1, \end{aligned}$$

and the claim follows. \square

The above lemma allows the following representation of $D(t)$ defined in (11).

$$\begin{aligned} D(t) &= \int_0^t r(s) ds + \int_0^t \sigma_2(s) dL_s^2 + \int_0^t \beta(s) dL_s^1 - \omega(t) - p(t) + \int_t^T f(t, s) ds - \delta T \\ &= -p(t) - \delta T + \int_0^T f(0, s) ds + \int_0^t A(s, T) ds - \int_0^t \Sigma(s, T) dL_s^1 \\ &\quad + \int_0^t \sigma_2(s) dL_s^2 + \int_0^t \beta(s) dL_s^1 - \omega(t). \end{aligned} \tag{A.1}$$

In addition, setting $t = T$ in the above lemma, we obtain

$$0 = \int_0^t r(s) ds - \int_0^t f(0, s) ds - \int_0^t A(s, t) ds + \int_0^t \Sigma(s, t) dL_s^1 \tag{A.2}$$

and hence, the bank account has the following well-known representation, see for example Eberlein and Raible (1999) or (13) in Eberlein and Rudmann (2018),

$$B(t) = \frac{1}{B(0, t)} \exp \left(\int_0^t A(s, t) ds - \int_0^t \Sigma(s, t) dL_s^1 \right). \tag{A.3}$$

APPENDIX B. PROOF OF THEOREM 4.2

Proof. By definition,

$$P^{\text{DB}} = \sum_{i=1}^N E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \text{DB}(\bar{t}_i) \right].$$

From equation (10), we obtain that

$$E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \text{DB}(\bar{t}_i) \right] = Q(\tau^m(x) \in [\bar{t}_{i-1}, \bar{t}_i]) E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \mathbf{1}_{\{\tau^s \geq \bar{t}_i\}} \max(IS_{\bar{t}_i}, G(\bar{t}_i)) \right].$$

We distinguish two cases: when $\bar{t}_i \leq t_1$ and when $t_1 < \bar{t}_i$. We start with the detailed description of the second case which is the more complex one. The first case is treated at the end.

For $j \in \{1, \dots, K-2\}$ and i such that $t_j < \bar{t}_i \leq t_{j+1}$, as well as for $j = K-1$ and i such that $t_{K-1} < \bar{t}_i \leq T$, we work along the same line as in the proof of Theorem 4.1, and get

$$\begin{aligned} & E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \mathbf{1}_{\{\tau^s \geq \bar{t}_i\}} \max(IS_{\bar{t}_i}, G(\bar{t}_i)) \right] \\ &= E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \mathbf{1}_{\{\tau^s \geq t_{j+1}\}} \max(IS_{\bar{t}_i}, G(\bar{t}_i)) \right] \\ &= E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} e^{-\int_0^{t_{j+1}} \lambda^s(u) du} \max(IS_{\bar{t}_i}, G(\bar{t}_i)) \right] \\ &= G(\bar{t}_i) E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} e^{-\int_0^{t_{j+1}} \lambda^s(u) du} \left(1 + \left(\frac{IS_{\bar{t}_i}}{G(\bar{t}_i)} - 1 \right)^+ \right) \right]. \end{aligned} \quad (\text{B.1})$$

We introduce the \bar{t}_i -forward measure $Q^{\bar{t}_i}$ defined by its Radon-Nikodym density

$$\frac{dQ^{\bar{t}_i}}{dQ} = \frac{1}{B(0, \bar{t}_i)B(\bar{t}_i)}. \quad (\text{B.2})$$

Denoting the expectation with respect to $Q^{\bar{t}_i}$ by $E^{\bar{t}_i}$ the quantity in equation (B.1) is

$$\begin{aligned} & G(\bar{t}_i)B(0, \bar{t}_i) \left(E^{\bar{t}_i} \left[e^{-\int_0^{t_{j+1}} \lambda^s(u) du} \right] + E^{\bar{t}_i} \left[e^{-\int_0^{t_{j+1}} \lambda^s(u) du} \left(\frac{IS_{\bar{t}_i}}{G(\bar{t}_i)} - 1 \right)^+ \right] \right) \\ &= G(\bar{t}_i)B(0, \bar{t}_i) (A_{j,i}^1 + A_{j,i}^2), \end{aligned} \quad (\text{B.3})$$

with an obvious notation in the last line. We note that

$$e^{C(t_{j+1}-t_1)} A_{j,i}^1 = E^{\bar{t}_i} \left[f(D(t_1), \dots, D(t_j)) \right], \quad (\text{B.4})$$

with $f(x_1, \dots, x_j) = \prod_{l=2}^{j+1} e^{-\beta \Delta t_l x_{l-1}^2}$, and $\Delta t_l = t_l - t_{l-1}$. As in (27) we obtain that the expectation in (B.4) can be represented as

$$\frac{1}{(2\pi)^j} \int_{\mathbb{R}^j} \tilde{M}_i^j(iu) \hat{f}(-u) du, \quad (\text{B.5})$$

with

$$\hat{f}(u_1, \dots, u_j) = \prod_{l=2}^{j+1} \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-u_{l-1}^2 / (4\beta \Delta t_l)},$$

and $\tilde{M}_i^j(iu)$ defined as

$$\tilde{M}_i^j(iu) = E^{\bar{t}_i} \left[e^{iu_1 D(t_1) + \dots + iu_j D(t_j)} \right].$$

By the representation of the bank account in (A.3), the Radon-Nikodym density (B.2) can be written as

$$\frac{dQ^{\bar{t}_i}}{dQ} = \exp \left(- \int_0^{\bar{t}_i} A(s, \bar{t}_i) ds + \int_0^{\bar{t}_i} \Sigma(s, \bar{t}_i) dL_s^1 \right). \quad (\text{B.6})$$

Consequently, from the representation of $D(t)$ in (A.1), it follows

$$\begin{aligned} \tilde{M}_i^j(iu) &= E^{\bar{t}_i} \left[e^{iu_1 D(t_1) + \dots + iu_j D(t_j)} \right] \\ &= E_Q \left[e^{iu_1 D(t_1) + \dots + iu_j D(t_j) - \int_0^{\bar{t}_i} A(s, \bar{t}_i) ds + \int_0^{\bar{t}_i} \Sigma(s, \bar{t}_i) dL_s^1} \right] \\ &= \exp \left(i \sum_{l=1}^j u_l (-p(t_l) - \delta T + \int_0^T f(0, s) ds + \int_0^{t_l} A(s, T) ds - \omega(t_l)) - \int_0^{\bar{t}_i} A(s, \bar{t}_i) ds \right) \\ &\quad \times E_Q \left[\exp \left(i \sum_{l=1}^j \left(\int_0^{t_l} u_l \sigma_2(s) dL_s^2 + \int_0^{t_l} u_l (\beta(s) - \Sigma(s, T)) dL_s^1 \right) + \int_0^{\bar{t}_i} \Sigma(s, \bar{t}_i) dL_s^1 \right) \right]. \end{aligned}$$

This last expectation is finite in virtue of (4), and returns

$$\begin{aligned} E_Q \left[\exp \left(\int_0^{\bar{t}_i} E_{j,i}(s, u, T) dL_s^1 + \int_0^{\bar{t}_i} F_j(s, u) dL_s^2 \right) \right] \\ = \exp \left(\int_0^{\bar{t}_i} \left(\theta_s^1(E_{j,i}(s, u, T)) + \theta_s^2(F_j(s, u)) \right) ds \right), \end{aligned}$$

in virtue of the definitions in (33) and equation (2). Therefore, with $D^{j,i}(u, T)$ defined in (33) we have

$$\tilde{M}_i^j(iu) = D^{j,i}(u, T) \exp \left(\int_0^{\bar{t}_i} \left(\theta_s^1(E_{j,i}(s, u, T)) + \theta_s^2(F_j(s, u)) \right) ds \right).$$

Finally, combining (B.4) with (B.5) and the definition of $M^{j,i}(u, T)$ in (33), we deduce that

$$A_{j,i}^1 = \frac{e^{-C(t_{j+1}-t_1)}}{(2\pi)^j} \int_{\mathbb{R}^j} M^{j,i}(u, T) du.$$

For the purpose of the computation of $A_{j,i}^2$, we replace $D(t)$ defined in (11) with the following quantity

$$D_{t,t'} = Y_t - p(t) + \int_t^{t'} f(t, s) ds - \delta t', \quad 0 \leq t \leq t', \quad (\text{B.7})$$

so that

$$\left(\frac{IS_{\bar{t}_i}}{G(\bar{t}_i)} - 1 \right)^+ = \left(\exp(D_{\bar{t}_i, \bar{t}_i} + p(\bar{t}_i)) - 1 \right)^+.$$

We note that

$$e^{C(t_{j+1}-t_1)} A_{j,i}^2 = E^{\bar{t}_i} \left[h(D(t_1), \dots, D(t_j), D_{\bar{t}_i, \bar{t}_i}) \right], \quad (\text{B.8})$$

for $h(x_1, \dots, x_{j+1}) := f(x_1, \dots, x_j)(e^{x_{j+1} + p(\bar{t}_i)} - 1)^+$, with f given in (B.4). In order to ensure integrability, let us define $H(x_1, \dots, x_{j+1}) := h(x_1, \dots, x_{j+1})e^{-rx_{j+1}}$, for some $1 < r < 2$,

and

$$H_{j+1}(x_{j+1}) := (e^{x_{j+1}+p(\bar{t}_i)} - 1)^+ e^{-rx_{j+1}}.$$

Then, $H_{j+1} \in L^1(\mathbb{R})$, and $H \in L^1(\mathbb{R}^{j+1})$. Moreover, elementary integration shows that for all $y \in \mathbb{R}$

$$\hat{H}_{j+1}(y) = \frac{\exp(-p(\bar{t}_i)(iy-r))}{(iy-r+1)(iy-r)}.$$

Observe that $|\hat{H}_{j+1}(y)|_{\mathbb{C}} = e^{rp(\bar{t}_i)}(((1-r)^2+y^2)(r^2+y^2))^{-1/2}$, thus, $\hat{H}_{j+1} \in L^1(\mathbb{R})$. Therefore, combining the last result with (26), we deduce that $\hat{H} \in L^1(\mathbb{R}^{j+1})$, and

$$\hat{H}(y_1, \dots, y_{j+1}) = \frac{\exp(-p(\bar{t}_i)(iy_{j+1}-r))}{(iy_{j+1}-r+1)(iy_{j+1}-r)} \prod_{l=2}^{j+1} \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-y_{l-1}^2/(4\beta \Delta t_l)}. \quad (\text{B.9})$$

As $H, \hat{H} \in L^1(\mathbb{R}^{j+1})$, it follows from Theorem 3.2 in Eberlein et al. (2010) that

$$E^{\bar{t}_i} [h(D(t_1), \dots, D(t_j), D_{\bar{t}_i, \bar{t}_i})] = \frac{1}{(2\pi)^{j+1}} \int_{\mathbb{R}^{j+1}} \tilde{N}_i^{j+1}(R+iu) \hat{h}(iR-u) du, \quad (\text{B.10})$$

for $R = (0, \dots, 0, r) \in \mathbb{R}^{j+1}$, $1 < r < 2$, and $\tilde{N}_i^{j+1}(R+iu)$ defined as

$$\tilde{N}_i^{j+1}(R+iu) := E^{\bar{t}_i} [e^{iu_1 D(t_1) + \dots + iu_j D(t_j) + (iu_{j+1}+r) D_{\bar{t}_i, \bar{t}_i}}]. \quad (\text{B.11})$$

Using (B.7) and (A.2), we get

$$\begin{aligned} D_{\bar{t}_i, \bar{t}_i} &= -p(\bar{t}_i) - \delta \bar{t}_i + Y_{\bar{t}_i} \\ &= -p(\bar{t}_i) - \delta \bar{t}_i + \int_0^{\bar{t}_i} r(s) ds + \int_0^{\bar{t}_i} \sigma_2(s) dL_s^2 + \int_0^{\bar{t}_i} \beta(s) dL_s^1 - \omega(\bar{t}_i) \\ &= -p(\bar{t}_i) - \delta \bar{t}_i + \int_0^{\bar{t}_i} f(0, s) ds + \int_0^{\bar{t}_i} A(s, \bar{t}_i) ds - \int_0^{\bar{t}_i} \Sigma(s, \bar{t}_i) dL_s^1 \\ &\quad + \int_0^{\bar{t}_i} \sigma_2(s) dL_s^2 + \int_0^{\bar{t}_i} \beta(s) dL_s^1 - \omega(\bar{t}_i) \\ &= -p(\bar{t}_i) - \delta \bar{t}_i - \omega(\bar{t}_i) + \int_0^{\bar{t}_i} f(0, s) ds + \int_0^{\bar{t}_i} A(s, \bar{t}_i) ds + \int_0^{\bar{t}_i} \sigma_2(s) dL_s^2 \\ &\quad + \int_0^{\bar{t}_i} (\beta(s) - \Sigma(s, \bar{t}_i)) dL_s^1. \end{aligned}$$

Plugging the last quantity in (B.11), in virtue of (A.1) and (B.6), we deduce that

$$\begin{aligned} &\tilde{N}_i^{j+1}(R+iu) \\ &= E_Q \left[e^{iu_1 D(t_1) + \dots + iu_j D(t_j) + (iu_{j+1}+r) D_{\bar{t}_i, \bar{t}_i} - \int_0^{\bar{t}_i} A(s, \bar{t}_i) ds + \int_0^{\bar{t}_i} \Sigma(s, \bar{t}_i) dL_s^1} \right], \end{aligned}$$

consequently

$$\begin{aligned}
& \tilde{N}_i^{j+1}(R + iu) \\
&= \exp\left(i \sum_{l=1}^j u_l (-p(t_l) - \delta T + \int_0^T f(0, s) ds + \int_0^{t_l} A(s, T) ds - \omega(t_l)) - \int_0^{\bar{t}_i} A(s, \bar{t}_i) ds\right) \\
&\quad \times \exp\left((iu_{j+1} + r)(-p(\bar{t}_i) - \delta \bar{t}_i - \omega(\bar{t}_i) + \int_0^{\bar{t}_i} f(0, s) ds + \int_0^{\bar{t}_i} A(s, \bar{t}_i) ds)\right) \\
&\quad \times E_Q \left[\exp\left(i \sum_{l=1}^j \left(\int_0^{t_l} u_l \sigma_2(s) dL_s^2 + \int_0^{t_l} u_l (\beta(s) - \Sigma(s, T)) dL_s^1 \right) \right. \right. \\
&\quad \left. \left. + i \left(\int_0^{\bar{t}_i} u_{j+1} \sigma_2(s) dL_s^2 + \int_0^{\bar{t}_i} u_{j+1} (\beta(s) - \Sigma(s, \bar{t}_i)) dL_s^1 \right) \right. \right. \\
&\quad \left. \left. + \int_0^{\bar{t}_i} r \sigma_2(s) dL_s^2 + \int_0^{\bar{t}_i} r (\beta(s) - \Sigma(s, \bar{t}_i)) dL_s^1 + \int_0^{\bar{t}_i} \Sigma(s, \bar{t}_i) dL_s^1 \right) \right].
\end{aligned}$$

Using the definitions from (33), the above can be rewritten as

$$\tilde{N}_i^{j+1}(R + iu) = \tilde{D}^{j,i}(u - iR, T) E_Q \left[\exp\left(\int_0^{\bar{t}_i} \tilde{E}_{j,i}(s, u - iR, T) dL_s^1 + \int_0^{\bar{t}_i} \tilde{F}_j(s, u - iR) dL_s^2 \right) \right].$$

Observe that, due to $1 < r < 2$, as well as (4) and (7), $r\sigma_2(s) \leq M_2$ and $|r\beta(s) + (1-r)\Sigma(s, T)| \leq (2r-1)\frac{M_1}{3} \leq M_1$. Thus, the above expectation exists. Using the independence of L^1 and L^2 and (2), we obtain that

$$\begin{aligned}
\tilde{N}_i^{j+1}(R + iu) &= \tilde{D}^{j,i}(u - iR, T) \\
&\quad \times \exp\left(\int_0^{\bar{t}_i} \theta_s^1(\tilde{E}_{j,i}(s, u - iR, T)) ds + \int_0^{\bar{t}_i} \theta_s^2(\tilde{F}_j(s, u - iR)) ds \right). \quad (\text{B.12})
\end{aligned}$$

On the other hand, we observe that for any $u \in \mathbb{R}^{j+1}$,

$$\hat{H}(u) = \int_{\mathbb{R}^{j+1}} e^{i\langle u, x \rangle} e^{-\langle R, x \rangle} h(x) dx = \hat{h}(u + iR).$$

Consequently, we deduce that

$$\hat{h}(iR - u) = \hat{H}(-u) = \frac{\exp(p(\bar{t}_i)(iu_{j+1} + r))}{(iu_{j+1} + r - 1)(iu_{j+1} + r)} \prod_{l=2}^{j+1} \sqrt{\frac{\pi}{\beta \Delta t_l}} e^{-u_{l-1}^2 / (4\beta \Delta t_l)}. \quad (\text{B.13})$$

Plugging (B.12) and (B.13) in (B.10), and using the definition of $N^{j,i}(u, T)$ in (33), it follows that

$$A_{j,i}^2 = \frac{e^{-C(t_{j+1}-t_1)}}{(2\pi)^{j+1}} \int_{\mathbb{R}^{j+1}} N^{j,i}(u, T) du.$$

We now consider the case $\bar{t}_i \leq t_1$ for $i \in \{1, \dots, N\}$. Using the same arguments as above, we derive

$$E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \mathbf{1}_{\{\tau^s \geq \bar{t}_i\}} \max(IS_{\bar{t}_i}, G(\bar{t}_i)) \right] = E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \max(IS_{\bar{t}_i}, G(\bar{t}_i)) \right],$$

from which it follows that

$$\begin{aligned}
& E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \mathbf{1}_{\{\tau^s \geq \bar{t}_i\}} \max(IS_{\bar{t}_i}, G(\bar{t}_i)) \right] \\
&= B(0, \bar{t}_i) G(\bar{t}_i) E^{\bar{t}_i} \left[\left(1 + \left(\frac{IS_{\bar{t}_i}}{G(\bar{t}_i)} - 1 \right)^+ \right) \right] \\
&= B(0, \bar{t}_i) G(\bar{t}_i) + B(0, \bar{t}_i) G(\bar{t}_i) E^{\bar{t}_i} \left[\left(\exp(D_{\bar{t}_i, \bar{t}_i} + p(\bar{t}_i)) - 1 \right)^+ \right] \\
&= B(0, \bar{t}_i) G(\bar{t}_i) + B(0, \bar{t}_i) G(\bar{t}_i) E^{\bar{t}_i} [h_1(D_{\bar{t}_i, \bar{t}_i})],
\end{aligned}$$

where $h_1(x) := (e^{x+p(\bar{t}_i)} - 1)^+$. For some $1 < r < 2$, we define the function H_1 as $H_1(x) := (e^{x+p(\bar{t}_i)} - 1)^+ e^{-rx}$. By Theorem 3.2 in [Eberlein et al. \(2010\)](#), we get

$$E^{\bar{t}_i} [h_1(D_{\bar{t}_i, \bar{t}_i})] = \frac{1}{2\pi} \int_{\mathbb{R}} \tilde{N}_i(r + iu) \hat{h}_1(ir - u) du, \quad (\text{B.14})$$

with

$$\tilde{N}_i(r + iu) := E^{\bar{t}_i} [e^{(r+iu)D_{\bar{t}_i, \bar{t}_i}}].$$

Using the same arguments as above, and the definitions in [\(33\)](#), we deduce that

$$\begin{aligned}
& \tilde{N}_i(r + iu) \\
&= \exp \left((r + iu)w_{\bar{t}_i} - \int_0^{\bar{t}_i} A(s, \bar{t}_i) ds \right) E_Q \left[\exp \left(\int_0^{\bar{t}_i} E_1(s, u) dL_s^1 + \int_0^{\bar{t}_i} F_1(s, u) dL_s^2 \right) \right] \\
&= \exp \left((r + iu)w_{\bar{t}_i} - \int_0^{\bar{t}_i} A(s, \bar{t}_i) ds \right) \exp \left(\int_0^{\bar{t}_i} \theta_s^1(E_1(s, u)) ds + \int_0^{\bar{t}_i} \theta_s^2(F_1(s, u)) ds \right).
\end{aligned} \quad (\text{B.15})$$

On the other hand, we have

$$\hat{h}_1(ir - u) = \hat{H}_1(-u) = \frac{\exp(p(\bar{t}_i)(iu + r))}{(iu + r - 1)(iu + r)}. \quad (\text{B.16})$$

Plugging [\(B.15\)](#) and [\(B.16\)](#) in [\(B.14\)](#), we get

$$E^{\bar{t}_i} [h_1(D_{\bar{t}_i, \bar{t}_i})] = \frac{1}{2\pi} e^{-\int_0^{\bar{t}_i} A(s, \bar{t}_i) ds} \int_{\mathbb{R}} N^i(u) du.$$

Therefore,

$$E_Q \left[e^{-\int_0^{\bar{t}_i} r(u) du} \mathbf{1}_{\{\tau^s \geq \bar{t}_i\}} \max(IS_{\bar{t}_i}, G(\bar{t}_i)) \right] = B(0, \bar{t}_i) G(\bar{t}_i) + B(0, \bar{t}_i) G(\bar{t}_i) A_{0,i}.$$

□

ONLINE SUPPLEMENTARY MATERIAL FOR
 VARIABLE ANNUITIES IN A LÉVY-BASED HYBRID MODEL WITH
 SURRENDER RISK

LAURA BALLOTTA^(a), ERNST EBERLEIN^{(b),(c),(d)}, THORSTEN SCHMIDT^{(b),(c),(d)}, AND
 RAGHID ZEINEDDINE^(d)

^(a) FACULTY OF FINANCE, CASS BUSINESS SCHOOL, CITY, UNIVERSITY OF LONDON

^(b) FREIBURG INSTITUTE FOR ADVANCED STUDIES (FRIAS)

^(c) DEPARTMENT OF MATHEMATICAL STOCHASTICS, UNIVERSITY OF FREIBURG

^(d) UNIVERSITY OF STRASBOURG INSTITUTE FOR ADVANCED STUDY (USIAS)

APPENDIX C. FURTHER NUMERICAL RESULTS: BENCHMARKING

In this Appendix, we provide full results from the benchmarking exercise of the Monte Carlo integration procedure.

From section 5, we recall that we consider contracts with short maturity, and surrender frequency $\Delta t_l = 1$ year. In addition, we assume half-annually spaced \bar{t}_i for the mortality monitoring.

Consequently, a sensible benchmarking exercise for the value of both the DB and the SB with deterministic quadrature methods can be achieved by considering the 4 year maturity contract, i.e. $K = 3$.

Starting with the value of the DB, the term $A_{0,i}$ in Theorem 4.2 is a 1-dimensional integral for $i = 1, \dots, N$, which can be obtained by direct quadrature, and therefore is not considered by this benchmarking exercise.

Monte Carlo integration is instead used to compute the remaining terms appearing in Theorem 4.2, i.e. $A_{1,i}^1$, $A_{1,i}^2$, $A_{2,i}^1$, which are two-dimensional integrals, and $A_{2,i}^2$ which is a three-dimensional integral, for all $i = 1, \dots, N$. These are benchmarked against the values obtained by deterministic quadrature procedures in MATLAB.

For importance sampling, we choose the same values of the variance of the importance sampling distribution as in the GMAB case, i.e. 0.25 for the first $K - 1$ dimensions and 1 for the final K^{th} dimension. The quality of the estimate is confirmed by the negligible bias and standard errors reported in Table C.1.

Concerning the value of the SB, we notice that the first term in the sum defining P^{SB} in Theorem 4.3 is composed by a constant (B_1^1) and a one-dimensional integral (B_1^2), which is obtained by deterministic quadrature; similarly to the previous case this term is not considered in this analysis. The second term in this sum is instead formed by a one- and a two-dimensional integral (B_2^1 and B_2^2 respectively) for which we deploy Monte Carlo integration. Benchmarking is performed against the corresponding quadrature routines in MATLAB.

For importance sampling, given the relatively simple forms of the integrand functions, we use the same variance fixed at 0.16 across all the K dimensions. The goodness of the estimate is confirmed by the negligible bias and standard errors shown in Table C.1.

TABLE C.1. Benchmarking Monte Carlo integration with importance sampling. Parameters: Table 1. ‘Quadrature’: MATLAB built-in functions `integral`, `integral2` and `integral3`. Bias/standard error expressed as percentage of the actual value. 100 batches of size 10^6 . CPU time expressed in seconds and referred to the average time of 1 batch of 10^6 iterations.

GMAB			Quadrature	Monte Carlo integration (Imp. Sampling)		
T	K		Value	Value	Bias (%)	Std. Error (%)
3 years	2	A_1	0.9867	0.9867	0.0035	0.0050
		A_2	0.1487	0.1482	0.3584	0.2683
		CPU	6.6323	31.2944		
			(A_1 : 0.1280) (A_2 : 6.5043)			
4 years	3	A_1	0.9703	0.9702	0.0139	0.0076
		A_2	0.1669	0.1669	0.0155	0.0647
		CPU	589.0926	51.6269		
			(A_1 : 1.3042) (A_2 : 587.7884)			
<hr/>						
DB			Quadrature	Monte Carlo integration (Imp. Sampling)		
T	K		Value	Value	Bias (%)	Std. Error (%)
4 years	3	$A_{1,1}^1$	0.9866	0.9865	0.0047	0.0051
		$A_{1,1}^2$	0.1122	0.1114	0.6452	2.0849
		CPU	9.0793	30.3230		
			($A_{1,1}^1$: 0.4369) ($A_{1,1}^2$: 8.6424)			
	$(j, i) = (1, 2)$	$A_{1,2}^1$	0.9866	0.9866	0.0040	0.0047
		$A_{1,2}^2$	0.1239	0.1247	0.6851	0.3349
		CPU	9.0793	30.2637		
			($A_{1,2}^1$: 0.4369) ($A_{1,2}^2$: 8.6424)			
	$(j, i) = (2, 1)$	$A_{2,1}^1$	0.9703	0.9704	0.0100	0.0074
		$A_{2,1}^2$	0.1349	0.1353	0.2873	0.2220
		CPU	834.0646	52.1560		
			($A_{2,1}^1$: 1.1743) ($A_{2,1}^2$: 832.8903)			
	$(j, i) = (2, 2)$	$A_{2,2}^1$	0.9703	0.9705	0.0159	0.0079
		$A_{2,2}^2$	0.1464	0.1461	0.2016	0.1338
		CPU	834.0646	52.0597		
			($A_{2,2}^1$: 1.1743) ($A_{2,2}^2$: 832.8903)			
	$(j, i) = (2, 3)$	$A_{2,3}^1$	0.9703	0.9703	0.0080	0.0069
		$A_{2,3}^2$	0.1570	0.1569	0.0378	0.0685
		CPU	834.0646	52.9188		
			($A_{2,3}^1$: 1.1743) ($A_{2,3}^2$: 832.8903)			
	$(j, i) = (2, 4)$	$A_{2,4}^1$	0.9703	0.9703	0.0012	0.0075
		$A_{2,4}^2$	0.1670	0.1669	0.0275	0.0588
		CPU	834.0646	52.9617		
			($A_{2,4}^1$: 1.1743) ($A_{2,4}^2$: 832.8903)			
<hr/>						
SB			Quadrature	Monte Carlo integration (Imp. Sampling)		
T	K		Value	Value	Bias (%)	Std. Error (%)
4 years	3	B_2^1	0.9871	0.9871	0.0030	0.0029
		B_2^2	0.9717	0.9717	0.0066	0.0041
		CPU	1.1314	31.7341		
			(B_2^1 : 0.1280) (B_2^2 : 1.0034)			

APPENDIX D. AN ALTERNATIVE SPECIFICATION FOR THE SURRENDER INTENSITY

The surrender intensity λ^s is defined in (12) by

$$\lambda^s(t) = \beta D^2(t_i) + C = W_1(D(t_i)), \quad t_i \leq t < t_{i+1} \quad (\text{D.1})$$

where $W_1(x) := \beta x^2 + C$. Consequently $W_1(x) \rightarrow \infty$ as $|x| \rightarrow \infty$. One might be interested in the behaviour of the contract value for a surrender intensity whose values do not become arbitrarily large. Therefore, in the following we study the valuation of the variable annuity's components under an alternative specification which keeps the surrender intensity bounded.

We start with an observation concerning the behaviour of $D(t_i)$, $i = 1, \dots, K-1$. The following lemma shows that these quantities stay in a bounded interval with high probability. Based on this fact, Proposition D.2 and its corollary show that the contract values obtained with the alternative surrender intensity approximate the values under the original assumption.

Lemma D.1. *For any $\epsilon > 0$, there exists $L > 0$ such that*

$$\begin{aligned} Q^T\left(\bigcup_{i=1}^{K-1} \{|D(t_i)| > L\}\right) &\leq \epsilon, \\ Q^{S,j}\left(\bigcup_{i=1}^j \{|D(t_i)| > L\}\right) &\leq \epsilon, \text{ for any } j \in \{1, \dots, K-1\}, \\ Q^{\bar{t}_i}\left(\bigcup_{l=1}^j \{|D(t_l)| > L\}\right) &\leq \epsilon, \end{aligned} \quad (\text{D.2})$$

where the last result holds for $j \in \{1, \dots, K-2\}$ and any i such that $t_j < \bar{t}_i \leq t_{j+1}$, and for $j = K-1$ and any i such that $t_{K-1} < \bar{t}_i \leq T$. $Q^{\bar{t}_i}$ is defined in (B.2), $Q^{S,j}$ is defined in (36), and Q^T is defined in (19).

Proof. Let L_1 denote a positive constant. Applying the Markov inequality and the inequality $e^{|x|} \leq e^x + e^{-x}$, we get

$$\begin{aligned} Q^T\left(\bigcup_{i=1}^{K-1} \{|D(t_i)| > L_1\}\right) &\leq \sum_{i=1}^{K-1} Q^T(\{|D(t_i)| > L_1\}) = \sum_{i=1}^{K-1} Q^T(\{e^{|D(t_i)|} > e^{L_1}\}) \\ &\leq e^{-L_1} \sum_{i=1}^{K-1} [E^T(e^{D(t_i)}) + E^T(e^{-D(t_i)})]. \end{aligned} \quad (\text{D.3})$$

Now we use the representation given in (A.1), the definition of Q^T in (19) and (A.3), to derive

$$\begin{aligned} E^T(e^{D(t_i)}) &= e^{-p(t_i) - \delta T + \int_0^T f(0,s)ds + \int_0^{t_i} A(s,T)ds - \omega(t_i)} E^T\left(e^{\int_0^{t_i} \sigma_2(s)dL_s^2 + \int_0^{t_i} (\beta(s) - \Sigma(s,T))dL_s^1}\right) \\ &= e^{-p(t_i) - \delta T + \int_0^T f(0,s)ds + \int_0^{t_i} A(s,T)ds - \omega(t_i) - \int_0^T A(s,T)ds} \\ &\quad \times E_Q\left(e^{\int_0^{t_i} \sigma_2(s)dL_s^2 + \int_0^{t_i} (\beta(s) - \Sigma(s,T))dL_s^1 + \int_0^T \Sigma(s,T)dL_s^1}\right) \end{aligned}$$

Observe that, by independence of L^1 and L^2 , we have

$$\begin{aligned} & E_Q(e^{\int_0^{t_i} \sigma_2(s) dL_s^2 + \int_0^{t_i} (\beta(s) - \Sigma(s, T)) dL_s^1 + \int_0^T \Sigma(s, T) dL_s^1}) \\ &= E_Q(e^{\int_0^{t_i} \sigma_2(s) dL_s^2 + \int_0^T ((\beta(s) - \Sigma(s, T)) \mathbf{1}_{\{0 \leq s \leq t_i\}} + \Sigma(s, T)) dL_s^1}) \\ &= E_Q(e^{\int_0^{t_i} \sigma_2(s) dL_s^2}) E_Q(e^{\int_0^T ((\beta(s) - \Sigma(s, T)) \mathbf{1}_{\{0 \leq s \leq t_i\}} + \Sigma(s, T)) dL_s^1}), \end{aligned}$$

where the last quantities are finite due to (4) and (7). Therefore, $C_i := E^T(e^{D(t_i)})$ is finite, and by similar arguments $C'_i := E^T(e^{-D(t_i)})$ is finite as well. Observe that $e^{-L_1} \sum_{i=1}^{K-1} (C_i + C'_i) \leq \epsilon$ is equivalent to $L_1 \geq -\log(\epsilon) + \log(\sum_{i=1}^{K-1} (C_i + C'_i))$. As a consequence of (D.3) and the last argument, we deduce that for $L_1 \geq -\log(\epsilon) + \log(\sum_{i=1}^{K-1} (C_i + C'_i))$, $Q^T(\bigcup_{i=1}^{K-1} \{|D(t_i)| > L_1\}) \leq \epsilon$.

By similar arguments, we can prove that for any $j \in \{1, \dots, K-1\}$, there exists $\hat{L}_j > 0$, such that $Q^{S,j}(\bigcup_{i=1}^j \{|D(t_i)| > \hat{L}_j\}) \leq \epsilon$. We can prove also that for $j \in \{1, \dots, K-2\}$ and any i such that $t_j < \bar{t}_i \leq t_{j+1}$, and for $j = K-1$ and any i such that $t_{K-1} < \bar{t}_i \leq T$, there exists $L_{i,j}$ such that $Q^{\bar{t}_i}(\bigcup_{l=1}^j \{|D(t_l)| > L_{i,j}\}) \leq \epsilon$. Therefore, by defining $\bar{L} := \max_{(i,j)} L_{i,j}$ and $L := \max(L_1, \hat{L}_1, \dots, \hat{L}_{K-1}, \bar{L})$, we deduce that (D.2) holds true. \square

Remark D.1. The constant L in Lemma D.1 depends on ϵ and should be denoted L_ϵ but we omit this notation for simplicity.

Based on the above, let us replace W_1 by another positive and continuous function W_2 which coincides with W_1 on $[-L, L]$, L as in Lemma D.1, is constant outside this compact set on the positive half line and converges to 0 as $x \rightarrow -\infty$. Specifically, we define W_2 as

$$W_2(x) = \begin{cases} W_1(L) & \text{for } x \in (L, \infty) \\ W_1(x) & \text{for } x \in [-L, L] \\ (\beta L^2 + C)e^{L+x} & \text{for } x \in (-\infty, -L) \end{cases},$$

with the constants β and C as in (12). The new surrender intensity $\bar{\lambda}^s$ is defined as

$$\bar{\lambda}^s(t) = W_2(D(t_i)), \quad t_i \leq t < t_{i+1}, \quad (\text{D.4})$$

for $i \in \{1, \dots, K-1\}$, and $\bar{\lambda}^s(t) = 0$ for $t \in [0, t_1] \cup [t_K, T]$. Then, as we will see in Corollary D.3, the corresponding values, denoted by \bar{P}^{GMAB} , \bar{P}^{DB} and \bar{P}^{SB} , approximate P^{GMAB} , P^{DB} and P^{SB} .

Starting with the value of the GMAB, we rewrite equation (21) as

$$\begin{aligned} \frac{\bar{P}^{\text{GMAB}}}{Q(\tau^m(x) > T)B(0, T)G(T)} &= E^T \left[e^{-\int_0^{t_K} \bar{\lambda}^s(u) du} \right] + E^T \left[e^{-\int_0^{t_K} \bar{\lambda}^s(u) du} \left(\frac{IS_T}{G(T)} - 1 \right)^+ \right] \\ &=: \bar{A}_1 + \bar{A}_2, \end{aligned}$$

with

$$\begin{aligned} \bar{A}_1 &= E^T \left[\prod_{i=2}^K (e^{-W_2(D(t_{i-1})) \Delta t_i}) \right], \\ \bar{A}_2 &= E^T \left[\prod_{i=2}^K (e^{-W_2(D(t_{i-1})) \Delta t_i}) (e^{D(T)} - 1)^+ \right]. \end{aligned}$$

Following the same calculations as in Theorem 4.2, we deduce that

$$\begin{aligned}\bar{P}^{\text{DB}} &= \sum_{i: \bar{t}_i \leq t_1} Q(\tau^m(x) \in [\bar{t}_{i-1}, \bar{t}_i]) (G(\bar{t}_i)B(0, \bar{t}_i) + G(\bar{t}_i)B(0, \bar{t}_i)A_{0,i}) \\ &+ \sum_{j=1}^{K-2} \sum_{i: \bar{t}_i \in (t_j, t_{j+1}]} Q(\tau^m(x) \in [\bar{t}_{i-1}, \bar{t}_i]) G(\bar{t}_i)B(0, \bar{t}_i) (\bar{A}_{j,i}^1 + \bar{A}_{j,i}^2) \\ &+ \sum_{i: \bar{t}_i \in (t_{K-1}, T]} Q(\tau^m(x) \in [\bar{t}_{i-1}, \bar{t}_i]) G(\bar{t}_i)B(0, \bar{t}_i) (\bar{A}_{K-1,i}^1 + \bar{A}_{K-1,i}^2),\end{aligned}$$

where, $A_{0,i}$ is the same as in Theorem 4.2, and for $j \in \{1, \dots, K-2\}$ and any i such that $t_j < \bar{t}_i \leq t_{j+1}$, and for $j = K-1$ and any i such that $t_{K-1} < \bar{t}_i \leq T$

$$\bar{A}_{j,i}^1 := E^{\bar{t}_i} \left[e^{-\int_0^{t_j+1} \bar{\lambda}^s(u) du} \right] = E^{\bar{t}_i} \left[\prod_{i=2}^{j+1} e^{-W_2(D(t_{i-1})\Delta t_i)} \right],$$

$$\bar{A}_{j,i}^2 := E^{\bar{t}_i} \left[e^{-\int_0^{t_j+1} \bar{\lambda}^s(u) du} \left(\frac{IS_{\bar{t}_i}}{G(\bar{t}_i)} - 1 \right)^+ \right] = E^{\bar{t}_i} \left[\prod_{i=2}^{j+1} e^{-W_2(D(t_{i-1})\Delta t_i)} \left(e^{D_{\bar{t}_i, \bar{t}_i + p(\bar{t}_i)} - 1} - 1 \right)^+ \right].$$

Finally, by the same argument as in Theorem 4.3, we deduce that

$$\begin{aligned}\bar{P}^{\text{SB}} &= I \sum_{j=1}^{K-1} P(t_j) Q(\tau^m(x) > t_j) E_{Q^{S,j}} \left[e^{-\int_0^{t_j} \bar{\lambda}^s(u) du} \right] \\ &\quad - I \sum_{j=1}^{K-1} P(t_j) Q(\tau^m(x) > t_j) E_{Q^{S,j}} \left[e^{-\int_0^{t_j+1} \bar{\lambda}^s(u) du} \right] \\ &= I \sum_{j=1}^{K-1} P(t_j) Q(\tau^m(x) > t_j) (\bar{B}_j^1 - \bar{B}_j^2),\end{aligned}$$

with $\bar{B}_1^1 = 1$, due to $\bar{\lambda}^s(u) = 0$, for $u \in [0, t_1)$ by construction, and

$$\begin{aligned}\bar{B}_j^1 &= E_{Q^{S,j}} \left[\prod_{l=2}^j (e^{-W_2(D(t_{l-1})\Delta t_l)}) \right], \text{ for } j \geq 2, \\ \bar{B}_j^2 &= E_{Q^{S,j}} \left[\prod_{l=2}^{j+1} (e^{-W_2(D(t_{l-1})\Delta t_l)}) \right].\end{aligned}$$

Given the above, the following result holds.

Proposition D.2. *We have*

- (1) $|\bar{A}_1 - A_1| \leq 2\epsilon,$
- (2) $|\bar{A}_2 - A_2| \leq 2C_2\epsilon^{1/2},$ with $C_2 := E^T [(e^{D(T)} - 1)^2]^{1/2},$
- (3) $|\bar{A}_{j,i}^1 - A_{j,i}^1| \leq 2\epsilon,$
- (4) $|\bar{A}_{j,i}^2 - A_{j,i}^2| \leq 2C_{2,i}\epsilon^{1/2},$ with $C_{2,i} := E^{\bar{t}_i} [(e^{D_{\bar{t}_i, \bar{t}_i + p(\bar{t}_i)} - 1} - 1)^2]^{1/2},$

where in (3) and (4) we have either $j \in \{1, \dots, K-2\}$ and i such that $t_j < \bar{t}_i \leq t_{j+1}$, or $j = K-1$ and i such that $t_{K-1} < \bar{t}_i \leq T$,

$$(5) \quad |\bar{B}_j^1 - B_j^1| \leq 2\epsilon, \text{ for } j \in \{2, \dots, K-1\},$$

$$(6) \quad |\bar{B}_j^2 - B_j^2| \leq 2\epsilon, \text{ for } j \in \{1, \dots, K-1\}.$$

A_1, A_2 are given in (21), $A_{j,i}^1, A_{j,i}^2$ in (B.3), and B_j^1, B_j^2 are given in (37) and (38).

Proof. It suffices to prove (2) which is representative for the degree of sophistication. The proofs of (1), (3), (4), (5) and (6) follow mutatis mutandis. From the definition of W_1 given in (D.1), we derive

$$\begin{aligned} \bar{A}_2 &= E^T \left[\prod_{i=2}^K (e^{-W_2(D(t_{i-1}))\Delta t_i}) (e^{D(T)} - 1)^+ \right] \\ &= E^T \left[\prod_{i=2}^K (e^{-W_2(D(t_{i-1}))\Delta t_i}) (e^{D(T)} - 1)^+ \mathbf{1}_{\{\cap_{i=2}^K \{|D(t_{i-1})| \leq L\}\}} \right] \\ &\quad + E^T \left[\prod_{i=2}^K (e^{-W_2(D(t_{i-1}))\Delta t_i}) (e^{D(T)} - 1)^+ \mathbf{1}_{\{\cup_{i=2}^K \{|D(t_{i-1})| > L\}\}} \right] \\ &= E^T \left[\prod_{i=2}^K (e^{-W_1(D(t_{i-1}))\Delta t_i}) (e^{D(T)} - 1)^+ \mathbf{1}_{\{\cap_{i=2}^K \{|D(t_{i-1})| \leq L\}\}} \right] \\ &\quad + a_1, \end{aligned}$$

with the obvious definition of a_1 in the last line. The third equality holds as W_1 and W_2 coincide on $[-L, L]$. Observe that

$$\begin{aligned} &E^T \left[\prod_{i=2}^K (e^{-W_1(D(t_{i-1}))\Delta t_i}) (e^{D(T)} - 1)^+ \mathbf{1}_{\{\cap_{i=2}^K \{|D(t_{i-1})| \leq L\}\}} \right] \\ &= E^T \left[\prod_{i=2}^K (e^{-W_1(D(t_{i-1}))\Delta t_i}) (e^{D(T)} - 1)^+ \right] \\ &\quad - E^T \left[\prod_{i=2}^K (e^{-W_1(D(t_{i-1}))\Delta t_i}) (e^{D(T)} - 1)^+ \mathbf{1}_{\{\cup_{i=2}^K \{|D(t_{i-1})| > L\}\}} \right] \\ &= A_2 - a_2, \end{aligned}$$

with the obvious notation in the last line. We deduce that

$$|\bar{A}_2 - A_2| \leq (a_1 + a_2).$$

As W_1 is a positive function, the following holds

$$\begin{aligned}
a_2 &\leq E^T \left[\left(e^{D(T)} - 1 \right)^+ \mathbf{1}_{\{\cup_{i=1}^{K-1} \{ |D(t_i)| > L \}\}} \right] \\
&\leq E^T \left[\left| e^{D(T)} - 1 \right| \mathbf{1}_{\{\cup_{i=1}^{K-1} \{ |D(t_i)| > L \}\}} \right] \\
&\leq E^T \left[\left(e^{D(T)} - 1 \right)^2 \right]^{1/2} Q^T \left(\bigcup_{i=1}^{K-1} \{ |D(t_i)| > L \} \right)^{1/2} \\
&\leq C_2 \epsilon^{1/2},
\end{aligned}$$

where the third line is a consequence of the Cauchy-Schwarz inequality. For the last inequality we use (D.2) together with the fact that $C_2 := E^T \left[\left(e^{D(T)} - 1 \right)^2 \right]^{1/2} < \infty$. This last fact is easy to verify from the definition of $D(T)$ in (A.1). A similar bound can be achieved for a_1 , therefore

$$|\bar{A}_2 - A_2| \leq 2C_2 \epsilon^{1/2},$$

which proves (2). \square

Based on the last proposition, the interested reader can prove the following corollary.

Corollary D.3. *We have*

- (1) $|\bar{P}^{\text{GMAB}} - P^{\text{GMAB}}| \leq 2Q(\tau^m(x) > T)B(0, T)G(T)[\epsilon + C_2\epsilon^{1/2}]$,
with $C_2 := E^T \left[\left(e^{D(T)} - 1 \right)^2 \right]^{1/2}$.
- (2) $|\bar{P}^{\text{DB}} - P^{\text{DB}}| \leq 2 \sum_{i: \bar{t}_i > t_1} Q(\tau^m(x) \in [\bar{t}_{i-1}, \bar{t}_i])B(0, \bar{t}_i)G(\bar{t}_i)[\epsilon + C_{2,i}\epsilon^{1/2}]$,
with $C_{2,i} := E^{\bar{t}_i} \left[\left(e^{D_{\bar{t}_i, \bar{t}_i + p(\bar{t}_i)}} - 1 \right)^2 \right]^{1/2}$.
- (3) $|\bar{P}^{\text{SB}} - P^{\text{SB}}| \leq 2IP(t_1)Q(\tau^m(x) > t_1)\epsilon + 4I\epsilon \sum_{j=2}^{K-1} P(t_j)Q(\tau^m(x) > t_j)$.