



City Research Online

City St George's, University of London

Citation: Hatch, S. L., Woodhead, C., Frissa, S., Fear, N. T., Verdecchia, M., Stewart, R., Reichenberg, A., Morgan, C., Bebbington, P. E., McManus, S., et al (2012). Importance of Thinking Locally for Mental Health: Data from Cross-Sectional Surveys Representing South East London and England. PLOS ONE, 7(12), e48012. doi: 10.1371/journal.pone.0048012

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/23641/>

Link to published version: <https://doi.org/10.1371/journal.pone.0048012>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Importance of Thinking Locally for Mental Health: Data from Cross-Sectional Surveys Representing South East London and England

Stephani L. Hatch^{1*}, Charlotte Woodhead¹, Souci Frissa¹, Nicola T. Fear², Maria Verdecchia¹, Robert Stewart³, Abraham Reichenberg¹, Craig Morgan⁴, Paul Bebbington⁵, Sally McManus⁶, Traolach Brugha⁷, Bwalya Kankulu¹, Jennifer L. Clark¹, Billy Gazard¹, Robert Medcalf¹, Matthew Hotopf¹, the SELCoH study team[†]

1 King's College London, Psychological Medicine, Institute of Psychiatry, London, United Kingdom, **2** King's College London, Academic Centre for Defence Mental Health, London, United Kingdom, **3** King's College London, Section of Epidemiology, Health Service and Population Research Department, Institute of Psychiatry, London, United Kingdom, **4** King's College London, Section of Social Psychiatry, Health Service and Population Research Department, Institute of Psychiatry, London, United Kingdom, **5** University College London, Department of Mental Health Sciences, London, United Kingdom, **6** National Centre for Social Research, London, United Kingdom, **7** Department of Health Sciences, New Academic Unit, Leicester General Hospital, Leicester, United Kingdom

Abstract

Background: Reliance on national figures may be underestimating the extent of mental ill health in urban communities. This study demonstrates the necessity for local information on common mental disorder (CMD) and substance use by comparing data from the South East London Community Health (SELCoH) study with those from a national study, the 2007 English Adult Psychiatric Morbidity Study (APMS).

Methodology/Principal Findings: Data were used from two cross-sectional surveys, 1698 men and women residing in south London and 7403 men and women in England. The main outcome, CMD, was indicated by a score of 12 or above on the Revised Clinical Interview Schedule. Secondary outcomes included hazardous alcohol use and illicit drug use. SELCoH sample prevalence estimates of CMD were nearly twice that of the APMS England sample estimates. There was a four-fold greater proportion of depressive episode in the SELCoH sample than the APMS sample. The prevalence of hazardous alcohol use was higher in the national sample. Illicit drug use in the past year was higher in the SELCoH sample, with cannabis and cocaine the illicit drugs reported most frequently in both samples. In comparisons of the SELCoH sample with the APMS England sample and the APMS sample from the Greater London area in combined datasets, these differences remained after adjusting for socio-demographic and socioeconomic indicators for all outcomes.

Conclusions/Significance: Local information for estimating the prevalence of CMD and substance use is essential for surveillance and service planning. There were similarities in the demographic and socioeconomic factors related to CMD and substance use across samples.

Citation: Hatch SL, Woodhead C, Frissa S, Fear NT, Verdecchia M, et al. (2012) Importance of Thinking Locally for Mental Health: Data from Cross-Sectional Surveys Representing South East London and England. PLoS ONE 7(12): e48012. doi:10.1371/journal.pone.0048012

Editor: Xiang Yang Zhang, Baylor College of Medicine, United States of America

Received: May 3, 2012; **Accepted:** September 20, 2012; **Published:** December 12, 2012

Copyright: © 2012 Hatch et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research was supported by the Biomedical Research Nucleus data management and informatics facility at South London and Maudsley NHS Foundation Trust, which is funded by the National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London and a joint infrastructure grant from Guy's and St Thomas' Charity and the Maudsley Charity. These funders had no involvement in study design, data collection, analysis or the decision to submit for publication. The Department of Health funded the APMS 2007 and had no involvement in study design, data collection, analysis or the decision to submit for publication.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: stephani.hatch@kcl.ac.uk

† Membership of the SELCoH study team is provided in the Acknowledgments.

Introduction

While national psychiatric epidemiological surveys are useful in setting macro-level policy agendas, combating inequalities in mental health requires knowledge generated from local data [1]. Epidemiological information allows high risk groups to be identified by services and also clarifies the adequacy of existing and planned service strategies. However, reliance on national figures may underestimate the extent of mental ill health in certain

communities – for example in those with high levels of deprivation [2–3]. In this context, variations in the characteristics both of individual members of the population and of the area as a whole may engender concomitant variation in morbidity. Moreover, differences in prevalence of mental ill health and related needs may change in the face of changes in the distribution of economic resources and the availability of social services; in demography as a result of shifting migration patterns; and in the built environment, particularly in urban settings [4–9]. The impact of persistent,

disabling and common mental disorders is most profound and costly in highly populated, urban communities [10–12]. However, data are lacking for such local communities, and there are methodological difficulties in extrapolating from national to local levels; for example national prevalence studies usually have insufficient sample size for robust analyses at the local level, and they may select only a proportion of the areas in their sampling frame [13–15]. Such difficulties are particularly acute if the major determinants of prevalence are characteristics of the area rather than of individual members of the population [13]. Thus, health planners need access to detailed local information in order to develop public mental health strategies. For this reason, we established the South East London Community Health Survey (SELCoH), which covers an inner-city population. Here we aimed to demonstrate the necessity for this level of information by comparing our data on the symptoms of common mental disorder (CMD) and the prevalence of common mental disorder (CMD) and substance use with those from a national study – the 2007 English Adult Psychiatric Morbidity Study (APMS). We hypothesise that the prevalence of CMD and substance use, including alcohol use, will be higher in the SELCoH sample in comparison to the APMS sample, and these differences will be explained by socio-demographic attributes and the socioeconomic status of the individual sample members.

Methods

Sample

The South East London Community Health (SELCoH) study is a local psychiatric and physical morbidity survey of 1698 adults, aged 16 years and over from 1075 randomly selected households in the South London boroughs of Southwark and Lambeth [16]. In the two boroughs, there is higher deprivation than the England average, but similar proportions of economically active and inactive residents in comparison to greater London [17–20]. The boroughs are also ethnically diverse, with a greater number of Black Caribbean residents but fewer South Asian residents than other areas of London [21]. The SELCoH sample resided in a community setting served by South London and Maudsley National Health Service Trust (SLaM) in the UK, and the partnership between King's College London and the SLaM NHS trust allows this and other research to inform and benefit clinical treatment.

Data were collected between June 2008 and December 2010 and the SELCoH study sampling strategy resulted in a 51.9% household participation rate and 71.9% participation within households. The study was conducted as a component of SLaM's NIHR Mental Health Biomedical Research Centre [16]. SELCoH aimed to provide updated local population data to inform the configuration of services.

The national data come from the Adult Psychiatric Morbidity Study (APMS) 2007 [22]. Data were collected in England between October 2006 and December 2007 from a random sample of private household residents aged 16 years and older. Using a multi-stage stratified probability sampling design [22], 7403 eligible individuals completed full interviews (57% of the total sampled).

SELCoH applied similar methods to the APMS [23], with a few notable exceptions. Both studies used the UK Small User Postcode Address File (PAF) for the sampling frames: this has near complete coverage of private households (defined as one person or group of people who have the accommodation as their only or main residence and for groups who either share at least one meal a day or share the living area). Trained interviewers conducted

interviews in participants' homes and administered structured assessments using laptops. The same measures were used for the principal outcomes, and many of the questions on demographic variables within SELCoH were based upon APMS methodology. Differences in methods included attempts in the SELCoH sample to interview all adults aged 16 years and over in each eligible household, whereas the APMS sample randomly selected one adult, aged 16 years and over to be interviewed in each eligible household using the Kish grid method. In addition, the APMS sample was stratified by region (Strategic Health Authorities) and manual and non-manual social class [22], whereas the SELCoH sample was stratified by borough.

Ethics statement

Ethical approval for the SELCoH study was received from the King's College London research ethics committee for non-clinical research populations; reference CREC/07/08-152. Ethical approval for APMS 2007 was obtained from the Royal Free Hospital and Medical School Research Ethics Committee, one of the Research Ethics Committees of the National Research Ethics Service for non-clinical populations.

Measures

Common mental disorder. Common mental disorder (CMD) in both samples was assessed by the Revised Clinical Interview Schedule (CIS-R) [24] - a structured interview that asks about 14 symptom domains (using skips to allow asymptomatic individuals to answer a minimum of 28 questions): fatigue, sleep problems, irritability, worry, depression, depressive ideas, anxiety, obsessions, subjective memory and concentration, somatic symptoms, compulsions, phobias, physical health worries and panic. A total CIS-R score of 12 or more is conventionally used to indicate the overall presence of CMD, with a total score of 18 or more that denotes a level of symptoms that are likely to require treatment. The CIS-R also provides ICD-10 diagnoses for six mental disorders through a standard algorithm (depressive episode, generalised anxiety disorder, panic disorder, phobias, obsessive-compulsive disorder and mixed anxiety and depressive disorder). Mixed anxiety and depressive disorder is a residual category, covering cases that have a total CIS-R score of 12 or more, but do not meet the specific criteria for the other five disorders. As such, it tends to be less severe, particularly in comparison to depressive episode, and less likely to be associated with treatment seeking and treatment receipt [25–26].

Substance use. In both samples, hazardous alcohol use was assessed by the Alcohol Use Disorders Identification Test (AUDIT) [27], developed by the World Health Organization, and comprising ten questions relating to alcohol consumption, symptoms of alcohol dependence and problems related to alcohol abuse within the last 12 months. Each item is scored 0–4 with a summed overall score ranging from 0–40. For this analysis, an AUDIT score of 8 or more has been used to define hazardous drinking; AUDIT scores were also recoded into four groups: an AUDIT score of 0 has been used to define non-drinkers, 1 to 7 indicated moderate drinkers, 8 to 15 defined hazardous alcohol use, and a score of 16 or more defined hazardous alcohol use that is harmful to health. Participants reported illicit drug use in the past year for the following drugs in both samples: cannabis, amphetamines, cocaine, crack, ecstasy, LSD, tranquilliser and heroin. *Any drug use* in the past year referred to use of at least one drug in the past year, while concurrent *poly drug use* (in the same time period) [28] was defined as the use of 2 or more drugs.

Socio-demographic and socio-economic factors. The following socio-demographic and socioeconomic indicators avail-

able in both samples were included: gender; the ethnic group categories of White British, Black Caribbean, Black African, South Asian or Other; age (years) both as a continuous variable and in the following categories: 16–24, 25–34, 35–44, 45–54, 55–64, and 65 and over; marital status; education level categorised as: no qualifications, qualifications up to GCSE or Ordinary Level, Advanced Level, and degree level or above; occupational social class (non-manual vs. manual); employment status categorised as: paid employment, unemployed, and economically inactive (i.e., student, permanent sick/disabled, temporary sick, retired, looking after the home and children); and housing tenure categorised as: own/mortgage, rented and rent free.

Analysis

Analyses were conducted in STATA 11 [29]. We used survey commands (svy) for estimates of prevalence and associations where appropriate to generate robust standard errors.

All analyses of SELCoH data accounted for clustering by household inherent in the study design and weighted for within household non-response, comparing all eligible household members (i.e., 16 years or older) by gender and age. As previously reported [16], the sample was similar to the most recent UK Census information in 2001 with regards to demographic and socioeconomic indicators for the catchment area under study, with the exception of the sample being slightly younger and having more students within the economically inactive group. However, there were no differences in the distribution by age across categories in comparisons of the SELCoH and APMS samples. In reference to the proportion of students, there were more students identified in the SELCoH sample (12.5%) in comparison to the APMS sample (3.3%); however, student status is likely to be underestimated in the APMS sample because the survey only inquired about student status as a response to a question asking about the main reason for being currently out of work.

All analyses of APMS data accounted for weighting, clustering and stratification built into the survey design. Weighting in the APMS data accounted for clusters by postcode sectors, stratification based on socio-economic status within regional areas and non-response based on differences between the sample and the mid-census estimates [22]. Analysis was conducted for those with complete data for all variables. We report the unweighted frequencies, and applied Pearson's χ^2 tests with Rao & Scott second-order corrections with 95 percent confidence intervals for categorical outcomes. Odds ratios (OR) with 95 percent confidence intervals (CI) were calculated for the associations of categorical outcomes with socio-demographic and socio-economic indicators. Models adjusted for gender and age in years are presented for all logistic regression models. To examine whether or not identified differences will be explained by socio-demographic attributes and the socioeconomic status of the individual sample members, data were combined to make direct comparisons across samples in fully adjusted multivariable logistic regression models for fully adjusted models. The definition of social class in SELCoH excludes participants without a current occupation; thus, social class is not included in the fully adjusted models for the combined dataset.

Results

Sample characteristics

Compared with the national survey data, the SELCoH sample had an over-representation of women, the youngest participants, Black Caribbean and Black African groups, never-married and

Table 1. Socio-demographic characteristics of the South East London Community Health (SELCoH) and Adult Psychiatric Morbidity Study (APMS) 2007 England samples.

		SELCoH	APMS 2007
Total		1698	7403
Gender	Female	959 (66.7)	4206 (51.4)
	Male	739 (33.3)	3197 (48.6)
Ethnic group	White British	1051 (63.5)	6499 (85.1)
	Black-Caribbean	143 (8.7)	104 (1.5)
	Black-African	234 (13.2)	78 (1.5)
	Asian	63 (3.5)	258 (5.0)
	Other	205 (11.2)	414 (6.9)
Age (years)	16–24	356 (18.2)	568 (14.2)
	25–39	572 (28.6)	1744 (26.1)
	40–54	432 (24.1)	1834 (25.9)
	55–64	163 (13.3)	1279 (14.8)
	65+	175 (15.9)	1978 (19.0)
Marital status	Never married	678 (35.7)	1428 (22.7)
	Married/cohabiting	786 (46.4)	4133 (62.9)
	Divorced/separated	181 (12.6)	893 (7.5)
	Widowed	53 (5.3)	949 (7.0)
Education levels	No qualifications	228 (16.9)	2278 (26.2)
	Up to GCSE level	332 (20.1)	2103 (30.9)
	Advanced level	426 (23.7)	938 (15.1)
	Higher degree or above	693 (39.3)	1916 (27.8)
Social class^a	Non-manual	703 (73.4)	4277 (60.8)
	Manual	244 (26.6)	2732 (39.2)
Employment status	Paid employment	921 (51.2)	3964 (60.4)
	Unemployed	170 (9.3)	164 (2.9)
	Economically inactive	598 (39.5)	3250 (36.7)
Housing tenure	Own/mortgage	525 (32.4)	5143 (69.8)
	Rented	1058 (61.7)	2077 (28.3)
	Rent free	112 (5.8)	117 (1.9)

Values are numbers (percentages) of respondents; weighted percentages to account for survey design; frequencies are unweighted and may not add up due to missing values.

^aSocial class is based on occupation and participants without a current occupation were excluded in both samples.

doi:10.1371/journal.pone.0048012.t001

divorced/separated groups, those with higher education levels, and the non-manual and the unemployed groups (Table 1).

Prevalence of CMD symptoms and diagnostic categories by gender

Table 2 compares the one-week prevalences and 95% confidence intervals (CI) for the distributions of CIS-R scores and psychiatric disorders across the two samples. A greater proportion of SELCoH sample met the criteria for CMD than the APMS sample. The differences in proportions was present for those with CIS-R scores at 12 to 17 and 18 and above, the latter denoting a level of symptoms that are likely to require treatment [17]. Overall, psychiatric diagnoses identified using CIS-R were significantly more common in the SELCoH sample than in the APMS sample ($p < 0.001$). There was a striking discrepancy in the prevalence of depressive episode. This, the most symptomatically

severe of the common mental disorders covered in the CIS-R, was four times more frequent in the SELCoH sample than in the national survey and was indeed the most prevalent mental disorder in the SELCoH participants. In contrast, the most common psychiatric diagnosis in the APMS sample was the residual diagnosis of mixed anxiety and depressive disorder, and there was no difference in the prevalence of this condition between the SELCoH and APMS samples. Obsessive compulsive disorder and panic disorder were the least common psychiatric diagnoses in both samples, but the prevalence of these disorders was greater in the APMS sample ($p < 0.001$ and $p = 0.03$, respectively).

The prevalence estimates for the CIS-R symptoms are shown in Figure 1. In both samples, fatigue was the most commonly reported symptom (33.7% in the SELCoH sample and 27.8% in the APMS sample), followed by sleep problems (32.5% in the SELCoH sample and 17.6% in the APMS sample) and worry (29.5% in the SELCoH sample and 18.7% in the APMS sample). Panic was the least common symptom reported (3.2% in the SELCoH sample and 2.5% in the APMS sample). An increase in symptom reporting was generally observed in the SELCoH sample; however, despite the disparity in the prevalence of a depressive episode diagnosis, depressed mood was one of the symptoms with a similar prevalence in the two surveys.

Socio-demographic and socio-economic factors and CMD

Table 3 compares the one-week prevalence for CMD by socio-demographic and socio-economic indicators across samples. There was a higher prevalence of CMD in the SELCoH sample within each stratum of all indicators, with the exception of comparisons between those in the Black African ethnic group. In general, there were similar associations between the majority of socio-demographic and socioeconomic indicators and CMD, with the exception of ethnicity and social class. There was no association between ethnicity and CMD in the SELCoH sample, although there was a trend towards lower prevalence of CMD in the Black African group than other groups (as previously noted in this area of south London [30]). In contrast, people in the Black African group in the APMS sample were more likely to meet the criteria for CMD than the White British group.

Prevalence of substance use by gender

As shown in Table 4, there was a higher prevalence of hazardous alcohol use in the APMS sample than in the SELCoH sample. In both samples, men reported more hazardous alcohol use than women ($p < 0.001$). In contrast to alcohol use, there was a two-fold higher prevalence of any illicit drug use in the past year in the SELCoH sample compared to the APMS sample. Further, there was a higher prevalence of concurrent poly drug use in the past year in the SELCoH sample. In both samples, cannabis was the most commonly used drug in the past year, but reported by a greater proportion of the SELCoH sample. Cocaine was the second most commonly reported drug in the past year in both samples. In both samples, men reported more cannabis and cocaine use in the past year than women.

Substance use by socio-demographic and socio-economic factors

Tables 5 and 6 compare the prevalence and confidence intervals for hazardous alcohol use and illicit drug use by socio-demographic and socio-economic indicators. There was a higher prevalence of hazardous alcohol use and a lower prevalence of illicit drug use in the APMS sample across all strata of indicators,

except for a lower proportion of hazardous alcohol use in the Asian ethnic group in the APMS sample. With few exceptions, there were similar associations between the socio-demographic indicators with both substance use outcomes across samples. However, those who were married or cohabitating and widowed in the APMS sample were less likely to report hazardous alcohol use in comparison to those in the never married group. In addition, those in the Black Caribbean group in the APMS sample were more likely to report illicit drug use than those in the White ethnic group, but there was no difference between the two ethnic groups in the SELCoH sample. Among the socio-economic indicators, those with a higher education level and in a non-manual social class were more likely to report hazardous alcohol use in the SELCoH sample, but there was no difference in these respects in the APMS sample. There was no association between employment status in the SELCoH sample, but those who were economically inactive in the APMS sample were less likely to report hazardous alcohol use. There were also greater odds of illicit drug use among those in manual social class and those in rented accommodation in the APMS sample, but not in the SELCoH sample.

Comparisons of CMD, hazardous alcohol use and illicit drug use in combined analysis

In Table 7 (full models presented in Table S1), the SELCoH sample was combined with data from the APMS England sample and the APMS sample from the Greater London area ($N = 792$) to determine whether or not differences in CMD, hazardous alcohol use and illicit drug use were explained by socio-demographic attributes and the socioeconomic status of the individual sample members. For the APMS London sample, the prevalence of CMD was 14.8% (95% CI 12.3–17.8), depressive episode was 2.6% (95% CI 1.7–4.1) and hazardous alcohol use was 22.6% (95% CI 19.4–26.3). Whereas, the prevalence of any drug use in the past year in the APMS London was greater than the national estimate and closer to the SELCoH sample estimate (APMS England = 8.9% (95% CI 8.1–9.8); APMS London = 13.1% (95% CI 9.9–17.2); SELCoH = 18.1% (95% CI 16.1–20.2). In the fully adjusted models, SELCoH participants had increased odds of CMD in comparison to both APMS England and APMS London participants after adjusting for socio-demographic and socio-economic indicators. SELCoH participants had lower odds of hazardous alcohol use than the APMS England and APMS London samples after adjusting for socio-demographic and socio-economic indicators. In contrast, SELCoH participants had more than twice the odds of reporting illicit drug use in the past year than the APMS England sample after adjusting for socio-demographic and socio-economic indicators in a combined dataset. This difference was weaker, but in the same direction in the comparison between the SELCoH sample and the APMS London sample in the fully adjusted model.

Discussion

Main findings

In comparing two household psychiatric morbidity surveys from South East London and a national sample in England, this study addressed two aims: to compare (1) prevalence estimates of common mental disorder (CMD) in the past week, alcohol and illicit drug use in the past year and (2) whether or not any identified differences could be explained by socio-demographic attributes and the socioeconomic status of the individual sample members. There were several notable differences: the prevalence of CMD in the SELCoH sample was more than 10% higher than

Table 2. Comparisons of prevalence estimates of common mental conditions and psychiatric diagnoses by gender.

	Total						Women			Men		
	SELCoH		APMS 2007		P value of difference		SELCoH		APMS 2007		P value of difference	
	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	P value of difference
CIS-R scores												
CMD 'case' (12+)	24.2 (21.9–26.5)	15.1 (14.1–16.0)	<0.001	27.3 (24.3–30.3)	18.4 (17.0–19.8)	<0.001	17.9 (15.0–20.9)	11.6 (10.3–12.8)	<0.001			
0–5	54.9 (52.3–57.6)	68.1 (66.7–69.4)	<0.001	50.3 (46.9–53.6)	62.3 (60.5–64.2)	<0.001	64.3 (60.7–67.9)	74.2 (72.3–75.9)	<0.001			
6–11	20.9 (18.8–22.9)	16.9 (15.8–17.9)		22.4 (19.7–25.1)	19.3 (17.9–20.7)		17.7 (14.8–20.5)	14.3 (12.8–15.7)				
12–17	11.7 (10.1–13.4)	7.5 (6.8–8.3)		12.5 (10.4–14.6)	9.1 (8.0–10.2)		10.2 (7.9–12.5)	5.9 (4.9–6.8)				
≥18	12.4 (10.7–14.2)	7.5 (6.9–8.2)		14.8 (12.5–17.1)	9.3 (8.4–10.2)		7.8 (5.7–9.8)	5.7 (4.8–6.5)				
Primary diagnosis												
Mixed anxiety and depressive disorder	7.7 (6.4–9.1)	8.4 (7.7–9.2)	0.4	8.9 (7.1–10.8)	10.3 (9.2–11.5)	0.2	5.3 (3.7–6.9)	6.4 (5.6–7.4)	0.3			
Depressive episode	11.9 (10.2–13.6)	2.9 (2.6–3.4)	<0.001	13.7 (11.4–15.9)	3.5 (3.0–4.1)	<0.001	8.4 (6.3–10.5)	2.4 (1.9–3.1)	<0.001			
Generalised anxiety disorder	4.4 (3.4–5.5)	4.4 (3.9–4.9)	0.9	4.8 (3.4–6.3)	5.3 (4.6–6.1)	0.6	3.7 (2.3–5.1)	3.4 (2.8–4.2)	0.7			
Obsessive-compulsive disorder	0.1 (0.0–0.3)	1.1 (0.9–1.4)	<0.001	0.1 (0.01–0.2)	1.3 (0.9–1.8)	<0.001	0.2 (0.2–0.6)	0.9 (0.6–1.4)	0.1			
Phobias	2.8 (1.9–3.6)	2.0 (1.7–2.4)	0.1	2.7 (1.7–3.7)	2.7 (2.2–3.4)	0.9	2.9 (1.7–4.2)	1.3 (0.9–1.8)	0.002			
Panic disorder	0.5 (0.1–0.9)	1.1 (0.9–1.4)	0.03	0.6 (0.1–1.1)	1.3 (0.9–1.7)	0.1	0.2 (0.1–0.6)	1.0 (0.7–1.5)	0.03			

Weighted percentages to account for survey design.
doi:10.1371/journal.pone.0048012.t002




Figure 1. Comparisons of weighted prevalence estimates of Revised Clinical Interview Schedule symptom scores (≥2).
doi:10.1371/journal.pone.0048012.g001

Table 3. One-week prevalence and adjusted odds ratios for common mental disorder (Revised Clinical Interview Schedule score ≥12) by socio-demographic and socio-economic indicators.

		SELCoH		APMS England 2007	
		% (95%CI)	Adjusted odds ratio ^a (95%CI), p-value	% (95%CI)	Adjusted odds ratio ^a (95%CI), p-value
Gender	Female	27.3 (24.3–30.2)	1.7 (1.4–2.2), p<0.001	18.4 (17.0–19.8)	1.8 (1.5–2.1), p<0.001
	Male	17.9 (15.0–20.8)	1.0	11.6 (10.3–12.9)	1.0
Ethnic group	White British	24.3 (21.5–27.3)	1.0	14.5 (13.5–15.5)	1.0
	Black-Caribbean	31.0 (22.3–39.8)	1.4 (0.9–2.1), NS	17.1 (9.9–27.9)	1.1 (0.5–2.2), NS
	Black-African	19.5 (13.8–25.1)	0.7 (0.5–1.1), NS	23.1 (16.1–31.9)	1.6 (1.0–2.6), p=0.04
	Asian	24.9 (15.9–33.9)	1.0 (0.6–1.7), NS	13.6 (9.8–18.6)	0.9 (0.6–1.3), NS
	Other	23.0 (16.8–29.3)	0.9 (0.6–1.3), NS	18.3 (14.8–22.4)	1.2 (0.9–1.6), NS
Age (years)	16–24	25.1 (20.2–30.0)	1.0	16.4 (13.4–19.9)	1.0
	25–39	21.9 (18.3–25.6)	0.8 (0.6–1.1), NS	16.9 (15.1–19.0)	1.0 (0.8–1.4), NS
	40–54	29.5 (24.8–34.1)	1.2 (0.9–1.7), NS	17.6 (15.8–19.5)	1.1 (0.8–1.4), NS
	55–64	25.4 (18.2–32.5)	1.0 (0.6–1.6), NS	13.2 (11.3–15.3)	0.8 (0.6–1.0), NS
	65+	18.3 (12.3–24.3)	0.7 (0.4–1.1), NS	9.5 (8.1–11.1)	0.5 (0.4–0.7), p<0.001
Marital status	Never married	26.5 (22.9–30.2)	1.0	18.2 (16.1–20.5)	1.0
	Married/cohabiting	19.8 (16.8–22.9)	0.7 (0.5–0.9), p=0.04	12.8 (11.8–13.9)	0.8 (0.7–1.0), p=0.04
	Divorced/separated	32.2 (25.1–39.4)	1.3 (0.9–2.0), NS	24.6 (21.7–27.9)	1.8 (1.4–2.3), p<0.001
	Widowed	27.1 (14.9–39.4)	1.1 (0.5–2.3), NS	14.8 (12.3–17.7)	1.2 (0.9–1.6), NS
Education levels	No qualifications	25.7 (19.8–31.6)	1.7 (1.1–2.5), p=0.01	17.4 (15.5–19.4)	2.0 (1.6–2.5), p<0.001
	Up to GCSE level	30.5 (25.2–35.8)	1.9 (1.4–2.6), p<0.001	16.3 (14.7–18.1)	1.5 (1.2–1.8), p<0.001
	Advanced level	25.6 (21.2–29.9)	1.4 (1.0–1.9), p=0.02	14.4 (12.1–17.0)	1.3 (1.0–1.6), NS
	Higher degree or above	19.2 (16.1–22.3)	1.0	11.3 (9.9–12.9)	1.0
Social class	Non-manual	19.9 (16.7–23.1)	1.0	13.6 (12.6–14.7)	1.0
	Manual	22.2 (16.4–28.0)	1.2 (0.8–1.8), NS	16.5 (15.0–18.2)	1.4 (1.2–1.6), p<0.001
Employment status	Paid employment	20.1 (17.3–22.9)	1.0	12.6 (11.5–13.8)	1.0
	Unemployed	35.4 (27.9–43.0)	2.2 (1.5–3.2), p<0.001	27.3 (19.4–37.1)	2.4 (1.5–3.7), p<0.001
	Economically inactive	26.9 (23.1–30.6)	1.5 (1.1–1.9), p=0.003	18.2 (16.6–19.8)	1.9 (1.6–2.2), p<0.001
Housing tenure	Own/mortgage	18.9 (15.1–22.7)	1.0	11.8 (10.8–12.9)	1.0
	Rented	27.9 (24.8–30.9)	1.6 (1.2–2.1), p=0.002	22.5 (20.5–24.3)	2.0 (1.7–2.4), p<0.001
	Rent free	14.2 (7.4–21.1)	0.6 (0.4–1.2), NS	16.9 (10.3–26.5)	1.5 (0.8–2.6), NS

NS=non-significant; Weighted percentages to account for survey design;
^a. Model adjusted for age and gender.
 doi:10.1371/journal.pone.0048012.t003

Table 4. Comparisons of prevalence estimates for substance use by gender between SELCoH and APMS England 2007.

	All		P value	Women		P value	Men		P value
	SELCoH	APMS 2007		SELCoH	APMS 2007		SELCoH	APMS 2007	
	% (95%CI)	% (95%CI)		% (95%CI)	% (95%CI)		% (95%CI)	% (95%CI)	
Alcohol Use									
Score 0	24.5 (22.0–27.0)	17.3 (16.2–18.4)		27.4 (24.3–30.5)	22.0 (20.6–23.4)		18.6 (15.4–21.8)	12.3 (10.9–13.8)	
Score 1–7	58.7 (56.0–61.4)	58.6 (57.3–59.9)		60.4 (57.1–63.6)	62.4 (60.6–64.1)		55.4 (51.5–59.3)	54.6 (52.6–56.6)	
Score 8–15	13.9 (12.1–15.6)	20.4 (19.3–21.6)		10.4 (8.5–12.4)	13.8 (12.6–15.0)		20.7 (17.7–23.8)	27.4 (25.6–29.2)	
Score 16–40	2.9 (2.1–3.7)	3.8 (3.3–4.4)		1.8 (1.0–2.6)	1.9 (1.5–2.5)		5.2 (3.6–6.9)	5.8 (4.8–6.8)	
Hazardous alcohol use (Score 8+)	17.5 (15.5–19.5)	24.2 (22.9–25.4)	<0.001	12.9 (10.8–15.1)	15.7 (14.4–16.9)	=0.04	26.7 (23.2–30.1)	33.2 (31.3–35.1)	=0.002
Past Year Drug Use									
Cannabis	15.2 (15.9–20.1)	7.6 (6.7–8.4)		12.1 (9.9–14.3)	5.0 (4.2–6.0)		21.4 (18.4–24.5)	10.1 (8.8–11.5)	
Amphetamine	1.4 (0.9–1.9)	0.7 (0.4–1.0)		0.8 (0.3–1.4)	0.4 (0.2–0.7)		2.5 (1.4–3.5)	1.1 (0.7–1.6)	
Cocaine	6.3 (5.1–7.5)	2.5 (2.1–2.9)		4.2 (2.9–5.5)	1.4 (1.0–1.9)		10.4 (8.2–12.7)	3.6 (2.9–4.5)	
Crack	0.2 (0.0–0.4)	0.2 (0.1–0.4)		0.1 (0.1–0.2)	–		0.4 (0.0–0.8)	0.4 (0.2–0.8)	
Ecstasy	3.8 (2.8–4.8)	1.2 (0.9–1.5)		2.6 (1.5–3.6)	0.6 (0.4–0.8)		6.3 (4.6–7.9)	1.9 (1.4–2.5)	
LSD	0.7 (0.3–1.0)	0.2 (0.1–0.3)		0.5 (0.1–0.9)	0.0 (0.0–0.2)		0.9 (0.3–1.6)	0.4 (0.2–0.7)	
Tranquilliser	1.7 (1.0–2.4)	0.7 (0.5–0.9)		1.9 (1.0–2.8)	0.5 (0.5–0.8)		1.2 (0.5–1.9)	0.9 (0.6–1.3)	
Heroin	0.1 (0.0–0.3)	0.2 (0.1–0.3)		0.1 (0.1–0.2)	–		0.3 (0.1–0.6)	0.3 (0.2–0.6)	
Any drugs (≥1 drugs)	18.1 (16.1–20.2)	8.9 (8.1–9.8)	<0.001	14.3 (11.9–16.6)	6.3 (5.5–7.3)	<0.001	25.8 (22.5–29.2)	11.6 (10.3–13.1)	<0.001
Poly drug use (≥2 drugs)	6.3 (5.1–7.5)	2.4 (2.0–2.8)	<0.001	4.5 (3.2–5.9)	1.1 (0.8–1.5)	<0.001	9.7 (7.6–1.9)	3.4 (3.0–4.6)	<0.001

Weighted percentages to account for survey design.
doi:10.1371/journal.pone.0048012.t004

that in the national sample; there was a four-fold greater proportion of depressive episodes in the SELCoH sample; the national sample had a higher prevalence of hazardous alcohol use; and SELCoH participants had a higher prevalence of illicit drug use (individual drugs and poly drug use) in the past year. However, in both samples, men reported more hazardous alcohol and drug use than women, and cannabis was the most commonly reported drug followed by cocaine. Comparisons made in combined datasets showed that these differences between the SELCoH sample and the APMS England sample persisted after adjusting for socio-demographic and socioeconomic indicators for all outcomes. Further, similar differences were also present in comparisons between the SELCoH sample and the APMS subsample from the Greater London area in the fully adjusted models.

Strengths and limitations

Among the strengths of this study are the similarities between the survey designs, the administration of the same validated structured clinical interview, and the representativeness of the APMS 2007 and the SELCoH samples. Specifically, the SELCoH sample was shown to be representative on most demographic and socioeconomic characteristics of the population in the study catchment area, the London boroughs of Southwark and Lambeth, according to the UK 2001 Census [16]. In terms of study limitations, non-response rates at the individual level in the national study and at the household level in the SELCoH sample may have resulted in participation bias, and the prevalence estimates should be considered with caution. A study linking a

Norwegian household survey with disability pension registry data illustrated that mental and substance use disorders were strongly associated with non-participation, and conventional surveys may under-estimate prevalence of mental disorder [31]. Further, different weighting procedures were utilised and it is likely that there is residual confounding in the analysis examining differences between the two samples. In terms of the timing of the studies, APMS data were collected in the year leading up to the UK economic recession (2006–2007), while SELCoH data were collected after the recession period began (2008–2010). Common mental disorders, increased substance use and substance disorder are among the mental health problems more likely to increase during an economic downturn, primarily as a result of an increase in the poor socioeconomic conditions included in this analysis [32–34]. Finally, we acknowledge concerns about the validity of measures, such as the CIS-R [35]. However, as with the national study, we enlisted experienced and trained lay interviewers to administer the CIS-R. Given these strengths and limitations, it is worthwhile considering the extent to which differences in prevalence estimates may be real, or have resulted from the design and conduct of the two studies. Low participation rates may be a factor in any psychiatric epidemiological survey, and the lower participation rate in the SELCoH sample compared with the APMS sample may have led to an under-estimate of differences in prevalence estimates [31]. We used appropriate statistical methods to apply weights for each sample and to control for the effect of clustering by household within the SELCoH study in all analyses, so believe it unlikely that this contributed to the observed differences.

Table 5. Prevalence estimates and adjusted odds ratios for hazardous alcohol use by socio-demographic and socio-economic indicators.

		SELCoH		APMS England 2007	
		% (95%CI)	Adjusted odds ratio ^a (95%CI), p-value	% (95%CI)	Adjusted odds ratio ^a (95%CI), p-value
Gender	Female	11.3 (9.2–13.3)	0.4 (0.3–0.5), p<0.001	15.7 (14.4–16.9)	0.4 (0.3–0.4), p<0.001
	Male	24.6 (21.2–27.9)	1.0	33.2 (31.3–35.1)	1.0
Ethnic group	White British	20.5 (17.9–23.2)	1.0	25.5 (24.2–26.8)	1.0
	Black-Caribbean	3.8 (0.8–6.8)	0.1 (0.1–0.3), p<0.001	17.5 (8.6–26.4)	0.6 (0.3–1.1), NS
	Black-African	3.4 (1.2–5.7)	0.1 (0.0–0.2), p<0.001	8.2 (0.7–15.7)	0.2 (0.1–0.5), p=0.001
	Asian	16.3 (6.9–25.6)	0.6 (0.3–1.3), NS	8.7 (5.1–12.4)	0.2 (0.1–0.3), p<0.001
	Other	11.8 (7.2–16.4)	0.4 (0.2–0.6), p<0.001	23.9 (19.0–28.8)	0.7 (0.6–1.0), NS
Age (years)	16–24	22.7 (17.8–27.7)	1.0	33.2 (32.6–41.9)	1.0
	25–39	22.5 (18.8–26.2)	1.0 (0.7–1.5), NS	29.3 (26.9–31.7)	0.7 (0.5–0.9), p=0.004
	40–54	13.4 (10.1–16.7)	0.5 (0.4–0.8), p=0.003	23.2 (21.1–25.3)	0.5 (0.4–0.6), p<0.001
	55–64	10.2 (5.6–14.8)	0.4 (0.2–0.7), p<0.001	19.3 (16.9–21.6)	0.4 (0.3–0.5), p<0.001
	65+	3.3 (0.8–5.7)	0.1 (0.0–0.3), p<0.001	12.4 (10.8–14.0)	0.2 (0.2–0.3), p<0.001
Marital status	Never married	21.4 (17.9–24.9)	1.0	36.7 (33.5–39.8)	1.0
	Married/cohabiting	12.9 (10.5–15.3)	0.9 (0.6–1.2), NS	21.3 (20.0–22.7)	0.7 (0.6–0.8), p<0.001
	Divorced/separated	14.1 (9.2–19.0)	1.5 (0.9–2.4), NS	24.2 (21.1–27.3)	1.0 (0.8–1.3), NS
	Widowed	4.8 (0.7–10.2)	1.0 (0.3–3.5), NS	8.6 (6.7–10.5)	0.5 (0.4–0.8), p<0.001
Education levels	No qualifications	9.9 (5.9–13.9)	0.7 (0.4–1.2), NS	17.4 (15.5–19.4)	0.9 (0.7–1.1), NS
	Up to GCSE level	9.3 (6.4–12.2)	0.3 (0.2–0.5), p<0.001	24.1 (21.9–26.2)	0.9 (0.7–1.0), NS
	Advanced level	15.6 (11.9–19.3)	0.5 (0.4–0.8), p<0.001	32.9 (29.4–36.5)	1.2 (0.9–1.4), NS
	Higher degree or above	21.7 (18.5–24.9)	1.0	26.0 (23.9–28.2)	1.0
Social class	Non-manual	19.5 (16.5–22.6)	1.0	22.9 (21.4–24.3)	1.0
	Manual	9.8 (5.9–13.6)	0.5 (0.3–0.7), p=0.002	26.8 (24.8–28.8)	1.1 (0.9–1.2), NS
Employment status	Paid employment	17.0 (14.5–19.6)	1.0	28.8 (27.1–30.4)	1.0
	Unemployed	20.6 (14.2–27.0)	1.2 (0.7–1.8), NS	33.4 (24.8–41.9)	0.9 (0.6–1.4), NS
	Economically inactive	12.9 (9.9–15.9)	0.9 (0.6–1.2), NS	15.8 (14.3–17.3)	0.7 (0.6–0.9), p<0.001
Housing tenure	Own/mortgage	13.9 (10.8–16.9)	1.0	23.3 (21.9–24.8)	1.0
	Rented	16.4 (13.9–18.9)	0.9 (0.7–1.4), NS	25.8 (23.6–28.3)	0.9 (0.8–1.1), NS
	Rent free	18.3 (10.1–26.5)	0.7 (0.4–1.4), NS	32.6 (22.6–44.4)	1.3 (0.7–2.2), NS

NS = non-significant; Weighted percentages to account for survey design;

^a. Model adjusted for age and gender.

doi:10.1371/journal.pone.0048012.t005

In theory these differences might have been a valid consequence of differences between the survey populations with respect to demographic profiles. The SELCoH sample was a somewhat younger and more female sample, with a much higher proportion of people with Black African or Caribbean ethnicity, and included more unemployed people and more manual workers, at the same time as having a better qualified population. Despite these differences in the demographic profile, there was a similar pattern of associations for the socio-demographic and socioeconomic indicators and the outcomes across the SELCoH and APMS samples. We argue that such differences in populations should not be seen as nuisance variables (i.e. confounders) but should be seen as plausible explanations for the differences observed. However, adjusting for these variables in analyses of combined datasets had no impact on the differences between the samples in frequency of disorders at the national and more local level. Thus, the differences are likely to be due to potential confounders not identified in this study. Further, it is notable that within virtually

each stratum of the analyses presented, the prevalence of CMD and illicit drug use was higher and the prevalence of hazardous alcohol use was lower in the SELCoH sample than in the APMS sample.

Comparisons with previous studies and implications

There were two main findings that deserve further discussion. First, to our knowledge there are no studies other than SELCoH that have compared UK inner city populations with national data in terms of prevalence of CMD or substance use. With the exception of more hazardous alcohol use in the national study, our findings support our hypotheses. Further, our findings are consistent with those from other national studies, such as the differences in CMD by urbanicity in the 1993 and 2000 British National Survey of Psychiatric Morbidity Surveys [11–12] and the graded increase in the prevalence of CMD and substance use disorders across multiple categories of urbanization in the Netherlands Mental Health Survey and Incidence Study [9].

Table 6. Comparisons of prevalence estimates and adjusted odds ratios for illicit drug use by socio-demographic and socio-economic indicators.

		SELCoH		APMS England 2007	
		% (95%CI)	Adjusted odds ratio ^a (95%CI), p-value	% (95%CI)	Adjusted odds ratio ^a (95%CI), p-value
Sample	SELCoH				
	APMS 2007				
Gender	Female	14.3 (11.9–16.6)	0.5 (0.4–0.6), p<0.001	6.3 (5.5–7.3)	0.5 (0.4–0.6), p<0.001
	Male	25.8 (22.5–29.2)	1.0	11.6 (10.3–13.1)	1.0
Ethnic group	White British	24.3 (21.5–27.3)	1.0	8.5 (7.6–9.3)	1.0
	Black-Caribbean	31.0 (22.3–39.8)	0.9 (0.5–1.5), NS	19.0 (10.1–27.9)	2.7 (1.4–5.2), p=0.003
	Black-African	19.5 (13.8–25.1)	0.1 (0.1–0.2), p<0.001	8.9 (1.0–16.8)	0.6 (0.2–1.6), NS
	Asian	24.9 (15.9–33.9)	0.2 (0.1–0.4), p<0.001	3.6 (0.9–6.3)	0.2 (0.1–0.5), p<0.001
	Other	23.0 (16.8–29.3)	0.6 (0.4–0.9), p=0.02	16.0 (11.6–20.4)	1.5 (1.0–2.1), p=0.03
Age (years)	16–24	32.5 (26.7–38.3)	1.0	23.7 (19.8–27.5)	1.0
	25–39	25.2 (21.3–29.1)	0.7 (0.5–0.9), p=0.05	14.9 (13.0–16.9)	0.6 (0.4–0.7), p<0.001
	40–54	14.6 (11.1–18.1)	0.4 (0.2–0.5), p<0.001	4.5 (3.5–5.5)	0.1 (0.1–0.2), p<0.001
	55–64	8.9 (4.6–13.3)	0.2 (0.1–0.4), p<0.001	2.1 (1.2–2.9)	0.1 (0.04–0.1), p<0.001
	65+	1.8 (0.3–3.9)	0.04 (0.0–0.1), p<0.001	0.8 (0.4–1.3)	0.03(0.02–0.05), p<0.001
Marital status	Never married	30.5 (26.5–34.5)	1.0	21.8 (19.2–24.4)	1.0
	Married/cohabiting	11.4 (9.0–13.8)	0.5 (0.4–0.7), p<0.001	4.9 (4.1–5.6)	0.5 (0.4–0.7), p<0.001
	Divorced/separated	14.8 (9.8–19.8)	1.1 (0.6–1.7), NS	10.0 (7.5–12.5)	1.8 (1.2–2.6), p=0.002
	Widowed	2.0 (0.8–4.9)	0.3 (0.1–1.4), NS	1.5 (0.5–2.5)	1.1 (0.5–2.3), NS
Education levels	No qualifications	9.7 (6.1–13.3)	1.0 (0.6–1.6), NS	5.3 (4.0–6.6)	1.2 (0.8–1.7), NS
	Up to GCSE level	19.3 (14.8–23.7)	0.9 (0.7–1.4), NS	10.2 (8.5–11.9)	0.9 (0.7–1.3), NS
	Advanced level	21.6 (17.2–25.9)	0.9 (0.6–1.3), NS	13.7 (10.9–16.4)	1.1 (0.8–1.6), NS
	Higher degree or above	19.4 (16.2–22.5)	1.0	8.5 (6.8–10.2)	1.0
Social class	Non-manual	22.3 (18.9–25.7)	1.0	7.4 (6.4–8.4)	1.0
	Manual	16.5 (11.7–21.3)	0.8 (0.5–1.1), NS	10.5 (9.1–11.9)	1.3 (1.1–1.7), p=0.02
Employment status	Paid employment	21.0 (18.1–23.9)	1.0	10.0 (8.9–11.2)	1.0
	Unemployed	26.2 (19.5–33.0)	1.2 (0.8–1.8), NS	25.1 (17.0–33.3)	1.7 (1.1–2.8), p=0.03
	Economically inactive	12.3 (9.51–15.1)	0.6 (0.4–0.8), p=0.001	5.7 (4.6–6.8)	1.1 (0.8–1.4), NS
Housing tenure	Own/mortgage	12.4 (9.4–15.4)	1.0	6.3 (5.5–7.2)	1.0
	Rented	20.1 (17.4–22.8)	1.4 (0.9–1.9), NS	15.3 (13.5–17.2)	1.8 (1.4–2.2), p<0.001
	Rent free	30.4 (20.8–39.9)	1.3 (0.8–2.3), NS	9.7 (4.7–18.8)	0.8 (0.3–1.9), NS

NS = non-significant; Weighted percentages to account for survey design;

^a. Model adjusted for age and gender.

doi:10.1371/journal.pone.0048012.t006

Second, the findings on illicit drug use indicated that a focus on individual drugs, such as cannabis use and cocaine, are necessary, but poly drug use in the local community should also be of concern. In particular, simultaneous drug and alcohol use is associated with poor social outcomes and mental ill health in a US national household survey [28]. More broadly, explanations for the differences in substance use between samples are likely to be rooted in the social environment. An international review of the social epidemiology of substance use suggested that type and density of social networks and neighbourhood-level indicators were associated with increased substance use, whereas findings on the relationship between socioeconomic status (SES) and substance use were mixed [36]. The latter is highlighted in further national and local comparisons; high SES groups were more likely to report hazardous alcohol use in the SELCoH sample, whereas there were

no differences by SES in the APMS or the British National Survey of Psychiatric Morbidity 2000 [37]. Further investigations of social and environmental factors that influence the population distribution of CMD and substance use are needed to better understand these differences between and within national and local community samples.

The findings of this study highlight the importance of considering differences between local and national public mental health that are needed to inform service provision and policy, specifically regarding the distribution of resources. While there are similarities in the social inequalities present at national and local levels that greatly contribute to the generation of mental health inequalities, differences in the prevalence of CMD and drug use justify the need to consider specific public mental health needs of those residing in urban environments. The findings resulting from

Table 7. Comparisons of adjusted odds ratios for common mental disorder, hazardous alcohol use and illicit drug use by socio-demographic and socio-economic indicators in combined data from both studies.

	Sample	Fully adjusted odds ratio ^a (95%CI), p-value	
Common Mental Disorder	SELCoH and APMS England 2007	SELCoH	1.4 (1.2–1.7), p<0.001
		APMS 2007	1.0
	SELCoH and APMS London 2007	SELCoH	1.7 (1.3–2.2), p<0.001
		APMS 2007	1.0
Hazardous Alcohol Use	SELCoH and APMS England 2007	SELCoH	0.8 (0.6–0.9), p = 0.001
		APMS 2007	1.0
	SELCoH and APMS London 2007	SELCoH	0.6 (0.4–0.8), p<0.001
		APMS 2007	1.0
Illicit Drug Use	SELCoH and APMS England 2007	SELCoH	2.1 (1.7–2.6), p<0.001
		APMS 2007	1.0
	SELCoH and APMS London 2007	SELCoH	1.6 (1.1–2.1), p = 0.008
		APMS 2007	1.0

NS = non-significant.

Weighted percentages to account for survey design.

^a. Fully adjusted model with combined comparable data from both studies; age entered as a continuous variable in fully adjusted model.

doi:10.1371/journal.pone.0048012.t007

comparisons between the SELCoH sample and the APMS London sample further highlight the need to drill down from the metropolitan area to the local area level. The growing number of individuals moving into urban environments, the impact of social relationships [36,38] and the urban context on health [39–40] suggests that there is a growing need for locally focused public mental health studies. A practical solution to this problem may be as simple as a joint investment by local and national public health and service provision authorities to collect comparable data at the local level in combination with monitoring inequalities by using secondary data sources such as electronic health records and case registers.

Supporting Information

Table S1 Comparisons of adjusted odds ratios for illicit drug use by socio-demographic and socio-economic indicators combined data from both studies (full models).
(DOCX)

Acknowledgments

The authors have the following disclaimers: SLH, SF, MH, MV, BK, JLC, BG, RM, RS, CM, AR receive salary support from the National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

References

- Jenkins R (2003) Making psychiatric epidemiology useful: the contribution of epidemiology to government policy. *Int Rev Psychiatry* 15: 188–200.
- Reijneveld SA, Schene AH (1998) Higher prevalence of mental disorders in socioeconomically deprived urban areas in the Netherlands: community or personal advantage? *J Epidemiol Community Health* 52: 2–7.
- Galea S, Ahern J, Nandi A, Tracy M, Beard J, et al. (2007) Urban neighbourhood poverty and the incidence of depression in a population-based cohort study. *Ann Epidemiol* 17:171–179.
- Galea S, Ahern J, Rudenstine S, Wallace Z, Vlahov D (2005) Urban built environment and depression: a multilevel analysis. *J Epidemiol Community Health* 59: 822–827.
- Galea S, Vlahov D (2005) Urban health: evidence, challenges and directions. *Annu Rev Public Health* 26:341–365.
- Weich S, Holt G, Twigg L, Jones K, Lewis G (2003) Geographical variation in the prevalence of common mental disorders in Britain: a multilevel investigation. *Am J Epidemiol* 157: 730–737.

SELCoH Study Team: Matthew Hotopf, Stephani L Hatch, Souci Frissa, Maria Verdecchia, Robert Stewart, Nicola T Fear, Abraham Reichenberg, Craig Morgan, Bwalya Kankulu, Jennifer Clark, Billy Gizard, Robert Medcalf, Alistair Baile, Elizabeth Doherty, Deborah Bekele, Joshua Buckman, Grant Henderson, Max Henderson and Benedict Weobong

Author Contributions

Analyzed the data: SLH CW MH. Contributed reagents/materials/analysis tools: SLH MH RS NTF CM AR PB SM TB BK JC BG RM. Wrote the paper: SLH MH CW PB. Designed the SELCoH study: MH. Sought and obtained funding: MH. Oversaw data collection and analysis, conceptualized and commented on all drafts of the paper: MH. Oversaw SELCoH data collection: SH. Conceptualised and conducted the analysis: SLH. Wrote all manuscript drafts: SLH. Had final responsibility for the decision to submit for publication: SLH. Conducted APMS analysis: CW. Conceptualized and commented on paper drafts: CW. Coordinated the SELCoH data collection and commented on all drafts of the paper: SF MV. All contributed to the design of the SELCoH study survey: RS NTF CM AR SF MV. Research assistants on the SELCoH team collected the data and commented on drafts of the paper: BK JC BG RM. Led the APMS data collection: SM, PB TB. Commented on drafts of the paper: RS NTF CM AR PB SM TB. Had access to study data (local data are available upon request and national data are publicly available): SLH CW SF NTF MV RS AR CM PB SM TB BK JLC BG RM. Full responsibility for the integrity of the data and the accuracy of the data analyses: SLH CW SF NTF MV RS AR CM PB SM TB BK JLC BG RM. Approved the final version of the paper: SLH CW SF NTF MV RS AR CM PB SM TB BK JLC BG RM. The guarantor for the study: MH.

7. Lorant V, Croux C, Weich S, Deliege D, Mackenbach J, et al. (2007) Depression and socio-economic risk factors: 7-year longitudinal population study. *Br J Psychiatry* 190:293–298.
8. Marmot M, Allen J, Goldblatt P, Boyce T, McNeish D, et al. (2010) Fair Society, Healthy Lives; 2010, available: <http://www.ucl.ac.uk/marmotreview>.
9. Peen J, Dekker J, Schoevers RA, ten Have M, de Graaf R, et al. (2007) Is the prevalence of psychiatric disorders associated with urbanization? *Soc Psychiatr Epidemiol* 42:984–989.
10. Eaton WW, Martins SS, Nestadt G, Bienvenu OJ, Clarke D, et al. (2008) The burden of mental disorders. *Epidemiol Rev* 30: 1–14.
11. Paykel E, Abbott R, Jenkins R, Brugha TS, Meltzer H (2000) Urban-rural mental health differences in Great Britain: findings from the National Morbidity Survey. *Psychol Med* 30:269–280.
12. Singleton N, Bumpstead R, O'Brien M, Lee A, Meltzer H (2003) Psychiatric morbidity among adults living in private households, 2000 (Reprinted from Psychiatric morbidity among adults living in private households, 2000: Summary report, 2001). *Int Rev Psychiatry* 15: 65–73.
13. Kirkbride JB, Fearon P, Morgan C, Dazzan P, Morgan K, et al. (2007) Neighbourhood variation in the incidence of psychotic disorders in Southeast London. *Soc Psychiatry Psychiatr Epidemiol* 42: 438–445.
14. Rezaeian M, Dunn G, St Leger S, Appleby L (2004) The production and interpretation of disease maps: A methodological case-study. *Soc Psychiatry Psychiatr Epidemiol* 39: 947–954.
15. Glover G, Arts G, Wooff D (2004) A needs index for mental health care in England based on updatable data. *Soc Psychiatry Psychiatr Epidemiol* 39: 730–738.
16. Hatch SL, Frissa S, Verdecchia M, Stewart R, Fear NT, et al. (2011) Identifying socio-demographic and socio-economic determinants of health inequalities in a diverse London community: the SELCoH study. *BMC Public Health* 11:861 doi:10.1186/1471-2458-11-861.
17. Department of Health website. Health profile 2011: Southwark <http://www.apho.org.uk/resource/view.aspx?RID=50215&SEARCH=Southwark&SPEAR>. Accessed 2012 May 1.
18. Department of Health website. Health profile 2011: Lambeth. Available: <http://www.apho.org.uk/resource/view.aspx?RID=50215&SEARCH=Lambeth&SPEAR>. Accessed: 2012 May 1.
19. Office of National Statistics website. NOMIS Labour market profile: Lambeth. Available: <http://www.nomisweb.co.uk/reports/Imp/la/2038431867/report.aspx?town=Lambeth>. Accessed: 2012 May 1.
20. Office of National Statistics website. NOMIS Labour market profile: Southwark. Available: <http://www.nomisweb.co.uk/reports/Imp/la/2038431870/report.aspx?town=Southwark>. Accessed: 2012 May 1.
21. Office of National Statistics website. Available: <http://www.ons.gov.uk/ons/taxonomy/index.html?nscl=Population>. Accessed: 2012 May 1.
22. McManus S, Meltzer H, Brugha T, Bebbington P, Jenkins R (2009) Adult Psychiatric Morbidity in England, 2007: Results of a Household Survey. National Health Service Information Centre for Health and Social Care.
23. Jenkins R, Bebbington P, Brugha T, Farrell M, Gill B, et al. (1997) The National Psychiatric Morbidity surveys of Great Britain-strategy and methods. *Psychol Med* 27: 765–774.
24. Lewis G, Pelosi AJ, Araya R, Dunn G (1992) Measuring psychiatric disorder in the community: a standardized assessment for use by lay interviewers. *Psychol Med* 22: 465–486.
25. Bebbington P, Meltzer H, Brugha T, Farrell M, Jenkins R, et al. (2000) Unequal access and unmet need: neurotic disorders and the use of primary care services. *Psychol Med* 30: 1359–1368.
26. Bebbington P, Brugha T, Meltzer H, Jenkins R, Ceresa C, et al. (2000) Neurotic disorders and the receipt of psychiatric treatment. *Psychol Med* 30: 1369–1376.
27. Babor TR, Higgins-Biddle JC, Saunders JB, Monteiro MG (2001) The Alcohol Use Disorders Identification Test: Guidelines for Use in Primary Care (Second ed.). Geneva: World Health Organization.
28. Midanik LT, Tam TW, Weisner C (2007) Concurrent and simultaneous drug and alcohol use: results of the 2000 national alcohol survey. *Drug Alcohol Depend* 90:72–80.
29. StataCorp (2009) Stata Statistical Software: Release 11. College Station, TX: StataCorp LP.
30. Bebbington PE, Hurry J, Tennant C (1981) Psychiatric disorders in selected immigrant groups in Camberwell. *Soc Psychiatry* 16: 43–51.
31. Knudsen AK, Hotopf M, Skogen JC, Overland S, Mykletun A (2010) The health status of nonparticipants in a population-based health study: the Hordaland health study. *Am J Epidemiol* 172:1306–1314.
32. World Health Organization website (2011) Impact of economic crises on mental health. Regional Office for Europe of the World Health Organization. Available: <http://www.euro.who.int/en/what-we-do/health-topics/noncommunicable-diseases/mental-health/publications/2011/impact-of-economic-crises-on-mental-health>. Accessed: 2012 Oct 1.
33. Stuckler D, Basu S, McDaid D (2011) Depression amidst depression: Mental health effects of the ongoing recession. World Health Organization website. Available: http://groups.stakes.fi/NR/rdonlyres/85320FDB-20EF-4673-A974-28681A4D7E01/0/Background_1_EconomicCrises_and_Mental_Health.pdf. Accessed 2012 Oct 1.
34. Uutela A (2010) Economic crisis and mental health. *Curr Opin Psychiatry* 23:127–130.
35. Brugha TS, Bebbington PE, Jenkins R (1999) A difference that matters: comparisons of structured and semi-structured psychiatric diagnostic interviews in the general population. *Psychol Med* 29: 1013–20.
36. Galea S, Nandi A, Vlahov D (2004) The social epidemiology of substance use. *Epidemiol Rev* 26:36–52.
37. Coulthard M, Farrell M, Singleton N, Meltzer H (2002) Tobacco, alcohol and drug use and mental health: London: TSO.
38. Das-Munshi J, Becares L, Dewey ME, Stansfeld SA, Prince MJ (2010) Understanding the effect of ethnic density on mental health: multi-level investigation of survey data from England. *BMJ* 341:c5367.
39. Dye C (2008) Health and urban living. *Science* 319:766–769.
40. Whitley R, Prince M, Cargo M (2005) Thinking inside the bubble: evidence for a new contextual unit in urban mental health. *J Epidemiol Community Health* 59:893–897.