



City Research Online

City, University of London Institutional Repository

Citation: Fring, A. ORCID: 0000-0002-7896-7161 and Taira, T. (2020). 't Hooft-Polyakov monopoles in non-Hermitian quantum field theory. Physics Letters B, 807, 135583.. doi: 10.1016/j.physletb.2020.135583

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/24440/>

Link to published version: <http://dx.doi.org/10.1016/j.physletb.2020.135583>

Copyright and reuse: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk



't Hooft-Polyakov monopoles in non-Hermitian quantum field theory

Andreas Fring*, Takanobu Taira

Department of Mathematics, City, University of London, Northampton Square, London EC1V 0HB, UK

ARTICLE INFO

Article history:

Received 9 June 2020

Received in revised form 18 June 2020

Accepted 20 June 2020

Available online 26 June 2020

Editor: N. Lambert

ABSTRACT

We construct exact 't Hooft-Polyakov monopole solutions in a non-Hermitian field theory with local non-Abelian $SU(2)$ gauge symmetry and a modified antilinear CPT symmetry. The solutions are obtained in a fourfold Bogomolny-Prasad-Sommerfield scaling limit giving rise to two different types of monopole masses that saturate the lower energy bound. These two masses only coincide in the Hermitian limit and in the limit in which the symmetry breaking vacuum tends to the trivial symmetry preserving vacuum. In the two theories corresponding to the two known Dyson maps these two masses are exchanged, unlike the Higgs and the gauge masses, which remain the same in both theories. We identify three separate regions in parameter space bounded by different types of exceptional points. In the first region the monopole masses are finite and tend both to zero at the boundary exceptional point, in the second the monopole masses become complex and in the third only one of the monopole masses becomes zero at the boundary exceptional point, whereas the other tends to infinity. We find a self-dual point in parameter space at which the gauge mass becomes exactly identical to the monopole mass.

© 2020 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

Ever since Dirac demonstrated 89 years ago [1] that the postulate of the existence of magnetic monopoles provides an explanation for the quantization of the electric charge, they have remained an appealing theoretical concept, despite the fact that up to now magnetic monopoles have not been observed in nature. See [2,3] for a very recent update on the experimental searches of magnetic monopoles in particle colliders and of cosmic origin that have been carried out, are currently performed and on future plans.

The concept of monopoles became an integral part of particle physics, notably of any theory aiming at the formulation of grand unification of all fundamental forces, after 't Hooft [4] and Polyakov [5] noticed that gauge theories almost inevitable contain monopole solutions. Subsequently the mechanism for the emergence of monopoles, together with their properties and relations to the internal structures of particular gauge theories have been extensively studied and many aspects are very well understood, see e.g. [6,7]. Here we are especially interested in the relation of their masses to the gauge particles and whether the intriguing features found in Hermitian theories also hold in non-Hermitian versions. Especially if the property found by Montonen and Olive [8] is still valid, that in non-Abelian gauge theories the soliton solutions become equivalent to gauge massive fields in a dual theory.

Thus our aim is to study the properties of the monopole solutions in a non-Hermitian field theory with local non-Abelian gauge symmetry and a modified antilinear CPT symmetry. We build on our previous investigations [9–11], and further elaborate on a particular model studied in [11] for which the Higgs masses have been identified in all PT -regimes. Variants of this model have also been investigated with different types of methods in [12–17].

We construct the monopole solutions in a fourfold Bogomolny-Prasad-Sommerfield (BPS) [18,19] scaling limit that has also been successful in the Hermitian setting. We shall investigate the properties of these solutions and in particular establish whether the Montonen-Olive strong-weak duality still holds.

Our manuscript is organised as follows: In section 2 we recall the non-Hermitian field theory previously studied in [11] and set up the equations of motion whose asymptotic solutions tend to the vacuum solutions. In sections 3 we discuss the Bogomolny energy bound for our theory. In section 4 we carry out the BPS-limit in form of a fourfold scaling limit obtaining two different types of masses from

* Corresponding author.

E-mail addresses: a.fring@city.ac.uk (A. Fring), takanobu.taira@city.ac.uk (T. Taira).

the monopole solutions. In section 5 we discuss the properties of these masses and identify physical regions in the parameter space. Our conclusions are stated in section 6.

2. Soliton solution in a non-Hermitian model with $SU(2)$ gauge symmetry

Here we consider the non-Hermitian $SU(2)$ gauge theory with matter fields in the adjoint representation. This is a non-Hermitian extension of the Lagrangian studied in [4,20] which is known to possess monopole solutions with finite energies

$$S = \int d^4x \frac{1}{2} \text{Tr}(D\phi_1)^2 + \frac{1}{2} \text{Tr}(D\phi_2)^2 - c_1 \frac{m_1^2}{2} \text{Tr}(\phi_1^2) + c_2 \frac{m_2^2}{2} \text{Tr}(\phi_2^2) - i\mu^2 \text{Tr}(\phi_1\phi_2) - \frac{g}{4} \left(\text{Tr}(\phi_1^2) \right)^2 - \frac{1}{4} \text{Tr}(F^2). \quad (2.1)$$

Here we take $g, \mu \in \mathbb{R}$, $m_i \in \mathbb{R}$ and discrete values $c_i \in \{-1, 1\}$. The two fields $\{\phi_i\}_{i=1,2}$ are Hermitian matrices $\phi_i(t, \vec{x}) \equiv \phi_i^a(t, \vec{x}) T^a$ where $\phi_i^a(t, \vec{x})$ is a real-valued field. The three generators $\{T^a\}_{a=1,2,3}$ of $SU(2)$ in the adjoint representation are defined by three Hermitian matrices of the form $(T^a)_{bc} = -i\epsilon_{abc}$, satisfying the commutation relation $[T^a, T^b] = i\epsilon^{abc} T^c$. One can check that $\text{Tr}(T^a T^b) = 2\delta^{ab}$. The field strength tensor is defined as $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - ie[A_\mu, A_\nu]$ where the gauge fields are $A_\mu = A_\mu^a T^a$.

This action is invariant under the following local $SU(2)$ transformation of the matter fields and gauge fields

$$\phi_i \rightarrow e^{i\alpha^a(x) T^a} \phi_i e^{-i\alpha^a(x) T^a}, \quad (2.2)$$

$$A_\mu \rightarrow e^{i\alpha^a(x) T^a} A_\mu e^{-i\alpha^a(x) T^a} + \frac{1}{e} \partial_\mu \alpha^a(x) T^a. \quad (2.3)$$

This action is also symmetric under modified $\mathcal{CP}\mathcal{T}$ symmetry where one of the matter fields ϕ_1 transforms as scalar field and the matter field ϕ_2 transforms as pseudo-scalar. This antilinear symmetry is an important indicator for the reality of the classical masses of the matter fields as extensively discussed in [9,17]

$$\mathcal{CP}\mathcal{T} : \phi_1(t, \vec{x}) \rightarrow \phi_1(-t, -\vec{x}), \quad \phi_2(t, \vec{x}) \rightarrow -\phi_2(-t, -\vec{x}), \quad i \rightarrow -i. \quad (2.4)$$

The equations of motion for the fields ϕ_i and A_μ are

$$(D_\mu D^\mu \phi_i)^a + \frac{1}{2} \frac{\delta V}{\delta \phi_i^a} = 0, \quad D_\nu F_a^{\nu\mu} - e\epsilon_{abc} \phi_1^b (D^\mu \phi)^c + e\epsilon_{abc} \phi_2^b (D^\mu \phi)^c = 0. \quad (2.5)$$

One of the problems often associated with non-Hermitian field theories is the incompatibility of the set of equations of motion [12]. To overcome these issues we employed the Pseudo-Hermitian method which consists of mapping the non-Hermitian model to a Hermitian model through a similarity transformation. This is a common procedure used in the \mathcal{PT} symmetric quantum mechanics [21,22], and in close analogy also their field-theoretic versions were studied for several different models [23,17]. The similarity transformations for our model eq. (2.1) have already been introduced in [11]

$$\eta_\pm = \prod_{a=1}^3 \exp\left(\pm \frac{\pi}{2} \int d^3x \Pi_2^a \phi_2^a\right). \quad (2.6)$$

The adjoint action of η_\pm maps the complex action in eq. (2.1) into the following real action

$$\begin{aligned} \eta_\pm S \eta_\pm^{-1} &= \int d^4x \frac{1}{2} \text{Tr}(D\phi_1)^2 - \frac{1}{2} \text{Tr}(D\phi_2)^2 - c_1 \frac{m_1^2}{2} \text{Tr}(\phi_1^2) - c_2 \frac{m_2^2}{2} \text{Tr}(\phi_2^2) - c_3 \mu^2 \text{Tr}(\phi_1\phi_2) - \frac{g}{4} \left(\text{Tr}(\phi_1^2) \right)^2 - \frac{1}{4} \text{Tr}(F^2) \\ &\equiv \int d^4x \frac{1}{2} \text{Tr}(D\phi_1)^2 - \frac{1}{2} \text{Tr}(D\phi_2)^2 - V - \frac{1}{4} \text{Tr}(F^2), \end{aligned} \quad (2.7)$$

where the parameter c_3 takes the value ± 1 for η_\pm respectively. Notice that this model is very similar to those with the actions considered in [4,20], but with second order coupling term $\mu^2 \text{Tr}(\phi_1\phi_2)$ and negative sign in the kinetic term of ϕ_2 .

Next one can use the simple scaling argument [24] to show that monopole solutions with finite energy require the monopole to asymptotically approach the vacuum solution $V[\phi^0] = 0$ (note that one can add a constant to the action so that this asymptotic condition is equivalent to $\delta V[\phi^0] = 0$). The explicit values of the vacuum solutions ϕ_α^0 and A_μ^0 are found by solving $\delta V = 0$ and $D_\mu \phi_\alpha = 0$ [25]

$$\begin{aligned} (\phi_1^0)^a &= \pm R \hat{r}^a \equiv h_1^{0\pm} \hat{r}^a, \quad (\phi_2^0)^a = \mp \frac{c_2 c_3 \mu^2}{m_2^2} R \hat{r}^a \equiv h_2^{0\pm} \hat{r}^a, \\ (A_i^0)^a &= -\frac{1}{e} \epsilon^{abc} \hat{r}^b \partial_i \hat{r}^c + \hat{r}^a A_i = -\frac{1}{er} \epsilon^{iaj} \hat{r}^j + \hat{r}^a A_i, \quad (A_0^0)^a = 0, \end{aligned} \quad (2.8)$$

where $r = (x, y, z)$, $R^2 \equiv (c_2 \mu^4 - c_1 m_1^2 m_2^2) / (2g m_2^2)$ and overhat indicate the normalisation $\hat{r} = r / \sqrt{x^2 + y^2 + z^2}$. The A_i are arbitrary functions of space-time. This vacuum solution is sometimes called the Higgs vacuum [26] to distinguish it from the usual much simpler vacuum solution where $A_\mu^0 = 0$. The asymptotic condition can be written more explicitly if we consider the spherical ansatz

$$(\phi_\alpha^0)^a(\vec{x}) = h_\alpha(r) \hat{r}^a, \quad (A_i^0)^a = \epsilon^{iaj} \hat{r}^j A(r), \quad (A_0^0)^a = 0, \quad (2.9)$$

where the subscript cl denotes the classical solutions to the equations of motion eq. (2.5). Here we are only considering the static ansatz to simplify our calculation, but one may of course also consider the time-dependent solution. For the monopole solution to have finite energy, we require the two matter fields of eq. (2.9) to approach the vacuum solutions in eq. (2.8) at spacial infinity

$$\lim_{r \rightarrow \infty} h_1(r) = h_1^{0\pm} = \pm R, \quad \lim_{r \rightarrow \infty} h_2(r) = h_2^{0\pm} = \mp \frac{c_2 c_3 \mu^2}{m_2^2} R. \tag{2.10}$$

Also notice that at some fixed value of the radius r , the vacuum solutions ϕ_α^0 and monopole solutions ϕ_α^{cl} both belongs to the 2-sphere in the field configuration space. For example, ϕ_1^0 belong to the 2-sphere with radius R because $(\phi_1^0)^2 = R^2$. Moreover, these vacuum and monopole solutions belong to the homotopy group $\pi^2(S^2) = \mathbb{Z}$. It maps the 2-sphere in the space-time to 2-sphere in the field configuration space, which implies that there are $n \in \mathbb{Z}$ many topologically inequivalent solutions that can be explicitly written by replacing \hat{r}^a in eq. (2.8), (2.9) with

$$\hat{r}_n^a = \begin{pmatrix} \sin(\theta) \cos(n\varphi) \\ \sin(\theta) \sin(n\varphi) \\ \cos(\theta) \end{pmatrix}. \tag{2.11}$$

Since we require the monopole and vacuum solutions to smoothly deform into each other at spacial infinity, both solutions need to share the same integer n usually referred to as the winding number. It is important to note that winding numbers of ϕ_1 and ϕ_2 need to be equal to satisfy $D\phi_1 = D\phi_2 = 0$ and therefore we will denote the winding numbers of ϕ_1 and ϕ_2 as n collectively. If they are not equal we would have $D\phi_1 = 0$ but $D\phi_2 \neq 0$. Next, let us insert our ansatz eq. (2.9) into the equations of motion eq. (2.5) by also making an explicit choice for $A(r)$

$$(\phi_\alpha^{cl})^a = h_\alpha(r) \hat{r}_{n_\alpha}^a, \quad (A_i^{cl})^a = \epsilon^{iaj} \hat{r}_n^j \left(\frac{u(r) - 1}{er} \right). \tag{2.12}$$

This ansatz is more in line with the original ansatz given in [18,19], compare to eq. (2.9). Inserting these expressions into the equations of motion eq. (2.5) we find

$$u''(r) + \frac{u(r)[1 - u^2(r)]}{r^2} + \frac{e^2 u(r)}{2} \{h_2^2(r) - h_1^2(r)\} = 0, \tag{2.13}$$

$$h_1''(r) + \frac{2h_1'(r)}{r} - \frac{2h_1(r)u^2(r)}{r^2} + g \left\{ c_1 \frac{m_1^2}{g} h_1(r) + c_3 \frac{\mu^2}{g} h_2(r) + 2h_1^3(r) \right\} = 0, \tag{2.14}$$

$$h_2''(r) + \frac{2h_2'(r)}{r} - \frac{2h_2(r)u^2(r)}{r^2} + c_2 m_2^2 \left\{ h_2(r) + c_3 \frac{\mu^2}{m_2^2} h_1(r) \right\} = 0. \tag{2.15}$$

Notice that these differential equations are similar to the ones discussed in [18,19], but with the extra field h_2 and extra differential equation eq. (2.15). In the Hermitian model the exact solutions to the differential equations were found by taking the parameter limit called the BPS limit [18,19] where parameters in the theory are taken to zero while keeping the vacuum solution finite. Here we will follow the same procedure and take the parameter limit where quantities in the curly brackets of eq. (2.14) and (2.15) vanish but keeping the vacuum solutions eq. (2.8) finite. We will see in section 4 that we also find the exact solutions in this limit. However, before we solve the differential equations, let us discuss the energy bound of the monopole.

3. The energy bound

The energy of the monopole can be found by inserting the monopole solution into the corresponding Hamiltonian of eq. (2.7).

$$\mathfrak{h} = \int d^3x \text{Tr} \left(E^2 \right) + \text{Tr} \left(B^2 \right) + \text{Tr} \left\{ (D_0 \phi_1)^2 \right\} + \text{Tr} \left\{ (D_i \phi_1)^2 \right\} - \text{Tr} \left\{ (D_0 \phi_2)^2 \right\} - \text{Tr} \left\{ (D_i \phi_2)^2 \right\} + V, \tag{3.1}$$

where E, B are $E_a^i = F_a^{0i}$, $B_a^i = -\frac{1}{2} \epsilon^{ijk} F_a^{jk}$, $i, j, k \in \{1, 2, 3\}$. The gauge is fixed to be the radiation gauge (i.e. $A_a^0 = 0$, $\partial_i A_a^i = 0$). Notice that our monopole ansatz eq. (2.12) is static with no electric charge $E_i^a = 0$ and therefore the above Hamiltonian simplifies to

$$E = \int d^3x \text{Tr} \left(B^2 \right) + \text{Tr} \left\{ (D_i \phi_1)^2 \right\} - \text{Tr} \left\{ (D_i \phi_2)^2 \right\} + V \\ = 2 \int d^3x B_i^a B_i^a + (D_i \phi_1)^a (D_i \phi_1)^a - (D_i \phi_2)^a (D_i \phi_2)^a + \frac{1}{2} V. \tag{3.2}$$

Here, we simplified our expression by dropping the subscripts $A_i^{cl} \rightarrow A_i$, $\phi_\alpha^{cl} \rightarrow \phi_\alpha$. We also keep in mind that these fields depend on the winding numbers $n \in \mathbb{Z}$. In the Hermitian model (i.e., when $\phi_2 = 0$) one can rewrite the kinetic term as $B^2 + D\phi^2 = (B - D\phi)^2 + 2BD\phi$ and find the lower bound to be $\int 2BD\phi$. Here we will follow the similar procedure but we introduce some arbitrary constant $\alpha, \beta \in \mathbb{R}$ such that $B^2 = \alpha^2 B - \beta^2 B$ where $\alpha^2 - \beta^2 = 1$. This will allow us to rewrite the above energy as

$$E = 2 \int d^3x \alpha^2 \left\{ B_i^a + \frac{1}{\alpha} (D_i \phi_1)^a \right\}^2 - \beta^2 \left\{ B_i^a + \frac{1}{\beta} (D_i \phi_2)^a \right\}^2 + 2 \left\{ \alpha B_i^a (D_i \phi_1)^a + \beta B_i^a (D_i \phi_2)^a \right\} + \frac{1}{2} V, \tag{3.3}$$

To proceed from here, we need to assume extra constraints on α and β such that the following is true

$$\int d^3x \alpha^2 \left\{ B_i^a + \frac{1}{\alpha} (D_i \phi_1)^a \right\}^2 - \beta^2 \left\{ B_i^a + \frac{1}{\beta} (D_i \phi_2)^a \right\}^2 \geq 0, \quad (3.4)$$

$$\int d^3x V \geq 0. \quad (3.5)$$

This allows us to write the lower bound on the energy as

$$\begin{aligned} E &\geq 2 \int d^3x \left\{ +\alpha B_i^a (D_i \phi_1)^a + \beta B_i^a (D_i \phi_2)^a \right\} \\ &= 2 \int d^3x \alpha \left\{ B_i^a \partial_i \phi_1^a + e B_i^a \epsilon^{abc} A_i^b \phi_1^c \right\} + \beta \left\{ B_i^a \partial_i \phi_2^a + e B_i^a \epsilon^{abc} A_i^b \phi_2^c \right\} \\ &= 2 \int d^3x \alpha \left\{ B_i^a \partial_i \phi_1^a + (-e \epsilon^{abc} A_i^b B_i^c) \phi_1^a \right\} + \beta \left\{ B_i^a \partial_i \phi_2^a + (-e \epsilon^{abc} A_i^b B_i^c) \phi_2^a \right\} \\ &= 2 \int d^3x \alpha \left\{ B_i^a \partial_i \phi_1^a + \partial_i B_i^a \phi_1^a \right\} + \beta \left\{ B_i^a \partial_i \phi_2^a + \partial_i B_i^a \phi_2^a \right\} \\ &= 2 \int d^3x \alpha \partial_i (B_i^a \phi_1^a) + \beta \partial_i (B_i^a \phi_2^a) \\ &= \lim_{r \rightarrow \infty} \left(2\alpha \int_{S_r} dS_i B_i^a \phi_1^a + 2\beta \int_{S_r} dS_i B_i^a \phi_2^a \right), \end{aligned} \quad (3.6)$$

where in the fourth line we used $D_i B_i^a = 0$ which can be shown from the Bianchi identity $D_\mu \epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}^a = 0$. The last line is obtained by using the Gauss theorem at some fixed value of the radius r . Since we are integrating over the 2-sphere with large radius, we can use the asymptotic conditions eq. (2.10) and replace the monopole solutions $\{\phi_\alpha^a, B_i^a\}$ with the Higgs vacuum $\{(\phi_\alpha^0)^a, (B_i^0)^a\}$

$$\begin{aligned} E &\geq \left(2\alpha \phi_1^{0a} + 2\beta \phi_2^{0a} \right) \lim_{r \rightarrow \infty} \int_{S_r} dS_i (B_i^0)^a \\ &= \left(\pm 2\alpha R \hat{r}_n^a \mp 2\beta \frac{\mu^2}{m_2^2} R \hat{r}_n^a \right) \lim_{r \rightarrow \infty} \int_{S_r} dS_i (B_i^0)^a, \end{aligned} \quad (3.7)$$

where the upper and lower signs of the above energy correspond to the upper and lower signs of the vacuum solutions in eq. (2.8). The explicit value of $(B_i^0)^a$ can be obtained by inserting eq. (2.8) into

$$(B_i^0)^a = -\frac{1}{2} \epsilon^{ijk} \left(\partial_j A_k^0 - \partial_k A_j^0 + e A_j^0 \times A_k^0 \right)^a. \quad (3.8)$$

After a lengthy calculation this expression can be simplified to $B_i^a = \hat{\phi}_1^{0a} B_i = \hat{r}_n^a B_i$ where B_i is defined as

$$B_i \equiv -\frac{1}{2} \epsilon_{ijk} \left\{ \partial^j A^k - \partial^k A^j + \frac{1}{e} \hat{r}_n \cdot \left(\partial^j \hat{r}_n \times \partial^k \hat{r}_n \right) \right\}. \quad (3.9)$$

Notice that integrating the first term over the 2-sphere gives zero by Stokes theorem $\int_S \partial \times A = \int_{\partial S} A = 0$ where one can show that Stokes's theorem on closed surface gives zero by dividing the sphere into two open surfaces. The second term is a topological term which can be evaluated [27] as

$$\int dS_i B_i = -\frac{4\pi n}{e}. \quad (3.10)$$

This is the magnetic charge of the monopole solutions. Notice that we could have chosen $B_i^a = \hat{\phi}_2^{0a} B_i$ instead, which also leads to $\int dS_i B_i = -\frac{4\pi n}{e}$ since we require winding numbers of ϕ_1 and ϕ_2 to be equal. Finally we find our lower bound of the monopole energy

$$E \geq 2R \left(\pm\alpha \mp \beta \frac{c_2 c_3 \mu^2}{m_2^2} \right) \hat{r}_n^a \hat{r}_n^a \left(\frac{-4\pi n}{e} \right) = \frac{-8\pi n}{e} R \left(\pm\alpha \mp \beta \frac{c_2 c_3 \mu^2}{m_2^2} \right). \quad (3.11)$$

Notice that we have some freedom to choose $\alpha, \beta \in \mathbb{R}$ as long as our initial assumptions eq. (3.4) are satisfied. We will see in the next section that we can take a parameter limit of our model which saturates the above inequality and gives exact values to α and β .

4. The fourfold BPS scaling limit

Our main goal is now to solve the coupled differential equations eq. (2.13)-(2.15). Prasad, Sommerfield and Bogomolny [19,18] managed to find the exact solution by taking the parameter limit which simplifies the differential equations. The multiple scaling limit is taken so that all the parameters of the model tend to zero with some combinations of the parameter remaining finite. The combinations are taken so that the vacuum solutions stay finite in this limit. Inspired by this, we will take here a fourfold scaling limit

$$g, m_1, m_2, \mu \rightarrow 0, \quad \frac{m_1^2}{g} < \infty, \quad \frac{\mu^2}{g} < \infty, \quad \frac{\mu^2}{m_2^2} < \infty. \tag{4.1}$$

This will ensure that the vacuum solutions eq. (2.8) stays finite, but crucially the curly bracket parts in eq. (2.14), (2.15) vanish. There is a physical motivation for this limit in which the mass ratio of the Higgs and gauge mass are taken to be zero (i.e. $m_{\text{Higgs}} \ll m_g$) as described in [28]. We will see in the next section that the same type of behaviour is present in our model, hence justifying eq. (4.1). The resulting set of differential equations, after taking the BPS limit is similar to the ones considered in [19,18] with the slightly different quadratic term in eq. (2.13). It is natural to consider a similar ansatz as given in [19,18]

$$u(r) = \frac{evr}{\sinh(evr)}, \tag{4.2}$$

$$h_1(r) = -\alpha \left(v \coth(evr) - \frac{1}{er} \right) \equiv -\alpha f(r), \tag{4.3}$$

$$h_2(r) = -\beta \left(v \coth(evr) - \frac{1}{er} \right) \equiv -\beta f(r). \tag{4.4}$$

where $\alpha, \beta \in \mathbb{R}$ were introduced in section 3 and $f(r) \equiv \left\{ v \coth(evr) - \frac{1}{er} \right\}$. One can check that this ansatz indeed satisfies differential equations eq. (2.13)-(2.15) in the BPS limit. We have decided to put a prefactor α and β in front of eq. (4.3), (4.4) to satisfy the differential equation eq. (2.13). Note that if we take $\alpha = 1$ we get exactly the same as given in [18,19], which is known to satisfy the first order differential equation called Bogomolny equation $B_i - D_i\phi = 0$. The ansatz eq. (4.2)-(4.4) only differs from the ones given in [18,19] by the prefactors α and β , and therefore our ansatz should satisfy Bogomolny equation with the appropriate prefactor to cancel the prefactor in eq. (4.3), (4.4)

$$B_i^b + \frac{1}{\alpha} (D_i\phi_1)^b = 0, \tag{4.5}$$

$$B_i^b + \frac{1}{\beta} (D_i\phi_2)^b = 0, \tag{4.6}$$

where $\phi_\alpha \equiv h_\alpha(r)\hat{r}_n$. If we compare these equations to the terms appearing in the energy of the monopole eq. (3.3), then we can saturate the inequality in eq. (3.11) by

$$E[\phi_1, \phi_2] = \frac{-8\pi nR}{e} \left(\pm\alpha \mp \beta \frac{c_2c_3\mu^2}{m_2^2} \right), \tag{4.7}$$

where upper and lower signs correspond to the vacuum solutions eq. (2.8). Note that this equality is true up to a constant given by $\int d^3xV$ which can be removed by introducing an appropriate constant in the action eq. (2.7). We can calculate the explicit forms of α and β by comparing the asymptotic conditions in eq. (2.10)

$$\lim_{r \rightarrow \infty} h_1^\pm = h_1^{0\pm} = \pm R, \quad \lim_{r \rightarrow \infty} h_2^\pm = h_2^{0\pm} = \mp \frac{c_2c_3\mu^2}{m_2^2} R, \tag{4.8}$$

with the asymptotic values of eq. (4.2)-(4.4)

$$\lim_{r \rightarrow \infty} u(r) = 0, \quad \lim_{r \rightarrow \infty} h_1^\pm(r) = -\alpha v, \quad \lim_{r \rightarrow \infty} h_2^\pm(r) = -\beta v. \tag{4.9}$$

Comparing the eq. (4.8) and eq. (4.9) we find the algebraic equations for α and β . Using $\alpha^2 - \beta^2 = 1$ and assuming $m_2^4 \geq \mu^4$, we find the two set of real solutions

$$\alpha = (\pm) \frac{m_2^2}{l}, \quad v = \mp(\pm) \frac{Rl}{m_2^2}, \quad \beta = \mp(\pm) \frac{c_2c_3\mu^2}{l}, \tag{4.10}$$

where $l = \sqrt{m_2^4 - \mu^4}$. The plus-minus signs in the brackets correspond to the two possible solutions to the algebraic equation of α . These need to be distinguished from the upper and lower signs of v and β which correspond to the vacuums solutions eq. (2.8). Inserting the explicit values of α and β to the energy eq. (4.7) we find

$$E[\phi_1, \phi_2] \equiv (\pm) \frac{8\pi nR}{em_2^2} \left(\frac{\mp m_2^4 - \mu^4}{l} \right), \tag{4.11}$$

where we observe two distinct energies for two vacuum solutions eq. (2.8). Notice that we need to take extra care when choosing the plus or minus sign in the brackets as wrong choices lead to negative energies. To see this explicitly let us consider the monopole solutions with vacuum solutions $\{h_1^{0+}, h_2^{0+}\}$ as boundary conditions. The corresponding energy is

$$E = (\pm) \frac{-8\pi nR}{em_2^2} \left(\frac{m_2^4 + \mu^4}{l} \right). \quad (4.12)$$

Then notice that the upper sign in the brackets leads to the positive energy only when $n < 0$ and the lower sign leads to the positive energy when $n > 0$. This allows us to choose the right sign for the α in eq. (4.10). This implies that the monopole solutions eq. (4.2)-(4.4) takes the following forms

Winding	$\{h_1, h_2\}$	u	Mass
$n > 0$	$\left\{ \frac{m_2^2}{l} f(r), -\frac{c_2 c_3 \mu^2}{l} f(r) \right\}$	$\frac{eRl}{m_2^2 \sinh(eRl/m_2^2)}$	$\frac{8\pi nR}{em_2^2} \left(\frac{m_2^4 + \mu^4}{l} \right)$
$n < 0$	$\left\{ -\frac{m_2^2}{l} f(r), \frac{c_2 c_3 \mu^2}{l} f(r) \right\}$	$\frac{-eRl}{m_2^2 \sinh(-eRl/m_2^2)}$	$\frac{8\pi nR}{em_2^2} \left(\frac{m_2^4 + \mu^4}{l} \right)$

(4.13)

where at rest the mass is equivalent to the energy of the corresponding monopole solutions. We can repeat the same analysis for the case when the monopole solutions asymptotically approach $\{h_1^{0-}, h_2^{0-}\}$ and find the different set of monopole solutions. We summarise here all possible solutions and its corresponding masses

Winding	$\{h_1^0, h_2^0\}$	$\{h_1, h_2\}$	u	Mass
$n > 0$	$\{h_1^{0\pm}, h_2^{0\pm}\}$	$\left\{ \frac{m_2^2}{l} f(r), \mp \frac{c_2 c_3 \mu^2}{l} f(r) \right\}$	$\frac{\pm eRl}{m_2^2 \sinh(\pm eRl/m_2^2)}$	M_{\pm}
$n < 0$	$\{h_1^{0\pm}, h_2^{0\pm}\}$	$\left\{ -\frac{m_2^2}{l} f(r), \pm \frac{c_2 c_3 \mu^2}{l} f(r) \right\}$	$\frac{\mp eRl}{m_2^2 \sinh(\mp eRl/m_2^2)}$	M_{\pm}

(4.14)

Where $M_{\pm} = 8\pi |n| R(m_2^4 \pm \mu^4)/em_2^2 l \geq 0$. We can combine these solutions and find

$\{h_1^0, h_2^0\}$	$\{h_1, h_2\}$	u	Mass
$\{h_1^{0\pm}, h_2^{0\pm}\}$	$\left\{ \text{Sign}(n) \frac{m_2^2}{l} f(r), \text{Sign}(n) \frac{\mp c_2 c_3 \mu^2}{l} f(r) \right\}$	$\frac{\pm \text{Sign}(n) eRl}{m_2^2 \sinh(\pm \text{Sign}(n) eRl/m_2^2)}$	M_{\pm}

(4.15)

So we find two monopoles with two distinct masses characterised by the upper and lower signs of the vacuum solutions eq. (2.8) which can be interchanged by choosing opposite sign for $c_2 c_3$. This is because changing the sign of $c_2 c_3$ exchanges the vacuum solutions eq. (2.8), resulting in swapping of the monopole solutions. Note that the two masses and solutions only coincide in the Hermitian limit $\mu = 0$.

5. Monopole mass and physical region

In this section, we will compare the monopoles masses with Higgs and massive gauge masses. We will analyse different versions of the model characterised by choosing different values of c_i . It was found in [11,16] that mapping the theories with respect to two different Dyson maps eq. (2.6) does not effect the Higgs and gauge masses as they depended on μ^4 . Here we find that different Dyson maps corresponds to exchanging the two monopole solutions eq. (4.15).

5.1. Higgs mass and exceptional points

Let us begin by recalling the results from our previous work [11]. The Higgs masses squared and gauge mass of our model defined in eq. (2.1) are

$$m_0^2 = -c_2 \frac{m_2^4 - \mu^4}{m_2^2}, \quad m_{\pm}^2 = K \pm \sqrt{K^2 - 2c_1 c_2 m_1^2 m_2^2 + 2\mu^4}, \quad m_g = e \frac{|R|}{m_2^2}, \quad (5.1)$$

where $K = -c_1 m_1^2 - c_2 \frac{m_2^2}{2} + \frac{3\mu^4}{2c_2 m_2^2}$. We recall that for some values of parameters, our masses can become complex, which is a common feature of non-Hermitian theories. The region where all Higgs masses are real were investigated in [11,16]. There are two non-overlapping physical regions for $c_1 = c_2 = \pm 1$ in our model eq. (2.1), which are equivalent to $-c_1 = c_2 = \pm 1$ in [11]. Notice that in the BPS limit we have $m_0 = m_{\pm} = 0$, but m_g and M_{\pm} stays finite, such that the ratios m_{Higgs}/m_g vanish in the BPS limit. This is in line with the Hermitian case [28], providing the physical interpretation $m_{\text{Higgs}} \ll m_g$ for the BPS limit.

We have observed that some boundaries of the physical region admit interesting behaviour where the gauge masses vanish. In particular, there are two distinct points (boundaries) called zero exceptional points of type *I* and *II* where the gauge masses vanish in each case. At the type *II* zero exceptional points, the vanishing of gauge masses is not surprising because the vacuum manifold collapse to a point and the broken symmetry is restored. However, at the type *I* zero exceptional point the vacuum manifold is finite and therefore the symmetry is still broken, nonetheless the gauge mass still loses its mass. The mass matrix for Higgs fields is also non-diagonalisable at this point, indicating that this point is a novel feature of the non-Hermitian theory. The type *I* zero exceptional point which we will call OEP, occurs when $m_2^4 = \mu^4$.

5.2. Mass swapping of monopoles with similarity transformations

The monopole solutions of the type eq. (4.2)-(4.4) can not exist in the physical region $c_1 = c_2 = 1$. To see this, recall that we assumed $m_2^4 - \mu^4 \geq 0$ in order to find the solutions to α and β in eq. (4.10). Since we want our Higgs masses to be positive, we require $m_2^4 - \mu^4 \leq 0$ for $c_1 = c_2 = 1$ and $m_2^4 - \mu^4 \geq 0$ for $c_1 = c_2 = -1$. Therefore we will only find the real solutions to α and β in eq. (4.10) when $c_1 = c_2 = -1$. This implies that the monopole solutions of type eq. (4.2)-(4.4) can only exist for the case $c_1 = c_2 = -1$. This restricts our monopole solutions eq. (4.15) to be

$\{h_1, h_2\}$	Mass
$\left\{ \frac{m_2^2}{l} f(r), c_3 \frac{\mu^2}{l} f(r) \right\}$	$M_+ = \frac{8\pi R}{em_2^2} \left(\frac{m_2^4 + \mu^4}{l} \right)$
$\left\{ \frac{m_2^2}{l} f(r), -c_3 \frac{\mu^2}{l} f(r) \right\}$	$M_- = \frac{8\pi R}{em_2^2} l$

(5.2)

here we took the winding number $n = 1$. Let us denote the solutions with the same signs to be $(+, +)$ solution and the opposite signs to be $(+, -)$ solution. Then for example, if $c_3 = 1$ then the first solution of above table is a $(+, -)$ solution. In summary we have

Solution	Mass when $c_3 = 1$	Mass when $c_3 = -1$
$(+, +)$ solution	M_+	M_-
$(+, -)$ solution	M_-	M_+

(5.3)

The Higgs masses and gauge masses are independent of the sign of the non-Hermitian coupling c_3 , but here we see that the monopole masses depend on the signs of the non-Hermitian coupling. This implies that the monopole mass depends on the similarity transformation in eq. (2.6), where choosing η_{\pm} corresponds to $c_3 = \pm 1$. So we observe that the different monopole masses, resulting from different solutions, can be exchanged by using different similarity transformations.

We started with the four different possible theories $S_{c_1 c_2}$ in eq. (2.1) which can be transformed into many different theories under appropriate similarity transformations. Here we considered two similarity transformations η_{\pm} which gives 8 possible theories $\eta_{\pm} S_{c_1 c_2} \eta_{\pm}^{-1} \equiv s_{c_1 c_2 \pm}$. We have observed in [11] that only s_{+++} and s_{---} admit positive gauge masses. This is somewhat reminiscent of how ϕ^4 theory can only admit symmetry breaking when the sign in the mass term is opposite to the sign in the ϕ^4 term. If we focus on the $(+, +)$ solution then it appears at first sight that the two theories s_{--+} and s_{---} are inequivalent as the mass of the $(+, +)$ solution changes. However, notice that the masses are swapped between $(+, +)$ and $(+, -)$ solutions, therefore the two theories s_{--+} and s_{---} are in fact equivalent with respect to the internal symmetry of exchanging the two monopole solutions. This internal symmetry can be seen in eq. (5.2) where choosing the different Dyson map (i.e. different values for c_3) leads to two monopole solutions to swap. Let us summaries this

Theories	Gauge mass	Mass of $(++)$ solution	Mass of $(+-)$ solution
s_{+++}	m_g	do not exist	do not exist
s_{--+}	m_g	M_+	M_-
s_{---}	m_g	M_-	M_+

(5.4)

5.3. Finite energy condition violation

As discussed in section 5.1, the gauge mass vanishes at the zero exceptional point $\mu^4/m_2^4 = 1$. We notice that in this parameter limit our two monopole masses eq. (5.3) vanish or diverge

$$\lim_{\frac{\mu^4}{m_2^4} \rightarrow 1} \frac{8\pi |n| R}{em_2^2} l = 0, \quad \lim_{\frac{\mu^4}{m_2^4} \rightarrow 1} \frac{8\pi |n| R}{em_2^2} \left(\frac{m_2^4 + \mu^4}{l} \right) \rightarrow \infty. \tag{5.5}$$

We have taken here $R = \sqrt{(m_1^2 m_2^2 - \mu^4)/2gm_2^2}$ since our monopole solution can only exist when $c_1 = c_2 = -1$. From now on we keep this choice and take $c_1 = c_2 = -1$. The monopole solutions eq. (5.2) also diverge at the zero exceptional point. This is a similar effect to how Higgs fields are no longer identifiable at the zero exceptional point because they diverge and the Hamiltonian is no longer diagonalisable. Let us denote the two monopole masses as $M_- \equiv M_{cov}$ and $M_+ \equiv M_{div}$ for converging and diverging masses at OEP limit. The converging mass can be written in terms of the gauge mass as

$$M_{cov} = \frac{8\pi |n| R l}{e m_2^2} = \frac{8\pi |n|}{e^2} m_g. \tag{5.6}$$

Therefore we see that only one of the monopole masses satisfies the Montonen-Olive duality [8], whereas the other mass does not satisfy this duality due to an additional factor $(m_2^4 + \mu^4)/(m_2^4 - \mu^4)$.

Let us rewrite the monopole and gauge masses in terms of the finite quantities in the BPS limit eq. (4.1)

$$m_g = \frac{e}{\sqrt{2}} \sqrt{(X - YZ)(1 - Z^2)}, \tag{5.7}$$

$$M_{cov} = \frac{8\pi |n|}{e} \sqrt{(X - YZ)(1 - Z^2)}, \tag{5.8}$$

$$M_{div} = \frac{8\pi |n|}{e} \sqrt{X - YZ} \frac{1 + Z^2}{\sqrt{1 - Z^2}}, \tag{5.9}$$




Fig. 1. Both panels are plotted for $X=1, Y=0.8, |n|=1$ with different values for e . The dashed lines in panel (b) indicates the self-dual points where the gauge mass and the monopole mass coincide at $Z_0^- = 0.689, Z_0^+ = 1.45$.

where $m_1^2/g \equiv X, \mu^2/g \equiv Y, \mu^2/m_2^2 \equiv Z$. Therefore the OEP corresponds to $Z=1$. One can view the parameters Y and Z as a measure of how strong the non-Hermitian term is coupled to the Hermitian theory. Notice that two monopole masses eq. (5.8), (5.9) are real and positive when the gauge mass eq. (5.7) is real and positive. This implies that the physical region of monopole masses and gauge mass coincide when $c_1 = c_2 = -1$ and $Z \neq 1$. Moreover, monopole masses and Higgs masses eq. (5.1) are both real and positive when $c_1 = c_2 = -1$ and $Z \neq 1$. This is because the physical region of Higgs masses is a subset of the physical region of the gauge mass as seen in [11].

Let us plot eq. (5.7)-(5.9) for $n=1, X > Y$ (i.e. $m_1^2 > \mu^2$) with a weak and a strong coupling $e=2, e=10$, respectively. The three regions are separated by the two types of exceptional points discussed in section 5.1. Region 1 is bounded between the Hermitian limit $Z \rightarrow 0$ and the above OEP of type I at $Z=1$, where the vacuum manifold stays finite. We see here that one of the monopole masses, M_{div} , diverges at the OEP. This violates our initial assumption eq. (2.10) of the finite energy of the monopole. The reason for this is because the monopole solution eq. (5.2) is ill-defined at the OEP, therefore, one can not continuously deform the monopole solutions to the vacuum solutions eq. (2.8). In fact, the finite energy condition eq. (2.10) is also violated in the region 2 as the monopole solutions are complex, but the vacuum solutions are real and therefore they can not continuously be deformed into each other by taking $r \rightarrow \infty$. In region 3, the finite energy condition is restored as both the monopole and the vacuum solutions are complex. In summary:

	Region 1	Region 2	Region 3
Monopole	Real	Complex	Complex
Vacuum	Real	Real	Complex

(5.10)

In the strong coupling case $e=10$, we find self-dual points, Z_0^\pm , at which the gauge mass m_g and the monopole mass M_{div} become equal and interchange their relative size in both regions 1 and 3. This phenomena only occurs when the gauge mass is bigger than the monopole mass in the Hermitian limit $Z \rightarrow 0$. To see this, consider the monopole masses when $Z=0$ $M_0 = M_{div}|_{Z=0} = M_{cov}|_{Z=0}$. Then one can show that $m_g < M_0$ implies $e^2 < 8\sqrt{2}|n|\pi$. On the other hand, solving $m_g = M_{div}$ one finds

$$(Z_0^\pm)^2 = \frac{e^2 \pm 8\sqrt{2}|n|\pi}{e^2 \mp 8\sqrt{2}|n|\pi} > 0 \implies e^2 > 8\sqrt{2}|n|\pi. \quad (5.11)$$

Therefore $m_g < M_0$ contradict $(Z_0^\pm)^2 > 0$. Note that, the existence of the self-dual points Z_0^\pm only depends on the values of e and $|n|$ and not on X and Y .

Next let us consider the $X < Y$ (i.e. $m_1^2 < \mu^2$) case. This case is equivalent to the $X > Y$ but with regions 1 and 3 exchanged. This means that the behaviour observed in the strongly non-Hermitian region in $X > Y$ of Fig. 1 is equivalent to the behaviour observed in the weakly non-Hermitian region in $X < Y$ of Fig. 2.

Finally let us look at the $X=Y$ (i.e. $m_1^2 = \mu^2$) case. We see that the regions 2 collapses as the two boundaries coincide (Fig. 3). The gauge and monopole masses m_g and M_{cov} behave similarly to the $X > Y$ case but we see an interesting behaviour of the monopole mass M_{div} at the OEP. As we discussed in previous cases, the finite energy condition fails at the OEP therefore one would expect to find unbounded energy. However, we observe that the monopole mass M_{div} is finite even at the OEP. This can be verified by looking at the asymptotic value of M_{div} at $Z=1$

$$\lim_{Z \rightarrow 1} (M_{div}|_{X=Y}) = \frac{8\pi\sqrt{X}}{e} \lim_{Z \rightarrow 1} \left(\frac{\sqrt{1-Z}(1+Z^2)}{\sqrt{1-Z^2}} \right) = \frac{8\pi\sqrt{X}}{e} \sqrt{2}. \quad (5.12)$$

At the OEP the monopole solutions (5.2) are ill-defined and violate the finite energy condition eq. (2.10). However for $X=Y=1$ they remain finite.

In the three cases we considered above, we see that the monopole mass can reach infinity with finite values of Z . For example, the region 1 of the Fig. 1 can not be physical as M_{div} diverges at $Z=1$. We see that the only physical region (i.e. no diverging mass with finite values of Z) is the region 1 of the case $X < Y$ and $X=Y$.




Fig. 2. Both panels are plotted for $X = 1, Y = 1.2$ with different values for e . The dashed lines in panel (b) indicates the self-dual points where the gauge mass and the monopole mass coincide at $Z_0^- = 0.689, Z_0^+ = 1.45$.




Fig. 3. Both panels are plotted for $X = 1, Y = 1$ with different values of e . The dashed lines in panel (b) indicates the self-dual points where the gauge mass and the monopole mass coincide at $Z_0^- = 0.689, Z_0^+ = 1.45$.

6. Conclusion

We found and analysed the 't Hooft-Polyakov monopole in the non-Hermitian model possessing $SU(2)$ symmetry and anti-linear CPT symmetry using the pseudo-Hermitian approach. Following the procedure outlined in [18,19] we have found two exact monopole solutions eq. (5.2) in the BPS limit which saturated the lower bound of the energy. These monopole solutions can only exist in one of the two physical regions characterised in [11] by c_1 and c_2 . We have considered four theories S_{c_1, c_2} eq. (2.1) which were mapped to 8 theories via two similarity transformations eq. (2.6). These transformations were previously found to not affect the Higgs masses and gauge masses as they have no dependencies on the similarity transformation parameter c_3 . Here we observed two distinct masses for each monopole solutions to be interchangeable under different similarity transformations, as they depend on c_3 .

At first sight that seemed to suggest that the two theories with different similarity transformations are not physically equivalent. This is a well known feature in the quantum mechanical setting, where different types of similarity transformations lead to different sets of observables, i.e. different physical theories [29]. Much to our surprise the interchange of the monopole masses for the different similarity transformations is, however, compensated by the internal structure that leads to different monopoles emerging from different soliton solutions. Thus, as discussed in section 5.2, overall the two theories resulting from the different similarity transformations can be seen as equivalent. It will be very interesting to investigate in future whether this feature survives when considering more complicated similarity transformations.

The behaviour of the gauge and monopole masses were investigated as function of the non-Hermitian coupling $Z = \mu^2/m_2^2$. Three disconnected regions were found in $Z \in (0, \infty)$ where finite energy condition failed in one of the regions, resulting in complex energies. At one of the boundary (OEP of type I) of the region, the monopole mass diverged to infinity, signalling that the theory is unbounded at this boundary. However, the monopole mass can have finite non-zero value even at OEP of type I when two boundaries (OEP of type I and II) coincide.

The issue of finite energy at the boundary requires further investigation as it is peculiar to observe a finite energy of the solution which is ill-defined at the boundary. Here we have considered here a classical theory and of course it will be very interesting to investigate its quantisation in detail.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] P.A.M. Dirac, Quantised singularities in the electromagnetic field, *Proc. R. Soc. Lond. A* 133 (821) (1931) 60–72.
- [2] V.A. Mitsou, The quest for magnetic monopoles—past, present and future, *Proc. Sci.* 188 (2018).
- [3] N.E. Mavromatos, V.A. Mitsou, Magnetic monopoles revisited: models and searches at colliders and in the cosmos, arXiv preprint, arXiv:2005.05100, 2020.
- [4] G. 't Hooft, Magnetic monopoles in unified theories, *Nucl. Phys. B* 79 (1974) 276–284 (CERN-TH-1876).
- [5] A.M. Polyakov, Spectrum of particles in quantum field theory, *JETP Lett.* 20 (1974) 430–433.
- [6] P. Goddard, D. Olive, Magnetic monopoles in gauge field theories, *Rep. Prog. Phys.* 41 (9) (1978) 1357.
- [7] E.J. Weinberg, *Classical Solutions in Quantum Field Theory: Solitons and Instantons in High Energy Physics*, Cambridge University Press, 2012.
- [8] C. Montonen, D.I. Olive, Magnetic monopoles as gauge particles?, *Phys. Lett. B* 72 (1977) 117–120 (CERN-TH-2391).
- [9] A. Fring, T. Taira, Goldstone bosons in different PT-regimes of non-Hermitian scalar quantum field theories, *Nucl. Phys. B* 950 (2020) 114834.
- [10] A. Fring, T. Taira, Pseudo-Hermitian approach to Goldstone's theorem in non-Abelian non-Hermitian quantum field theories, *Phys. Rev. D* 101 (4) (2020) 045014.
- [11] A. Fring, T. Taira, Massive gauge particles versus Goldstone bosons in non-Hermitian non-Abelian gauge theory, arXiv preprint, arXiv:2004.00723, 2020.
- [12] J. Alexandre, P. Millington, D. Seynaeve, Symmetries and conservation laws in non-Hermitian field theories, *Phys. Rev. D* 96 (6) (2017) 065027.
- [13] J. Alexandre, D. Seynaeve, P. Millington, Consistent description of field theories with non-Hermitian mass terms, *J. Phys. Conf. Ser.* 952 (2017) 012012.
- [14] J. Alexandre, J. Ellis, P. Millington, D. Seynaeve, Gauge invariance and the Englert-Brout-Higgs mechanism in non-Hermitian field theories, *Phys. Rev. D* 99 (7) (2019) 075024.
- [15] J. Alexandre, J. Ellis, P. Millington, D. Seynaeve, Spontaneous symmetry breaking and the Goldstone theorem in non-Hermitian field theories, *Phys. Rev. D* 98 (4) (2018) 045001.
- [16] J. Alexandre, J. Ellis, P. Millington, D. Seynaeve, Spontaneously breaking non-Abelian gauge symmetry in non-Hermitian field theories, *Phys. Rev. D* 101 (3) (2020) 035008.
- [17] P.D. Mannheim, Goldstone bosons and the Englert-Brout-Higgs mechanism in non-Hermitian theories, *Phys. Rev. D* 99 (4) (2019) 045006.
- [18] E.B. Bogomolny, The stability of classical solutions, *Sov. J. Nucl. Phys. (Engl. Transl.)* 24 (4) (1976).
- [19] M.K. Prasad, C.M. Sommerfield, Exact classical solution for the 't Hooft monopole and the Julia-Zee dyon, *Phys. Rev. Lett.* 35 (12) (1975) 760.
- [20] A.M. Polyakov, Particle spectrum in quantum field theory, in: *30 Years of the Landau Institute — Selected Papers*, World Scientific, 1996, pp. 540–541.
- [21] C.M. Bender, *PT Symmetry: In Quantum and Classical Physics*, World Scientific Publishing, 2018.
- [22] A. Mostafazadeh, Pseudo-Hermitian representation of quantum mechanics, *Int. J. Geom. Methods Mod. Phys.* 7 (7) (2010) 1191–1306.
- [23] C.M. Bender, H.F. Jones, R.J. Rivers, Dual PT-symmetric quantum field theories, *Phys. Lett. B* 625 (3–4) (2005) 333–340.
- [24] G.H. Derrick, Comments on nonlinear wave equations as models for elementary particles, *J. Math. Phys.* 5 (9) (1964) 1252–1254.
- [25] E. Corrigan, D.I. Olive, J. Nuyts, D.B. Fairlie, Magnetic monopoles in SU(3) gauge theories, *Nucl. Phys.* 106 (1975) 475–492 (CERN-TH-2102).
- [26] P. Goddard, D.I. Olive, Magnetic monopoles in gauge field theories, *Rep. Prog. Phys.* 41 (9) (1978) 1357.
- [27] J. Arafune, P.G.O. Freund, C.J. Goebel, Topology of Higgs fields, *J. Math. Phys.* 16 (2) (1975) 433–437.
- [28] T.W. Kirkman, C.K. Zachos, Asymptotic analysis of the monopole structure, *Phys. Rev. D* 24 (4) (1981) 999.
- [29] F.G. Scholtz, H.B. Geyer, F.J.W. Hahne, Quasi-Hermitian operators in quantum mechanics and the variational principle, *Ann. Phys.* 213 (1992) 74–101.