



# City Research Online

## City St George's, University of London

**Citation:** Rawson, T. M., Moore, L. S. P., Castro-Sanchez, E., Charani, E., Davies, F., Satta, G., Ellington, M. J. & Holmes, A. H. (2020). COVID-19 and the potential long-term impact on antimicrobial resistance. *Journal of Antimicrobial Chemotherapy*, 75(7), pp. 1681-1684. doi: 10.1093/jac/dkaa194

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/24498/>

**Link to published version:** <https://doi.org/10.1093/jac/dkaa194>

**Copyright and Reuse:** Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

# 1 COVID-19 and the potential long-term impact on Antimicrobial 2 Resistance

3

4 Timothy M Rawson,<sup>1,2,3</sup> Luke SP Moore,<sup>1,4</sup> Enrique Castro-Sanchez,<sup>1</sup> Esmita Charani,<sup>1,5</sup> Frances Davies,<sup>1,3</sup>  
5 Giovanni Satta,<sup>1,3</sup> Matthew J Ellington,<sup>5</sup> Alison H Holmes<sup>1,2,3</sup>

6

7

## 8 Affiliations:

- 9 1. *National Institute for Health Research Health Protection Research Unit in Healthcare Associated Infections and*  
10 *Antimicrobial Resistance, Imperial College London, Hammersmith Campus, Du Cane Road, London. W12 0NN.*  
11 *United Kingdom.*
- 12 2. *Centre for Antimicrobial Optimisation, Hammersmith Hospital, Imperial College London, Du Cane Road, London,*  
13 *W12 0NN. United Kingdom.*
- 14 3. *Imperial College Healthcare NHS Trust, Hammersmith Hospital, Du Cane Road, London. W12 0HS. United*  
15 *Kingdom*
- 16 4. *Chelsea & Westminster NHS Foundation Trust, 369 Fulham Road. London. SW10 9NH. United Kingdom.*
- 17 5. *Public Health England, National Infection Service, 61 Colindale Avenue, London, NW9 5EQ. United Kingdom*

18

19

20

## 21 Corresponding author:

22 Professor Alison Holmes, Health Protection Research Unit in Healthcare Associated Infections & Antimicrobial Resistance,  
23 Hammersmith Hospital, Du Cane Road, London. W12 0NN. United Kingdom. Email: [alison.holmes@imperial.ac.uk](mailto:alison.holmes@imperial.ac.uk)

24 Telephone: +44 (0) 2033132732.

25

26

27

28 **Summary**

29 The emergence of the SARS-CoV-2 respiratory virus has required an unprecedented  
30 response to control the spread of the infection and protect the most vulnerable within  
31 society. Whilst the pandemic has focused society on the threat of emerging infections and  
32 hand hygiene, certain infection control and antimicrobial stewardship policies may have  
33 to be relaxed. It is unclear whether the unintended consequences of these changes will  
34 have a net-positive or -negative impact on rates of antimicrobial resistance. Whilst the  
35 urgent focus must be on allaying this pandemic, sustained efforts to address the longer-  
36 term global threat of antimicrobial resistance should not be overlooked.

37

38

39

40

41

42

43

44

45

46 Summary: 98 words

47 Text: 1178 words

48 **Main text**

49 The emergence of, and subsequent pandemic caused by the Severe Acute Respiratory  
50 Syndrome Coronavirus 2 (SARS-CoV-2) has placed an immense strain on healthcare  
51 systems.<sup>1,2</sup> This has required unprecedented response(s) to control the spread of  
52 infection and protect the most vulnerable.

53 In response to the pandemic, healthcare systems have rapidly adapted infection control  
54 policies to ensure adequate capacity to isolate patients with potential SARS-CoV-2  
55 infection. Societal focus on the threat from this emerging infectious disease has driven a  
56 heightened awareness of the importance of personal hygiene, particularly hand hygiene,  
57 environmental contamination, and increased use of personal protective equipment (PPE).  
58 The pandemic is also likely to require the relaxing of measures to prevent the spread of  
59 multi-drug resistant organisms (MDRO), such as screening and isolation in single rooms,  
60 and antimicrobial stewardship.

61 The paucity of available data makes it difficult to predict the impact that this pandemic  
62 may have on antimicrobial stewardship programs and long-term rates of antimicrobial  
63 resistance (AMR). On one hand, the increased focus on hand hygiene, attempts to limit  
64 patient contact, and social distancing may lead to reductions in healthcare associated  
65 transmission of disease. On the other hand, due to the prioritization of isolation rooms to  
66 COVID patients, the cohorting and/or management in open bays of patients colonized  
67 with CPE/VRE/MRSA/*Clostridium difficile* (*C.difficile*), and the inevitable higher workload  
68 of healthcare workers may potentially lead to a higher number of hospital transmissions.

69 The potential propagation of AMR may also be exacerbated by increasing rates of  
70 antimicrobial prescribing and potential breakdown in well-established stewardship  
71 programs. For example, despite few reports of bacterial co-infection, 62% of patients with  
72 COVID-19 had received antimicrobial therapy in the recent International Severe Acute  
73 Respiratory and Emerging Infections Consortium (ISARIC) report.<sup>3</sup> These prescriptions  
74 tend to be broad-spectrum in nature.<sup>4</sup> In addition to excessive and inappropriate  
75 antimicrobial prescribing, the spread of other pathogens and MDRO's may also be  
76 affected by day-to-day practicalities of an emergency focus on a single primary pathogen  
77 which may affect the depth of sampling for other organisms. The redeployment of  
78 antimicrobial stewardship teams and laboratory capacity to support the workload  
79 associated with SARS-CoV-2 is likely to compound this further.

80 With predictions that the current pandemic could continue to consume the focus of  
81 individual national healthcare systems such as the United Kingdom and United States for  
82 up to 18 months,<sup>5</sup> urgent analysis of its impact on AMR is required. This will support the  
83 development of contingency interventions to mitigate the potential impact of the pandemic  
84 on rates and transmission of AMR. Learning early lessons from countries currently  
85 affected will be important in supporting evidence-based guidance for those regions not  
86 yet burdened by an exponential rise in COVID-19 cases. In particular, the unintended  
87 consequences, whether positive or negative, of these health system changes need to be  
88 described, and where negative impacts are identified these must be mitigated against to  
89 ensure that sustained efforts to address the long-term and devastating threat of AMR.

90 **Table 1** outlines some of the core antimicrobial stewardship efforts affected by the SARS-  
91 CoV-2 pandemic, suggesting potential interventions to help mitigate the impact of the  
92 pandemic response on AMR.

93 With the rapid redeployment of side rooms and PPE to protect patients and staff from  
94 respiratory viral infections and saturation of any isolation capacity, the ability to adhere to  
95 isolation policies normally deployed to prevent the spread of MDROs is likely to be  
96 challenged.(REF) Whilst increased hand hygiene may help prevent transmission<sup>6</sup> -  
97 pragmatic interventions are required to ensure sustained surveillance for MDRO's in both  
98 SARS-CoV-2 positive and negative patient cohorts.

99 With the disruption to routine services and redeployment of staff to alternative areas, such  
100 as critical care, is likely to cause disruption to engrained, top down antimicrobial  
101 stewardship programs within hospitals. With the requirement for increased infectious  
102 disease support for managing the response to the pandemic, stewardship teams may be  
103 redeployed from their primary roles leading to reduced opportunities to optimize  
104 antimicrobial therapy in patients. Loss of expert support for antimicrobial decision making,  
105 a paucity of evidence-based guidelines for antimicrobial prescribing in SARS-CoV-2, and  
106 anxiety of medical colleagues in front of deteriorating patients, may lead to further  
107 inappropriate use of antimicrobials. Urgent steps are needed to develop consensus on  
108 empirical use of antimicrobials together with clarification on the role of *Watch* and *Reserve*  
109 agents in the WHO essential medicines AWaRe criteria.<sup>7</sup> To support the redeployment of  
110 stewardship teams, leadership must be developed within local teams managing SARS-  
111 CoV-2 patients, supported by the development of evidence-based guidelines for the role  
112 of diagnostic tests, such a procalcitonin, to inform appropriate empirical treatment.(REF)

113 Institutions must also focus on behavioral factors and team dynamics that will come under  
114 immense pressure as healthcare services respond to the increase in demand associated  
115 with the pandemic. Team dynamics, fear, and specialty level cultural norms of practice  
116 are major drivers of inappropriate antimicrobial prescribing.<sup>8</sup> These factors must rapidly  
117 be assessed and stabilised in the face of a large variations in staffing levels within  
118 healthcare systems in the coming months. This will require effective leadership, clear  
119 communication across professions, and realigning chains of command in order to  
120 accommodate staff from multiple professions and experience. We must learn from  
121 existing social science research to be responsive and adaptive to the changing priorities  
122 and clinical needs. This cannot be achieved without engagement from across the  
123 healthcare multi-professional teams; the very personnel the interventions will target.

124 The planned implementation of large field-hospitals with a major focus on a primary viral  
125 pathogen also presents potential negative drivers for control of AMR. Uncertainty  
126 regarding the levels of staffing and support services for such facilities, as well the  
127 pressures of clinical practice and patient culture in such a setting do present opportunities  
128 for many of relevant MDRO's to spread with or without disease presentation.

129 In community practice, primary and secondary care has rapidly shifted towards  
130 telemedicine.<sup>9</sup> This is a vital step in protecting both healthcare workers and patients, but  
131 currently has limited data to support its potential to reduce or propagate suboptimal  
132 antimicrobial prescribing, and therefore AMR.<sup>10,11</sup> Limited evidence suggests that  
133 telemedicine is associated with increased rates of antimicrobial use.<sup>10,11</sup> Currently, there  
134 is little guidance and support for colleagues practicing telemedicine for the management  
135 of infection. With primary care being the biggest prescriber of antimicrobials, there is an

136 urgent need for education and training, development of risk stratification, and guidance  
137 that is specific for telemedicine consultations. Community pharmacy roles must be  
138 urgently developed to support appropriate stewardship of antimicrobials prescribed by  
139 this route.

140 In summary, before the outbreak of SARS-CoV-2 in December 2019, major national and  
141 international interventions had begun to focus on reducing the potential future impact of  
142 AMR on society. The consequences of the current pandemic have the potential to stretch  
143 well into the post-COVID19 era. With increased societal sensitization towards emerging  
144 threats from infectious diseases and the concept of transmission and acquisition of  
145 disease, this may subsequently drive greater engagement with the problem of AMR.  
146 However, the current pandemic may also have a greater impact on society through the  
147 unintended propagation of AMR. Whilst, undoubtedly the main focus of healthcare must  
148 be on controlling the spread of SARS-CoV-2 and mitigating immediate impact on  
149 individual patients, we must not lose sight of the longer-term threat of AMR if our current  
150 structures and stewardship programs are completely disrupted during this unprecedented  
151 time.

152

## **Additional information**

### **Transparency declarations**

LSPM reports personal fees from bioMérieux, personal fees from DNAelectronics, personal fees from Dairy Crest, personal fees from Profile Pharma, grants and personal fees from Pfizer, grants from Leo Pharma, grants from CW+ Charity. MJE is a member of PHE's AMRHAI Reference lab which has received financial support for conference attendance, lectures, research projects or contracted evaluations from numerous sources, including: Accelerate Diagnostics, Achaogen Inc., Allegra Therapeutics, Amplex, AstraZeneca UK Ltd, AusDiagnostics, Basilea Pharmaceutica, Becton Dickinson Diagnostics, bioMérieux, Bio-Rad Laboratories, BSAC, Cepheid, Check-Points B.V., Cubist Pharmaceuticals, Department of Health, Enigma Diagnostics, ECDC, Food Standards Agency, GlaxoSmithKline Services Ltd, Helperby Therapeutics, Henry Stewart Talks, IHMA Ltd, Innovate UK, Kalidex Pharmaceuticals, Melinta Therapeutics, Merck Sharpe & Dohme Corp., Meiji Seika Pharma Co., Ltd, Mobidiag, Momentum Biosciences Ltd, Neem Biotech, NIHR, Nordic Pharma Ltd, Norgine Pharmaceuticals, Rempex Pharmaceuticals Ltd, Roche, Rokitan Ltd, Smith & Nephew UK Ltd, Shionogi & Co. Ltd, Trius Therapeutics, VenatoRx Pharmaceuticals, Wockhardt Ltd and WHO. All other authors have no conflicts of interest to declare

### **Funding**

ECS is an NIHR Senior Nurse and Midwife Research Leader, and acknowledges the support of the NIHR Biomedical Research Centre

### **Acknowledgements**

The authors would like to thank members of Imperial College NHS Healthcare Trust who participated in the study. The authors would also like to acknowledge 1) the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Healthcare Associated Infection and Antimicrobial Resistance at Imperial College London in partnership with Public Health England and the NIHR Imperial Patient Safety Translational Research Centre and 2) The Department for Health and Social Care funded Centre for Antimicrobial Optimisation (CAMO) at Imperial College London. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research or the UK Department of Health.

## References

1. Zhu N, Zhang D, Wang W, *et al.* A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 2020; **382**: 727–33.
2. Phelan AL, Katz R, Gostin LO. The Novel Coronavirus Originating in Wuhan, China: Challenges for Global Health Governance. *JAMA - J Am Med Assoc* 2020; **323**: 709–10.
3. International Severe Acute Respiratory and Emerging Infection Consortium. *COVID-19 Report: 08 April 2020*. 2020. <https://isaric.tghn.org/about/>.
4. Chen N, Zhou M, Dong X, *et al.* Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020; **395**: 507–13.
5. Ferguson NM, Laydon D, Nedjati-Gilani G, *et al.* Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand. 2020 <https://doi.org/10.25561/77482>.
6. Erasmus V, Daha TJ, Brug H, *et al.* Systematic Review of Studies on Compliance with Hand Hygiene Guidelines in Hospital Care. *Infect Control Hosp Epidemiol* 2010; **31**: 283–94.
7. Sharland M, Pulcini C, Harbarth S, *et al.* Classifying antibiotics in the WHO Essential Medicines List for optimal use-be AWARe. *Lancet Infect Dis* 2018; **18**: 18–20.
8. Charani E, Ahmad R, Rawson TM, Castro-Sanchèz E, Tarrant C, Holmes AH. The Differences in Antibiotic Decision-making Between Acute Surgical and Acute Medical Teams: An Ethnographic Study of Culture and Team Dynamics. *Clin Infect Dis* 2019; **69**: 12–20.
9. Hollander JE, Carr BG. Virtually Perfect? Telemedicine for Covid-19. *N Engl J Med* 2020: NEJMp2003539.
10. Ray KN, Shi Z, Gidengil CA, Poon SJ, Uscher-Pines L, Mehrotra A. Antibiotic prescribing during pediatric direct-to-consumer telemedicine visits. *Pediatrics* 2019; **143**.
11. Martinez KA, Rood M, Jhangiani N, Kou L, Boissy A, Rothberg MB. Association Between Antibiotic Prescribing for Respiratory Tract Infections and Patient Satisfaction in Direct-to-Consumer Telemedicine. *JAMA Intern Med* 2018; **178**: 1558.

**Table 1. Potential impacts of healthcare system adaption during the COVID-19 pandemic on antimicrobial resistance.**

<b>Effected area</b>	<b>Potential impact</b>	<b>Potential interventions</b>
<b>Increased focus on hand hygiene in hospitals</b>	Reduction in the spread of AMR within healthcare settings.	<p>Ensure adequate resources and equipment available to support increase in demand (e.g. hand sanitizer).</p> <p>Ensure that routine surveillance systems remain in place to monitor rates of AMR within healthcare settings.</p>
<b>Social distancing in the community</b>	Reduction in antimicrobial seeking behaviours by members of the public, leading to reductions in antimicrobial prescribing.	Reinforcement through public engagement.
<b>Less opportunity for isolation of infective / MDRO patients</b>	<p>Potential spread of MDRO.</p> <p>Potential for suboptimal management of other public health challenges (e.g. tuberculosis).</p>	<p>Hand hygiene and barrier nursing.</p> <p>Sustaining MDRO surveillance.</p> <p>Staff and patient education &amp; training.</p> <p>Clustered cohorting of patients by risk-factor (e.g. COVID-19 &amp; CPE, COVID-19 &amp; MRSA).</p>
<b>Pre-emptive discharge of patients and cancellation of routine procedures to enhance bed capacity</b>	Reduction in patients carrying MDRO, such as CPE, within the hospital environment.	Stringent surveillance systems to detect and track the spread of AMR on reintroduction of these patients to healthcare services.
<b>Diversion of all PPE for SARS-CoV-2 patients</b>	Potential spread of MDRO.	<p>Hand hygiene and barrier nursing.</p> <p>Sustaining MDRO surveillance.</p> <p>Staff and patient education &amp; training.</p> <p>Appropriate stratification of PPE for different indications in line with evidence-based guidelines.</p>
<b>Increased rates of empirical antimicrobial therapy for patients presenting with respiratory symptoms</b>	Potential of AMR.	<p>Clear guidelines for empirical therapy in suspected SARS-CoV-2 patients. Specifically delineating the requirement for anti-pseudomonal and / or atypical coverage.</p> <p>Education and emphasis on local stewardship within all healthcare workers.</p> <p>Re-establishment of AMS oversight as soon as possible.</p> <p>Upskilling of staff within the organisation (e.g nurses and</p>

		<p>pharmacy technicians) to take on broader roles and responsibilities.</p> <p>Development of rapid diagnostics to support prescribing decisions. Including a clear role for the use of procalcitonin to detect bacterial infection.</p> <p>Ensuring that pandemic preparedness is part of future IPC and AMS strategy.</p>
<b>Increased rate of telemedicine within primary and secondary care and outpatient services</b>	<p>Possible increase in community rates of antimicrobial prescribing as part of safety netting.</p> <p>Possible reduction in community antimicrobial prescriptions due to social distancing and reduced access to pharmacies.</p>	<p>Need for education and specialist support to develop AMS strategies for telemedicine.</p> <p>Engagement with community-based pharmacies, who may not be confident in screening secondary care medicines.</p> <p>Need for development in technology to support risk stratification.</p>
<b>Redeployment of antimicrobial stewardship teams to deal with healthcare strain due to pandemic</b>	<p>Loss of developed stewardship frameworks within local healthcare environments.</p>	<p>Focus on education and responsibility of individual teams for promotion of appropriate antimicrobial usage.</p> <p>Addressing current social hierarchies within healthcare and upskilling of staff within the organisation (e.g nurses and pharmacy technicians) to provide routine AMS services.</p> <p>Integration of AMS / IPC teams.</p>
<b>Maintenance of institutional memory and team dynamics within organisations experiencing rapid reorganization and recruitment of staff</b>	<p>Loss of best practice and leadership within local team environments.</p>	<p>Education and training.</p> <p>Focus on fostering positive behaviours towards antimicrobials and infection control.</p> <p>Ensuring that structures are agile enough to absorb new individuals with minimal impact of process and patient care.</p>
<b>Overcrowding associated with overloading of healthcare systems</b>	<p>Major driver for the transmission of AMR.</p>	<p>Stringent surveillance systems to detect and track the spread of AMR.</p> <p>Ensuring that routine MDRO screening still takes place in the face of increased viral screening.</p> <p>Cohorting of high-risk patients.</p> <p>Contingency plans for rapidly responding to detected outbreaks.</p>

**Depletion of structural resources**

Loss of side room capacity leading to propagation of SARS-CoV-2 infection due to cohorting of positive and negative patients.

Stringent pathways for segregation of cases.

Rapid diagnostics to facilitate rapid identification.

**Legend:** AMR = antimicrobial resistance; AMS = antimicrobial stewardship; CPE = carbapenemase producing enterobacteriaceae; COVID-19 = novel coronavirus 2019; IPC = infection prevention and control; MDRO = multi-drug resistant organism; MRSA = methicillin resistant *Staphylococcus aureus*; PPE = personal protective equipment; SARS-COV-2 = severe acute respiratory distress coronavirus 2