



# City Research Online

## City St George's, University of London

**Citation:** Mena-Tudela, D., Gonzalez-Chorda, V. M., Soriano-Vidal, F. J., Bonanad-Carrasco, T., Centeno-Rico, L., Vila-Candel, R., Castro-Sanchez, E. & Gasch, A. C. (2020). Changes in health sciences students' perception of obstetric violence after an educational intervention. *Nurse Education Today*, 88, 104364. doi: 10.1016/j.nedt.2020.104364

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/24499/>


**Link to published version:** <https://doi.org/10.1016/j.nedt.2020.104364>

**Copyright and Reuse:** Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

## Journal Pre-proof

Changes in health sciences STUDENTS' perception of obstetric violence after an educational intervention

Desirée Mena-Tudela, Víctor Manuel González-Chordá, Francisco Javier Soriano-Vidal, Teresa Bonanad-Carrasco, Laura Centeno-Rico, Rafa Vila-Candel, Enrique Castro-Sánchez, Águeda Cervera Gasch



PII: S0260-6917(19)31136-0

DOI: <https://doi.org/10.1016/j.nedt.2020.104364>

Reference: YNEDT 104364

To appear in: *Nurse Education Today*

Received date: 24 July 2019

Revised date: 18 December 2019

Accepted date: 7 February 2020

Please cite this article as: D. Mena-Tudela, V.M. González-Chordá, F.J. Soriano-Vidal, et al., Changes in health sciences STUDENTS' perception of obstetric violence after an educational intervention, *Nurse Education Today*(2020), <https://doi.org/10.1016/j.nedt.2020.104364>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## CHANGES IN HEALTH SCIENCES STUDENTS' PERCEPTION OF OBSTETRIC VIOLENCE AFTER AN EDUCATIONAL INTERVENTION

Word count: 4,658 words.

Desirée Mena-Tudela<sup>1</sup>, Víctor Manuel González-Chordá<sup>1</sup>, Francisco Javier Soriano-Vidal<sup>2,3,4</sup>, Teresa Bonanad-Carrasco<sup>1</sup>, Laura Centeno-Rico<sup>1</sup>, Rafa Vila-Candel<sup>2,5</sup>, Enrique Castro-Sánchez<sup>6</sup>, Águeda Cervera Gasch<sup>1</sup>

<sup>1</sup> Nursing Department, Universitat Jaume I, Spain.

<sup>2</sup> Department of Nursing, Universidad Católica de Valencia "San Vicente Mártir", Valencia, Spain.

<sup>3</sup> Department of Obstetrics and Gynecology, Hospital Lluís Alcanyis, Foundation for the Promotion of Health and Biomedical Research in the Valencian Region (FISABIO), Valencia, Spain.

<sup>4</sup> Department of Nursing, University of Alicante, San Vicente del Raspeig, Alicante, Spain.

<sup>5</sup> Department of Obstetrics and Gynecology, Hospital Universitario de la Ribera, Foundation for the Promotion of Health and Biomedical Research in the Valencian Region (FISABIO), Valencia, Spain

<sup>6</sup> National Institute for Health Research, Health Protection Research Unit (NIHR HPRU) in Healthcare-Associated Infection and Antimicrobial Resistance at Imperial College London, United Kingdom

Corresponding author: Desirée Mena-Tudela. PhD

## CHANGES IN HEALTH SCIENCES STUDENTS' PERCEPTION OF OBSTETRIC VIOLENCE AFTER AN EDUCATIONAL INTERVENTION

### Abstract

**Background:** Obstetric violence is a type of gender-based violence that is presented structurally. This type of violence has physical and psychological consequences for both the women who experience it and health professionals. The World Health Organization adds that health professionals need training to ensure that pregnant women are treated with compassion and dignity.

**Objectives:** The objective of the study was to evaluate health sciences students' perception of obstetric violence and to identify possible changes after an educational intervention.

**Design:** A pre-post quasi-experimental study was carried out between January and June 2019.

**Settings and participants:** Students of medicine and nursing from Jaume I University (Universitat Jaume I) (Spain).

**Methods:** An ad hoc scale comprising 33 items was designed to measure the students' perceptions. In addition, sociodemographic and control variables were collected. Descriptive analyses of the sample and the scale were carried out, and a bivariate analysis was performed.

**Results:** Of the students surveyed, 89.7% were women, and the majority were nursing students. Of the 33 items, 28 (84.84%) showed statistically significant changes in the pre-post-intervention measurement. Twenty-five of the 33 items (75.75%) showed a relationship with the sociodemographic variables of gender, field, course and ever having been pregnant.

**Conclusion:** This study shows the change in health sciences students' perceptions of obstetric violence after an educational intervention. In addition, the normalization of this type of violence was observed with the progression of training and with personal obstetric experience.

**Keywords:** Obstetric Violence; Nursing students; Medical students; Perception

## **Introduction**

The World Health Organization (WHO) states that "*all women have the right to receive the highest level of health care, which includes the right to dignified and respectful care in pregnancy and childbirth, and the right not to suffer violence or discrimination*" (WHO, 2014). In 1985, the European regional office of the WHO, the Pan American Health Organization and the regional office of the WHO for the Americas, at a conference on appropriate technology for childbirth, created a series of consensus recommendations among obstetricians, paediatricians, midwives, psychologists, epidemiologists, mothers and other professionals. The result was the "Declaration of Fortaleza" (World Health Organization, 1985), which the WHO considers applicable to all perinatal services throughout the world.

## **Background**

Some definitions of obstetric violence (OV) exist. Specifically, the "Organic Law on the Right of Women to a Life Free of Violence", published in March 2007 in Venezuela, defines this term as "*...the appropriation of the body and reproductive processes of women by health personnel, which is expressed as dehumanized treatment, an abuse of medication, and the conversion of natural processes into pathological ones, bringing with it loss of autonomy and the*

*ability to decide freely about their bodies and sexuality, negatively impacting the quality of life of women"* (Diaz-Tello, 2016; Pérez D'gregorio, 2010).

The WHO warns that an increasing number of studies on the experiences of women during pregnancy and, in particular, during childbirth present an alarming scenario, indicating that many women around the world experience disrespectful, offensive or negligent treatment during labour (WHO, 2014). In addition, it describes the practices that make OV visible: *disrespectful and offensive treatment during childbirth, physical abuse, profound humiliation and verbal abuse, medical procedures performed without consent or under coercion (including sterilization), lack of confidentiality, failure to obtain the complete informed consent, refusal to administer analgesics, flagrant violations of privacy, refusal of admission to a health centre, negligence towards women during childbirth and the retention of women and new-borns in health centres due to their inability to pay* (WHO, 2014), among others.

Concerning this type of violence, several conjectures have been raised regarding possible variables that favour its social stratification (Castro and Frías, 2019): lower socioeconomic level (Brandão et al., 2018; Santiago et al., 2018); youth, race, poor economic status and women's ignorance of their rights (Perera et al., 2018); or having dark skin (Grilo Diniz et al., 2018). However, an analysis of the main reasons and places of occurrence of this type of violence can reveal that OV is a type of structural violence. Structural violence is *one way of describing social arrangements that put individuals and populations in harm's way [...]; the arrangements are structural because they are embedded in the political and economic organization of our social world* (Farmer et al., 2006).

The main reason for OV is gender bias, in which women's right to choose is nullified and replaced (Jardim and Modena, 2018). Regarding sites of occurrence, OV occurs throughout the world. Evidence shows that it exists in countries such as Mexico (Castro and Savage, 2019; Castro and Frías, 2019; Santiago et al., 2018), Venezuela, Argentina, Brazil (Ishola et al., 2017; Jardim and Modena, 2018), India (Bhattacharya and Sundari Ravindran, 2018), Tanzania (Mselle et al., 2018), the Czech Republic (Begley et al., 2018), Ecuador (Brandão et al., 2018), Italy (Castro and Frías, 2019; Raval di et al., 2018; Scambia et al., 2018), the United States (Perera et al., 2018), and Nigeria (Ishola et al., 2017), among other places. The structural nature of OV makes the health professional who exercises it unaware of it and even normalizes this practice (Borges, 2018).

Practices characterized by OV have physical and psychological consequences for both the women who experience them and the health professionals who practise or witness them. Women have shown how their physical, sexual and psychological health has been negatively affected (Chattopadhyay et al., 2018), and a very meaningful experience in their lives has been transformed into a violent and negative one (Borges, 2018; McGarry et al., 2017). On the other hand, the literature suggests that personnel who witness this type of violence during childbirth may suffer from compassion fatigue or secondary traumatic stress in response to observing the traumas the woman experiences first-hand (Sadler et al., 2016). The WHO states that it is necessary to generate data related to respectful and disrespectful care practices, responsibility systems and valuable professional support, adding that health professionals need support

and training to ensure that pregnant women are treated with compassion and dignity (WHO, 2014).

It is important to note that no similar studies have been found in the literature; more specifically, there are no studies analysing health sciences students' perspectives regarding OV. Therefore, the main objective of this study was to evaluate the perceptions of health sciences students at Jaume I University (Universitat Jaume I) (Spain) regarding OV and to detect possible changes in these perceptions after an educational intervention.

## **Methods**

### **Design and sample**

A pre-post quasi-experimental study was conducted among health sciences students at Jaume I University (Universitat Jaume I) (Spain) between January and June 2019.

Returned questionnaires with up to 10% of items incomplete were excluded from the study analysis. A sample size calculation was performed using the GRANMO programme, which determined that a sample of 99 subjects was sufficient. The values considered for the calculation of the sample size included a confidence interval of 95%, highlighting an initial proportion of events of 0.1 percentage points and a loss to follow-up of 20%.

### **Variables and instruments**

The sociodemographic variables that were considered were age, gender, field (medicine, nursing), course, health experience in gynaecology and obstetrics services (yes, no), duration of experience (less than 1 year, between 1 and 4

years, more than 4 years), having been present at a birth (yes, no), duration of experience being present at births (less than 1 year, between 1 and 4 years, more than 4 years), personal experience with pregnancies and births (yes, no), time since pregnancy and birth (less than 1 year, between 1 and 4 years, more than 4 years).

The perception of OV was measured with an ad hoc questionnaire composed of 33 items that referred to OV practices and were divided into 4 key moments (before delivery, during delivery, in case of caesarean section and after delivery). These items were measured using a 5-point Likert scale (1 strongly disagree - 5 strongly agree). The questionnaire was developed by a group of 3 experts and was based on the Guía de Práctica Clínica de Atención al Parto Normal (Clinical Practice of Normal Birth Care Guide) (Ministerio de Sanidad, 2014). The internal consistency of the scale, measured with Cronbach's alpha, was 0.922 for the pre-intervention measurement and 0.975 for the post-intervention measurement.

The intervention consisted of an 8-hour seminar. This activity was composed of a theatrical performance on OV in the delivery room performed by "*The Other Part of the Theatre*" (1 hour); a master class on legal aspects presented by a lawyer specializing in health law (2 hours); a round table composed of professionals from the different fields, who contributed their experiences (4 hours); and another round table in which four volunteer mothers narrated their experiences of childbirth (1 hour). The session with the theatrical performance and the master class on legal issues was conducted on 03/07/2019. The round tables of the professionals and the mothers was held at a second session on 03/12/2019.

**Data collection**

Fieldwork was conducted in March 2019 after the launch of a seminar related to OV, in which students voluntarily enrolled. Data were collected through a self-completed survey administered before the students entered the seminar on 03/07/2019 and after the activity on 03/12/2019. This survey was accompanied by an explanation of the study objective and an explanation of its voluntary and anonymous nature.

**Analysis**

A descriptive analysis of the data was performed considering means, standard deviations and 95% confidence intervals for the quantitative variables and the distribution of frequencies and percentages were taken into account for the qualitative variables. For the bivariate analysis, applicability was determined using parametric tests, the Kolmogorov-Smirnov normality test and Levene test for the homogeneity of variances. After these conditions were confirmed as acceptable, Student's T test was applied for paired data, with the intention of detecting the effect of the change in the different measurements, and the Mann Whitney U and Kruskal Wallis tests were used to determine the relationships of the sociodemographic variables and perceptions of OV with the responses on the pre-intervention measurement. The analysis was carried out with the statistical package Statistical Package for Social Sciences (SPSS), version 21. A statistical significance level of  $p < 0.05$  was established.

**Ethical considerations**

The study was approved by the management of the Jaume I University (Universitat Jaume I) Nursing Research Group. The intervention was approved by the directorate of the Nursing Department and the dean of the School of Health Sciences of Jaume I University (Universitat Jaume I). Before data collection, the students received information about the objectives of the study as well as its methodology and the voluntary and anonymous nature of participation. The data collection tool did not include any personal data that could compromise the identity of the participants. The project was designed in accordance with the December 5 Protection of Personal Data and Guarantee of Digital Rights Organic Law 03/2018. In addition, the principles of the Declaration of Helsinki (charity, nonmaleficence, autonomy and justice) were respected. To respect the anonymity of the data and to match the first and second measurements, an ID was created consisting of the last two digits of the student's cell phone number, the last two digits of his or her National ID and his or her initials.

## Results

A total of 107 questionnaires were collected. The mean age of the students was 22.5 years ( $\pm 5.87$ ). Women represented 89.7% of the sample ( $n = 96$ ). A total of 86.9% ( $n = 93$ ) of the students belonged to the degree programme in nursing, and 28% ( $n = 30$ ) of the students had completed clinical practice in the gynaecology and obstetrics departments. Of the sample, 20.6% ( $n = 22$ ) had been present at a birth; only 4.7% ( $n = 5$ ) had been pregnant, and 2.8% ( $n = 3$ ) had given birth (Table 1).

The Kolmogorov-Smirnov test ( $p < 0.01$ ) and the Levene test ( $p < 0.05$ ) verified the appropriateness of the Student's t-test for paired data. Table 2 shows the descriptive analysis of the variables for the pre- and post-intervention measurements and the bivariate analysis. All analyses of the OV perception scale by paired data showed statistically significant differences between the pre- and post-intervention measures, except on the items related to performing a pelvic examination without consent ( $p = 0.368$ ); not preserving the privacy of the woman ( $p = 0.389$ ); not considering the woman's decision ( $p = 0.086$ ); taking pictures without permission ( $p = 0.379$ ); saying "*Stop complaining, it is not that bad*" ( $p = 0.181$ ); Separating the mother and new-born ( $p = 1.00$ ); and giving formula to the baby without the mother's consent ( $p = 0.320$ ).

Of the 33 items on the OV perception scale, only 24.24% ( $n = 8$ ) had no relationship with the sample's sociodemographic and control variables. The variables with the most statistically significant differences in relation to the OV perception scale were gender, course, and having been pregnant. Field (nursing vs. medicine) showed statistically significant differences on the items related to not offering measures for pain (nursing:  $m = 4.43$ ,  $SD = 0.85$ , medicine:  $m = 4.86$ ,  $SD = 0.36$ ,  $p = 0.047$ ); performing the Kristeller manoeuvre (nursing:  $m = 4.60$ ,  $SD = 0.75$ ; medicine:  $m = 3.92$ ,  $SD = 1.17$ ,  $p = < 0.01$ ); performing an episiotomy without anaesthesia (nursing:  $m = 4.46$ ,  $SD = 1.01$ ; medicine:  $m = 3.89$ ,  $SD = 1.05$ ;  $p = 0.033$ ); allowing skin-to-skin contact after the paediatric examination (nursing:  $m = 3.93$ ,  $SD = 1.35$ ; medicine:  $m = 3.15$ ,  $SD = 1.52$ ,  $p = 0.049$ ) and taking the new-born to the nursery (nursing:  $m = 3.85$ ,  $SD = 1.12$ ; medicine:  $m = 3.15$ ,  $SD = 1.14$ ,  $p = 0.025$ ) (Table 3).

The completion of rotations in obstetrics-gynaecology was statistically significant related to the variables on the OV perception scale: directing the position of the woman in labour (yes:  $m = 2.57$ ,  $SD = 1.33$ ; no:  $m = 1.95$ ,  $SD = 1.33$ ;  $p = 0.025$ ); performing genital shaving (yes:  $m = 3.80$ ,  $SD = 1.29$ ; no:  $m = 3.16$ ,  $SD = 1.46$ ;  $p = 0.041$ ) and convincing the woman to undergo a caesarean section to end the labour quickly and without pain (yes:  $m = 4.70$ ,  $SD = 0.79$ ; no:  $m = 4.38$ ,  $SD = 0.93$ ;  $p = 0.021$ ). Having been present at a childbirth was statistically significantly related to the following variable on the OV perception scale: Saying "you do not know how to push" (yes:  $m = 4.45$ ,  $SD = 1.01$ ; no:  $m = 4.84$ ,  $SD = 0.65$ ;  $p = <0.01$ ).

Table 4 shows the descriptive and comparative results for the control variables with respect to the items on the OV perception scale. Having given birth was statistically significantly related to the following items on the scale: performing routine genital shaving (yes:  $m = 1.00$ ,  $SD = 0.00$ ; no:  $m = 3.41$ ,  $SD = 1.40$ ,  $p = <0.01$ ), performing routine episiotomy (yes:  $m = 2.67$ ,  $SD = 0.57$ ; no:  $m = 4.44$ ,  $SD = 0.95$ ;  $p = <0.01$ ); saying "you do not know how to push" (yes:  $m = 2.33$ ,  $SD = 1.15$ ; no:  $m = 4.82$ ,  $SD = 0.62$ ;  $p = <0.01$ ); performing a caesarean section due to slow dilation (yes:  $m = 2.00$ ,  $SD = 1.00$ ; no:  $m = 4.09$ ,  $SD = 1.07$ ;  $p = <0.01$ ) and not allowing company in cases of instrumentation or caesarean section (yes:  $m = 3.33$ ,  $SD = 0.57$ ; no:  $m = 4.09$ ,  $SD = 1.07$ ;  $p = 0.02$ ). Having been pregnant was statistically significantly related to the following items: performing routine genital shaving (yes:  $m = 1.80$ ,  $SD = 1.78$ ; no:  $m = 3.42$ ,  $SD = 1.38$ ;  $p = 0.028$ ); performing a pelvic exam without consent (yes:  $m = 4.20$ ,  $SD = 1.30$ ; no:  $m = 4.83$ ,  $SD = 0.64$ ,  $p = 0.026$ ); encouraging the use of an epidural (yes:  $m = 4.40$ ,  $SD = 0.54$ ; no:  $m = 4.81$ ,  $SD = 0.68$ ,  $p = <0.01$ ); not

considering the woman's decision (yes:  $m = 4.40$ ,  $SD = 0.54$ ; no:  $m = 4.79$ ,  $SD = 0.62$ ;  $p = 0.01$ ); taking pictures without permission (yes:  $m = 4.00$ ,  $SD = 1.73$ ; no:  $m = 4.85$ ,  $SD = 0.62$ ;  $p = 0.015$ ); performing routine episiotomy (yes:  $m = 3.00$ ,  $SD = 0.70$ ; no:  $m = 4.46$ ,  $SD = 0.94$ ;  $p = <0.01$ ); saying "you do not know how to push" (yes:  $m = 3.40$ ,  $SD = 1.67$ ; no:  $m = 4.82$ ,  $SD = 0.62$ ;  $p = <0.01$ ); saying "stop complaining, it is not that bad" (yes:  $m = 3.80$ ,  $SD = 1.78$ ; no:  $m = 4.83$ ,  $SD = 0.62$ ;  $p = 0.034$ ); and performing a caesarean due to slow dilation (yes:  $m = 2.60$ ;  $SD = 1.51$ ; no:  $m = 4.10$ ,  $SD = 1.05$ ;  $p = 0.022$ ) (Table 4).

## Discussion

First, it is important to emphasize that students' participation in the proposed teaching activity was much greater than initially expected. The students seemed very motivated by the central theme, and their involvement was notable; therefore, the capture of individuals from the sample exceeded the calculated sample size. The high percentage of women in the sample (89.7%) should be noted; it may have occurred because women were especially motivated by the issue or because female representation is increasing in the health sciences (Bernalte-Martí, 2015). It is noteworthy that students in different years of their programme were equitably represented in the seminar, although more second- and fourth-year students than students of other years were in attendance. The representation of medical students was low; some possible reasons for this low attendance may be low dissemination of the activity among these students or the possibility that nursing students feel more linked to this type of practice (Olza-Fernández and Ruiz-Berdún, 2015). Because our sample was young in relation to the mean age (32.58 years) for maternity in Spain according to data

from the National Institute of Statistics (Instituto Nacional de Estadística - INE) («Edad Media a la Maternidad por orden del nacimiento según nacionalidad (española/extranjera) de la madre(1579)», s. f.), very few participants had a personal medical history of pregnancy or birth, although 30% of the sample had experience in gynaecology and obstetrics.

Regarding the comparison of the pre- and post-intervention measures by paired data, it is noteworthy that with the exception of a few items, all measures presented statistically significant differences. Even so, it is worth highlighting the low results found in the pre-intervention measurement. The Fortaleza document (World Health Organization, 1985) states that a family member chosen by the mother may accompany her during childbirth and throughout the postnatal period to promote her well-being; the new-born should stay with his/her mother whenever possible; immediate breastfeeding should be promoted; the dorsal position of the woman in lithotomy during dilation is not recommended; and the shaving of pubic hair, the administration of enemas, the systematic use of episiotomy and the early artificial rupture of membranes should not be performed as routine procedures. All these procedures were included in the questionnaire, and of them, the only one that was identified as violence pre-intervention was "*giving formula without the mother's consent*".

It should be noted that health science students should be trained in the latest available evidence (Aglen, 2016). Apparently, this does not occur in the field of obstetrics in Spain, highlighting the need for all women of child-bearing age to receive evidence-based care that is applied respectfully without neglecting the woman's opinions and preferences (Begley et al., 2018). Along the same lines, the Fédération Internationale de Gynécologie et d'Obstétrique (FIGO) adds that

*“Every woman and every baby should be protected from unnecessary interventions, practices and procedures that are not evidence-based, and any practices that are not respectful of their culture, bodily integrity, and dignity”* (International Federation of Gynaecology and Obstetrics et al., 2014). Special focus falls on the Kristeller manoeuvre, which, despite being contraindicated, continues to be put into practice (Borges, 2018; Fritz et al., 2017; Rubashkin et al., 2019) and was not recognized by the students as OV.

It is evident that there is a close relationship between ideological representations of the female gender and the existence of OV. The cultural image of women as reproductive and submissive serves as a precedent for the domination, control and abuse they experience in relation to their bodies and sexuality. As a consequence, women are nullified, and their rights to choose are replaced (Jardim and Modena, 2018). These assertions are corroborated by the results obtained in this study; when the responses to the OV scale were compared by gender, a large number of variables presented statistically significant differences, and in all of them, the perception of OV was higher among females.

Another feature of this OV is that it is rooted in a system that stands in the way of optimal health outcomes (Castro and Savage, 2019); thus, it also has a structural nature (Bhattacharya and Sundari Ravindran, 2018). In this way, the researchers assumed two facts that were confirmed through this study: a) the normalization of this type of violence according to the student's year of study, i.e., a lower perception of OV among more advanced students and a relationship between perceptions of OV and having participated in obstetric practices during study; and b) the normalization of this type of obstetric

practices in relation to the participant's personal experience with pregnancies and births (a decreased perception of OV after having been pregnant or given birth). A larger study is necessary to determine the degree of normalization and the normalization process; however, given these preliminary data, it is essential to change the training of health personnel, who should have a solid foundation in ethics and gender and human rights because emotional factors or burnout may be among the reasons for practising OV (Olza Fernández, 2013). The strain on health personnel is so high that many professionals have to abandon their job and even their profession (Beck and Gable, 2012). Therefore, public policies must direct attention towards humane and respectful treatment that is based on and supported by the latest available evidence. However, in order for this to happen, health personnel who work with pregnant women must abandon the traditional hierarchy and structure in which medical supervision implies a subordination of women's bodies and sexuality. This fact is further aggravated when all attention is paid to techniques, and the value of how people are treated is lost (Grilo Diniz et al., 2018; Mselle et al., 2018). Education that promotes respect and informs and raises awareness among future professionals, along with policies, guides, protocols and education, will eradicate OV (Brandão et al., 2018; Diaz-Tello, 2016; Grilo Diniz et al., 2018; Mselle et al., 2018; Sen et al., 2018). Education is a fundamental aspect for ending the normalization of OV in society; it approaches the problem from the root and will evolve until the rights of women are respected. The results should be taken with caution because there are some limitations. This is a quasi-experimental study without a control group, carried out in a single institution and, also, the post-test data were collected immediately after the intervention. In spite of these limitations, the

results obtained are interesting because they highlight the consideration of OV by health students and add value because they show that it is possible to change this perception with an educational intervention directed specifically at OV. Among the future lines of research, the authors propose designs that overcome these limitations, for example, considering a control group and long-term monitoring of students.

### **Conclusions**

According to the results of this study, health sciences students integrate the normalizations of OV during their studies. A formative activity aimed at making this type of violence visible and reflecting on OV helps to create awareness among students, making it possible for them to notice this type of violence and be able to identify it. It is noteworthy that from the beginning, the women in the study have perceived all the points raised on the OV scale as having higher OV; additionally, OV becomes normalized as a result of being present at a delivery, the progression of training (depending on the course) and obstetric experience itself, including pregnancies and births.

### **References**

- Aglen, B., 2016. Pedagogical strategies to teach bachelor students evidence-based practice: A systematic review. *Nurse Educ. Today* 36, 255-263. <https://doi.org/10.1016/j.nedt.2015.08.025>
- Beck, C.T., Gable, R.K., 2012. A Mixed Methods Study of Secondary Traumatic Stress in Labor and Delivery Nurses. *JOGNN - J. Obstet. Gynecol. Neonatal Nurs.* 41, 747-760. <https://doi.org/10.1111/j.1552->

6909.2012.01386.x

- Begley, C., Sedlicka, N., Daly, D., 2018. Respectful and disrespectful care in the Czech Republic: An online survey. *Reprod. Health* 15, 198. <https://doi.org/10.1186/s12978-018-0648-7>
- Bernalte-Martí, V., 2015. The minority of male in the nursing profession. Reflexions of their history, image and their evolution in Spain. *Enfermería Glob.* 14, 328-334.
- Bhattacharya, S., Sundari Ravindran, T.K., 2018. Silent voices: Institutional disrespect and abuse during delivery among women of Varanasi district, northern India. *BMC Pregnancy Childbirth* 18, 338. <https://doi.org/10.1186/s12884-018-1970-3>
- Borges, M.T., 2018. A Violent Birth: Reframing Coerced Procedures During Childbirth as Obstetric Violence. *Duke Law J.* 67, 827-862.
- Brandão, T., Cañadas, S., Galvis, A., de los Ríos, M.M., Meijer, M., Falcon, K., 2018. Childbirth experiences related to obstetric violence in public health units in Quito, Ecuador. *Int. J. Gynecol. Obstet.* 143, 84-88. <https://doi.org/10.1002/ijgo.12625>
- Castro, A., Savage, V., 2019. Obstetric Violence as Reproductive Governance in the Dominican Republic. *Med. Anthropol. Cross Cult. Stud. Heal. Illn.* 38, 123-136. <https://doi.org/10.1080/01459740.2018.1512984>
- Castro, R., Frías, S.M., 2019. Obstetric Violence in Mexico: Results From a 2016 National Household Survey. *Violence Against Women.* <https://doi.org/10.1177/1077801219836732>
- Chattopadhyay, S., Mishra, A., Jacob, S., 2018. 'Safe', yet violent? Women's experiences with obstetric violence during hospital births in rural Northeast

India. Cult. Heal. Sex. 20, 815-829.

<https://doi.org/10.1080/13691058.2017.1384572>

Diaz-Tello, F., 2016. Invisible wounds: obstetric violence in the United States.

Reprod. Health Matters 24, 56-64.

<https://doi.org/10.1016/j.rhm.2016.04.004>

Edad Media a la Maternidad por orden del nacimiento según nacionalidad

(española/extranjera) de la madre (1579) [WWW Document], s. f. URL

<http://www.ine.es/jaxiT3/Datos.htm?t=1579> (Accessed: 03.05.2019).

Farmer, P.E., Nizeye, B., Stulac, S., Keshavjee, S., 2006. Structural violence

and clinical medicine. PLoS Med. 3, e449.

<https://doi.org/10.1371/journal.pmed.0030449>

Fritz, J., Walker, D.M., Cohen, S., Angeles, G., Lamadrid-Figueroa, H., 2017.

Can a simulation-based training program impact the use of evidence based routine practices at birth? Results of a hospital-based cluster randomized

trial in Mexico. PLoS One 12, e0172623.

<https://doi.org/10.1371/journal.pone.0172623>

Grilo Diniz, C.S., Rattner, D., Lucas d'Oliveira, A.F.P., de Aguiar, J.M., Niy,

D.Y., 2018. Disrespect and abuse in childbirth in Brazil: social activism, public policies and providers' training. Reprod. Health Matters 26, 19-35.

<https://doi.org/10.1080/09688080.2018.1502019>

International Federation of Gynecology and Obstetrics, International

Confederation of Midwives, White Ribbon Alliance, International Pediatric Association, World Health Organization, 2014. Mother-baby friendly

birthing facilities. Int. J. Gynecol. Obstet.

<https://doi.org/10.1016/j.ijgo.2014.10.013>

- Ishola, F., Owolabi, O., Filippi, V., 2017. Disrespect and abuse of women during childbirth in Nigeria: A systematic review. *PLoS One* 12, e0174084. <https://doi.org/10.1371/journal.pone.0174084>
- Jardim, D.M.B., Modena, C.M., 2018. Obstetric violence in the daily routine of care and its characteristics. *Rev. Lat. Am. Enfermagem* 26, e3069. <https://doi.org/10.1590/1518-8345.2450.3069>
- McGarry, J., Hinsliff-Smith, K., Watts, K., McCloskey, P., Evans, C., 2017. Experiences and impact of mistreatment and obstetric violence on women during childbearing: A systematic review protocol. *JBISRIR-2016-Implement. Reports* 15, 620-627. <https://doi.org/10.11124/JBISRIR-2016-002950>
- Ministerio de Sanidad, 2014. Guía de Práctica Clínica sobre la Atención al Parto Normal, Internet. [https://doi.org/http://www.guiasalud.es/GPC/GPC\\_472\\_Part0\\_Normal\\_Osteba\\_compl.pdf](https://doi.org/http://www.guiasalud.es/GPC/GPC_472_Part0_Normal_Osteba_compl.pdf)
- Mselle, L.T., Kohi, T.W., Dol, J., 2018. Barriers and facilitators to humanizing birth care in Tanzania: Findings from semi-structured interviews with midwives and obstetricians. *Reprod. Health* 15, 137. <https://doi.org/10.1186/s12978-018-0583-7>
- Olza-Fernández, I., Ruiz-Berdún, D., 2015. Midwives experiences regarding obstetric violence, en: *Archives of Women's Mental Health*. <https://doi.org/10.1007/s00737-014-0488-6>
- Olza Fernández, I., 2013. PTSD and obstetric violence. *Midwifery Today Int. Midwife Spring*, 48-49.
- Perera, D., Lund, R., Swahnberg, K., Schei, B., Infanti, J.J., Darj, E., Lukasse,

- M., Bjørngaard, J.H., Joshi, S.K., Rishal, P., Koju, R., Pun, K.D., Wijewardena, K., Muzrif, M.M., Campbell, J.C., 2018. «When helpers hurt»: Women's and midwives' stories of obstetric violence in state health institutions, Colombo district, Sri Lanka. *BMC Pregnancy Childbirth* 18, 211. <https://doi.org/10.1186/s12884-018-1869-z>
- Pérez D'gregorio, R., 2010. Obstetric violence: A new legal term introduced in Venezuela. *Int. J. Gynecol. Obstet.* 111, 201-202. <https://doi.org/10.1016/j.ijgo.2010.09.002>
- Ravaldi, C., Skoko, E., Battisti, A., Cericco, M., Vannacci, A., 2018. Sociodemographic characteristics of women participating to the LOVE-THEM (Listening to Obstetric Violence Experiences THrough Enunciations and Measurement) investigation in Italy. *Data Br.* 19, 226-229. <https://doi.org/10.1016/j.dib.2018.04.146>
- Rubashkin, N., Torres, C., Escuriet, R., Dolores Ruiz-Berdún, M., 2019. “Just a little help”: A qualitative inquiry into the persistent use of uterine fundal pressure in the second stage of labor in Spain. *Birth.* <https://doi.org/10.1111/birt.12424>
- Sadler, M., Santos, M.J., Ruiz-Berdún, D., Rojas, G.L., Skoko, E., Gillen, P., Clausen, J.A., 2016. Moving beyond disrespect and abuse: addressing the structural dimensions of obstetric violence. *Reprod. Health Matters* 24, 47-55. <https://doi.org/10.1016/j.rhm.2016.04.002>
- Santiago, R.V., Monreal, L.A., Rojas Carmona, A., Domínguez, M.S., 2018. «If we're here, it's only because we have no money.» discrimination and violence in Mexican maternity wards. *BMC Pregnancy Childbirth* 18, 244. <https://doi.org/10.1186/s12884-018-1897-8>

Scambia, G., Viora, E., Chiantera, A., Colacurci, N., Vicario, M., 2018.

“Obstetric violence”: Between misunderstanding and mystification. Eur. J.

Obstet. Gynecol. Reprod. Biol. Sep, 331.

<https://doi.org/10.1016/j.ejogrb.2018.06.012>

Sen, G., Reddy, B., Iyer, A., 2018. Beyond measurement: the drivers of

disrespect and abuse in obstetric care. Reprod. Health Matters 26, 6-18.

<https://doi.org/10.1080/09688080.2018.1508173>

World Health Organization, 2014. The prevention and elimination of disrespect and abuse during facility-based childbirth., World Health Organization.

<https://doi.org/WHO/RHR/14.23>

World Health Organization, 1985. Appropriate technology for birth. Lancet 24,

436-437.

**Table 1. Sociodemographic and control variables of the sample.**

	Total		Nursing		Medicine	
	n	%	n	%	n	%
Gender						
Male	11	10.3	10	9.3	1	0.9
Female	96	89.7	83	77.6	13	12.1
Course						
First	10	9.3	8	7.5	2	1.9
Second	46	43.0	40	37.4	6	5.6
Third	10	9.3	7	6.5	3	2.8
Fourth	40	37.4	37	34.6	3	2.8
Health experience in gynaecology and obstetrics services						
Yes	30	28.0	25	23.5	5	4.7
No	77	72.0	68	63.6	9	8.4
Duration of experience						
Less than 1 year	16	15.0	12	11.2	4	3.7
Between 1 and 4 years	13	12.1	12	11.2	1	0.9
More than 4 years	1	0.9	1	0.9	-	-
Having been present at a birth						
Yes	22	20.6	20	18.7	2	1.9
No	85	79.4	73	68.2	12	11.2
Personal experience with pregnancies						
Yes	5	4.7	5	4.7	-	-
No	102	95.3	88	82.2	14	13.1
Time since pregnancy						
Between 1 and 4 years	2	1.9	2	1.9	-	-
More than 4 years	3	2.8	3	2.8	-	-
Personal experience with birth						
Yes	3	2.8	3	2.8	-	-
No	104	97.2	90	84.1	14	13.1
Time since birth						
More than 4 years	3	2.8	3	2.8	-	-

**Table 2. Results of student perceptions of obstetric violence.**

Item no	Variable	Pre-intervention measure				Post-intervention measure				p-value <sup>a</sup>
		n	M	SD	IC 95%	n	M	SD	IC 95%	
1	Inserting an intravenous channel	105	2.15	1.18	1.92-2.38	98	4.00	1.00	3.80-4.20	<0.01
2	Directing the woman's position	105	2.12	1.35	1.86-2.39	98	3.97	1.21	3.73-4.21	<0.01
3	Accelerate the birthing process artificially	105	3.34	1.31	3.09-3.60	98	4.61	0.83	4.45-4.78	<0.01
4	Administering routine enemas	106	3.12	1.37	2.86-3.39	98	4.65	0.79	4.50-4.81	<0.01
5	Performing routine amniorrhexis	106	4.08	0.93	3.90-4.27	98	4.82	0.66	4.68-4.95	<0.01
6	Performing routine genital shaving	105	3.34	1.44	3.06-3.62	98	4.57	0.76	4.42-4.72	<0.01
7	Immobilizing the woman	106	4.49	0.85	4.33-4.65	98	4.87	0.62	4.74-4.99	<0.01
8	Performing a pelvic exam without consent	106	4.8	0.69	4.67-4.94	98	4.87	0.62	4.74-4.99	0.368
9	Not offering measures for pain	106	4.49	0.81	4.33-4.65	98	4.72	0.80	4.56-4.88	<0.01
10	Encouraging the use of an epidural	106	3.94	0.99	3.75-4.13	98	4.58	0.73	4.44-4.73	<0.01
11	Not preserving privacy	106	4.79	0.68	4.66-4.92	98	4.84	0.63	4.71-4.96	0.389
12	Convincing the woman to undergo a c-section to end labour quickly and without pain	106	4.47	0.91	4.30-4.65	98	4.83	0.65	4.69-4.96	<0.01
13	Not considering the woman's decision	106	4.77	0.62	4.65-4.89	98	4.88	0.59	4.76-5.00	0.086
14	Taking pictures without permission	106	4.81	0.71	4.67-4.95	98	4.86	0.71	4.71-5.00	<0.01
15	Enforcing the lithotomy position	103	3.27	1.30	3.02-3.53	98	4.53	0.84	4.36-4.70	0.379
16	Allowing accompaniment during the second stage	105	2.95	1.55	2.65-3.25	95	4.12	1.27	3.86-4.38	<0.01
17	Performing routine episiotomy	102	4.39	0.98	4.20-4.59	98	4.81	0.68	4.67-4.94	<0.01
18	Saying "You do not know how to push"	105	4.75	0.75	4.61-4.90	98	4.88	0.64	4.75-5.01	0.05
19	Performing the Kristeller manoeuvre	104	4.52	0.83	4.36-4.68	98	4.85	0.63	4.72-4.97	<0.01
20	Performing an episiotomy without anaesthesia	101	4.41	1.02	4.20-4.61	98	4.76	0.71	4.61-4.90	<0.01

21	Prohibiting eating and drinking	105	3.51	1.17	3.29-3.74	98	4.62	0.76	4.47-4.78	<0.01
22	Not providing covering/heating during delivery	105	4.43	0.85	4.26-4.59	97	4.70	0.85	4.53-4.87	0.01
23	Saying "Stop complaining, it is not that bad"	105	4.78	0.73	4.64-4.92	98	4.88	0.64	4.75-5.01	0.181
24	Not letting the woman shout	104	4.51	0.89	4.34-4.68	98	4.78	0.76	4.62-4.93	0.012
25	Performing a caesarean section due to slow dilation	105	4.03	1.12	3.81-4.25	98	4.76	0.62	4.63-4.88	<0.01
26	Performing an emergency caesarean section without consent	104	4.38	1.07	4.17-4.58	98	4.77	0.63	4.64-4.89	<0.01
27	Not allowing accompaniment in cases of instrumentation or caesarean section	105	4.49	0.84	4.32-4.65	98	4.82	0.69	4.68-4.96	<0.01
28	Immediately cutting the cord	104	3.51	1.30	3.26-3.76	98	4.35	0.89	4.17-4.53	0.018
29	Suturing a tear without anaesthesia	104	4.39	0.91	4.22-4.57	98	4.74	0.80	4.58-4.91	<0.01
30	Separating the mother and new-born	105	4.90	0.57	4.78-5.01	98	4.86	0.65	4.73-4.99	1.00
31	Allowing skin-to-skin contact after the paediatric examination	105	3.84	1.38	3.57-4.11	98	4.66	0.77	4.51-4.82	<0.01
32	Taking the baby to the nursery	105	3.76	1.14	3.54-3.98	97	4.58	0.85	4.41-4.75	<0.01
33	Giving formula without the mother's consent	105	4.74	0.77	4.59-4.89	98	4.82	0.67	4.68-4.95	0.320

<sup>a</sup> T of Student for related samples.

**Table 3. Descriptive and comparative results of the sociodemographic variables and the obstetric violence scale.**

Item no	Gender				p-value <sup>a</sup>	Field				p-value <sup>a</sup>	Course								p-value <sup>b</sup>
	Male		Female			Nursing		Medicine			First		Second		Third		Fourth		
	M	S	M	S		M	S	M	S		M	S	M	S	M	S	M	S	
2	1.90	1.37	2.58	1.18	-	2.99	1.33	2.66	1.55	-	1.30	0.48	2.44	1.44	2.40	1.46	1.98	1.52	-
6	2.70	1.33	3.44	1.44	-	3.42	1.44	2.73	1.36	-	3.44	1.73	3.40	1.40	3.00	1.82	3.52	1.26	-
8	4.40	1.38	4.85	0.48	-	4.78	0.77	4.92	0.22	-	5.00	0.08	4.07	0.87	4.08	0.48	4.86	0.62	0.02
9	4.00	1.25	4.57	0.45	-	4.48	0.88	4.83	0.30	0.04	4.74	0.43	4.88	0.88	4.84	0.58	4.87	0.82	0.04
10	3.00	0.51	4.07	0.97	0.02	3.99	0.93	4.03	0.30	-	3.06	0.86	3.06	0.60	4.09	0.88	3.82	0.22	-
11	4.30	1.09	4.85	0.43	-	4.77	0.70	5.00	0.00	-	4.88	0.48	4.07	0.93	4.09	0.37	4.77	0.77	-
12	3.80	1.03	4.88	0.80	0.04	4.95	0.64	4.99	0.49	-	4.51	0.58	4.81	0.74	4.44	0.40	4.08	0.80	-
13	4.40	0.98	4.85	0.60	0.03	4.76	0.92	4.86	0.26	-	4.74	0.86	4.86	0.84	4.04	0.76	4.89	0.69	-
14	4.40	1.03	4.86	0.51	-	4.88	0.60	5.00	0.00	-	5.00	0.08	4.09	0.90	4.01	0.56	4.65	0.01	0.05
16	4.40	1.05	3.50	1.55	-	2.99	1.57	2.66	1.33	-	2.66	1.66	2.55	2.55	2.55	3.44	1.43	0.04	0.06
17	3.20	1.03	4.88	0.80	0.01	3.53	1.54	4.07	1.07	-	4.19	0.59	4.99	0.82	4.03	1.30	4.07	0.70	-
18	3.00	1.05	4.86	0.76	0.01	4.72	0.80	5.00	0.00	-	4.93	0.86	4.90	0.65	4.27	0.69	4.89	0.99	-
19	4.40	0.88	4.88	0.58	-	4.67	0.91	3.91	1.01	<0.01	4.17	0.39	4.95	0.75	4.85	0.55	4.85	0.50	<0.01
20	3.40	1.03	4.00	0.40	0.00	4.13	1.31	3.91	1.01	0.00	4.18	0.68	4.06	0.62	4.34	1.41	4.01	0.99	-

0	6	5	4	9	03	4	0	8	0	03	5	7	4	9	3	9	4	0	
	7	8	8	3	5	6	1	9	5	3	6	2	0	7	8	1	8	3	
2	2.	0.	3.	1.	<0	3.	1.	3.	1.	-	2.	0.	3.	1.	4.	1.	3.	1.	0.
1	3	6	6	1	.0	5	1	3	5		7	9	2	1	1	1	8	0	01
	0	7	4	4	1	4	2	1	4		8	7	2	8	0	0	5	9	1
2	4.	0.	4.	0.	0.	4.	0.	5.	0.	-	4.	0.	4.	0.	4.	0.	4.	0.	-
2	0	9	4	8	04	4	7	0	0		2	8	3	9	9	3	4	8	
	0	4	7	3	8	0	7	0	0		2	3	8	1	0	1	3	7	
2	4.	1.	4.	0.	-	4.	0.	5.	0.	-	4.	0.	4.	0.	5.	0.	4.	0.	0.
3	4	2	8	6		7	7	0	0		8	3	8	6	0	0	7	7	02
	0	6	2	5		5	7	0	0		9	3	2	8	0	0	5	0	4
2	4.	1.	4.	0.	0.	4.	0.	4.	0.	-	4.	0.	4.	0.	4.	0.	4.	0.	-
4	0	2	5	8	04	4	9	9	2		4	7	5	9	6	6	4	9	
	0	4	6	9	0	6	3	2	9		4	2	7	2	0	9	8	3	
2	3.	1.	4.	1.	-	4.	1.	4.	0.	-	3.	0.	3.	1.	3.	1.	4.	1.	-
5	4	5	0	0		0	1	2	9		8	9	9	1	7	4	2	0	
	0	7	9	5		0	4	3	2		9	2	8	3	0	1	3	7	
2	3.	1.	4.	0.	0.	3.	0.	4.	1.	-	4.	0.	4.	0.	4.	0.	4.	0.	-
7	8	4	5	7	04	5	7	0	1		3	8	3	9	5	7	7	6	
	0	0	6	4	0	4	9	8	1		3	6	6	8	0	0	0	4	
2	3.	1.	4.	0.	<0	4.	0.	4.	0.	-	4.	1.	4.	0.	4.	0.	4.	0.	-
9	6	1	4	8	.0	3	9	5	6		4	0	4	7	6	6	3	9	
	0	7	8	5	1	7	4	8	6		4	1	5	9	0	9	5	4	
3	4.	1.	4.	0.	0.	4.	0.	5.	0.	-	5.	0.	4.	0.	5.	0.	4.	0.	-
0	5	2	9	4	01	8	6	0	0		0	0	8	6	0	0	8	6	
	0	6	4	3	7	8	0	0	0		0	0	7	2	0	0	8	4	
3	3.	1.	3.	1.	-	3.	1.	3.	1.	0.	3.	0.	4.	1.	3.	1.	3.	1.	0.
1	1	5	9	3		9	3	1	5	04	0	8	0	4	3	4	8	3	01
	0	9	2	5		3	5	5	2	9	0	6	9	1	0	9	5	6	8
3	3.	0.	3.	1.	0.	3.	1.	3.	1.	0.	3.	0.	3.	0.	3.	1.	3.	1.	-
2	0	9	8	1	01	8	1	1	1	02	5	5	9	9	7	2	6	3	
	0	4	4	4	2	5	2	5	4	5	6	2	6	7	0	5	0	9	
3	4.	1.	4.	0.	-	4.	0.	4.	0.	-	4.	0.	4.	0.	5.	0.	4.	0.	0.
3	3	3	7	6		7	8	9	2		7	6	8	6	0	0	6	8	01
	0	4	9	8		2	2	2	7		8	6	4	3	0	0	5	0	5

<sup>a</sup> Mann-Whitney U test

<sup>b</sup> Kruskal-Wallis test

**Table 4. Descriptive and comparative results of the control variables and the obstetric violence scale.**

Item no	Personal experience with pregnancies					Personal experience with birth				
	Yes		No		p-value <sup>a</sup>	Yes		No		p- value <sup>a</sup>
	M	SD	M	SD		M	SD	M	SD	
2	2.80	1.78	2.12	1.14	-	3.00	2.00	2.13	1.15	-
6	1.80	1.78	3.42	1.38	0.028	1.00	0.00	3.41	1.40	<0.01
8	4.20	1.30	4.83	0.64	0.026	3.67	1.52	4.83	0.64	-
9	4.00	1.22	4.51	0.79	-	3.67	1.52	4.51	0.79	-
10	3.20	1.64	3.98	0.94	-	3.00	1.73	3.97	0.96	-
11	4.40	0.54	4.81	0.68	<0.01	4.33	0.57	4.81	0.68	-
12	4.20	1.30	4.49	0.89	-	4.00	1.73	4.49	0.88	-
13	4.40	0.54	4.79	0.62	0.010	4.33	0.57	4.79	0.62	-
14	4.00	1.73	4.85	0.62	0.015	3.33	2.08	4.85	0.61	-
16	2.40	1.94	2.98	1.54	-	2.00	1.73	2.98	1.55	-
17	3.00	0.70	4.46	0.94	<0.01	2.67	0.57	4.44	0.95	<0.01
18	3.40	1.67	4.82	0.62	<0.01	2.33	1.15	4.82	0.62	<0.01
19	4.00	1.00	4.55	0.82	-	4.00	1.00	4.53	0.83	-
20	3.80	1.78	4.44	0.97	-	3.00	2.00	4.45	0.96	-
21	2.80	1.64	3.55	1.14	-	2.67	1.15	3.54	1.17	-
22	4.20	1.30	4.44	0.83	-	3.67	1.52	4.45	0.82	-
23	3.80	1.78	4.83	0.62	0.034	3.00	2.00	4.83	0.61	-
24	4.00	1.41	4.54	0.86	-	3.33	1.52	4.54	0.85	-
25	2.60	1.51	4.10	1.05	0.022	2.00	1.00	4.09	1.07	<0.01
27	4.00	1.00	4.51	0.83	-	3.33	0.57	4.52	0.82	0.020
29	3.80	1.64	4.42	0.87	-	3.33	2.08	4.43	0.86	-
30	4.80	0.44	4.90	0.57	-	4.67	0.57	4.90	0.57	-
31	4.00	1.73	3.83	1.37	-	3.67	2.30	3.84	1.37	-
32	4.20	0.83	3.74	1.16	-	4.00	1.00	3.75	1.15	-
33	4.00	1.73	4.78	0.69	-	3.33	2.08	4.78	0.68	-

<sup>a</sup> Mann-Whitney U test