



City Research Online

City St George's, University of London

Citation: Hochberg, M. E. & Noble, R. (2017). A framework for how environment contributes to cancer risk. *Ecology Letters*, 20(2), pp. 117-134. doi: 10.1111/ele.12726

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/24706/>

Link to published version: <https://doi.org/10.1111/ele.12726>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

A framework for how novel environments contribute to cancer risk

Michael E. Hochberg^{1,2} & Robert J. Noble¹

1 Institut des Sciences de l'Évolution de Montpellier, Université de Montpellier, Place E. Bataillon, CC065, 34095 Montpellier Cedex 5, France

2 Santa Fe Institute, 1399 Hyde Park Rd., Santa Fe, NM 87501, USA

Type of article: Ideas and Perspectives

Running title: Novel environments and cancer

Keywords: Aging, anthropogenic impact, cancer risk, environment, epidemiology, evolutionary mismatch, longevity, body size, global change, modern lifestyles, mutagens, pathogens

Statement of authorship: MEH conceived the study, wrote the manuscript, designed the figures, and contributed to the figures and boxes; RJN performed the modelling work, designed and produced the figures, and contributed to the boxes and revisions.

Data accessibility statement: We confirm that, should our manuscript be accepted, the data supporting our results will be archived in an appropriate public repository.

Content: Abstract (192 words); Main text (7836 words); 236 References; 1 Glossary; 4 Figures; 4 Boxes

Author for correspondence: Michael Hochberg, ISEM - UMR 5554, Université de Montpellier, CC065, Place E. Bataillon, 34095 Montpellier Cedex 5, France. Telephone: +33 4 67143667. Email: mhochber@univ-montp2.fr

Abstract

Evolutionary theory explains why metazoan species are largely protected against the negative fitness effects of cancers. Nevertheless, cancer is often observed at high incidence across a range of species. Although there are many challenges to quantifying cancer epidemiology and assessing its causes, we claim that most modern-day cancer in animals--and humans in particular-- is due to environments deviating from central tendencies of distributions that have prevailed during cancer resistance evolution. Such novel environmental conditions may be natural and/or of anthropogenic origin, and may interface with cancer risk in numerous ways, broadly classifiable as those: increasing organism body size and/or life span, disrupting processes within the organism, and changing germline. We argue that anthropogenic influences, in particular, explain much of the present-day cancer risk across life, including in humans. Based on a literature survey of animal species and a parameterized mathematical model for humans, we suggest that combined risks of all cancers in a population beyond $c.$ 5% can almost invariably be explained by the influence of novel environments. Our framework provides a basis for understanding how natural environmental variation and human activity impact cancer risk, with potential implications for species ecology.

Introduction

Cancer is a pervasive threat to many multicellular organisms, and vulnerable populations are expected to evolve cancer prevention and suppression (hereafter ‘resistance’). Indeed, a large array of resistance mechanisms exist, including tissue architecture, cell cycle regulation and differentiation, DNA mismatch repair, apoptosis, immune responses, and replicative aging (Greaves 2000; DeGregori 2011; Reinhardt and Schumacher 2012; Campisi 2013). Despite the likely deep evolutionary history in the establishment and reinforcement of these processes, cancers occur across the tree of life, and sometimes at high incidence (Aktipis et al. 2015). These contradictory observations beg for explanation, specifically regarding the role of environments in promoting or suppressing cancer, and the implications of dynamic cancer risks for species ecology and evolution.

Much of our understanding of cancer comes from human tumour cells, either isolated *de novo* from biopsies, or studied on well characterized cell lines *in vitro*, or as allografts in immunodeficient laboratory mice. Only recently has increased emphasis been given to investigating cancers over a broad range of taxa (comparative oncology, see Glossary, (Schiffman et al. 2015; Nunney et al. 2015)). Although our knowledge remains largely restricted to humans, most cancers are thought to emerge through a stepwise mutational process, resulting in the Hallmarks of Cancer (Hanahan and Weinberg 2000; Hanahan and Weinberg 2011). A central prediction of the multistage theory of carcinogenesis (Glossary) is that, all else being equal, lifetime risk of any form of cancer (hereafter ‘cancer risk’; Glossary) should correlate with the lifetime number of stem cell divisions (Box 1). Perhaps surprisingly, cancer risk across mammal species shows no significant trend with longevity or with body mass (Peto et al. 1975; Peto 1977). There are numerous explanations for what has come to be known as Peto’s paradox (Glossary) (e.g., (Nunney 1999; Leroi et al. 2003; Nagy

et al. 2007; Caulin and Maley 2011; DeGregori 2011; Nunney 2013; Maciak and Michalak 2015)), most being based on differential selection for phenotypic traits that reduce cancer risk in more massive and/or longer-lived species.

Whereas considerable study addresses Peto's paradox, few authors have recognized an additional feature of the cancer risk data: most surveyed species and taxonomic groups have lifetime cancer risks of less than *c.* 5%, and for many, cancer is very rarely observed, if at all (e.g., (Effron et al. 1977; Varki 2000; Abegglen et al. 2015)). To the best of our knowledge and when sufficiently investigated, high lifetime cancer risk (conservatively defined here as greater than *c.* 5% of a population suffering cancer-related negative effects on performance, health, survival, and/or fitness during their lifetimes) is observed only in populations that live in captivity or in polluted areas, or that are subject to cancer-promoting infections, founder effects, inbreeding, or selective breeding. Even in the human population where cancers account for *c.* 11-25% of mortality (Ferlay et al. 2015), and as will be described below, the majority of cases are associated with present-day human ecology, including survival to older ages, modern lifestyles, parasitic infections, and mutagenic exposure.

Previous study has discussed how evolutionarily novel environments (Glossary) are a likely source of many human diseases (Gluckman and Hanson 2004; Gluckman et al. 2011), and associations between environments and cancer across species (e.g., (Newman and Smith 2006; McAloose and Newton 2009)) and in humans (Irigaray et al. 2007). We go significantly beyond these by proposing a framework for how environments, and specifically novel environments, affect cancer risk across species and in the human population. We argue that many, probably most cancers across animal species, and particularly in humans, result from environmental states that increase cancer risk above baseline levels occurring in native habitats (Figure 1). These baseline levels reflect long-term adaptations to prevailing

exogenous (e.g., temperature, physical habitat and biotic interactions) and endogenous (e.g., behaviour and physiology) environmental conditions. Environments obviously vary over different spatial and temporal scales, and evolutionary theory predicts how a species adapts to varying selection pressures (e.g., (Chevin et al. 2010; Tuomainen and Candolin 2011)). Importantly, trait evolution can be slow relative to the time scales of environmental variation and behavioural responses (Tuomainen and Candolin 2011), meaning that the absolute fitness associated with a trait may vary considerably without a significant evolutionary response (so called “evolutionary mismatches”). This is particularly relevant to cancers, where specific costly adaptations are unlikely to obtain in response to the diversity of possible transient and spatially limited environmental insults. Moreover, due to the complexity of cancer genetics (Blekhman et al. 2008; Hindorff et al. 2011), and weak selection against certain cancer sensitivity mutations (Frank 2004a), the fixation of resistance across all cancer-risk loci is unlikely, even in the long term (Maher et al. 2013).

Deviations from native habitat conditions may take the form of idiosyncratic pulses, sustained temporal variation, directional trends, or shifts to new states. Given heterogeneity at different spatial and temporal scales, environmental exposure and associated cancer risk are expected to vary within a population. Although it is difficult to accurately estimate baseline cancer risks and the environments under which these prevailed, we *can* identify putative environmental drivers of increased risk by focussing on species (or populations) showing unexpectedly high overall levels of cancer (i.e., greater than the conservative benchmark level of 5%). Humans have influenced ecosystems since the Late Pleistocene (Boivin et al. 2016), and we claim that as a result of the intensification and globalization of these effects, the environments experienced by many present-day metazoan species differ significantly from their pre-anthropogenic states (Eriksson 2013), and consequently many, possibly most instances of high cancer risk are attributable to human impacts.

We begin by reviewing the theoretical and empirical basis for how evolutionary adaptations limit cancer as a life threatening and fitness reducing disease. This sets the stage for understanding how environmental deviations from native conditions, typically acting on shorter time scales than significant evolutionary responses, affect individual biology and ecology, which in turn, impacts individual cancer risks. There are nevertheless numerous hurdles in quantifying cancer epidemiology and assessing relationships between environment and cancer, and we discuss five such limitations. We then present a framework where novel environments impact cancer risk via one or more of three basic biological levels: those increasing body size and/or longevity, those disrupting within-organism processes, and those resulting in heritable genetic change. We finally develop and analyse an empirically parameterized mathematical model to deduce overall cancer incidence in recent and ancient humans, which also allows estimation of how different components of modern (novel) environments have increased cancer risk in our species. Our framework provides a basis for understanding the relevance of environments to cancer across life, uncovers a number of important questions regarding the implications of cancer dynamics for ecological communities and evolutionary change, and makes predictions for future research.

The evolutionary theory of cancer resistance

Somatic cells in metazoans exhibit an array of cooperative behaviours that contribute to the fitness of the whole organism. Obtaining the benefits of cooperation without paying the costs would have provided the context for cells to effectively cheat (Glossary). Cancer is a prime example of cheating in the metazoa (Aktipis et al. 2015), whereby somatic cells grow and invade local and distant tissues (invasive carcinoma and metastasis, respectively). Failure of higher-level physiological systems to prevent this process or keep it in check would have had

substantial fitness consequences for the host organism, resulting in selection for costly cancer resistance mechanisms (Crespi and Summers 2005; Merlo et al. 2006; Casás-Selves and Degregori 2011; Greaves and Maley 2012; Aktipis et al. 2015). The evolution of some of these processes appears to have occurred deep in the metazoan phylogeny, where cancer risk and its reduction were associated with the emergence of “gatekeeper” genes (Domazet-Loso and Tautz 2010), possibly accompanied (and followed) by purifying selection against cancer-prone genotypes (Thomas et al. 2003) (but see (Blekhman et al. 2008)). With subsequent selection for larger body plans and longer life spans, additional, sometimes apparently species-specific, mechanisms have emerged (Gorbunova et al. 2014).

The evolutionary theory of cancer resistance makes two central predictions. First, *taxonomic-group or species-specific* cancer suppression has evolved in response to environmental challenges (e.g., UV radiation, temperature, diet, and exposure to competitors, predators and parasites) and coevolved with metabolic processes, mutation rates, characteristics of tissue and organ microenvironments (Glossary), and life history traits such as body size and life span. Theory suggests that if there is positive selection acting on traits that also render individuals more susceptible to disease, then the positive trait is more likely to evolve if the marginal costs of augmented disease (cancer) resistance are sufficiently low (Hochberg et al. 1992; Boddy et al. 2015; Kokko and Hochberg 2015).

Most empirical support for this first prediction comes from studies on the body size or longevity extremes that generate Peto’s paradox. For example, carcinoma in long-lived captive naked mole rats has been observed in only a few individuals out of thousands examined (Buffenstein 2008; Taylor et al. 2016; Delaney et al. 2016), and it remains to be determined whether inbreeding or laboratory conditions may have contributed to these cases, which were exclusively middle-aged and older individuals. Study indicates that this species

has reinforced resistance against various diseases, including cancers (e.g., (Tian et al. 2013)), and that occasional cancer may have both an evolutionary and ecological basis (Hochberg et al. 2016). Other work on captive African and Asian elephants reveals that they have 20 copies of the TP53 tumour suppressor gene, suggesting that the total stem cell population in an elephant can tolerate numerous mutations with little risk of any single stem cell lineage transforming into a fitness-threatening neoplasm (Abegglen et al. 2015; Sulak et al. 2016). Lifetime cancer risk in captive elephants is estimated at 3-5% (Elephant Encyclopedia 1995-2012. www.elephant.se) (Abegglen et al. 2015).

Although the evolutionary theory of cancer resistance is consistent with observations of low incidence across species and with reinforced protective mechanisms in those species that tend to generate Peto's paradox (i.e., the most massive and longest lived; (Seluanov et al. 2009; Gorbunova et al. 2014; Abegglen et al. 2015; MacRae et al. 2015), but see e.g., (Rozhok and DeGregori 2016)), complementary approaches such as phylogenetic analyses are uncommon (Gomes et al. 2011; Keane et al. 2015). For example, Gomes and colleagues (Gomes et al. 2011) compared and contrasted telomere length and telomerase activity across more than 60 mammal species. They propose that the capacity of ancestral adult mammals to suppress telomerase has been lost in certain short-lived, small mammal species. Their results are consistent with the prediction that replicative aging (shorter telomeres and repressed telomerase activity) tends to protect the organism from the fitness-reducing effects of cancer as a correlated response to larger body sizes and longer life spans (Weinstein and Ciszak 2002) (but see (Seluanov et al. 2007; Eisenberg 2011)).

The second prediction of evolutionary theory is that, whereas no correlation is expected between species over the full range of metazoan taxa, variation *within a species* should reveal positive associations between life history traits (particularly longevity) and cancer risk (Box

1). This is because selection on life history - cancer resistance tradeoffs is expected to be too weak (due, for example, to cancer emerging late in life) over ecological time scales of a few generations or less, or there is simply little or no additive genetic variation upon which selection can work. Limited support for this prediction comes from the correlation of cancer incidence with body mass across dog breeds (Fleming et al. 2011) and with human height (Shors et al. 2001; Green et al. 2011; Kabat et al. 2014; Jiang et al. 2015). An alternative, untested explanation for these patterns is that life history traits are in linkage disequilibrium with specific mutations increasing (or decreasing) cancer risk.

Similar to body size within a species, cancer incidence and mortality tend to correlate with age (in humans (Martincorena and Campbell 2015); in dogs (Fleming et al. 2011); but see (Li et al. 1996; Pompei and Wilson 2001)). This is not surprising since classic theory of the evolution of aging and senescence (Medawar 1952; Williams 1957; Hamilton 1966) leads to the prediction that cancer resistance will be generally favoured in an inverse age-dependent manner (e.g., (DeGregori 2011; Hochberg et al. 2013)), but this effect (increasing cancer risk for longer lived individuals within a species) is also predicted to be a straightforward consequence of more time for (multistage) cancer to obtain, and for emerging tumours to grow and constitute risks to individual performance and survival. Another explanation for within-species patterns is that cancer risk is somehow pleiotropically associated with genetically heritable differences in height or longevity (e.g., (Cournil and Kirkwood 2001)), although we are not aware of any supporting evidence.

Challenges to how we perceive and quantify cancer

Despite a plethora of anecdotal observations of cancer in animals and, more rarely, epidemiological studies in natural populations, the actual importance of cancers (altered

behaviours, reduced reproduction, synergisms with other diseases, morbidity, mortality) to demography, population dynamics, intra- and interspecific interactions, spatial distributions, and/or genomic evolution has seldom been demonstrated for any species in the wild[1]. When cancer is assessed on wild animals, it is usually semi-quantitative, recorded from individuals in captivity (Ratcliffe 1933), and/or typically for only one sampled population at a single time point (but see (Lair et al. 2015)).

We present five non-mutually exclusive factors that limit our ability to accurately assess the causes of cancer, and the importance of cancer to the performance, survival or Darwinian fitness of the individual, and by extension, the ecological and evolutionary dynamics of populations.

Sampling. Numerous biases exist in interpreting epidemiological data on cancer (e.g., for humans see (Doll and Peto 1981)), that also limit our ability to generalize findings about cancer in the wild. First and foremost, accurate field sampling in natural, near-pristine habitats is highly unusual (McAloose and Newton 2009), and may even be infeasible should individuals harbouring a cancer have altered behaviours (Vittecoq et al. 2013; Vittecoq et al. 2015). In addition, quantification of epidemiological parameters such as incidence, morbidity and mortality is potentially biased by (1) small sample sizes (eTable 1 in (Abegglen et al. 2015)), (2) reports with no follow-up or associated study of hypothetical explanations (Daoust et al. 1991), (3) the underreporting of studies revealing little or no observed cancer, (4) oversampling of species that are relatively easy to sample and examine (e.g., humans, domestic and captive mammals), (5) oversampling of emblematic species (Aguirre and Lutz 2004), (6) populations suspected to be exposed to carcinogens (Hueper 1963), (7) oversampling of species showing unexpectedly high (Lair et al. 2015) or low (Buffenstein 2008) cancer incidence, (8) oversampling of animals attracting attention through aberrant

behaviours, external signs of cancer, or found in atypical locations (e.g., (Newman and Smith 2006)), and (9) paleopathological analyses indicating high incidence, but that cannot accurately ascribe associated health impacts (Nerlich et al. 2006).

Another form of sampling bias stems from potentially irrelevant phylogenetic and ecological comparisons, such as between mice, humans, naked mole rats and whales. A more insightful approach is to make comparisons within phylogenetic groups, ecological niches, or life histories (Abegglen et al. 2015; Faulkes et al. 2015; Keane et al. 2015; Sanchez et al. 2015). Thus, for example, rather than compare mice and naked mole rats, which have little in common excepting body size, comparisons should begin within their taxonomic groups, e.g., Hystricomorpha (including *Heterocephalus glaber*) and Myomorpha (including *Mus musculus*). In contrast, some examples of elevated cancer risk in captivity are associated with phylogenetically-related taxa (e.g., (Canfield et al. 1990; Owston et al. 2008)), but it is unknown to what extent these patterns are related to similar types of environmental forcing, similar ecology or biology, or to other, unexplained factors.

The differences between disease, individual condition, population limitation and natural selection. Establishing benchmarks for studying cancer (i.e., simple presence vs. importance to the functioning of the host organism) is highly challenging. Even should stem cells obtain one or more cancer driver mutations (Glossary), the affected tissue may not become invasive (Martincorena et al. 2015), and should a lesion progress to invasive cancer, it may never present health (Greaves 2014) or fitness (DeGregori 2011; Hochberg et al. 2013) consequences. As with highly virulent microparasitic diseases, the symptoms associated with many cancers – especially those producing external tumours – elicit the intuitive reaction that they are likely to have individual health and population consequences. However, the severity of a disease on individuals does not necessarily predict how the disease will affect population size (Anderson 1979), and disease-driven population limitation need not be associated with

strong selection for disease resistance (Holt and Hochberg 1997). Rather, the strength of selection integrates reduced fitness, allelic dominance (in diploids), the frequency of alternative fitter genotypes, and population structure over spatio-temporally heterogeneous environments (e.g. (Orr 2009)).

Cancer emergence, health effects, and fitness consequences are each sequentially probabilistic. This means that adaptations to minimize the fitness impacts of cancers should increase with extended pre-reproductive and reproductive life spans (Kokko and Hochberg 2015), but also that cancer will be expressed more often with longer periods of aging and senescence, as is the case, for example, in humans (DeGregori 2011; Hochberg et al. 2013). Thus, cancer morbidity and mortality at older, post-reproductive ages are generally not expected to produce a substantial evolutionary response ((Kirkwood 2005), but see (Brown and Aktipis 2015)). Species with short pre-reproductive and reproductive life spans and little senescence (e.g., many small mammals and birds (Ricklefs 1998)) will tend to perish before succumbing to a threatening neoplasm, and are expected to have less reinforced adaptation to preventing cancer. Limited data concur with these basic predictions. For example, elephants have long pre-reproductive and reproductive life spans, show little senescence (Promislow 1991), have reinforced cancer suppression, and their lifetime cancer rates in captivity are less than 5% (Abegglen et al. 2015); field mice show high cancer rates, but only when reared beyond their short, natural life spans (Andervont and Dunn 1962; Schug et al. 1991; Pompei et al. 2001); naked mole rats have long reproductive life spans, show little senescence (e.g., Buffenstein 2008), and have reinforced cancer protection (e.g., Tian et al. 2013).

Real antagonistic pleiotropy? Positive selection on certain phenotypic traits due, for example, to antagonistic coevolution may be associated with increased cancer risks (Crespi and Summers 2006). Antagonistic pleiotropy (Glossary) is often advanced as an explanation

for correlations between positively selected traits and the occurrence of diseases (Carter and Nguyen 2011), but evidence for its specific role in cancer is largely indirect, particularly in humans (Leroi et al. 2005; Grimes and Chandra 2009; Giaimo and d’Adda di Fagagna 2012). For example, studies of human female breast cancers suggest a pleiotropic association between fecundity and the presence of BRCA1/2 mutations (Smith et al. 2012; Kwiatkowski et al. 2015). Although many factors may be involved, da Silva (da Silva 2012) used mathematical models to show that considerable augmentations in fecundity associated with weak negative impacts on fitness (resource transfers or the “grandmother effect”; (Croft et al. 2015; Brown and Aktipis 2015); see also (Pavard and Metcalf 2007)) could explain the current low frequencies of BRCA alleles, but only in conjunction with low general female fertility in our distant ancestors (by which time much of the evolution leading to the present-day frequencies would have occurred). Alternatively, the association between BRCA mutations and increased fecundity may be real, but not genetically linked. This could occur, for example, if there are associations between the mutated BRCA gene and particular ethnic groups (Fackenthal and Olopade 2007) or families (Kwiatkowski et al. 2015).

Perhaps the most convincing evidence for positive selection on a trait linked with cancer susceptibility comes from male swordtail fish (*Xiphophorus*). Hybrid males harbouring the *Xmrk* oncogene are more susceptible to melanomas in later life, but are also larger and better at competing for mating opportunities (Fernandez and Morris 2008). Limited evidence comes from two additional sources. First, in free-living populations of *X. variatus* subsequently reared in the laboratory, older, non-hybrid males express melanosis or nodular melanomas (e.g., (Schartl et al. 1995; Fernandez and Bowser 2010)). Second, Fernandez and Bowser (Fernandez and Bowser 2010) observed that 8 out of 52 non-hybrid males of *X. cortezi* at one field site showed signs of melanoma stemming from *Xmrk* genes. However, closer examination of these fish did not reveal adverse impacts of the neoplasms, and the authors

did not note whether the remaining field sites were similarly inspected. Nevertheless, the fact that not all fish harbour the *Xmrk* gene suggests some form of frequency dependent or fluctuating habitat selection (see (Fernandez and Bowser 2010) for discussion), indicative of antagonistic pleiotropy.

Inferring and untangling causal factors. In species where cancer epidemiology has been quantified, there is almost invariably a hypothesized contributing novel environment (Vittecoq et al. 2015; Ujvari et al. 2016a)). However, inferring environmental influence is often based on probable or possible cause. For example, Daoust et al. (1991) notes the stark contrast in multicentric mesenchymal tumour incidence between samples of White-fronted geese (*Anser albifrons*), (23%, n=30) and three other goose species: *Branta canadensis* (0.9%, n=117), *Chen caerulescens caerulescens* (0%, n=594), and *Chen rossii* (0%, n=77). The cause (s) of these striking differences, although suggestive of environmental impacts on White-fronted geese, would be difficult if not impossible to either retrospectively evaluate, or to retest should present-day populations show little or no cancer.

Indeed, determining contribution and, more specifically, causality in disease and mortality is notoriously difficult (Rothman and Greenland 2005). Cancers are no exception, in part because one or more etiological agents may be contribute (for certain bivalves, see (Carballal et al. 2015)) and interact with one another in a causal web (for humans (Galea et al. 2010)). Moreover, cancer may be only one of a myriad of diseases and dysfunctions affecting the organism. This means that it will often be difficult to establish the causal and/or contributing role of cancer in individual health, reproduction and survival, and by extension, whether *cancer itself* will impact natural selection. Consider the following hypothetical but realistic scenario. In studying samples of a population over many generations it is discovered that individuals exposed to a toxin show one or more cellular and tissue abnormalities (including

cancers) that are statistically associated with mortality risk. Should resistance evolve, will it (1) detoxify the chemical, and/or (2) result in the avoidance of habitats where the chemical is found, and/or (3) act downstream and suppress cancers and/or other diseases caused by toxin exposure? Assuming that a unique set of cancer driver mutations are obtained sequentially, different environmental (causal) factors may each differentially be responsible for each driver mutation as part of a multistage process. Determining how natural selection proceeds in complex situations such as this is largely unexplored.

The 'Primacy of Mortalities'. All else being equal, changes in the incidence of factors that contribute to mortality over a given age range will (obviously) alter the incidence of other factors that act at the same or at later ages. In contrast, changes in the incidence of late acting factors are less likely to affect the dynamics of those acting at younger ages. This well known demographic effect is potentially important for cancer risks that increase precipitously with age (e.g., human prostate cancers). For example, improvements in preventing and treating early-onset cardiovascular diseases have not only increased life span, but also appear to have shifted deaths to other later-occurring causes, including cancer (e.g., (Davis 1994)). Moreover, different diseases show contrasting age-specific patterns in multi-stage acceleration (Frank 2004b), suggesting that complex patterns in age-specific cancer risk may emerge due to changes in the probabilities of other diseases occurring (e.g., through more effective prevention or individual treatment). As discussed in more detail below, many cancers observed in captive animals and in humans are an indirect consequence of reducing mortality in young and mid-aged individuals through, for example, animal protection in households and zoos, and, for humans in particular, improved sanitation and health care.

A framework for novel environmental impacts on cancer

Despite biases in quantification and difficulties in ascribing cause, as discussed below, there is considerable evidence that novel environments contribute to explaining the high levels of present-day cancer in many animal species, and the majority of cancer risk in humans. What is currently lacking is a framework for understanding the causal web of how environments impact cancer risk. We take first steps towards a more general theory of cancer risk by proposing a framework for how environments influence one or more of three distinct biological levels (*The organism, Within-organism processes, The genome*; Figure 2) and how this, in turn, leads to cancer (Figure 3). We then illustrate how our framework can be used to gain quantitative insights into overall cancer risk in the human population.

1. *The organism: longer life span and larger body size.* Most cancers in humans and the few other animal species studied in detail are associated with longevity and aging (Campisi 2013; de Magalhães 2013). Whereas survival to ages at which cancer is a health threat is probably rare in natural habitats (Finch 1990; Kirkwood 2005), modern-day transformations have made survival to cancer-prone ages the rule for many species living in protected environments, particularly humans.

Improved survival increases cancer risk in three interactive ways. First, because cancer is a multi-stage mutational process, longer life means *more time to obtain* all necessary mutations to produce a cancer and for the cancer to grow, spread, and impact performance and health. Second, the *aging process itself* favours both genomic instability (Vijg and Suh 2013) and the likelihood that tissue microenvironmental conditions will be hospitable to the expression of deleterious mutations leading to tumour development and invasion (DeGregori 2011; DeGregori 2012; Campisi 2013). Third, longer life means *more potential exposure* to environmental factors (e.g., mutagens) that increase the probability of multi-stage carcinogenesis obtaining (see *Within-organism processes* below).

One prime example of increases in health and longevity is the confinement of wildlife in protected habitats (e.g., laboratories, nature reserves, zoos, homes; but see (Clubb and Mason 2003)). As alluded to above, studies from zoo animals indicate that most species have cancer incidences lower than 5% (Effron et al. 1977; Abegglen et al. 2015), though according to the data compiled by Abegglen and coworkers (2015) a significant minority (30% or 11/37 species) exhibit incidences (based on necropsied animals) above 5%. Beyond the reasonable hypothesis that many observed cancers in laboratory and captive populations are associated with longer life spans compared to field situations, and notwithstanding potential biases due to small sample sizes, to the best of our knowledge, many of the cases greater than *c.* 5% lifetime risk reported in the literature can be explained by other novel environmental effects as contributing factors, for example, viral infections or the use of chemical contraceptives (e.g., (McAloose et al. 2007; Ewald and Swain Ewald 2015) (see also *Within-organism processes* below).

The life spans of modern humans and their ancestors have increased over time scales of millions (Caspari and Lee 2004; Finch 2007), tens of thousands and thousands (Kaplan et al. 2000; Caspari and Lee 2004; Burger et al. 2012), and hundreds and tens (Fogel 2012; Burger et al. 2012; NCD Risk Factor Collaboration (NCD-RisC) 2016) of years. Modern medicine promotes longevity (increasing cancer risk), but also lowers cancer mortality at younger ages and shifts some cancer effects on health and survival to older ages. The net overall pattern is that overall cancer risk in Western society is less than 1% by age 20, about 2% by age 40, and less than 10% at 50 (Martincorena and Campbell 2015). These low figures, despite numerous mutagenic exposures (e.g., smoking, UV radiation), can be explained both by highly effective cancer resistance mechanisms at younger ages, and by the many years typically required to complete the multistage process and for the resultant cancer to have health consequences (Jones et al. 2008; Yachida et al. 2010).

Similar to life span, body mass is expected to correlate with the lifetime number of stem cell divisions in the organism, and therefore cancer risk (Peto 1977; Nunney 1999; Noble et al. 2015). Although body mass itself is associated with life span between species, the correlations differ between taxa (Healy et al. 2014), and the within species correlation is more complex (Speakman 2005). Indirect evidence for associations between increased body size and cancer comes from how improved nutrition and health care have contributed to general increases in human stature and life span over recorded history ((Floud et al. 2011), but see e.g., (Formicola and Gianecchini 1999) for variation over much longer time spans). Human height is correlated with the masses of many tissues and organs, such as the liver, kidneys and the heart (Heymsfield et al. 2007). Height is therefore also expected to correlate with the number of cell divisions required to establish these organs and maintain their function (Floud et al. 2011; Zhao et al. 2011), which in turn is expected to correlate with cancer risk (Albanes and Winick 1988; Noble et al. 2015). Limited support for this complex chain of causality comes from study of human height (Shors et al. 2001; Green et al. 2011; Kabat et al. 2014; Jiang et al. 2015) and the size of certain organs (Silva et al. 2011) correlating with cancer risk.

2. *Within-organism processes: disrupted cells, tissues and support systems.* Even with the establishment of highly effective cancer resistance mechanisms, natural selection is not expected to achieve zero risk (see above and, more generally, (Lynch 2012)). Our hypothesis is that sufficient perturbations with respect to the native states of environmental variables affecting cancer risk will increase the probability of obtaining somatic mutations leading to cancer. This may be associated with, for example, injury, disease, or exposure to mutagens (e.g., (Grivennikov et al. 2010; Dang 2015)). Gluckman and Hanson (2004) have argued that environmental conditions during periconceptual, fetal and infant periods of life can have significant impacts on disease risk. However, the implications of these effects for cancers are

little understood.

There are numerous examples in humans and other metazoans of suspected or demonstrated associations between environmental impacts on individual physiology and cancer (Newman and Smith 2006; McAloose and Newton 2009), such as chemical exposure (Martineau et al. 2002; Lerebours et al. 2014), contraceptives (Harrenstien et al. 1996), and UV radiation (Fernandez et al. 2012; Leiter et al. 2014). Whereas humans often have a causal role in cancers stemming from these and other abiotic or physical environmental variables, human implication is usually more difficult to demonstrate in the many cancers associated with extra- and intra-cellular organisms. Indeed, extra- and intra-cellular organisms are widely considered to be a major factor associated with certain cancers in animals and in humans (Ewald 2009; McAloose and Newton 2009; zur Hausen 2009; Coffee et al. 2013; Ewald and Swain Ewald 2015) (Box 2). This may involve changes to the microbiome (Dalmaso et al. 2014; Vogtmann and Goedert 2016), but is more usually associated with parasitic infections. According to de Martel and colleagues (de Martel et al. 2012) pathogens are estimated to account for 16.1% of human cancer globally, with considerable variation between cancer types and geographical regions. Evaluating causality between pathogens and cancer is challenging, in part because of potential delays between the presence of the parasite and cancer detection (Ewald and Swain Ewald 2015). An alternative approach to linking infection with cancer is to study temporal correlations in parasite infection dynamics and cancer. Examples include the spread of HPV (e.g., (Baseman and Koutsky 2005; Ramqvist and Dalianis 2010; Chaturvedi et al. 2011)), HIV (Robbins et al. 2014) and Hepatitis C virus (Perz et al. 2006).

In addition to impacts mediated by the external environment, humans, in particular, influence their own cancer risks through certain lifestyle behaviours. These include alcohol

consumption, UV exposure, smoking (Irigaray et al. 2007), but also certain foods (Soto and Sonnenschein 2010), and lack of physical activity and obesity (Vucenik and Stains 2012). Moreover, there is some evidence that other behaviours such as sleep patterns (Blask 2009), reproductive biology (Aktipis et al. 2014), and psychological states (Reiche et al. 2004), stresses and trauma (Reiche et al. 2004; Antoni et al. 2006) influence cancer risk.

3. *The genome: the emergence of cancer sensitivity genes.* Changes to the heritable genome can have considerable impacts on cancer. These range from very simple to highly complex predispositions (for humans, see (Frank 2004a)). Certain cancers are promoted by homozygosity or drift due to genetic bottlenecks, selective breeding, or by founder effects. For instance, in humans, founder effects and preferential association within groups may favour the persistence of cancer sensitivity genes (de la Chapelle and Wright 1998; Rudan et al. 2003; Fackenthal and Olopade 2007).

In animals, the emergence and persistence of some cancers, in particular transmissible cancers (Box 2), may be promoted by low genetic diversity or heritable sensitivity genes (Fredrickson 1987; Siddle et al. 2007; Miller et al. 2011; Murchison et al. 2012; Murchison et al. 2014; Browning et al. 2014). For example, the Santa Catalina Island fox shows a cancer rate of *c.* 50%, and this elevated level could be explained by some combination of mite infections often associated with these cancers, and the accumulation of deleterious mutations in this highly inbred, low genetic diversity population (Vickers et al. 2015; Robinson et al. 2016). Other fox populations on two other of the Channel Islands are also inbred, also subject to mite infections, but show no cancer (Vickers et al. 2015). Thus, mites and genetic effects may be causal, but additional factors appear necessary to explain cancers on Santa Catalina Island only.

Human-induced population bottlenecks and selective breeding appear to contribute to many

cancers in domestic animals (Vail and MacEwen 2000; Dobson 2013). For example, some cancers in domestic dogs have a genetic basis, stemming from selective breeding (Dobson 2013; Karyadi et al. 2013; Davis and Ostrander 2014; Schiffman and Breen 2015). Similarly, strong selection for specific traits may increase cancer risk, such as more frequent egg laying in farmed chickens leading to increased age-related incidence of ovarian cancer (Johnson et al. 2015). Finally, a recent study on *Hydra* suggests that prolonged breeding in captivity fostered the emergence of tumours after approximately 50 asexual generations (Domazet-Lošo et al. 2014); Alexander Klimovich, pers comm), suggestive of epigenetic alterations, emerging pleiotropy, or mutation accumulation.

Cancer risk in past and present human populations

As indicated above, cancer risk in the human population is subject to alterations in all three biological levels. Not surprisingly, many if not most of these effects are associated with impacts from humans themselves, and in particular extended life spans, environmental mutagens, and lifestyle behaviours. These risks (and by extension, their environmental drivers) are the basis for the current debate of the relative importance of random mutation and environmental drivers in human cancers (Box 3). Here, we take initial steps towards understanding the roles of environment in cancer risk by considering some of the general features of human life history and demography past and present. In particular, we employ information about hunter-gatherers currently and in the recent past to calibrate a mathematical model and conduct a sensitivity analysis with the goal of predicting what overall cancer risks may have been in our recent and more distant ancestors.

Estimates vary, but generally indicate expected hunter-gatherer life spans of 30-40 years (Gurven et al. 2007). Tuljapurkar and colleagues (2007) used demographic models of men

and women to estimate the relative force of selection (Glossary) as a function of age. Their results indicate that the fitness effect of a mutant allele expressed beyond 40 years of age is 20% of what it would be if it were expressed beyond 20 years of age. Selection on an individual at 50 and 60 years of age is 5% and 1%, respectively, of selection on a 20 year old. If this applies generally to pre-modern humans, then it suggests that there would have been selection against germline cancer sensitivity, but with a pronounced decline for those cancers with health effects after approximately 40 years of age. Supporting this prediction, modern day age-specific cancer risk (e.g. (Howlader et al. 2013)) recapitulates the inverse of the predicted force of selection in early *Homo sapiens* (Tuljapurkar et al. 2007) and in modern hunter-gatherers (Lee 2003).

Additional evidence for the hypothesis of less cancer in our recent and distant ancestors comes from mortality data in hunter-gatherer groups prior to outside contact. Data from pre-contact (before 1960) Hiwi indicate low levels (4/86 or 5% of young and older adult deaths) of “other organic/pathological” causes of mortality (including cancers), whereas post-contact was considerably higher (26/69 or 38%) (Hill et al. 2007). However, it is not known with certainty how cancers may have contributed to these or other mortalities. Similar assessments (lower life expectancy and lower risks for certain cancers compared to Western society) hold for the Tsimané Amerindians (excluding deaths from unknown causes, 1.3% (16/1276) of deaths attributable to cancers for all ages, and 7.8% (7/90) for individuals 60 years old or greater) (Gurven et al. 2007), and Inuits (Wilkins et al. 2008; Boysen et al. 2008) for which rapid social and environmental changes may explain why the incidence of many cancers has recently increased (Boysen et al. 2008; Young et al. 2016), whereas high incidence, endemic nasopharyngeal cancers appear to be linked to infectious agents (Friborg and Melbye 2008).

A mathematical model of cancer risk in humans

To examine the relationship between longevity and cancer risk, we use a simple mathematical model (Supplementary methods) that accounts for the probability of acquiring cancer for the first time, the mortality rate due to cancer, and the rate of background mortality (that is, death from causes other than cancer).

However, in reality, populations vary in cancer risk and cancer death rate, as well as background mortality. Based on modern US life expectancy and cancer incidence curves, our model predicts, as expected, a lifetime cancer risk of approximately 40% (Fig. 4C). A population that is identical except that it has 30% lower cancer incidence in adults is predicted to have approximately 30% lower lifetime risk of cancer (Fig. 4C). In pre-modern humans, the model estimates a lifetime cancer risk between 0.6% and 10% (for those alive and cancer-free at age 5). The higher estimate (Fig. 4C) assumes that background mortality in pre-modern humans was similar to what is observed in modern hunter-gatherers (Gurven et al. 2007), and that adult cancer risk per year was 30% lower in pre-modern humans than in the modern US population (based on epidemiological studies that estimate 30%-50% of risk in modern Western populations is attributable to lifestyle (Doll and Peto 1981; Parkin et al. 2011; Song and Giovannucci 2016); see also Supplementary Fig. 1)). The lower estimate assumes a much higher background mortality rate deduced from prehistoric skeletal remains (Gage 1998) – which most likely overestimates adult mortality (O’Connell et al. 1999) – and assumes that cancer risk per year was 80% lower than in the US currently (due to changes in both environment and stature). The model results indicate that most of the difference in lifetime cancer risk between pre-modern humans and the modern US population is due to lower background mortality in the latter, which enables more individuals to attain the advanced ages at which cancer risk is substantially increased. If modern US cancer incidence applied to modern-day hunter gatherers and to Sweden in 1861, then lifetime cancer risks in the latter populations would be 13% and 24%, respectively (in individuals alive and cancer-

free at age 5, assuming the same stature and environmental contributors as in the US currently).

Contrasting patterns emerge when we examine cancer risk over the age range 5-40 years old, which is when the force of selection is expected to be strongest (Fig. 4A). The model indicates that cancer risk by 40 years old was between 0.4% and 1% in pre-modern humans, whereas it is *c.* 2% currently in the US population (Fig. 4D). Most of this difference is predicted to be due to increased environmental forcing in the modern population. In particular, once life expectancy reaches 75 years, the background mortality rate before age 40 is already very low. Therefore further reductions in background mortality (which will increase life expectancy beyond 75 years) are predicted to have almost no effect on cancer risk by 40 years old (Fig. 4D), whereas not surprisingly they have a dramatic effect on cancer risk over a lifetime (Fig. 4C).

Note that these results for the human population concern the lifetime risk of acquiring cancer, irrespective of possible effects on health or survival. Thus, the age-specific incidence data used in the model is likely to overestimate actual cancer risks, in part due to advances in detecting cancers, many of which have no subsequent effects on health or survival. In populations with high background mortality rates (e.g., ancient hunter-gatherers), cancer would less often have negative consequences due to the primacy of mortalities.

Discussion

We argue that elevated cancer risks across the tree of life – particularly observations of greater than *c.* 5% lifetime risk – are largely attributable to novel environments, that is deviations in certain environmental conditions experienced over evolutionary time scales in

native habitats (Fig. 1). This is not to say that resistance to certain cancers in certain species does not continue to evolve, but rather that over ecologically relevant time spans of less than a few generations, environmental drivers and not evolutionary responses dominate variation in cancer epidemiology. We stress that *c.* 5% is a proposed benchmark, based on our literature survey showing no indiscutable examples of cancer beyond this level that cannot be explained by influences of novel environments, and our case study of ancient humans suggestive of similar levels. Inversely, the benchmark of *c.* 5% is a prediction of the limits of evolution in limiting cancer risks across life, and we expect that the actual figure may be much lower for many species. Supporting this, Abegglen et al. (2015) found that 51% of the captive species in their data set showed *no* cancer. Although these cases may be explained by e.g., small sample sizes--taken at face value they suggest that baseline levels in native habitats are likely to be extremely low as well. First-principles theory would be valuable to test our prediction as a general limit across species, and to discover the limits of selection in preventing cancer across different species ecologies.

Our proposal that anthropogenic influences explain much of the novel environmental variation responsible for present-day cancer risks, is in line with previous study suggesting that cancer was of lower importance in free-living animal populations and in humans in the (distant) past (e.g., (Capasso 2005; David and Zimmerman 2010)). We contend, nevertheless, that cancer *has been and still is occasionally important* for certain populations in the wild that are relatively untouched by human activity. Arguably the most prevalent contributors are parasitic infections (Ewald 2009; Ewald and Swain Ewald 2015), but more research is needed to evaluate the extent to which these deviate from evolutionary norms (Fig. 1), and if so, the potential indirect causal role of humans (e.g., contaminants the increase parasitic infection and cancer).

Although environmental impacts on cancer epidemiology are attracting increased interest, the

relevance for ecological and evolutionary dynamics remains underexplored. Similar to certain parasites, transmissible cancers and cancers associated with parasitic infections can have significant population and evolutionary impacts (Box 2), and possibly epidemic-like behaviour (McCallum et al. 2009; Ramqvist and Dalianis 2010). Epidemic-like phenomena could also occur in species vulnerable to endogenous cancers, but aside from correlations between environmental factors and cancer incidence, the implications for species ecology is unknown. The implications of cancer epidemiology for species ecology will be challenging to study, given the typical delays between environmental causes and the effect of the disease on individual performance and survival (e.g., (Ewald and Swain Ewald 2015; Lair et al. 2015)).

Beyond the obvious need for more studies explaining cancer risks in free-living species, both in near-pristine systems and in those perturbed either naturally or by human activity, several important questions merit attention. (1) To what extent are cancers distinct from infectious and noninfectious diseases in how they are affected by environments (Eisenberg et al. 2007) and, inversely, in how they affect species ecology and evolution (Vittecoq et al. 2013; Ujvari et al. 2016a)? (2) For the many cancers associated with viral or parasitic infections, will selection for resistance be directed at the parasite, or at the subsequent induction of cancer? Is the epidemiology of cancer-associated parasites distinct from other parasitic organisms? (3) Can we detect contemporary signals of selection on cancer genes (either directly or linked to other traits under selection) due to altered environments? (4) We actually do not know cancer risks as a function of environmental conditions (Fig. 1) for any species or cancer type. How sensitive are different cancer types on the basis of the tissues they affect to different environmental drivers? (5) Our study considered the risk of *any* cancer affecting individual performance, health, survival and fitness. How do the risk levels of different cancer types (e.g., cancer cell phenotype; tissues or organs affected) vary with environments and phylogenetically across species? (6) Are there any phylogenetic signals of cancer resistance

evolution and natural cancer risk? That is, to what extent do novel genes associated with cancer risk appear in species lineages, or rather is most evolution due to modifications to existing genes (e.g., copy number (Abegglen et al. 2015))? And finally, (7) to what extent can cancer be employed as an indicator of more general environmental impacts, such as global climate change, species invasions or infectious disease epidemics? Linking specific cancers with a single or small number of causal factors could be useful in certain wildlife conservation efforts (McAloose and Newton 2009) and in cancer prevention in the human population (Box 3).

In conclusion, the framework presented here focuses on how deviations from native environmental conditions increase cancer risk at (1) the whole-organism level, by enabling larger body size and longer life span, (2) the within-organism level, by disrupting intra-individual physiology and (3) the germline level, by altering genomes and cancer sensitivity[2]. Supporting the generality of this approach, each of these three levels can be linked to more distal contributions of species ecology (e.g., environmental change leads to heightened susceptibility to a pathogen associated with cancer, or to population contraction and a genetic bottleneck; the removal of a predator results in longer life spans), and to the more proximal process of multistage carcinogenesis (the probability of which depends on lifetime numbers of stem cell divisions and mutation rates). Nevertheless, given the complexity of the causal network proposed here (Fig. 3), testing predictions (Box 4) and dissecting impacts of the various contributors to cancer will be challenging, requiring theoretical, comparative and experimental approaches. For example, more rigorous tests of the effects of captivity on cancer in zoo animals (Ratcliffe 1933), or proposals that reductions in predation or parasite pressure should result in increased cancer risk (Vittecoq et al. 2013), would require contrasts with data from natural habitats and communities with predators/parasites, respectively. For some model species, this could be accomplished with

the employment of field enclosures to monitor all individuals and control species composition (Legrand et al. 2012).

Acknowledgements. We thank Manfred Scharl, Daniel Martineau, Ula Hibner, Hanna Kokko, Paul Ewald, Michael Gurven, Gillian Bentley, Robert Wayne, Alexander Klimovich, and James Sherry for discussions, and the helpful comments from three anonymous reviewers. MEH acknowledges support from the Agence National de la Recherche (EvoCan ANR-13-BSV7-0003-01), the Institut National du Cancer (2014-1-PL-BIO-12-IGR-1), the McDonnell Foundation (Studying Complex Systems Research Award No. 220020294), and the Wissenschaftskolleg zu Berlin.

Footnotes

[1] The most notable exception is transmissible facial tumours in Tasmanian devils (e.g., (McCallum et al. 2009; Hollings et al. 2016; Epstein et al. 2016)). However, transmissible cancers differ importantly from endogenously emerging cancers, in that the population and evolutionary dynamics of the former are more akin to certain infectious diseases.

[2] Evidently, environments approaching a species' native conditions will result in lowered overall cancer risk.

References

- Abegglen LM, Caulin AF, Chan A, et al (2015) Potential Mechanisms for Cancer Resistance in Elephants and Comparative Cellular Response to DNA Damage in Humans. *JAMA* 314:1.
- Aguirre AA, Lutz P (2004) Marine Turtles as Sentinels of Ecosystem Health: Is Fibropapillomatosis an Indicator? *Ecohealth* 1:275–283.
- Aktipis CA, Boddy AM, Jansen G, et al (2015) Cancer across the tree of life: cooperation and cheating in multicellularity. *Philos Trans R Soc Lond B Biol Sci* 370:20140219–20140219.
- Aktipis CA, Ellis BJ, Nishimura KK, Hiatt RA (2014) Modern reproductive patterns associated with estrogen receptor positive but not negative breast cancer susceptibility. *Evol Med Public Health* 2015:52–74.
- Albanes D, Winick M (1988) Are Cell Number and Cell Proliferation Risk Factors for Cancer? 1. *J Natl Cancer Inst* 80:772–775.
- Anderson RM (1979) Parasite pathogenicity and the depression of host population equilibria. *Nature* 279:150–152.
- Andervont HB, Dunn TB (1962) Occurrence of tumors in wild house mice. *J Natl Cancer Inst* 28:1153–1163.
- Antoni MH, Lutgendorf SK, Cole SW, et al (2006) The influence of bio-behavioural factors on tumour biology: pathways and mechanisms. *Nat Rev Cancer* 6:240–248.
- Armitage P, Doll R (1954) The age distribution of cancer and a multi-stage theory of carcinogenesis. *Br J Cancer* 8:1–12.
- Ashby B, Boots M (2015) Coevolution of parasite virulence and host mating strategies. *Proc Natl Acad Sci U S A* 112:13290–13295.
- Barber BJ (2004) Neoplastic diseases of commercially important marine bivalves. *Aquat Living Resour* 17:449–466.
- Baseman JG, Koutsky LA (2005) The epidemiology of human papillomavirus infections. *J Clin Virol* 32 Suppl 1:S16–24.
- Blask DE (2009) Melatonin, sleep disturbance and cancer risk. *Sleep Med Rev* 13:257–264.
- Blekhman R, Man O, Herrmann L, et al (2008) Natural Selection on Genes that Underlie Human Disease Susceptibility. *Curr Biol* 18:883–889.
- Boddy AM, Kokko H, Breden F, et al (2015) Cancer susceptibility and reproductive trade-offs: a model of the evolution of cancer defences. *Philos Trans R Soc Lond B Biol Sci* 370:20140220.
- Boivin NL, Zeder MA, Fuller DQ, et al (2016) Ecological consequences of human niche construction: Examining long-term anthropogenic shaping of global species distributions. *Proc Natl Acad Sci U S A* 113:6388–6396.
- Boysen T, Friborg J, Andersen A, et al (2008) The Inuit cancer pattern—The influence of migration. *Int J Cancer* 122:2568–2572.
- Bozic I, Antal T, Ohtsuki H, et al (2010) Accumulation of driver and passenger mutations during tumor progression. *Proc Natl Acad Sci U S A* 107:18545–18550.
- Branzei D, Foiani M (2005) The DNA damage response during DNA replication. *Curr Opin Cell Biol* 17:568–575.
- Browning HM, Acevedo-Whitehouse K, Gulland MD, et al (2014) Evidence for a genetic basis of urogenital carcinoma in the wild California sea lion. *Proc Biol Sci* 281:20140240.
- Brown JS, Aktipis CA (2015) Inclusive fitness effects can select for cancer suppression into old age. *Philos Trans R Soc Lond B Biol Sci* 370:20150160.
- Buffenstein R (2008) Negligible senescence in the longest living rodent, the naked mole-rat: Insights from a successfully aging species. *J Comp Physiol B* 178:439–445.
- Burger O, Baudisch A, Vaupel JW (2012) Human mortality improvement in evolutionary context. *Proc Natl Acad Sci U S A* 109:18210–18214.
- Cairns J (2002) Somatic stem cells and the kinetics of mutagenesis and carcinogenesis. *Proc Natl Acad Sci U S A* 99:10567–10570.
- Calabrese P, Shibata D (2010) A simple algebraic cancer equation: calculating how cancers may arise with normal mutation rates. *BMC Cancer* 10:3.
- Campisi J (2013) Aging, cellular senescence, and cancer. *Annu Rev Physiol* 75:685–705.
- Canfield PJ, Hartley WJ, Reddacliff GL (1990) Spontaneous proliferations in Australian marsupials--

- a survey and review. 2. Dasyurids and bandicoots. *J Comp Pathol* 103:147–158.
- Capasso LL (2005) Antiquity of cancer. *Int J Cancer* 113:2–13.
- Carballal MJ, Barber BJ, Iglesias D, Villalba A (2015) Neoplastic diseases of marine bivalves. *J Invertebr Pathol* 131:83–106.
- Carter AJR, Nguyen AQ (2011) Antagonistic pleiotropy as a widespread mechanism for the maintenance of polymorphic disease alleles. *BMC Med Genet* 12:160.
- Casás-Selves M, Degregori J (2011) How cancer shapes evolution, and how evolution shapes cancer. *Evolution* 4:624–634.
- Caspari R, Lee S-H (2004) Older age becomes common late in human evolution. *Proc Natl Acad Sci U S A* 101:10895–10900.
- Caulin AF, Maley CC (2011) Peto's Paradox: Evolution's prescription for cancer prevention. *Trends Ecol Evol* 26:175–182.
- Chaturvedi AK, Engels EA, Pfeiffer RM, et al (2011) Human papillomavirus and rising oropharyngeal cancer incidence in the United States. *J Clin Oncol* 29:4294–4301.
- Chevin L-M, Lande R, Mace GM (2010) Adaptation, plasticity, and extinction in a changing environment: towards a predictive theory. *PLoS Biol* 8:e1000357.
- Clubb R, Mason G (2003) Animal welfare: captivity effects on wide-ranging carnivores. *Nature* 425:473–474.
- Coffee LL, Casey JW, Bowser PR (2013) Pathology of tumors in fish associated with retroviruses: a review. *Vet Pathol* 50:390–403.
- Colditz GA, Sutcliffe S (2016) The Preventability of Cancer: Stacking the Deck. *JAMA Oncology* 19–20.
- Cournil A, Kirkwood TB (2001) If you would live long, choose your parents well. *Trends Genet* 17:233–235.
- Crespi BJ, Summers K (2006) Positive selection in the evolution of cancer. *Biol Rev Camb Philos Soc* 81:407–424.
- Crespi B, Summers K (2005) Evolutionary biology of cancer. *Trends Ecol Evol* 20:545–552.
- Croft DP, Brent LNJ, Franks DW, Cant MA (2015) The evolution of prolonged life after reproduction. *Trends Ecol Evol* 30:407–416.
- Dalmasso G, Cougnoux A, Delmas J, et al (2014) The bacterial genotoxin colibactin promotes colon tumor growth by modifying the tumor microenvironment. *Gut Microbes* 5:675–680.
- Dang CV (2015) A metabolic perspective of Peto's paradox and cancer. *Philos Trans R Soc Lond B Biol Sci*. doi: 10.1098/rstb.2014.0223
- Daoust PY, Wobeser G, Rainnie DJ, Leighton FA (1991) Multicentric intramuscular lipomatosis/fibromatosis in free-flying white-fronted and Canada geese. *J Wildl Dis* 27:135–139.
- da Silva J (2012) BRCA1/2 mutations, fertility and the grandmother effect. *Proc Biol Sci* 279:2926–2929.
- David AR, Zimmerman MR (2010) Cancer: an old disease, a new disease or something in between? *Nat Rev Cancer* 10:728–733.
- Davis BW, Ostrander EA (2014) Domestic dogs and cancer research: a breed-based genomics approach. *ILAR J* 55:59–68.
- Davis DL (1994) Decreasing cardiovascular disease and increasing cancer among whites in the United States from 1973 through 1987. Good news and bad news. *JAMA* 271:431–437.
- DeGregori J (2012) Challenging the axiom: does the occurrence of oncogenic mutations truly limit cancer development with age? *Oncogene* 32:1869–1875.
- DeGregori J (2011) Evolved tumor suppression: Why are we so good at not getting cancer? *Cancer Res* 71:3739–3744.
- de la Chapelle A, Wright FA (1998) Linkage disequilibrium mapping in isolated populations: the example of Finland revisited. *Proc Natl Acad Sci U S A* 95:12416–12423.
- Delaney MA, Ward JM, Walsh TF, et al (2016) Initial Case Reports of Cancer in Naked Mole-rats (*Heterocephalus glaber*). *Vet Pathol* 53:691–696.
- de Magalhães JP (2013) How ageing processes influence cancer. *Nat Rev Cancer* 13:357–365.
- de Martel C, Ferlay J, Franceschi S, et al (2012) Global burden of cancers attributable to infections in 2008: a review and synthetic analysis. *Lancet Oncol* 13:607–615.

- DevCan: Probability of Developing or Dying of Cancer Software (2005) Statistical Research and Applications Branch, National Cancer Institute
- Dobson JM (2013) Breed-predispositions to cancer in pedigree dogs. *ISRN Vet Sci* 2013:941275.
- Doll R, Peto R (1981) The causes of cancer: quantitative estimates of avoidable risks of cancer in the United States today. *J Natl Cancer Inst* 66:1191–1308.
- Domazet-Lošo T, Klimovich A, Anokhin B, et al (2014) Naturally occurring tumours in the basal metazoan Hydra. *Nat Commun* 5:4222.
- Domazet-Lošo T, Tautz D (2010) Phylostratigraphic tracking of cancer genes suggests a link to the emergence of multicellularity in metazoa. *BMC Biol* 8:66.
- Effron M, Griner L, Benirschke K (1977) Nature and rate of neoplasia found in captive wild mammals, birds, and reptiles at necropsy. *J Natl Cancer Inst* 59:185–198.
- Eisenberg DTA (2011) An evolutionary review of human telomere biology: the thrifty telomere hypothesis and notes on potential adaptive paternal effects. *Am J Hum Biol* 23:149–167.
- Eisenberg JNS, Desai MA, Levy K, et al (2007) Environmental determinants of infectious disease: a framework for tracking causal links and guiding public health research. *Environ Health Perspect* 115:1216–1223.
- Epstein B, Jones M, Hamede R, et al (2016) Rapid evolutionary response to a transmissible cancer in Tasmanian devils. *Nat Commun*. doi: 10.1038/ncomms12684
- Eriksson O (2013) Species pools in cultural landscapes – niche construction, ecological opportunity and niche shifts. *Ecography* 36:403–413.
- Ewald PW (2009) An Evolutionary Perspective on Parasitism as a Cause of Cancer. *Adv Parasitol* 68:21–43.
- Ewald PW, Swain Ewald HA (2015) Infection and cancer in multicellular organisms. *Philos Trans R Soc Lond B Biol Sci* 370:20140224.
- Fackenthal JD, Olopade OI (2007) Breast cancer risk associated with BRCA1 and BRCA2 in diverse populations. *Nat Rev Cancer* 7:937–948.
- Faulkes CG, Davies KTJ, Rossiter SJ, Bennett NC (2015) Molecular evolution of the hyaluronan synthase 2 gene in mammals: implications for adaptations to the subterranean niche and cancer resistance. *Biol Lett* 11:20150185.
- Fay MP (2004) Estimating age conditional probability of developing disease from surveillance data. *Popul Health Metr* 2:6.
- Ferlay J, Soerjomataram I, Dikshit R, et al (2015) Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 136:E359–86.
- Fernandez AA, Bowser PR (2010) Selection for a dominant oncogene and large male size as a risk factor for melanoma in the Xiphophorus animal model. *Mol Ecol* 19:3114–3123.
- Fernandez A a., Morris MR (2008) Mate choice for more melanin as a mechanism to maintain a functional oncogene. *Proc Natl Acad Sci U S A* 105:13503–13507.
- Fernandez AA, Paniker L, Garcia R, Mitchell DL (2012) Recent advances in sunlight-induced carcinogenesis using the Xiphophorus melanoma model. *Comp Biochem Physiol C Toxicol Pharmacol* 155:64–70.
- Finch CE (2007) The biology of human longevity: Inflammation, Nutrition, and Aging in the Evolution of Lifespans. Academic Press
- Finch CE (1990) Longevity, senescence, and the genome. University of Chicago Press, Chicago, USA
- Fleming JM, Creevy KE, Promislow DEL (2011) Mortality in north american dogs from 1984 to 2004: an investigation into age-, size-, and breed-related causes of death. *J Vet Intern Med* 25:187–198.
- Floud R, Fogel RW, Harris B, Hong SC (2011) The Changing Body: Health, Nutrition, and Human Development in the Western World since 1700. Cambridge University Press
- Fogel RW (2012) Explaining Long-Term Trends in Health and Longevity. Cambridge University Press
- Formicola V, Giannecchini M (1999) Evolutionary trends of stature in upper Paleolithic and Mesolithic Europe. *J Hum Evol* 36:319–333.
- Frank SA (2004a) Genetic predisposition to cancer—insights from population genetics. *Nat Rev Genet* 5:764–772.
- Frank SA (2004b) A multistage theory of age-specific acceleration in human mortality. *BMC Biol*

2:16.

- Frank SA (2010) *Dynamics of Cancer: Incidence, Inheritance, and Evolution*. Princeton University Press, Princeton (NJ)
- Frank SA (1996) Models of Parasite Virulence. *Q Rev Biol* 71:37–78.
- Frank SA (2004c) Age-specific acceleration of cancer. *Curr Biol* 14:242–246.
- Fredrickson TN (1987) Ovarian tumors of the hen. *Environ Health Perspect* 73:35–51.
- Friborg JT, Melbye M (2008) Cancer patterns in Inuit populations. *Lancet Oncol* 9:892–900.
- Gage TB (1998) The comparative demography of primates: with some comments on the evolution of life histories. *Annu Rev Anthropol* 27:197–221.
- Galea S, Riddle M, Kaplan GA (2010) Causal thinking and complex system approaches in epidemiology. *Int J Epidemiol* 39:97–106.
- Giaimo S, d’Adda di Fagagna F (2012) Is cellular senescence an example of antagonistic pleiotropy? *Aging Cell* 11:378–383.
- Gluckman PD, Hanson MA (2004) Living with the past: evolution, development, and patterns of disease. *Science* 305:1733–1736.
- Gluckman PD, Low FM, Buklijas T, et al (2011) How evolutionary principles improve the understanding of human health and disease. *Evol Appl* 4:249–263.
- Gomes NMV, Ryder OA, Houck ML, et al (2011) Comparative biology of mammalian telomeres: Hypotheses on ancestral states and the roles of telomeres in longevity determination. *Aging Cell* 10:761–768.
- Gorbunova V, Seluanov A, Zhang Z, et al (2014) Comparative genetics of longevity and cancer: insights from long-lived rodents. *Nat Rev Genet* 15:531–540.
- Greaves M (2000) *Cancer: The Evolutionary Legacy*. Oxford University Press
- Greaves M (2014) Does everyone develop covert cancer? *Nat Rev Cancer* 14:209–210.
- Greaves M, Maley CC (2012) Clonal evolution in cancer. *Nature* 481:306–313.
- Green J, Cairns BJ, Casabonne D, et al (2011) Height and cancer incidence in the Million Women Study: prospective cohort, and meta-analysis of prospective studies of height and total cancer risk. *Lancet Oncol* 12:785–794.
- Grimes A, Chandra SBC (2009) Significance of cellular senescence in aging and cancer. *Cancer Res Treat* 41:187–195.
- Grivennikov SI, Greten FR, Karin M (2010) Immunity, Inflammation, and Cancer. *Cell* 140:883–899.
- Gurven M, Kaplan H, Supa AZ (2007) Mortality experience of Tsimane Amerindians of Bolivia: regional variation and temporal trends. *Am J Hum Biol* 19:376–398.
- Hamilton WD (1966) The moulding of senescence by natural selection. *J Theor Biol* 12:12–45.
- Hanahan D, Weinberg R a. (2000) The hallmarks of cancer. *Cell* 100:57–70.
- Hanahan D, Weinberg RA (2011) Hallmarks of cancer: the next generation. *Cell* 144:646–674.
- Harrenstien LA, Munson L, Seal US, The American Zoo and Aquarium Association Mammary Cancer Study Group (1996) Mammary Cancer in Captive Wild Felids and Risk Factors for Its Development: A Retrospective Study of the Clinical Behavior of 31 Cases. *J Zoo Wildl Med* 27:468–476.
- Healy K, Guillerme T, Finlay S, et al (2014) Ecology and mode-of-life explain lifespan variation in birds and mammals. *Proc Biol Sci* 281:20140298.
- Heymsfield SB, Gallagher D, Mayer L, et al (2007) Scaling of human body composition to stature: new insights into body mass index. *Am J Clin Nutr* 86:82–91.
- Hill K, Hurtado a. M, Walker RS (2007) High adult mortality among Hiwi hunter-gatherers: Implications for human evolution. *J Hum Evol* 52:443–454.
- Hindorff LA, Gillanders EM, Manolio TA (2011) Genetic architecture of cancer and other complex diseases: lessons learned and future directions. *Carcinogenesis* 32:945–954.
- Hochberg ME, Michalakis Y, de Meeus T (1992) Parasitism as a constraint on the rate of life-history evolution. *J Evol Biol* 5:491–504.
- Hochberg ME, Noble RJ, Braude S (2016) A Hypothesis to Explain Cancers in Confined Colonies of Naked Mole Rats. *bioRxiv* 079012.
- Hochberg ME, Thomas F, Assenat E, Hibner U (2013) Preventive Evolutionary Medicine of Cancers. *Evol Appl* 6:134–143.
- Hollings T, Jones M, Mooney N, McCallum H (2016) Disease-induced decline of an apex predator

- drives invasive dominated states and threatens biodiversity. *Ecology* 97:394–405.
- Holt RD, Hochberg ME (1997) When is Biological Control Evolutionarily Stable (or is it)? *Ecology* 78:1673.
- Howlander N, Noone AM, Krapcho M, et al (2013) SEER cancer statistics review, 1975–2010. 2013.
- Hueper WC (1963) Environmental carcinogenesis in man and animals. *Ann N Y Acad Sci* 108:963–1038.
- Irigaray P, Newby JA, Clapp R, et al (2007) Lifestyle-related factors and environmental agents causing cancer: an overview. *Biomed Pharmacother* 61:640–658.
- Jiang Y, Marshall RJ, Walpole SC, et al (2015) An international ecological study of adult height in relation to cancer incidence for 24 anatomical sites. *Cancer Causes Control* 26:493–499.
- Johnson PA, Stephens CS, Giles JR (2015) The domestic chicken: Causes and consequences of an egg a day. *Poult Sci* 94:816–820.
- Jones S, Chen W-DD, Parmigiani G, et al (2008) Comparative lesion sequencing provides insights into tumor evolution. *Proc Natl Acad Sci U S A* 105:4283–4288.
- Kabat GC, Kim MY, Hollenbeck AR, Rohan TE (2014) Attained height, sex, and risk of cancer at different anatomic sites in the NIH-AARP diet and health study. *Cancer Causes Control* 25:1697–1706.
- Kaplan H, Hill K, Lancaster J, Hurtado AM (2000) A theory of human life history evolution: diet, intelligence, and longevity. *Evolutionary Anthropology Issues News and Reviews* 9:156–185.
- Karyadi DM, Karlins E, Decker B, et al (2013) A copy number variant at the KITLG locus likely confers risk for canine squamous cell carcinoma of the digit. *PLoS Genet* 9:e1003409.
- Keane M, Semeiks J, Webb AE, et al (2015) Insights into the evolution of longevity from the bowhead whale genome. *Cell Rep* 10:112–122.
- Kirkwood TBL (2005) Understanding the odd science of aging. *Cell* 120:437–447.
- Knudson AG (2001) Two genetic hits (more or less) to cancer. *Nat Rev Cancer* 1:157–162.
- Kokko H, Hochberg ME (2015) Towards cancer-aware life-history modelling. *Philos Trans R Soc Lond B Biol Sci* 370:20140234.
- Kwiatkowski F, Arbre M, Bidet Y, et al (2015) BRCA Mutations Increase Fertility in Families at Hereditary Breast/Ovarian Cancer Risk. *PLoS One* 10:e0127363.
- Lafferty KD, Kuris AM (1999) How environmental stress affects the impacts of parasites. *Limnol Oceanogr* 44:925–931.
- Lair S, Measures LN, Martineau D (2015) Pathologic Findings and Trends in Mortality in the Beluga (*Delphinapterus leucas*) Population of the St Lawrence Estuary, Quebec, Canada, From 1983 to 2012. *Veterinary Pathology* 53:2–36.
- Lee RD (2003) Rethinking the evolutionary theory of aging: transfers, not births, shape senescence in social species. *Proc Natl Acad Sci U S A* 100:9637–9642.
- Legrand D, Guillaume O, Baguette M, et al (2012) The Metatron: an experimental system to study dispersal and metaecosystems for terrestrial organisms. *Nat Methods* 9:828–833.
- Leiter U, Eigentler T, Garbe C (2014) Epidemiology of skin cancer. *Adv Exp Med Biol* 810:120–140.
- Lerebours A, Stentiford GD, Lyons BP, et al (2014) Genetic alterations and cancer formation in a European flatfish at sites of different contaminant burdens. *Environ Sci Technol* 48:10448–10455.
- Leroi AM, Bartke A, De Benedictis G, et al (2005) What evidence is there for the existence of individual genes with antagonistic pleiotropic effects? *Mech Ageing Dev* 126:421–429.
- Leroi AM, Koufopanou V, Burt A (2003) Cancer selection. *Nat Rev Cancer* 3:226–231.
- Li Y, Deeb B, Pendergrass W, Wolf N (1996) Cellular proliferative capacity and life span in small and large dogs. *J Gerontol A Biol Sci Med Sci* 51:B403–B408.
- Lynch M (2012) Evolutionary layering and the limits to cellular perfection. *Proc Natl Acad Sci U S A* 109:18851–18856.
- Maciak S, Michalak P (2015) Cell size and cancer: a new solution to Peto's paradox? *Evol Appl* 8:2–8.
- MacRae SL, Croken MMK, Calder RB, et al (2015) Dna Repair in Species With Extreme Lifespan Differences. *Aging* 7:1171–1184.
- Maher MC, Uricchio LH, Torgerson DG, Hernandez RD (2013) Population Genetics of Rare Variants and Complex Diseases. *Hum Hered* 74:118–128.

- Martincorena I, Campbell PJ (2015) Somatic mutation in cancer and normal cells. *Science* 349:1483–1489.
- Martincorena I, Roshan A, Gerstung M, et al (2015) High burden and pervasive positive selection of somatic mutations in normal human skin. *Science* 348:880–886.
- Martineau D, Lemberger K, Dallaire A, et al (2002) Cancer in wildlife, a case study: beluga from the St. Lawrence estuary, Quebec, Canada. *Environ Health Perspect* 110:285–292.
- McAloose D, Munson L, Naydan DK (2007) Histologic features of mammary carcinomas in zoo felids treated with melengestrol acetate (MGA) contraceptives. *Vet Pathol* 44:320–326.
- McAloose D, Newton AL (2009) Wildlife cancer: a conservation perspective. *Nat Rev Cancer* 9:517–526.
- McCallum H, Jones M, Hawkins C, et al (2009) Transmission dynamics of Tasmanian devil facial tumor disease may lead to disease-induced extinction. *Ecology* 90:3379–3392.
- Medawar PB (1952) An unsolved problem of biology: an inaugural lecture delivered at University College, London, 6 December, 1951. HK Lewis and Company
- Merlo LMF, Pepper JW, Reid BJ, Maley CC (2006) Cancer as an evolutionary and ecological process. *Nat Rev Cancer* 6:924–935.
- Metzger MJ, Reinisch C, Sherry J, Goff SP (2015) Horizontal transmission of clonal cancer cells causes leukemia in soft-shell clams. *Cell* 161:255–263.
- Metzger MJ, Villalba A, Carballal MJ, et al (2016) Widespread transmission of independent cancer lineages within multiple bivalve species. *Nature* 1–11.
- Michalakis Y, Olivieri I, Renaud F, Raymond M (1992) Pleiotropic action of parasites: How to be good for the host. *Trends Ecol Evol* 7:59–62.
- Miller W, Hayes VM, Ratan A, et al (2011) Genetic diversity and population structure of the endangered marsupial *Sarcophilus harrisii* (Tasmanian devil). *Proc Natl Acad Sci U S A* 108:12348–12353.
- Murchison EP, Schulz-Trieglaff OB, Ning Z, et al (2012) Genome sequencing and analysis of the Tasmanian devil and its transmissible cancer. *Cell* 148:780–791.
- Murchison EP, Wedge DC, Alexandrov LB, et al (2014) Transmissible [corrected] dog cancer genome reveals the origin and history of an ancient cell lineage. *Science* 343:437–440.
- Murgia C, Pritchard JK, Kim SY, et al (2006) Clonal origin and evolution of a transmissible cancer. *Cell* 126:477–487.
- Muttray A, Reinisch C, Miller J, et al (2012) Haemocytic leukemia in Prince Edward Island (PEI) soft shell clam (*Mya arenaria*): spatial distribution in agriculturally impacted estuaries. *Sci Total Environ* 424:130–142.
- Nagy JD, Victor EM, Cropper JH (2007) Why don't all whales have cancer? A novel hypothesis resolving Peto's paradox. *Integr Comp Biol* 47:317–328.
- NCD Risk Factor Collaboration (NCD-RisC) (2016) A century of trends in adult human height. *Elife* 5:e13410.
- Nerlich AG, Rohrbach H, Bachmeier B, Zink A (2006) Malignant tumors in two ancient populations: An approach to historical tumor epidemiology. *Oncol Rep* 16:197–202.
- Newman SJ, Smith SA (2006) Marine mammal neoplasia: a review. *Vet Pathol* 43:865–880.
- Noble RJ, Kaltz O, Nunney L, Hochberg ME (2016) Overestimating the role of environment in cancers. *Cancer Prev Res* . doi: 10.1158/1940-6207.CAPR-16-0126
- Noble R, Kaltz O, Hochberg ME (2015) Peto's paradox and human cancers. *Philos Trans R Soc Lond B Biol Sci* 370:20150104.
- Nunney L (2013) The real war on cancer: The evolutionary dynamics of cancer suppression. *Evol Appl* 6:11–19.
- Nunney L (1999) Lineage selection and the evolution of multistage carcinogenesis. *Proc Biol Sci* 266:493–498.
- Nunney L, Maley CC, Breen M, et al (2015) Peto's paradox and the promise of comparative oncology. *Philos Trans R Soc Lond B Biol Sci* 370:20140177.
- O'Connell JF, Hawkes K, Jones N (1999) Grandmothering and the evolution of *Homo erectus*. *J Hum Evol* 36:461–485.
- Orr HA (2009) Fitness and its role in evolutionary genetics. *Nat Rev Genet* 10:531–539.
- Ostrander EA, Davis BW, Ostrander GK (2016) Transmissible Tumors: Breaking the Cancer

- Paradigm. *Trends Genet* 32:1–15.
- Owston MA, Ramsay EC, Rotstein DS (2008) Neoplasia in felids at the Knoxville Zoological Gardens, 1979–2003. *J Zoo Wildl Med* 39:608–613.
- Parkin DM, Boyd L, Walker LC (2011) The fraction of cancer attributable to lifestyle and environmental factors in the UK in 2010. *Br J Cancer* 105 Suppl 2:S77–81.
- Pavard S, Metcalf CJE (2007) Negative selection on BRCA1 susceptibility alleles sheds light on the population genetics of late-onset diseases and aging theory. *PLoS One* 2:e1206.
- Perz JF, Armstrong GL, Farrington LA, et al (2006) The contributions of hepatitis B virus and hepatitis C virus infections to cirrhosis and primary liver cancer worldwide. *J Hepatol* 45:529–538.
- Peto R (1977) Epidemiology, multistage models, and short-term mutagenicity tests. In: Hiatt HH, Watson JD, WJA (ed) *The Origins of Human Cancer*, Cold Spring Harbor Conferences on Cell Proliferation. Cold Spring Harbor Laboratory, New York, pp 1403–1428
- Peto R, Roe FJ, Lee PN, et al (1975) Cancer and ageing in mice and men. *Br J Cancer* 32:411–426.
- Pompei F, Polkanov M, Wilson R (2001) Age distribution of cancer in mice: the incidence turnover at old age. *Toxicol Ind Health* 17:7–16.
- Pompei F, Wilson R (2001) Age Distribution of Cancer: The Incidence Turnover at Old Age. *Human and Ecological Risk Assessment: An International Journal* 7:1619–1650.
- Promislow DEL (1991) Senescence in Natural Populations of Mammals: A Comparative Study. *Evolution* 45:1869–1887.
- Ramqvist T, Dalianis T (2010) Oropharyngeal cancer epidemic and human papillomavirus. *Emerg Infect Dis* 16:1671–1677.
- Ratcliffe HL (1933) Incidence and Nature of Tumors in Captive Wild Mammals and Birds. *Am J Cancer* 17:116–135.
- Reiche EMV, Nunes SOV, Morimoto HK (2004) Stress, depression, the immune system, and cancer. *Lancet Oncol* 5:617–625.
- Reinhardt HC, Schumacher B (2012) The p53 network: cellular and systemic DNA damage responses in aging and cancer. *Trends Genet* 28:128–136.
- Ricklefs RE (1998) Evolutionary theories of aging: confirmation of a fundamental prediction, with implications for the genetic basis and evolution of life span. *Am Nat* 152:24–44.
- Robbins HA, Shiels MS, Pfeiffer RM, Engels EA (2014) Epidemiologic contributions to recent cancer trends among HIV-infected people in the United States. *AIDS* 28:881–890.
- Robinson JA, Ortega-Del Vecchyo D, Fan Z, et al (2016) Genomic Flatlining in the Endangered Island Fox. *Curr Biol* 26:1183–1189.
- Rothman KJ, Greenland S (2005) Causation and Causal Inference in Epidemiology. *Am J Public Health* 95:S144–S150.
- Rozhok AI, DeGregori J (2016) The Evolution of Lifespan and Age-Dependent Cancer Risk. *Trends Cancer Res.* doi: 10.1016/j.trecan.2016.09.004
- Rudan I, Rudan D, Campbell H, et al (2003) Inbreeding and risk of late onset complex disease. *J Med Genet* 40:925–932.
- Sanchez JR, Milton SL, Corbit KC, Buffenstein R (2015) Multifactorial processes to slowing the biological clock: Insights from a comparative approach. *Exp Gerontol* 71:27–37.
- Schartl A, Malitschek B, Kazianis S, et al (1995) Spontaneous melanoma formation in nonhybrid *Xiphophorus*. *Cancer Res* 55:159–165.
- Schiffman JD, Breen M (2015) Comparative oncology: what dogs and other species can teach us about humans with cancer. *Philos Trans R Soc Lond B Biol Sci.* doi: 10.1098/rstb.2014.0231
- Schiffman J, Maley CC, Nunney L, et al (2015) Cancer across life: Peto’s paradox and the promise of comparative oncology. *Philos Trans R Soc Lond B Biol Sci* 370:1673.
- Schottenfeld D, Beebe-Dimmer JL, Buffler PA, Omenn GS (2013) Current perspective on the global and United States cancer burden attributable to lifestyle and environmental risk factors. *Annu Rev Public Health* 34:97–117.
- Schug MD, Vessey SH, Korytko AI (1991) Longevity and Survival in a Population of White-Footed Mice (*Peromyscus leucopus*). *J Mammal* 72:360–366.
- Seluanov A, Chen Z, Hine C, et al (2007) Telomerase activity coevolves with body mass, not lifespan. *Aging Cell* 6:45–52.

- Seluanov A, Hine C, Azpurua J, et al (2009) Hypersensitivity to contact inhibition provides a clue to cancer resistance of naked mole-rat. *Proc Natl Acad Sci U S A* 106:19352–19357.
- Shors AR, Solomon C, McTiernan A, White E (2001) Melanoma risk in relation to height, weight, and exercise (United States). *Cancer Causes Control* 12:599–606.
- Siddle HV, Kaufman J (2013) A tale of two tumours: comparison of the immune escape strategies of contagious cancers. *Mol Immunol* 55:190–193.
- Siddle HV, Kreiss A, Eldridge MDB, et al (2007) Transmission of a fatal clonal tumor by biting occurs due to depleted MHC diversity in a threatened carnivorous marsupial. *Proc Natl Acad Sci U S A* 104:16221–16226.
- Silva AS, Wood SH, van Dam S, et al (2011) Gathering insights on disease etiology from gene expression profiles of healthy tissues. *Bioinformatics* 27:3300–3305.
- Smith KR, Hanson H a., Mineau GP, Buys SS (2012) Effects of BRCA1 and BRCA2 mutations on female fertility. *Proc Biol Sci* 279:1389–1395.
- Song M, Giovannucci E (2016) Preventable Incidence and Mortality of Carcinoma Associated With Lifestyle Factors Among White Adults in the United States. *JAMA Oncology*. doi: 10.1001/jamaoncol.2016.0843
- Soto AM, Sonnenschein C (2010) Environmental causes of cancer: endocrine disruptors as carcinogens. *Nat Rev Endocrinol* 6:363–370.
- Sottoriva A, Kang H, Ma Z, et al (2015) A Big Bang model of human colorectal tumor growth. *Nat Genet* 47:209–216.
- Speakman JR (2005) Body size, energy metabolism and lifespan. *J Exp Biol* 208:1717–1730.
- St-Jean SD, Stephens RE, Courtenay SC, Reinisch CL (2005) Detecting p53 family proteins in haemocytic leukemia cells of *Mytilus edulis* from Pictou Harbour, Nova Scotia, Canada. *Can J Fish Aquat Sci* 62:2055–2066.
- Strakova A, Ni Leathlobhair M, Wang G-D, et al (2016) Mitochondrial genetic diversity, selection and recombination in a canine transmissible cancer. *Elife*. doi: 10.7554/eLife.14552
- Sulak M, Fong L, Mika K, et al (2016) TP53 copy number expansion is associated with the evolution of increased body size and an enhanced DNA damage response in elephants. *Elife*. doi: 10.7554/eLife.11994
- Taylor KR, Milone NA, Rodriguez CE (2016) Four Cases of Spontaneous Neoplasia in the Naked Mole-Rat (*Heterocephalus glaber*), A Putative Cancer-Resistant Species. *J Gerontol A Biol Sci Med Sci* 00:glw047.
- Thomas MA, Weston B, Joseph M, et al (2003) Evolutionary dynamics of oncogenes and tumor suppressor genes: Higher intensities of purifying selection than other genes. *Mol Biol Evol* 20:964–968.
- Tian X, Azpurua J, Hine C, et al (2013) High-molecular-mass hyaluronan mediates the cancer resistance of the naked mole rat. *Nature* 499:346–349.
- Tomasetti C, Marchionni L, Nowak M a., et al (2015) Only three driver gene mutations are required for the development of lung and colorectal cancers. *Proc Natl Acad Sci U S A* 112:118–123.
- Tomasetti C, Vogelstein B (2015) Cancer etiology. Variation in cancer risk among tissues can be explained by the number of stem cell divisions. *Science* 347:78–81.
- Tuljapurkar SD, Puleston CO, Gurven MD (2007) Why men matter: Mating patterns drive evolution of human lifespan. *PLoS One* 2:e785.
- Tuomainen U, Candolin U (2011) Behavioural responses to human-induced environmental change. *Biol Rev Camb Philos Soc* 86:640–657.
- Ujvari B, Beckmann C, Biro PA, et al (2016a) Cancer and life-history traits: lessons from host-parasite interactions. *Parasitology* 143:533–541.
- Ujvari B, Gatenby RA, Thomas F (2016b) The evolutionary ecology of transmissible cancers. *Infect Genet Evol* 39:293–303.
- Vail DM, MacEwen EG (2000) Spontaneously occurring tumors of companion animals as models for human cancer. *Cancer Invest* 18:781–792.
- Varki A (2000) A chimpanzee genome project is a biomedical imperative. *Genome Res* 10:1065–1070.
- Vickers TW, Clifford DL, Garcelon DK, et al (2015) Pathology and epidemiology of ceruminous gland tumors among endangered santa catalina island foxes (*Urocyon littoralis catalinae*) in the

- channel Islands, USA. *PLoS One* 10:1–18.
- Vijg J, Suh Y (2013) Genome instability and aging. *Annu Rev Physiol* 75:645–668.
- Vineis P, Wild CP (2014) Global cancer patterns: causes and prevention. *Lancet* 383:549–557.
- Vittecoq M, Ducasse H, Arnal A, et al (2015) Animal behaviour and cancer. *Anim Behav* 101:19–26.
- Vittecoq M, Roche B, Daoust SP, et al (2013) Cancer: A missing link in ecosystem functioning? *Trends Ecol Evol* 28:628–635.
- Vogtmann E, Goedert JJ (2016) Epidemiologic studies of the human microbiome and cancer. *Br J Cancer* 114:237–242.
- Vucenik I, Stains JP (2012) Obesity and cancer risk: evidence, mechanisms, and recommendations. *Ann N Y Acad Sci* 1271:37–43.
- Weinstein BS, Ciszek D (2002) The reserve-capacity hypothesis: Evolutionary origins and modern implications of the trade-off between tumor-suppression and tissue-repair. *Exp Gerontol* 37:615–627.
- White MC, Holman DM, Boehm JE, et al (2014) Age and cancer risk: A potentially modifiable relationship. *Am J Prev Med* 46:S7–S15.
- Wilkins R, Uppal S, Finès P, et al (2008) Life expectancy in the Inuit-inhabited areas of Canada, 1989 to 2003. *Health Rep* 19:7–19.
- Williams GC (1957) Pleiotropy, Natural Selection, and the Evolution of Senescence. *Evolution* 11:398–411.
- Wirth T, Falush D, Lan R, et al (2006) Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Mol Microbiol* 60:1136–1151.
- Wu S, Powers S, Zhu W, Hannun YA (2016) Substantial contribution of extrinsic risk factors to cancer development. *Nature* 529:43–47.
- Yachida S, Jones S, Bozic I, et al (2010) Distant metastasis occurs late during the genetic evolution of pancreatic cancer. *Nature* 467:1114–1117.
- Young TK, Kelly JJ, Friberg J, et al (2016) Cancer among circumpolar populations: an emerging public health concern. *Int J Circumpolar Health* 75:29787.
- Zhao B, Tumaneng K, Guan K-L (2011) The Hippo pathway in organ size control, tissue regeneration and stem cell self-renewal. *Nat Cell Biol* 13:877–883.
- zur Hausen H (2009) The search for infectious causes of human cancers: where and why (Nobel lecture). *Angew Chem Int Ed Engl* 48:5798–5808.

Glossary

Antagonistic pleiotropy. Expression of a single gene affecting two or more phenotypic traits, where at least one trait is beneficial to fitness, and another trait has negative fitness consequences.

Cancer driver mutations. Epigenetic alterations or mutations to key genes that result in the Hallmarks of Cancer (Hanahan and Weinberg 2000; Hanahan and Weinberg 2011), including unchecked cellular proliferation, tissue invasion and metastasis. In addition to those necessary for the Hallmarks of Cancer, other driver mutations may increase the selective growth advantage of cancer subclones resulting in accelerated disease spread (Bozic et al.

2010) (but see (Sottoriva et al. 2015)).

Cancer risk. The probability over an individual's lifetime of cancer emerging and having performance, health, survival or reproduction consequences.

Cellular cheating. Achieving one or more of the Hallmarks of Cancer, resulting in deregulation of proliferation inhibition, cell death, division of labour, resource allocation and extracellular environment maintenance (Aktipis et al. 2015).

Comparative oncology. The study of cancer defences and carcinogenesis between species, individuals within a species, or between tissues within individuals.

Force of selection. The fitness of an allele expressed at age x , measured as remaining fecundity and contributions (e.g., resource transfers) to the survival and fecundities of close kin.

Invasive carcinoma. A growing population of epithelial cells that exhibit all of the Hallmarks of Cancer (Hanahan and Weinberg 2000; Hanahan and Weinberg 2011), infiltrate or destroy surrounding tissue, and possibly metastasize.

Microenvironment. The immediate environment of a cell, including other cells, the vascular system, nutrients, immune factors, and chemical signals.

Multistage theory of carcinogenesis. The theory that cancers proceed through successive stages, corresponding to one or more mutations (or epigenetic alterations) activating oncogenes and disabling tumour suppressor genes (Armitage and Doll 1954; Frank 2010). Some of these mutations may be inherited (Frank 2004a), increasing the risk of certain cancer types obtaining .

Novel environment. The full set of environmental conditions, including those stemming from behaviour and ecology, that tend to differ from the conditions under which a species evolved, and (in the context of this study) which result in a significant effect on cancer risk (Figure 1). Novel environments can be either quantitative (e.g., increased temperatures,

increased caloric intake), or both quantitative and qualitative (e.g., pollutants, infectious disease, selective breeding). They may also be distal in causation, for example life in captivity, healthcare, habitat restriction leading to bottlenecks and inbreeding. We hypothesize that novel environments will usually increase overall cancer risk, but it is possible that some cancers will decrease (e.g., the invasion of a generalist predator that tends to kill older individuals before cancer typically obtains).

Peto's paradox. The empirical observation that neither species longevity nor body size correlates with cancer risk. The same observation has been identified for tissue families within organisms (Noble et al. 2015).

Figure 1

Figure 2

Figure 3

Figure 4

Figure Legends

Figure 1. Schematic representation of how a novel environmental variable is associated with increases in cancer risk above levels established by long-term evolution. The long-term exposure to an environmental variable (e.g., UV radiation) (black solid line) is expected to select for adaptations (see Introduction in main text) that form a norm of reaction in cancer risk (red line) across the distribution of the environmental variable. Novel environments shift the frequency distribution of the environmental variable (black dashed line) and associated cancer risks. Shaded red area around mean risk line either indicates variability among

individuals within a population, or between populations of a given species. Note that novel environments may result in *decreases* in cancer risk, as for example, following the invasion of a predator or parasite, that shifts species demography to younger (less cancer prone) ages.

Figure 2. Hypothetical effects on cancer risk of variation in *The Organism, Within-Organism, and The Genome*. Specific abiotic and biotic environment variables (including lifestyle behaviours) will alter one or more of these biological levels and in so doing, increase cancer beyond baseline risks associated with environments experienced over evolutionary time (“native” environments) (see Fig. 1). Whereas we expect interactions between whole organism and within-organism effects (left panel) on cancer risk, the interactive effects of changes to genomes (right panel) will depend, in part, on whether genetic effects selectively increase cancer risk in young individuals (e.g., mutations in *RB* and retinoblastoma in humans), or rather, are (partially) age related (e.g., *BRCA1* mutations and breast cancer in humans).

Figure 3. A conceptual framework for how environmental factors influence cancer risk over ecological and evolutionary timescales. Intrinsic and environmental/behavioural factors (green boxes) ultimately determine cancer risk (red boxes) through their effects on the proximate mechanism of somatic evolution (blue boxes). In general, a population of stable size in a stable environment is expected to evolve low cancer risk, because cancer-affected individuals have lower fitness. Environmental change or variation (yellow boxes) can increase cancer risk if it outpaces organismal evolution. Within-species variation in cancer risk results from genetic diversity, environmental heterogeneity, and from stochasticity in the mutational process (“bad luck”; see also Box 3).

Figure 4. Relationship between longevity and cancer risk in humans, based on a mathematical model using empirical data. (A) Relative force of selection (red line, qualitative approximation based on Lee (Lee 2003); for illustrative purposes; not used in the model) and relative risk of acquiring cancer for the first time (blue line, used as model input, estimated using the DevCan software package (Fay 2004; DevCan: Probability of Developing or Dying of Cancer Software 2005), based on US data for 2010-2012). (B) Survivorship curves generated by the model (thin lines), with empirical curves for hunter-gatherers, Sweden in 1861, and the US in 2011 (thick lines) for comparison. Mortality in children less than 5 years old is excluded so as to better reflect typical life expectancy among individuals who are at substantial risk of cancer. Model results are shown for low cancer risk (left panel, using as input the US cancer incidence curve rescaled by 0.4 for adults) and high cancer risk (right panel, using as input the US cancer incidence curve). In both cases, the death rate of individuals with cancer in the model is assumed to be 0.1 per year. (C) Lifetime cancer risk (for those alive and cancer-free at age 5) versus life expectancy at age 5, according to the model. Each lifetime risk curve shows the effect of varying the parameters of background mortality, such that the survivorship curve varies within a family of curves resembling those shown in the previous panel. Crosses mark model predictions for the modern US population; a population identical to the modern US population except that the incidence of cancer per year is 30% lower in adults; and a population with the background mortality of hunter-gatherers and with 30% lower incidence of cancer per year in adults, which is assumed to provide an upper bound for cancer risk in pre-modern humans. A 30% difference in cancer incidence can be accounted for by lifestyle factors, or by a difference in stature of

approximately 19 cm (Supplementary methods). (D) Cancer risk by age 40 (for those alive and cancer-free at age 5) versus life expectancy at age 5, according to the model. Crosses are as in the previous panel. When cancer incidence and the cancer death rate remain constant, cancer risk by age 40 is positively correlated with life expectancy. This is because higher life expectancy corresponds to less background mortality, which results in more individuals being alive and susceptible to cancer.

Box 1. Relative contributions of life span and body mass to cancer risk

The multistage theory of carcinogenesis as envisaged by Armitage and Doll (1954) posits that several independent mutations are typically required to initiate cancer. This process implies that, all else being equal, the more stem cell divisions occurring over a lifetime, the higher the probability of a given cancer being obtained (Albanes and Winick 1988), since cells are most vulnerable to mutation during the cell cycle (Cairns 2002; Branzei and Foiani 2005). The lifetime number of stem cell divisions (and therefore cancer risk) is expected to correlate with stem cell number (body size), stem cell division rate, and life span (Nunney 1999; Noble et al. 2015). Focusing on body size and life span, theory predicts that the former should have a smaller effect on cancer risk than the latter (Nunney 1999). To see this, consider a very simple multi-stage model for cancer risk, R

TO BE ADDED (1)

where s is the number of stem cells, u is the mutation rate at any one of M genes necessary for cancer, and d is the number of divisions per stem cell over a lifetime. **Rob: other assumptions? This simple model assumes that $ud < 1$.** Interestingly, these cellular processes

can be reinterpreted as life history traits, whereby s correlates with body mass and d with life span. To see the difference in the relative importance of body size (s) and life span (d), consider a cancer requiring three mutations ($M=3$) (Tomasetti et al. 2015). Here, increasing d by a factor of 10 changes R as much as multiplying s by a factor of 1000.

Equation 1 and related formulae (e.g., (Calabrese and Shibata 2010)) characterize R as cellular transformation to invasive carcinoma (Glossary), and not the related definition employed in our study, the added risk of subsequent effects (e.g., tissue invasion and metastasis) on performance or health. A more complete model would incorporate the possibility that larger body sizes require more time for cancer foci to have impacts on individual performance or health. Moreover, whether modelling cancer emergence only, or both emergence and progression, Equation 1 neglects age-related changes in parameters, which has been hypothesized to explain differences in cancers typically occurring before and during reproduction ages with those usually associated with the aging process (e.g., (Rozhok and DeGregori 2016)).

Box 2. Cancer and parasites

Although cancer cells are comparable to some parasitic organisms in that grow and may spread and multiply within the host, there is one fundamental difference: unlike free-living parasites, cancer emerges from self, and is not transmitted between individual hosts (but see exceptions below). Sporadic somatic cancer is therefore an evolutionary dead end. Cancer genes, however, can evolve (and be positively selected (Crespi and Summers 2006)) beyond the host individual as vertically transmitted allelic variants within the genome. Evolutionary theory predicts that selection on vertically transmitted parasites typically results in lowered virulence (Frank 1996). Virulence in the context of cancer genes (i.e., probability of cancer

expression, rates of tumour progression and metastasis) would reduce the survival and reproduction of hosts, resulting in purifying selection on such genes and/or positive selection on other genes contributing to cancer suppression. Another possibility, similar to certain parasites (Michalakis et al. 1992)), is that cancer is antagonistically pleiotropic on positively genes (Crespi and Summers 2006) (see also discussion in main text in section on *Challenges to how we perceive and quantify cancer*).

Transmissible cancers share features with certain horizontally transmitted parasites (Ostrander et al. 2016; Ujvari et al. 2016b), including intra-specific (Murgia et al. 2006; Siddle et al. 2007; Metzger et al. 2015) and interspecific (Metzger et al. 2016) infection. Transmissible cancers are also similar to some parasites in that emergence and spread may depend on environmental conditions (Lafferty and Kuris 1999). For cancers, this includes low genetic diversity in Tasmanian devils and in dogs (Murgia et al. 2006; Siddle et al. 2007; Miller et al. 2011), and pollutants in molluscs (J. Sherry, pers comm: (St-Jean et al. 2005; Muttray et al. 2012) but see (Barber 2004) for both examples and counterexamples)).

There is also recent evidence that (similar to many parasites) canine transmissible cancers may have evolved via recombination (Strakova et al. 2016). However, it is not known for this or other transmissible cancers whether evolutionary rates are similar to parasitic species with comparable life cycles and life histories. Although not experimentally demonstrated, the low virulence of transmissible cancers in dogs (evolved over thousands of years) and the high virulence of facial tumours in Tasmanian devils (tens of years) ((Murgia et al. 2006; Siddle and Kaufman 2013); but see (Epstein et al. 2016)) are consistent with predictions of the evolutionary attenuation of initially high virulence (Frank 1996; Wirth et al. 2006; Ashby and Boots 2015).

Box 3: Cancer and “bad luck” in humans

Tomasetti and Vogelstein (Tomasetti and Vogelstein 2015) compiled data on 31 cancer types in humans to assess the relative contributions of a random factor (“bad luck”) and environmental and inherited factors to cancer incidence. They proposed that the random factor should correlate with the total lifetime number of stem cell divisions per tissue or organ type, assuming that cancer-causing mutations have a fixed probability of emerging per stem cell division. Stem cells divide during ontogeny and in tissue repair, the latter being associated with, for example, injury, parasitism, mutagenic exposure, cellular aging and cellular diseases other than cancer. Their analysis uncovered a strong positive association between cancer incidence and the lifetime number of stem cell divisions, which they interpreted as indicating that random mutations due to replication and repair errors (“bad luck”) explain much of the variance in risk among cancer types in humans. In a subsequent study, Wu et al. (Wu et al. 2016) claimed, on the contrary, that extrinsic environments play a much larger role in human cancer risk than does “bad luck”. Such contrasting results highlight the challenges in untangling the effects of intrinsic processes and extrinsic environments on cancer risk (Noble et al. 2015; Noble et al. 2016). In the more complete framework proposed here, environmental and intrinsic factors interactively determine the parameters of somatic evolution, which, in turn, contributes to cancer risk (Fig. 2). Some of these factors (e.g., those due to lifestyle) are preventable, and account for many, possibly the majority of present-day human cancers (Doll and Peto 1981; Schottenfeld et al. 2013; White et al. 2014; Colditz and Sutcliffe 2016; Song and Giovannucci 2016), whereas others (e.g., inherited cancer risks, old age) magnify within-organism disruption (shape of risk isoclines in Fig. 2), and together with particular risks of certain cancers across ethnic groups and geographical regions (Vineis and Wild 2014), should be taken into consideration in determining risk status.

Box 4: Predictions

Below, we list several predictions based on one or more of the following: Equation 1 (Box 1), interpretations of how environment influences cancer risk (Fig. 3), empirical examples (main text), and the parameterised model for the human population (Supp. info.).

1. Populations with greater than *c.* 5% overall cancer risk are notably impacted by novel environments. Environmental drivers—not necessarily stemming from human influences—include parasites, increased senescence following predator extinction, genetic bottlenecks (e.g., island species), and founder effects. All else being equal, for taxonomic or interpopulation comparisons, reduced performance due to physiological aging (or senescence) should correlate with increased lifetime cancer risks in either native or novel environments.
2. The effect of disrupted bodies on cancer risk relative to the baseline risk is larger when life expectancy is lower, but the absolute effect is greater in longer-lived populations. For example, if the lifetime cancer risk in the native environment is 1% and in the novel environment is 1.3%, then the relative change is a 30% increase, and the absolute change is a 0.3 percentage points increase. Therefore, the effects of disrupted bodies are more likely to be observed in species with longer life spans and senescence.
3. Marginal increases in survival, especially post-reproduction, will have a greater marginal impact on cancer risk than marginal increases in body size (see also (Frank 2004c)).
4. To the extent that cancer reduces the performance or survival of senescing individuals, it will (not surprisingly) have less impact on the ecology and evolution of the species. Notable exceptions are social species in which older individuals play important roles in group dynamics (e.g. many primates).

5. Based on a first approximation of linear marginal effects of environmental change on each parameter in Eqn. 1, Box 1, and the realistic assumption that most cancers require two or more mutations (in humans, where this is best understood (Knudson 2001)), we predict that environmental impacts on mutation rate (u) or life span (as a proxy for total stem cell divisions per lineage, d) will have exponential implications for cancer risk, whereas the influence on body size (as a proxy for total number of stem cells s) will be linear. Whether environmental perturbations alter the number of mutations required for a cancer to obtain (M) is an open question, but if they do, then their effects may be highly non-linear depending on the baseline level of M .

6. Similar to prediction 5, the relation between life span d and body size s : (Speakman 2005) leads to the prediction that when selection for larger body size coevolves with cancer resistance (Kokko and Hochberg 2015) so as to maintain a constant risk level, (correlated) longer life spans will have a greater marginal effect than evolving larger body size alone, but only when $M > 5$. In contrast, for $M < 5$, body size will have a greater marginal effect than correlated changes in life span. Note that this prediction assumes that coevolution will follow the same relation as identified by Speakman (2005) for interspecific comparisons.

7. Responses to selection of environmentally-driven cancers will frequently be slow or insignificant due to the long generation times of many vulnerable species, the low selective value of older individuals typically impacted by cancers, the probabilistic nature of tumorigenesis, the complexity of the genetic basis for cancer sensitivity, and/or the low population frequency of sensitivity genes.

8. Species with few and/or weak mechanisms of cancer resistance (which nevertheless are effective in native environments; Fig. 1), are particularly vulnerable to cancers stemming from environmental vagaries.

9. Increasing US life expectancy by one year (through decreases in background mortality) is predicted to increase lifetime cancer risk by *c.* 1%.