



# City Research Online

## City St George's, University of London

**Citation:** Moghadam, K., Chehrzad, M., Masouleh, S., Mardani, A., Maleki, M., Akhlaghi, E. & Harding, C. (2021). Nursing workload in intensive care units and the influence of patient and nurse characteristics. *Nursing In Critical Care*, 26(6), pp. 425-431. doi: 10.1111/nicc.12548

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/25079/>

**Link to published version:** <https://doi.org/10.1111/nicc.12548>

**Copyright and Reuse:** Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

1 **Title: Nursing Workload in ICUs and the Influence of Patient and Nurse**  
2 **Characteristics**

3 **Kolsoom Nasirizad Moghadam MScN | Minoo Mitra Chehrzad PhD | Shademan Reza**  
4 **Masouleh MScN | Abbas Mardani MScN | Maryam Maleki MScN | Elham Akhlaghi MScN**  
5 **| Celia Harding PhD**

6 *Journal of Nursing in Critical Care*

7

8 **Abstract**

9 **Background:** Understanding factors that can potentially influence patient care and nursing  
10 workload in intensive care units is important. Previous studies have shown contradictory outcomes  
11 about the relationships between nursing workload and patient and nurse characteristics.

12 **Aims and objectives:** This study aimed to investigate nursing workload in intensive care units  
13 and examine the association between this in relation to patient and nurse characteristics.

14 **Design:** A cross-sectional design was conducted.

15 **Methods:** All nurses who were working in the intensive care units of five hospitals and met the  
16 study criteria were enrolled in the study. Two demographic questionnaires collected nurse and  
17 patient demographic information. The Nursing Activities Score was applied to determine nursing  
18 workload in three shifts (morning, evening, night) for each nurse. Data were analyzed using the  
19 independent sample t-test, one-way analysis of variance, and multivariable linear regression  
20 analysis.

21 **Results:** The Nursing Activities Score was calculated for 509 patients who were under the care of  
22 the 105 intensive care unit nurses. The mean (SD) nursing activities score was 72.84% (22.07%).  
23 Morning shifts, male patients, medical treatments, and referred patients from the emergency ward  
24 and other intensive care units imposed a higher workload for nurses. Specifically, female nurses,  
25 increased number of patients receiving care, and increased patient length of intensive care unit  
26 stay were directly associated with increased nursing activities scores. Work in surgical and burn  
27 intensive care units were inversely associated with the nursing activities score.

28 **Conclusion:** This study suggests that the workload of nurses in intensive care units can be affected  
29 by both nurse and patient characteristics.

30 **Relevance to clinical practice:** The findings can be used to ensure appropriate staffing of  
31 intensive care units by nurses. However, nurse and patient characteristics should not be considered  
32 as the only factors which influence nursing workload in intensive care units.

33 **Key words:** *Critical care; ICU; Intensive care units; Nursing; Workload*

34

## 35 1. INTRODUCTION

36 Health-care workers are crucial in supporting and caring for patients (Heckemann et al., 2015).  
37 Within typical health organizations, nurses are the largest workforce and play a key role in the  
38 quality of care and health promotion (Maenhout and Vanhoucke, 2013). They include 62% of all  
39 the hospital staff (Momennasab et al., 2017). According to the National Association of Safety  
40 Professionals in the United States, nursing is one of the forty professions with a high prevalence  
41 of work-related stress (Yusefi et al., 2019). For that reason, workload is known to influence the  
42 behavior and performance of nurses in the workplace (Holland et al., 2019). Generally, the nursing  
43 workload is determined by time spent on patient care, nursing activities, and the skills needed to  
44 care for the patient (Myny et al., 2012). The nursing staff workload is clearly related to patient  
45 safety, quality of care, and cost of health care (Swiger et al., 2016).

46 The Intensive Care Unit (ICU) is an environment that provides care for patients with severe clinical  
47 conditions which require ventilation and acute medical clinical care (Ferreira et al., 2017). Nurses  
48 in ICUs have extensive and intense duties with limited variation in tasks, exposing them to  
49 extremely high workloads both physically and mentally (Mohammadi et al., 2015). There can be  
50 a need to react to both the demands of patients as well as their families. In addition, they are  
51 frequently making many decisions in relation to the urgent and critical conditions of patients' lives  
52 (Mohammadi et al., 2015). Abbey et al. (2012) report that nurses in the ICU perform 3,081  
53 different activities during the day, 43% of which are performed simultaneously. Consequently,  
54 nurses may be at risk of making more errors with ICU patient care procedures, thus impacting on  
55 patient safety. In the ICU, the high workload and the low number of nurses per patient increases  
56 the risk of nosocomial infections in patients and mortality. Therefore, the ICU in the hospital is a

57 stressful work setting due to the complexity of patients and the high need for direct patient care  
58 (Hoogendoorn et al., 2019).

59 Given the importance of nursing workload in ICU and the factors that can affect it, this study aimed  
60 to determine the nursing workload in the ICUs and examine the association between nursing  
61 workload and patient and nurse characteristics.

## 62 **2. METHODS**

### 63 **2.1. Design and Participants**

64 A cross-sectional study was conducted on nurses and their patients under their care in the ICUs of  
65 five teaching hospitals affiliated to the ××× University of Medical Sciences, Rasht, Iran in 2015-  
66 2016.

67 In this study inclusion criteria for the recruitment of participants included: having an  
68 undergraduate/postgraduate degree in nursing, being responsible for the direct care of at least one  
69 patient alone during the shift, having at least one year's experience working in an ICU, having  
70 direct face to face care of a patient at least 4 hours in ICU per shift, and provision of signed consent  
71 to participate in the study. The exclusion criteria in this study included nurses who did not meet  
72 the inclusion criteria.

73 For sampling, the researchers contacted the study ICUs, then used face to face interviews to inform  
74 124 nurses of the project and the participant inclusion criteria. Nurses responded to the researchers  
75 if they were interested. The nurses were screened by the study team to ensure they met the inclusion  
76 criteria.

### 77 **2.2. Measures**

#### 78 **2.2.1. Demographic Data Questionnaires**

79 Two demographic data questionnaires were developed based on the review of the literature to  
80 collect demographic characteristics of nurses and the patients in their care. Nurses' demographic  
81 data questionnaires included questions about a nurse's gender, marital status, educational level,  
82 work experience, weekly working hours, employment at the other health care centers, resting  
83 before and after the work shift and type of ICU worked in. Patients' demographic data  
84 questionnaires contained questions about patients' gender, age, treatment types, referral sources,  
85 and length of ICU stay. Content and face validity methods were applied to confirm the validity of  
86 these questionnaires.

### 87 **2.2.2. Nursing Activities Score**

88 The Nursing Activities Score (NAS) was used in this study to determine nursing workload. The  
89 NAS designed by Miranda et al. (2003) measures the nursing workload in ICUs. This instrument  
90 consists of 23 items which are divided into seven major categories including basic measures,  
91 ventilation support, cardiovascular support, renal support, neurological support, metabolic support,  
92 and specific intervention. Each item is scored based on the amount of time the nurse spends on  
93 patient care. Although the original version of the NAS was validated to measure the workload of  
94 nursing over a 24 hour period, the modified version of this tool was validated by Debergh et al.  
95 (2012) to measure nursing workload per shift. In fact, for each patient, the scores of the NAS were  
96 calculated during each shift unless the time spent with a patient was less than 4 hours. The NAS  
97 ranges from 0 to 177 % that indicates the sum of scores of 23 items which demonstrates the  
98 percentage of time spent by a nurse on care-related tasks per ICU patient per 24 hours. Each NAS  
99 point corresponds to 14.4 minutes. Ideally, one nurse who provides care for two patients will obtain  
100 a score of 50% for each patient. A NAS of 100% is the ideal score for a nurse per shift in a 24-h  
101 period and would demonstrate that the nurse spent 100% of the working time caring directly for

102 patients. A NAS of above 100% indicates that more than one nurse is needed to perform caring  
103 activities. Therefore, a NAS of 177% also would equal 1.77 full-time equivalent ICU nurse time  
104 (Miranda et al., 2003, Debergh et al., 2012).

105 To evaluate the validity of the NAS, the original version with the available Farsi version (translated  
106 by Alizadeh et al., (2015)) was provided to 10 faculty members of the nursing school who  
107 specialize in this area. After collecting their opinions, corrective feedback was considered and the  
108 results (content validity ratio (CVR) = 93% and content validity ratio (CVI) = 90%) indicated a  
109 high validity of the Farsi version of the NAS. In addition, the equivalent reliability method was  
110 used to determine the reliability of the NAS. The NAS was completed by two evaluators  
111 simultaneously and separately for 17 nurses who were providing nursing care for ICU patients.  
112 The results showed there was a high correlation between the scores of two evaluators based on the  
113 Pearson correlation coefficient ( $r = 0.81$  and  $p < 0.001$ ). In addition, there was no significant  
114 difference between the mean of the two evaluator scores based on a paired t-test ( $p = 0.72$ ).

### 115 **2.3. Data Collection**

116 Data were collected in six adult ICUs (two general ICUs with 19 beds, one neurosurgical ICU with  
117 8 beds, one cardiac surgery ICU with 6 beds, one obstetric surgery ICU with 3 beds, and one burn  
118 ICU with 4 beds) of 5 hospitals during a fourth-month period from November 2015 to February  
119 2016. The demographic questionnaire was completed on paper by the nurses and the patient  
120 demographic questionnaire was completed by the researchers during the study. The NAS was  
121 completed on paper for each nurse in three shifts (morning, evening, night). The researchers  
122 completed the NAS instrument in three mentioned shifts for each patient under the care of the  
123 nurses participating in the study based on direct observation of nursing activities, the information  
124 in the patient records and nursing reports. If the nurses were responsible for taking care of two

125 patients, the NAS was completed for each patient separately. Next, to obtain the score of the  
126 workload per shift, the scores of the NAS for each patient were summed together. Finally, the  
127 mean scores of the NAS in three shifts (morning, evening and night) were considered in relation  
128 to nursing workload.

#### 129 **2.4. Ethical Considerations**

130 ××× University of Medical Sciences approved the study protocol which was assigned the following  
131 code: IR.GUMS.REC.1394.286. The necessary permissions were obtained from the hospital  
132 authorities prior to the sampling. Next, information about the study was given to study participants  
133 (nurses and their patients/ patient companions). Then, the informed consent form was signed by  
134 participants who agreed to participate in this study. They were informed that they could opt-out of  
135 the study at any time without being penalized.

#### 136 **2.5. Data Analysis**

137 Data were analyzed using the SPSS v. 25 software (IBM, Armonk, NY, USA) through descriptive  
138 and inferential statistics. Frequency and percent for qualitative variables and mean and standard  
139 deviation for quantitative variables were used to describe participants' characteristics. Nursing  
140 workload was described using means and standard deviations. In addition, an independent sample  
141 t-test and One-Way ANOVA were performed to compare nursing workload according to working  
142 shifts, patient genders, patient treatment types, and patient admission referrers. Moreover,  
143 multivariable linear regression analysis using the SATA software (Version 15, Stata Corporation,  
144 and College Station, TX, USA) was conducted to investigate nursing workload association with  
145 characteristics of nurses and patients. P-value < 0.05 was considered as a significant level.

### 146 **3. RESULTS**

147 Out of 124 nurses who work in six ICUs 7 nurses (5.6%) declined to participate in the study and  
148 12 nurses (9.7%) including head nurses and assistant head nurses were excluded because they were  
149 not directly looking after patients. A total of 105 nurses (84.7%) who met the inclusion criteria  
150 were enrolled in the study. The majority of nurses in this study were female (96.2%), between the  
151 ages of 30-40 years old (61%), and married (59%). Most of them had a bachelor's degree in nursing  
152 (94.3%). Only 17.1% of nurses were employed at the other healthcare centers. In addition, 45.7%  
153 of nurses worked in general ICUs, 19% in a neurosurgical ICU, 20% in a cardiac surgery ICU,  
154 10.5% in a obstetric surgery ICU, and 4.8% in a burns ICU were employed. The burns ICU had  
155 14 nurses but due to a lack of patients at the time of the study only 5 nurses participated. Moreover,  
156 the mean (SD) of nurses' resting before and after the work shift was 33.43 (13.44) and 30.91  
157 (11.73) hours, respectively. The mean number of patients under the care of nurses was 1.61 (0.40).  
158 Also, the mean of nurses' work experiences was 8.14 (4.72) years and the mean weekly working  
159 of them was 48.21 (8.59) hours.

160 Furthermore, 509 patients were under the care of nurses during the study period. The majority of  
161 patients in this study were male (56.6%). The mean age of patients participating in the study was  
162 48.85 (20.31) years, and the mean length of stay on ICU for all participants was 6.31 (5.51) days.  
163 Most referrals to the ICU came from the operating theatre (52.3%) and emergency ward (34.6%).  
164 Of this sample, 63.1% of patients needed surgery and 36.9% required medical treatment only.

165 The mean (SD) of the nursing workload in three shifts (morning, evening, night) was 72.84%  
166 (22.07%). According to Table 2, there was a significant difference between the workload  
167 experienced by nurses on different work shifts ( $p = 0.001$ ). It was noted that there were significant  
168 differences between the workload in morning shift in comparison with the evening shift (mean  
169 difference (MD) = 10.85,  $p = 0.002$ ) and night shift (MD = 13.37,  $p = 0.001$ ). Male patients

170 produced a significantly higher workload for nurses ( $p < 0.001$ ) in contrast to female patients.  
171 Patients who received medical treatments caused a significantly higher workload for nurses than  
172 patients who had received surgery ( $p < 0.001$ ). Moreover, a statistically significant association was  
173 identified between the nursing workload and patient referral sources ( $p < 0.001$ ). Based on the post  
174 hoc test, there was a significant difference between the patients admitted from the operating theatre  
175 with patients admitted from the emergency ward (MD = -4.80,  $p < 0.001$ ) and patients admitted  
176 from other ICUs (MD = -7.99,  $p = 0.02$ ).

177 Based on the multivariable linear regression analysis, a positive association was found between  
178 nursing workload and female nurses ( $b = 6.27$ , 95% CI = 1.41, 11.13). Also, there was a direct  
179 association between workload of nursing and number of patients under their care ( $b = 25.04$ , 95%  
180 CI = 15.13, 34.96) and the length of patients' ICU stay ( $b = 0.75$ , 95% CI = 0.53, 0.97). Moreover,  
181 an inverse relationship was seen between nursing workload and working in neurosurgical ICU ( $b$   
182 = -7.91, 95% CI = -12.91, -2.91), cardiac surgery ICU ( $b = -14.58$ , 95% CI = -20.87, -8.29),  
183 obstetric ICU ( $b = -31.91$ , 95% CI = -39.15, -24.67), and burns ICU ( $b = -31.18$ , 95% CI = -41.22,  
184 -21.14) (Table 2).

#### 185 **4. DISCUSSION**

186 The purpose of this study was to determine the nursing workload in ICUs and investigate the  
187 association between nursing workload and patients' and nurses' characteristics. In this study, the  
188 mean NAS of the nursing workload was reported at 72.84%. The average nursing workload in  
189 ICUs reported in other studies shows a wide variation (Esmaili et al., 2015). A study of  
190 multidisciplinary ICUs in Norway showed that the nursing workload was 96.2% using the NAS  
191 tool (Stafseth et al., 2011). The findings of a recent study conducted in Brazil revealed that the  
192 nursing workload using the NAS in ICUs is lower at 63.5% (Strazzieri-Pulido et al., 2019). The

193 discrepancy of nursing workload in these studies could be due to different nurse staff frameworks  
194 and organizations, differences in the types of patients admitted to ICUs and the technical and  
195 practical differences in the use of the NAS tool (Stafseth et al., 2011, Strazzieri-Pulido et al., 2019,  
196 Padilha et al., 2015).

197 To our knowledge, this study is one of the limited studies (Bruyneel et al., 2019, Debergh et al.,  
198 2012) that evaluated the NAS for individual nurses across all three shifts. In comparison to  
199 evaluating the NAS over 24 hours, evaluation using the NAS per shift can possibly provide better  
200 data to determine a precise nurse-patient ratio. The findings of the current study indicated that ICU  
201 nurses experience a higher workload in the morning shift than in the evening and night shifts.,  
202 despite having a higher number of nurses in the morning shift (Ducci et al., 2008). Most patient  
203 admissions, patients discharges, patient treatment procedures and follow-up administration are  
204 performed during the morning shift (Armstrong et al., 2015). In addition, the number of nurses  
205 does not decrease significantly in evening and night shifts in ICUs despite the possibility of new  
206 patient admissions or the deterioration of patients' conditions (Debergh et al., 2012). Findings of  
207 a study undertaken in 16 hospitals in Belgium demonstrated that ICU nurse workload in the  
208 morning shift was considerably higher than evening and night shifts (Bruyneel et al., 2019).  
209 Debergh et al.'s (2012) study also showed that workload in the night shift is lower than evening  
210 and night shifts.

211 The results of the present study identified that male patients were associated with a higher  
212 workload for ICU nurses. Similarly, findings of another study also showed that male trauma  
213 patients in ICUs create a higher workload for nurses (Nogueira et al., 2014). Two other studies  
214 also found a positive association between male patients and increased nursing workload in ICUs  
215 (Padilha et al., 2008, Samuelsson et al., 2015). Although evidence is currently limited, it has been

216 suggested by some previous studies that male patients are more likely to be hospitalized in ICUs  
217 and as a result of severe injuries, are likely to receive aggressive life support (Mahmood et al.,  
218 2012, Fowler et al., 2009).

219 In our study, patients admitted with specific medical diagnoses ' were associated with a higher  
220 workload than surgical admissions. In addition, there was a direct association between having a  
221 higher nursing workload for nurses working in general ICUs. Similarly, Padilha et al.'s (2007)  
222 study results highlighted that nursing workload in general ICUs was higher than surgical and burn  
223 ICUs. Lucchini et al. (2015) used the NAS to evaluate nursing workload on ICUs and found  
224 general ICUs in contrast with surgical ICUs imposed a higher workload for nurses. The higher  
225 nursing workload of general ICUs may be due to the increased complexity and intensity of patient  
226 care such as the need for hygiene procedures, mobilization, and positioning (Reich et al., 2015,  
227 Lucchini et al., 2015). Furthermore, Padilha et al. (2008) documented that patients receiving  
228 medical treatments in comparison with those who have had surgery created a higher nursing  
229 workload in ICUs. Although according to our data, the type of patients' treatments and type of  
230 ICU can be considered one of the factors that may influence the workload of nursing in different  
231 ICUs, the previous studies have demonstrated that the intensity of the patient's condition more  
232 important predictor of nursing workload in clinical practice (Altafin et al., 2014, Nogueira et al.,  
233 2014, Romano et al., 2019). For instance, Altafin et al. (2014) found patients who died required a  
234 higher workload for ICU nurses than those who survived.

235 In our study, a significant association was identified between nursing workload and patient referral  
236 source. The referral sources of ICU patients can help predict nursing workload (Padilha et al.,  
237 2008). Consistent with our results, another study found that the mean percentage NAS for patients  
238 admitted from the emergency department was higher than for those admitted from the operating

239 theatre and general ward (Armstrong et al., 2015). However, this study also reported that the  
240 workload of patients who were admitted from other ICUs was not high which is in conflict with  
241 our results (Armstrong et al., 2015). Romano et al. (2019) in their study concluded the type of  
242 admission to ICUs provides no valuable guidance to predict the required nursing workload for  
243 patient care. This is inconsistent with our study and other published studies (Coelho et al., 2011,  
244 Padilha et al., 2008).

245 Our study found a direct association between nursing workload and the number of patients under  
246 the care of nurses which would be expected. Similarly, previous studies also showed a significant  
247 relationship between the number of patients and the nursing workload (Penz et al., 2007, Coventry  
248 et al., 2015). In fact, when the number of patients increase, the responsibility of nurses and the  
249 time spent on direct patient care increases (Yusefi et al., 2019, Hugonnet et al., 2004, Halwani et  
250 al., 2006).

251 We showed that there was a direct association between the patients' length of stay on ICU and  
252 nursing workload. The relationship between NAS and length of ICU stay in the literature is  
253 conflicting. Consistent with our findings, Padilha et al. (2008) reported that patients who remain  
254 longer in the ICUs, for each extra ICU-day, have a 7.0% more probability of increasing the NAS.  
255 Lucchini et al. (2014) during a five-year study with 5856 patients found the NAS was high in  
256 patients with the increasing length of ICU stay. However, findings of another study revealed that  
257 the NAS decreased with increasing patient length of stay in the ICU (Altafin et al., 2014).

258 In the present study, we found a direct association between the nursing workload and female  
259 nurses. Consistent with our findings, Hoonakker et al. (2011) and Mirzaei et al. (2015) found  
260 female nurses in ICUs experienced a higher workload. Female nurses may, in addition to the  
261 stressful work environment in ICU face personal stressors including daily housework, parenting,

262 and childcare demands, and challenges to the work-life balance that lead to perceived high  
263 workload and burnout (Papazian et al., 2018).

#### 264 **4.1. Implications and Recommendations for Practice**

265 Our study extends the previous knowledge about nursing workload in ICUs and their related  
266 factors. The findings of this study can be used by nursing and hospital managers to provide an  
267 appropriate allocation of nursing staff on ICUs with additional consideration given to nurse and  
268 patient characteristics, and can therefore improve the quality of care in ICUs. However, nurse and  
269 patient characteristics should not be considered as the only predictive factors of nursing workload  
270 in ICUs.

#### 271 **4.2. Strengths and Limitations**

272 Our study investigated and measured the workload of each nurse in the morning, evening and  
273 night shifts, therefore differing from the other studies cited in our literature review. Our study has  
274 several limitations that should be considered. Firstly, a cross-sectional design was used in this  
275 study; therefore, the results should not be considered causal. Secondly, this study was conducted  
276 in Iran, which possibly decreases the generalizability of the results to other countries. Thirdly, the  
277 NAS tool was originally developed to be applied over a 24-hour period. Although Debergh et al.  
278 (Debergh et al., 2012) demonstrated that it could be used per shift, it seems that using this tool per  
279 shift requires further validation in larger studies. Fourthly, although the results showed a  
280 significant association between workload and nursing gender, this finding may be due to a greatly  
281 uneven number of male nurses in our sample. Finally, the data for this study were collected about  
282 4 years ago which could generate bias.

#### 283 **Conclusion**

284 Our study shows that the nursing workload in ICUs was associated with both nurse and patient  
285 characteristics. Type of work shifts, nurse gender, type of ICU which the nurses worked in, the  
286 number of patients under a nurse's care, patient gender, medical or surgical treatment for patient,  
287 patient referral source, and the length of patient ICU stay influenced nursing workload. Future  
288 research should focus on gathering more clinical patient data in order to extend our study findings  
289 regarding the relationship between nursing workload, patient characteristics and nurse  
290 characteristics.

291 **Impact**

292 **WHAT IS KNOWN ABOUT THIS TOPIC**

- 293 • Some of the previous studies about nursing workload in an ICU and the associations with  
294 nurse and patient characteristics present conflicting outcomes.

295 **WHAT THIS PAPER ADDS**

- 296 • Type of work shifts, nurse gender, type of ICU which nurses work in and the number of  
297 patients under care were nurse characteristics that influenced workload.

298 Patient gender, patient ICU requirements, patient referral source, the length of patient ICU stay  
299 were patient characteristics that influenced nursing workload.

300

301

302

303 **Table 1.** Comparison of nursing workload according to work shift, patient gender, patient treatment  
 304 types, and patient referral sources

Variables		Workload Mean (SD)	Test	p-value
Work shift (workload per shift)	morning	81.46 (25.94)	F-value = 6.94	<b>0.001*</b>
	evening	70.61 (26.01)		
	night	68.09 (23.17)		
Patient gender (workload per patient)	male	47.78 (12.23)	t-value = 5.77 Df = 507	< <b>0.001**</b>
	female	41.55 (11.85)		
Patient treatment types (workload per patient)	surgical	42.89 (11.04)	t-value = 5.30 Df = 507	< <b>0.001**</b>
	medical	48.80 (13.79)		
Patient referral sources (workload per patient)	Operation theatre	42.93 (11.16)	F-value = 6.42	< <b>0.001*</b>
	Emergency ward	47.73 (10.36)		
	General wards	45.57 (9.97)		
	Other ICUs	50.92 (41.04)		

\* One-Way ANOVA

\*\* Independent samples t-test

Df : degree of freedom.

305

306 Table 2. Association between nursing mean workload with individual and occupational characteristics of  
 307 ICU nurses and their patients' characteristics

<b>Nurses characteristics</b>	<b>b*</b>	<b>SE**</b>	<b>95% CI***</b>	<b>p-value</b>
Female	6.27	3.96	1.41, 11.13	<b>0.01</b>
Age (ref # to < 30)				
30-40	2.59	2.93	-3.16, 8.35	0.37
> 40	2.78	7.44	-11.81, 17.37	0.70
Married	-2.98	1.86	-6.62, 0.66	0.10
Educational level (ref to Bachelor)				
Master	0.20	3.73	-7.10, 7.51	0.95
Work experience	0.42	0.34	-0.24, 1.09	0.21
Weekly working hours	0.31	0.11	-0.12, 0.74	0.08
Being employed at the other health care centers	3.02	2.46	-1.80, 7.84	0.21
Resting before the work shift	0.18	0.07	-0.07, 0.43	0.07
Resting after the work shift	0.01	0.07	-0.13, 0.16	0.83
Number of patients under care	25.04	5.05	15.13, 34.96	<b>&lt; 0.001</b>
Type of ICU (ref # to General ICUs)				
Neurosurgical	-7.91	2.55	-12.91, -2.91	<b>0.002</b>
Cardiac surgery	-14.58	3.20	-20.87, -8.29	<b>&lt; 0.001</b>
Obstetric surgery	-31.91	3.69	-39.15, -24.67	<b>&lt; 0.001</b>
Burn	-31.18	5.12	-41.22, -21.14	<b>&lt; 0.001</b>
<b>Patients characteristics</b>				
Age of patients	-0.02	0.12	-0.26, 0.21	0.82
Length of ICU stay	0.75	0.23	0.53, 0.97	<b>0.02</b>

\* b coefficient was obtained according to the multivariable linear regression  
 \*\* Standard error  
 \*\*\* Confidence interval  
 # Reference group

309 **References**

- 310 Abbey M, Chaboyer W & Mitchell MJACC (2012). Understanding the work of intensive care  
311 nurses: a time and motion study. *Aust Crit Care*; **25**: 13-22.
- 312 Alizadeh M, Heidari Gorji MA, Khalilian AR & Esmaeili R (2015). Assessment of nursing  
313 workload and related factors in intensive care units using the nursing activities score.  
314 *JMUMS*; **24**: 147-157.
- 315 Altafin JAM, Grion CMC, Tanita MT, Festti J, Cardoso LTQ, Veiga CFF, Kamiji D, Barbosa  
316 ÁRG, Matsubara CCT & Lara ABJRBDti (2014). Nursing Activities Score and workload  
317 in the intensive care unit of a university hospital. *Rev Bras Ter Intensiva*; **26**: 292.
- 318 Armstrong E, de Waard MC, de Grooth H-JS, Heymans MW, Miranda DR, Girbes AR, Spijkstra  
319 JJJA & Analgesia (2015). Using nursing activities score to assess nursing workload on a  
320 medium care unit. *Anesth Analg*; **121**: 1274-1280.
- 321 Bruyneel A, Tack J, Droguet M, Maes J, Wittebole X, Miranda DR & Di Pierdomenico L (2019).  
322 Measuring the nursing workload in intensive care with the Nursing Activities Score (NAS):  
323 A prospective study in 16 hospitals in Belgium. *J Crit Care*; **54**: 205-211.
- 324 Coelho FUDa, Queijo AF, Andolhe R, Gonçalves LA & Padilha KG (2011). Carga de trabalho de  
325 enfermagem em unidade de terapia intensiva de cardiologia e fatores clínicos associados.  
326 *Texto e Contexto Enferm*; **20**: 735-741.
- 327 Coventry TH, Maslin-Prothero SE & Smith G (2015). Organizational impact of nurse supply and  
328 workload on nurses continuing professional development opportunities: an integrative  
329 review. *J Adv Nurs*; **71**: 2715-2727.
- 330 Debergh DP, Myny D, Van Herzele I, Van Maele G, Miranda DR & Colardyn FJlcm (2012).  
331 Measuring the nursing workload per shift in the ICU. *Intensive Care Med*; **38**: 1438-1444.
- 332 Ducci AJ, Zanei SSV & Whitaker IY (2008). Nursing workload to verify nurse/patient ratio in a  
333 cardiology ICU. *Rev esc enferm USP*; **42**: 673-680.
- 334 Esmaeili R, Moosazadeh M, Alizadeh M & Afshari M (2015). A systematic review of the workload  
335 of nurses in intensive care units using NAS. *Acta Med Mediterr*; **31**: 1455-1460.
- 336 Ferreira P, Machado R, Martins Q & Sampaio SJRGE (2017). Classification of patients and  
337 nursing workload in intensive care: comparison between instruments. *Revista gaucha  
338 deenfermagem*; **38**: e62782.
- 339 Fowler RA, Filate W, Hartleib M, Frost DW, Lazongas C & Hladunewich M (2009). Sex and  
340 critical illness. *Curr Opin Crit Care*; **15**: 442-449.
- 341 Halwani M, Solaymani-Dodaran M, Grundmann H, Coupland C & Slack RJJoHI (2006). Cross-  
342 transmission of nosocomial pathogens in an adult intensive care unit: incidence and risk  
343 factors. *J Hosp Infect*; **63**: 39-46.
- 344 Heckemann B, Zeller A, Hahn S, Dassen T, Schols J & Halfens RJNet (2015). The effect of  
345 aggression management training programmes for nursing staff and students working in an

- 346 acute hospital setting. A narrative review of current literature. *Nurse Educ Today*; **35**: 212-  
347 219.
- 348 Holland P, Tham TL, Sheehan C & Cooper B (2019). The impact of perceived workload on nurse  
349 satisfaction with work-life balance and intention to leave the occupation. *Appl Nurs Res*;  
350 **49**: 70-76.
- 351 Hoogendoorn ME, Margadant CC, Brinkman S, Haringman JJ, Spijkstra JJ & de Keizer NFJJons  
352 (2019). Workload scoring systems in the Intensive Care and their ability to quantify the  
353 need for nursing time: a systematic literature review. *Int J Nurs Stud*: 103408.
- 354 Hoonakker P, Carayon P, Gurses AP, Brown R, Khunlertkit A, McGuire K & Walker JM (2011).  
355 Measuring workload of ICU nurses with a questionnaire survey: the NASA Task Load  
356 Index (TLX). *IIE Trans Healthc Syst Eng*; **1**: 131-143.
- 357 Hugonnet S, Harbarth S, Sax H, Duncan RA & Pittet DJCOiID (2004). Nursing resources: a major  
358 determinant of nosocomial infection? *Curr Opin Infect Dis*; **17**: 329-333.
- 359 Lucchini A, De Felippis C, Elli S, Schifano L, Rolla F, Pegoraro F & Fumagalli R (2014). Nursing  
360 Activities Score (NAS): 5 years of experience in the intensive care units of an Italian  
361 University hospital. *Intensive Crit Care Nurs*; **30**: 152-158.
- 362 Lucchini A, Elli S, Bambi S, Becattini G, Vanini S, Piantanida C, Iacca C & Sartori D (2015).  
363 Nursing activities score: differences in nursing workload in three intensive care units.  
364 *Assist Inferm Ric*; **34**: 6-14.
- 365 Maenhout B & Vanhoucke MJO (2013). An integrated nurse staffing and scheduling analysis for  
366 longer-term nursing staff allocation problems. *Omega*; **41**: 485-499.
- 367 Mahmood K, Eldeirawi K & Wahidi MM (2012). Association of gender with outcomes in critically  
368 ill patients. *Crit Care*; **16**: R92.
- 369 Miranda DR, Nap R, de Rijk A, Schaufeli W & Iapichino GJCcm (2003). Nursing activities score.  
370 *Crit Care Med*; **31**: 374-382.
- 371 Mirzaei S, Zamanian Z & Hasan Zade J (2015). Effects of work shifts and mental workload on  
372 chronic fatigue among female nurses in intensive care units. *Journal of Health Sciences &*  
373 *Surveillance System*; **3**: 113-118.
- 374 Mohammadi M, Mazloumi A, Kazemi Z & Zeraati HJHpp (2015). Evaluation of mental workload  
375 among ICU ward's nurses. *Health Promot Perspect*; **5**: 280.
- 376 Momennasab M, Karimi F, Dehghanrad F & Zarshenas L (2017). Evaluation of nursing workload  
377 and efficiency of staff allocation in a trauma Intensive Care Unit. *Trauma Mon*; **23**:  
378 e58161-e58161.
- 379 Myny D, Van Hecke A, De Bacquer D, Verhaeghe S, Gobert M, Defloor T & Van Goubergen  
380 DJIjns (2012). Determining a set of measurable and relevant factors affecting nursing  
381 workload in the acute care hospital setting: a cross-sectional study. *Int J Nurs Stud*; **49**:  
382 427-436.

- 383 Nogueira dSL, de Alencar Domingues C, Poggetti RS & de Sousa RMC (2014). Nursing workload  
384 in intensive care unit trauma patients: analysis of associated factors. *PLoS one*; **9**: e112125.
- 385 Padilha KG, de Sousa RMC, Queijo AF, Mendes AM & Miranda DR (2008). Nursing Activities  
386 Score in the intensive care unit: analysis of the related factors. *Intensive Crit Care Nurs*;  
387 **24**: 197-204.
- 388 Padilha KG, Sousa RMC, Kimura M, Miyadahira AMK, da Cruz DALM, de Fátima Vattimo M,  
389 Fusco SRG, de Campos MEF, Mendes EMT & Mayor ERC (2007). Nursing workload in  
390 intensive care units: a study using the Therapeutic Intervention Scoring System-28 (TISS-  
391 28). *Intensive Crit Care Nurs*; **23**: 162-169.
- 392 Padilha KG, Stafseth S, Solms D, Hoogendoorn M, Monge FJC, Gomaa OH, Giakoumidakis K,  
393 Giannakopoulou M, Gallani MC & Cudak E (2015). Nursing Activities Score: an updated  
394 guideline for its application in the Intensive Care Unit. *Rev esc enferm USP*; **49**: 131-137.
- 395 Papazian L, Sylvestre A & Herridge M (2018). Should all ICU clinicians regularly be tested for  
396 burnout? Yes. *Intensive Care Med*; **44**: 681-683.
- 397 Penz K, D'Arcy C, Stewart N, Kosteniuk J, Morgan D & Smith B (2007). Barriers to participation  
398 in continuing education activities among rural and remote nurses. *The J Contin Educ Nurs*;  
399 **38**: 58-66.
- 400 Reich R, Vieira DFVB, Lima LBd & Rabelo-Silva ERJRgde (2015). Nursing workload in a  
401 coronary unit according to the Nursing Activities Score. *Rev Gaúcha Enferm*; **36**: 28-35.
- 402 Romano JL, Garcia PC, Silva DV, Moura BRS & de Souza Nogueira L (2019). Type of admission  
403 and nursing workload of critical patients: a cross-sectional study. *Nurs Crit Care*; **24**: 387-  
404 391.
- 405 Samuelsson C, Sjöberg F, Karlström G, Nolin T & Walther SM (2015). Gender differences in  
406 outcome and use of resources do exist in Swedish intensive care, but to no advantage for  
407 women of premenopausal age. *Crit Care*; **19**: 129.
- 408 Stafseth SK, Solms D & Bredal IS (2011). The characterisation of workloads and nursing staff  
409 allocation in intensive care units: a descriptive study using the Nursing Activities Score for  
410 the first time in Norway. *Crit Care Nurs*; **27**: 290-294.
- 411 Strazzieri-Pulido KC, S. González CV, Nogueira PC, Padilha KG & G. Santos VL (2019). Pressure  
412 injuries in critical patients: Incidence, patient-associated factors, and nursing workload. *J  
413 Nurs Manag*; **27**: 301-310.
- 414 Swiger PA, Vance DE & Patrician PAJNo (2016). Nursing workload in the acute-care setting: A  
415 concept analysis of nursing workload. *Nurs outlook*; **64**: 244-254.
- 416 Yusefi AR, Ebrahim Z, Zargar Balaye Jame S, Bastani PJJorDiN & Midwifery (2019). Workload  
417 and its Associated Factors among Nurses in Teaching Hospitals of Shiraz University of  
418 Medical Sciences in 2017. *J Res Dev Nurs Midw*; **16**: 82-92.

419