



# City Research Online

## City St George's, University of London

**Citation:** Douglas, R. H. (2017). A review of the nature & role of pre-receptor, wavelength-selective, ocular filters in vertebrates. *Acta Ophthalmologica*, 95(S259), doi: 10.1111/j.1755-3768.2017.02161

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/25425/>

**Link to published version:** <https://doi.org/10.1111/j.1755-3768.2017.02161>

**Copyright and Reuse:** Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

**Douglas, R., 2017. A review of the nature & role of pre-receptor, wavelength-selective, ocular filters in vertebrates. *Acta Ophthalmologica*, 95.**

Abstracts from the 2017 European Association for Vision and Eye Research Conference

First published: 07 September 2017

<https://doi.org/10.1111/j.1755-3768.2017.02161>

## **Summary**

Such filters are widespread in all vertebrates groups, most notably fish. Several species have pigmented corneas, whose density may alter with illumination. Filters within lenses are common, occurring in animals that inhabit well-lit environments as well as the deep-sea. Pigmented humours are rare but have been reported in a few species. Various layers within the retina also contain pigments, including the primate macular pigment, that absorb short wavelength light and the inner segments of many animals, such as birds, contain coloured oil droplets.

While many of these filters are various carotenoids, other substances such as mycosporine-like amino acids, are common. The variety of filtering pigments indicates pigmentation has evolved independently on a number of occasions; a clear testament to its importance.

Most pigments serve to reduce short wavelength light and have a variety of functions including; protection of the eye by removing the most damaging photons, acting as antioxidants, and improving image quality by reducing scatter. In deep-sea fish they enhance the visibility of bioluminescence and in animals with oil droplets they refine the spectral properties of photoreceptors.