



# City Research Online

## City St George's, University of London

**Citation:** Ctori, I. & Huntjens, B. (2017). Does ethnicity and foveal morphology play a role in the spatial distribution of macular pigment?. *Acta Ophthalmologica Scandinavica*, 95(S259), doi: 10.1111/j.1755-3768.2017.02163

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/25428/>

**Link to published version:** <https://doi.org/10.1111/j.1755-3768.2017.02163>

**Copyright and Reuse:** Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

# European Association for Vision and Eye Research Conference (EVER) abstract 2017

Title:

**Does ethnicity and foveal morphology play a role in the spatial distribution of macular pigment?**

Authors:

Irene Ctori <sup>1</sup>

Byki Huntjens <sup>1</sup>

<sup>1</sup> Applied Vision Research Centre, School of Health Sciences, City, University of London, UK

## **Abstract**

While macular pigment optical density typically follows an exponential decline with eccentricity from the fovea, *atypical* profiles have also been described, including reports of increased prevalence of ring-like and central dip profiles among non-white ethnicities. However, classification of macular pigment (MP) profile phenotypes varies and is often based on subjective visualisation. We present data showing that compared to visual assessment, objective MP profiling (taking into account measurement repeatability) is a more reliable method that should be considered in future observational and interventional studies. We will also describe ethnic variations in MP spatial distribution (among white, black and South Asian individuals) in relation to variations in foveal architecture, including data regarding individual retinal layers, foveal width and pit slope taken from SD-OCT images. Our data suggests that although ethnicity explains around 10% of the variance in MP profile, gender plays no significant role. In addition, although white subjects demonstrate thicker retinal layers than non-whites, the significant differences in the amount and distribution of MP between ethnicities is not explained by variations in foveal morphology.