



City Research Online

City, University of London Institutional Repository

Citation: Hofstad, T., Hampton, J. A. & Hofmann, B. (2020). What Makes Some Diseases More Typical than Others? A Survey on the Impact of Disease Characteristics and Professional Background on Disease Typicality. *INQUIRY: The Journal of Health Care Organization, Provision, and Financing*, 57, pp. 1-9. doi: 10.1177/0046958020972813

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/25950/>

Link to published version: <https://doi.org/10.1177/0046958020972813>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

What Makes Some Diseases More Typical than Others? A Survey on the Impact of Disease Characteristics and Professional Background on Disease Typicality

INQUIRY: The Journal of Health Care
Organization, Provision, and Financing
Volume 57: 1–9
© The Author(s) 2020
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/0046958020972813
journals.sagepub.com/home/inq



Tore Hofstad¹ , James A. Hampton² , and Bjørn Hofmann^{1,3}

Abstract

Health professionals tend to perceive some diseases as more typical than others. If disease typicalities have implications for health professionals or health policy makers' handling of different diseases, then it is of great social, epistemic, and ethical interest. Accordingly, it is important to find out what makes health professionals rank diseases as more or less typical. This study investigates the impact of various factors on how typical various diseases are perceived to be by health professionals. In particular, we study the influence of broad disease categories, such as somatic versus psychological/behavioral conditions, and a wide range of more specific disease characteristics, as well as the health professional's own background. We find that professional background strongly impacted disease typicality. All professionals (MD, RN, physiotherapists and psychologists) considered somatic conditions to be more typical than psychological/behavioral. As expected, psychologists also found psychological/behavioral conditions to be more typical than did other groups. Professions of respondents could be well predicted from their individual typicality judgments, with the exception of physiotherapists and nurses who had very similar judgment profiles. We also demonstrate how various disease characteristics impact typicality for the different professionals. Typicality showed moderate to strong positive correlations with condition severity and mortality, and only non-severe conditions were rated as atypical. Hence, studying how different disease characteristics and occupational background influences health professionals' perception of disease typicality is the first and important step toward a more general study of how typicality influences disease handling.

Keywords

disease, typical, severity, profession, health policy, ethics, disease characteristics, judgment, respondents, survey

What do we already know about this topic?

Health professionals ascribe different prestige to various diseases and they consider certain diseases to be more typical than others, but little or nothing is known about why they do so.

How does this research contribute to the field?

This is the first study that investigates what makes health professionals consider some diseases to be more typical than others.

What are the implications for practice and policy?

How typical professionals find diseases can influence how they treat these diseases and persons having them. Knowing what drives this differentiation is a prerequisite for handling the problem.

Highlights

- Health professionals tend to perceive some diseases as more typical than others.
- This is of great social, epistemic, and ethical interest in cases where disease typicalities have implications for health professionals or health policy makers' handling of different diseases.
- We find that professional background impacted disease typicality.
- All professionals (MD, RN, physiotherapists and psychologists) considered somatic conditions to be more typical than psychological/behavioral ones.
- Severity and mortality also influence whether professionals consider conditions to be typical diseases.



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

- Studying how different disease characteristics and occupational background influences health professionals' perception of disease typicality is the first and important step toward a more general study of how typicality influences disease handling.

Introduction

The concept of disease is crucial to healthcare—both theoretically and practically. It defines the goal of medicine: detecting, understanding, treating, curing, and/or palliating disease. Moreover, it has a normative function in demarcating who is entitled to attention, care and treatment.¹ Disease also directs the action of health professionals. Moreover, it decides a person's social role and status,^{2–5} for example, who gets sick leave and sick pay,^{6,7} and it can determine a person's moral and legal accountability.^{1,8–10} Hence, disease is a core concept in health care, in health policy making, and in society at large.^{11–13}

Despite the crucial role of the concept of disease, it is notoriously difficult to define.^{10,11,13–19} From a theoretical point of view it can be argued that the concept of disease is defined by its constituents (according to *classical concept theory*), by its relation to the world (*conceptual atomism*), by its description (*descriptivist theories*), by theories (*theory theory of concepts*), by perceptual and motor states (*neo-empiricist theories*), by natural kinds (*natural kind theory*),²⁰ by actual exemplars in memory (*exemplar theory*),^{21,22} by abstract average prototypes (*prototype theory*),^{23–27} or as fuzzy sets (*fuzzy theory of concepts*).²⁸ Despite long and thorough debates, no agreements are obtained.

From an empirical perspective, diseases have been found to make up a prestige hierarchy.^{29,30} Organ specific, acute, and actionable conditions where you can use advanced technologies have high prestige among health professionals. Correspondingly diseases have been shown to be more or less typical.^{31–33} According to prototype theory our concepts are structured on the basis of their similarity to some central prototype representation.^{34–36}

If such prestige hierarchies or disease typicalities have implications for health professionals or health policy makers' handling of different diseases, then it is of great social, epistemic, and ethical interest.^{37–39} From other areas we know that typicality affects what people consider to be trustworthy and good.⁴⁰ Hence, if some diseases are considered

as better or deserve more attention due to their typicality, this has implications for patients, professionals, and policy makers.

In a previous study³³ among Norwegian health professionals, it has been shown that some conditions are considered to be very typical diseases. Examples are lung cancer, leukemia, colon cancer, myocardial infarction, AIDS, diabetes, multiple sclerosis, and schizophrenia. Others score very low on how typical they are perceived to be, for example caffeine-dependency, kleptomania, and infertility.³³

The objective of this study was to investigate what makes professionals perceive some diseases as more typical than others. In particular we wanted to assess the importance of specific characteristics of the health professionals (profession) and some features of the conditions (somatic vs psychological/ behavioral; mortality; prevalence; severity; media coverage; curability; etiology).

Our specific research questions were:

1. What is the impact of broad disease category on perceived typicality among different professional groups?
2. Can different respondent professions be discriminated based on their perceived typicality of diseases?
3. What are the individual and combined effects of disease characteristics (such as mortality; prevalence; severity; media coverage; curability; etiology) on perceived typicality?

Methods

To answer our research questions, we used the dataset from an earlier study investigating how typical health professionals considered various conditions to be,³³ and we compiled a data set with information about each condition. Conditions were included partly based on a study of health professionals' conceptions of prestige related to diseases³⁰ where diseases were selected to represent a broad range of conditions handled by a variety of specialities. The respondents were asked to rate whether they found the diseases more or less typical.

Typicality was explained in terms of how people would find different birds, such as sparrows versus ostriches, more or less typical of birds. Answers were given from 0 to 10,

¹The University of Oslo, Oslo, Norway

²City, University of London, England

³The Norwegian University of Science and Technology (NTNU), Gjøvik, Norway

Received 17 July 2020; revised 6 October 2020; revised manuscript accepted 13 October 2020

Corresponding Author:

Bjørn Hofmann, Department in Health Sciences, Faculty of Medicine and Health Sciences, the Norwegian University of Science and Technology (NTNU), PO Box 191, Gjøvik, N-2801, Norway.

Email: bjoern.hofmann@ntnu.no

where 0 was atypical, and 10 was very typical. The final survey, which was in Norwegian, was printed on white paper (1 page, 2-sided). More background details are available in Hofmann.³³ The diseases used may also be seen displayed in Figure 1.

One of the authors (BH) designed the survey and distributed it to 254 health related professionals while they were attending university courses or seminars at the Faculty of Medicine, University of Oslo, during the spring terms of 2010 to 2014. The respondents were categorized as MD (42), RN (67), physiotherapists (22), psychologists (13), and others (52). The latter category encompassed dentists, biologists, epidemiologists, social scientists, philosophers, and professionals with backgrounds from other humanities. Forty-seven respondents had missing information on professional background, and their data were only included in the average scores. The respondents either had education at Masters (135) or Bachelors (119) level.

The data were registered manually in Microsoft Excel 2010. Statistical analyses were performed with R version 3.6.3,⁴¹ and IBM-SPSS, and plots were created with ggplot2 3.2.1⁴² and sjPlot.⁴³ Minor misprints in the data were corrected and 11 respondents with more than 20% missing responses were removed, in addition to one disease (neuralgia) with 26% missing data.

To investigate the importance of type of disease, we categorized the diseases based on the presence or absence of *predominantly* psychological/behavioral components. While all internal, psychological states and conditions are likely to show some form of external correlate in the body, such as altered brain waves or heightened blood pressure, the following taxonomies were based on what is emphasized in the current understanding of the etiology. Similarly, the diseases we categorized as somatic may well involve a change in individual behavior and psychological well-being, but neither of them is assumed to originate in the psyche. This left us with a handful of diseases/conditions that are less easily confined under these two labels, and they made out the ambiguous class.

“Somatic”: myocardial infarction, chlamydia pneumoniae, ruptured spleen, prolapse, candida, infertility, pollen allergy, kidney stone, cataract, colon cancer, diabetes, lumbago, angina pectoris, asthma, MS, psoriasis, lung cancer, cirrhosis, pregnancy, leukemia, rheumatoid arthritis, ankylosing spondylitis, AIDS, kidney failure, appendicitis, miscarriage, stomach ulcer, ectopic pregnancy, hip fracture, menopause, dementia.

“Psychological/Behavioral”: nostalgia, homosexuality, kleptomania, dissidence, media victim syndrome, anxiety neurosis, borderline personality disorder, electromagnetic hypersensitivity, gambling addiction, porn addiction, opioid addiction, shopping addiction, cola addiction, caffeine addiction, alcoholism, self-harm, body dysmorphic disorder, anorexia, Gulf war syndrome, agoraphobia, drapetomania (runaway disease for slaves⁴⁴).

“Ambiguous status”: schizophrenia, PMS, AD/HD, Asperger’s syndrome, erectile dysfunction, obesity, ME/CFS, IBS, fibromyalgia.

Research Question 1: In order to assess the impact of disease category on perceived typicality among different professional groups, we calculated the average and standard deviations of the typicality scores within each category, separately for each profession. Reliability of typicality ratings across categories within each group was high (Cronbach $\alpha > .86$).

Research Question 2: To determine whether different professional groups perceive typicality in characteristically different ways, we employed linear discriminant analysis, using the typicality ratings of disease categories for each individual in each of the four defined groups, MDs, RNs, physiotherapists and psychologists to construct linear functions to maximally separate the groups from each other.

Research Question 3: In order to test why perceived typicality varies among the different diseases, we conducted a correlational analysis. A data set was compiled containing a variety of theoretically relevant factors pertaining to each disease. We classified the diseases on the following variables: severity, based on disability weights from the Global Health Estimate⁴⁵; global prevalence rates in 2010⁴⁶; global and Norwegian mortality in 2010 (ibid.); fame, operationalized as number of mentions in newspaper articles online and in print in Norway in 2010 to 2015, and whether the conditions were incurable or infectious. (Detailed references to the disease characteristics can be found in Appendix A.) The presence of an infectious agent was coded as a four-level scale: no infectious agent/infections may cause the disease/infections the most common cause but not necessary/infection is a necessary cause. The categories (Somatic, Psychological/Behavioral, Ambiguous) were dummy coded when calculating correlations, where 1 indicates membership and 0 non-membership in the category.

Given non-normal distributions, we report Spearman’s rank order correlation coefficients. Correlations between perceived typicality for each respondent and disease characteristics were transformed using Fisher’s Z transformation for the purpose of averaging across respondents. Table 1 displays the inverted Z-transformed average correlations.

In order to assess the combined impact of disease characteristics on typicality we conducted mixed effects multilevel regression analyses⁴⁷ using the *lme4* package.⁴⁸ We estimated several models with varying intercepts for each respondent and disease. This was done in order to account for dependency in the data, since each typicality score can be considered to be nested within diseases and respondents. The regression coefficients for the disease characteristics are assumed to be the same across respondents and diseases. Models were fitted using

restricted maximum likelihood, and model diagnostics were checked using the *performance* package.⁴⁹ Some variables were strongly right skewed (mortality, prevalence, fame), and these were log transformed prior to running the regressions.

Finally, in order to inspect non-linear relationships among the strongest predictors, we plotted mean typicality scores (separately for each profession) versus severity, by disease category and mortality.

Results

The role of occupational background and disease category

Table 2 shows the impact of disease category on perceived typicality among different professional groups (research question 1). All professionals considered somatic conditions to be more typical as diseases than psychological/behavioral

conditions, with the ambiguous category in between. There was also evidence of an interaction effect, such that different professions rated different categories of disease as most typical. On average MDs considered somatic conditions to be more typical diseases than any other professions. Psychologists on the other hand considered both the psychological/behavioral conditions, as well as those falling in the ambiguous category, to be more typical than any other profession. Figure 2 shows a graphical illustration of perceived typicality according to category and profession.

To test whether different professions consider different broad categories of disease to be more or less typical, a two-way ANOVA was run with factors of Profession (4 levels, between-subjects) and Disease Category (3 levels, somatic, psychological/behavioral, and ambiguous, within-subjects). The main effect of Profession was not significant ($F(3,140) = 2.05, P = .11, \eta_p^2 = .042$). The effect of Disease Category was strong ($F(2,280) = 464.4, P < .001, \eta_p^2 = .768$), with somatic





Figure 1. Scatterplot of mean typicality of diseases by severity, grouped by mortality (in Norway), and category (somatic, psychological/behavioral, ambiguous).

Table 1. Bivariate Correlations. Spearman's Rho: Typicality vs Disease Characteristics by Profession.

Profession	Medical doctor	Psychologist	Registered nurse	Physiotherapist
Severity	0.60	0.57	0.58	0.63
Category: Somatic	0.58	0.34	0.51	0.50
Category: Behavioral	-0.52	-0.34	-0.50	-0.47
Mortality: Norway	0.47	0.47	0.49	0.51
Mortality: Global	0.46	0.38	0.45	0.46
Fame	0.39	0.34	0.41	0.42
Infectious	0.30	0.21	0.29	0.28
Incurable	0.17	0.16	0.20	0.20
Category: Ambiguous	-0.12	0.02	-0.05	-0.06
Prevalence	-0.02	0.00	-0.02	-0.02

Table 2. Mean Typicality of Diseases and Standard Deviation by Profession and Broad Disease Category.

Profession	Category of disease		
	Somatic	Ambiguous	Psychological/behavioral
Medical Doctor	8.11 (0.80)	5.51 (1.42)	3.64 (1.18)
Registered Nurse	7.81 (1.00)	5.86 (1.45)	3.49 (1.26)
Physiotherapist	7.59 (0.98)	5.62 (1.64)	3.48 (1.27)
Psychologist	7.53 (0.99)	6.69 (1.72)	5.08 (1.28)

**Figure 2.** Mean typicality of diseases by profession and category. Note. Error bars 95% CI.

diseases ($M = 7.84$, $SD = .95$) more typical than ambiguous ($M = 5.79$, $SD = 1.51$) which in turn were more typical than psychological/behavioral conditions ($M = 3.66$, $SD = 1.30$).

Interestingly, the interaction between the factors was significant ($F(6,280) = 5.97$, $P < .001$, $\eta_p^2 = .113$), as seen in Figure 2. The source of the interaction was the group of psychologists, who judged the psychological/behavioral and ambiguous disease categories to be clearly more typical than did the other groups. In confirmation, a rerun of the ANOVA without the psychologist group led to the interaction ceasing to be significant ($F(4, 258) = 2.19$, $P = .071$, $\eta_p^2 = .033$).

Differences between professions across different individual diseases

Research question 2 asked whether, given the difference between professions in how typical broad categories of disease were seen (see above), is it possible to look in more

detail at where the professional groups differed most strongly?

A MANOVA analysis with the four professions (MD, RN, Psychologist, and Physiotherapist) as the Independent factor and the rated typicalities of the 61 diseases within each profession as dependent variables confirmed a significant effect of Profession, with partial η^2 (similar to R^2) of .58 using Wilks Lambda. Over half the diseases (about 38) showed significant univariate effects of profession ($P < .05$). It thus appears that respondents generally agree on which diseases are typical vs non-typical, but disagree on how typical the psychological/behavioral diseases should be understood. Profession seems to explain parts of this variation.

Linear discriminant analysis (LDA) takes a set of measures (in this case the rated typicality of diseases for each respondent) and aims to construct a set of dimensions (as linear functions of the measures) that will enable one to predict which profession a person belongs to, based on their judgments. In the case of our professional groups, three

independent functions were created, placing each individual in a 3D space, with clusters in the space corresponding to each of the four professions. The functions were optimal in maximizing the distance between members of different groups and minimizing distance within groups. The aim was to see how well the analysis would be able to separate the professions. This was done by leaving each respondent out of the analysis in turn and predicting their professional group based on functions calculated from the remaining data (a cross-validation, or jack-knife procedure). A fully successful analysis would predict the correct professional group for each individual, just based on their judgments of typicality.

To avoid over-fitting of the model, a stepwise procedure was employed, adding in diseases to the analysis one at a time until no significant improvement in fit was seen. Two functions were significant and captured 97% of the variance. Figure 3 shows how the individual respondents were clustered on these functions. Function 1 distinguished between the psychologists and doctors to the right and the other professional groups to the left. Function 2 then differentiated psychologists from the rest. The cross-validated classifications showed that 62% of doctors, 84% of nurses and 58% of psychologists were correctly predicted to fall in their respective groups. In contrast, none of the physiotherapists were

correctly classified as such, with 86% predicted to be nurses, and 14% to be doctors.

Disease characteristics associated with typicality

Research question 3 concerned how disease characteristics correlate with perceived typicality. Table 1 shows how various characteristics of the diseases correlated with perceived typicality for the different professions. In general, the correlations in Table 1 differed little between professions, with correlations among psychologists somewhat lower on most of the characteristics.

We observed a moderate to strong positive correlation between the typicality scores and severity, category of disease, and mortality (national and global). There was also a low to moderate positive correlation between typicality and fame and infectiousness. There was no correlation between typicality and prevalence, and only a small positive correlation with respect to whether the condition is incurable. Finally, typicality scores showed a strong negative correlation with whether the condition is categorized as psychological/behavioral.

Figure 4 shows plotted coefficients from the full hierarchical model (see Appendix B for coefficients). The plot shows the standardized regression coefficients for comparison between predictors and confirms the strong effect of severity and mortality on typicality.

Table 1 and Figure 4 show that severity was the strongest predictor of disease typicality, with correlations of around 0.6 within each profession. To understand this effect, Figure 1 shows a scatter plot of average typicality vs severity, grouped by disease category and mortality. The colors differ by disease category and display the patterns clearly. Somatic diseases, in black are seen to be more typical than ambiguous diseases, in yellow, which were more typical than psychological/behavioral ones in blue. All categories of disease showed a similar relation between severity and typicality. Above a severity of 0.2 on the scale, typicality was not correlated with severity. Below this level of severity, there was a wide range of typicalities, and almost all diseases with lower typicality (<5) were at the




Figure 3. Plot showing individual professionals mapped on the first two functions of the linear discriminant analysis.




Figure 4. Plotted coefficients from hierarchical linear model.

least severe end of the scale. Virtually all of these atypical diseases are either psychological/behavioral, or are related to sex or reproducibility: pregnancy, menopause, PMS, miscarriage and infertility.

The effect of severity on typicality can be captured by noting the absence of any diseases in the lower right quadrant. Typical diseases could vary in severity across the whole scale, but there were no atypical severe diseases. If a disease was severe, it scored at least 4.5 on the typicality scale, and if a disease was atypical it scored less than 0.2 on the severity scale.

Discussion

In this study we considered three research questions. First, we considered how the four groups of MD, RN, physiotherapists and psychologists differed in their perceptions of disease typicality. Here we found that while there was a high level of consensus across groups, there were many diseases where differences appeared. In an overall analysis, we identified the key difference as relating to either fully or ambiguously psychological/behavioral conditions, which were rated as more typical by the psychologists in comparison to the other groups.

One explanation for the fact that psychologists consider psychological diseases to be far more typical than other professions, while medical doctors are more likely to perceive somatic conditions as more typical, is that members of these professions are more frequently exposed to individuals exhibiting these problems, both from clinical experience and from the literature. Regardless of explanation, the difference is striking and could influence the way patients with various diseases are met and treated, depending on the profession of the therapist. However, there were only 13 psychologists in the sample, which urges cautiousness with making strong claims of an effect that is generalizable beyond the sample. Moreover, the generally lower typicality of psychological/behavioral diseases for other professions may also be related to the low prestige of these diseases among professionals.^{29,30}

The second research question concerned whether the profile of a professional's typicality judgments was sufficient to identify their professional background. Using LDA we were able to show that a substantial majority of the MD, RN and psychologist groups could be differentiated on the basis of their typicality judgments. RNs were a particularly well-clustered group. However, physiotherapists were almost completely contained within the RN group cluster. It is interesting how MDs had a very distinct view of disease typicality compared to the RN and physiotherapist groups, which themselves did not differ.

The third and final research question asked what characteristics of diseases lead them to being considered typical. The professional groups hardly differed in which characteristics were most predictive. In addition to the already described

effect of somatic versus non-somatic diseases, the most important aspect of a disease driving typicality was its severity, followed by mortality and infectiousness. Curability turned out to have only a low correlation with typicality.

A closer look at how severity affected typicality revealed that the only diseases considered atypical tended to have low severity. Put another way, any disease that was above a certain threshold of severity was considered more or less typical. This pattern suggests that, logically, severity is a sufficient (but not necessary) condition for a disease to be considered typical.

Another characteristic—fame—was poorly correlated with typicality. This fact may in part be due to the inclusion of concepts such as pregnancy, homosexuality and nostalgia, which are not typically considered diseases, but often show up in the media. The search terms used for measuring this variable required the inclusion of additional terms like disease, sickness or illness. Still they are likely to include articles where the terms are not understood as diseases.

There are of course many ways to categorize diseases. For example, whether they are chronic is, no doubt, an important characteristic. While we had included chronicity as a characteristic, we abandoned it in the end. One reason was that it overlapped with incurability. Moreover, there exist many definitions of chronicity that are quite different in scope.

It is also noteworthy that conditions related to female reproduction, such as infertility, miscarriage, pregnancy, PMS, and menopause came out low in typicality although most of them are characterized as somatic. Hence, there may be a gender issue to address in further research.

Measures of disease characteristics such as mortality and prevalence were collected in July 2019 and represent global numbers for 2010. Clearly, updated estimates are likely to become available later. However, our ambition was to identify general trends in correlations, rather than specific magnitude, so minor deviations are not likely to have a significant impact.

It is also very important to acknowledge that this study was performed in a specific cultural setting. The results are by no means generalizable to an Asian or African setting. Here we can only encourage other researchers to supplement our study. In a similar vein, it would be interesting and informative to learn how the typicality of diseases is understood by people who are not health professionals.

Conclusion

The conclusions of the research were, first, that while different health professions agreed well on which diseases are typical, their judgments could also be clearly differentiated. Psychologists, naturally, considered psychological/behavioral conditions to be more typical, but in addition, MDs could be distinguished from RN and physiotherapists by their profile of disease typicality judgments. The second conclusion was that looking at a number of disease characteristics

that may influence typicality, there were two that stood out. The involvement of somatic disease clearly raised typicality over either partly or fully psychological/behavioral conditions, and there was a strong relation between severity and typicality, such that any severe disease would necessarily be considered typical, while any atypical disease would necessarily be non-severe. Hence, studying how different disease characteristics and occupational background influences health professionals' perception of disease typicality is the first and crucial step toward a more general study of how typicality influences disease handling.

Authors' Note

Data are available in appendixes and further information on request. No patients were involved in this study and no patient consent was applied. All data are anonymous, and no ethics approval was necessary for collecting anonymous data in Norway. Participants were informed about the study and consented by filling in the questionnaire.

Author Contributions

BH designed the questionnaire and collected the data. TH and BH designed this study and TH gathered data for analysis parameters. TH designed the data analysis and performed these in collaboration with JH. BH made the first draft of the manuscript. TH, JH and BH revised the manuscript in several rounds. All authors have approved the final manuscript.

Declaration of Conflicting Interests


The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: TH and BH received no financial support for the research, authorship, and/or publication of this article. JH has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme grant agreement Nos. 681422 and 742204.

ORCID iDs

Tore Hofstad  <https://orcid.org/0000-0003-0718-7931>

James A. Hampton  <https://orcid.org/0000-0002-0363-8232>

Bjørn Hofmann  <https://orcid.org/0000-0001-6709-4265>

Supplemental Material

Supplemental material for this article is available online.

References

- Nordenfelt L. On the relevance and importance of the notion of disease. *Theor Med.* 1993;14(1):15-26.
- Parsons T. Definitions of health and illness in the light of American values and social structure. In: Jaco EG, ed. *Patients, Physicians and Illness: Sourcebook in Behavioral Science and Medicine*. Free Press; 1958:165-187.
- Twaddle A. *Influence and Illness: Definitions and Definers of Illness Behavior Among Older Males in Providence, Rhode Island*. Brown University; 1968.
- Twaddle A. Disease, illness and sickness revisited. In: Twaddle A, Nordenfelt L, eds. *Disease, Illness and Sickness: Three Central Concepts in the Theory of Health*. Linköping University Press; 1994: 1-18.
- Jutel A. Classification, disease, and diagnosis. *Perspect Biol Med.* 2011;54(2):189-205.
- Hofmann B. On the dynamics of sickness in work absence. In: Nordby H, Rønning R, Tellnes G, eds. *Social Aspects of Illness, Disease and Sickness Absence*. Unipub Forlag; 2011: 287.
- Hofmann B. The concepts of sickness and illness in the transformation of sick leave. In: *24th International Conference of the European Society for Philosophy of Medicine and Healthcare: "Human Nature, Medicine and Healthcare"*, University of Zagreb, Croatia, Book of Abstracts. European Society for Philosophy of Medicine and Healthcare; 2010: 30.
- Sisti D, Caplan AL. The concept of disease. *Routledge Companion Philos Med.* 2017;2017:5-15.
- Merskey H. Variable meanings for the definition of disease. *J Med Philos.* 1986;11(3):215-232.
- Hofmann B. Complexity of the concept of disease as shown through rival theoretical frameworks. *Theor Med Bioethics.* 2001;22(3):211-236.
- Kovacs J. The concept of health and disease. *Med Health Care Philos.* 1998;1(1):31-39.
- Schramme T. The significance of the concept of disease for justice in health care. *Theor Med Bioethics.* 2007;28(2):121-135.
- Hofmann B. The concept of disease-vague, complex, or just indefinable? *Med Health Care Philos.* 2010;13(1):3-10.
- Boorse C. Concepts of health and disease. In: Gifford F, ed. *Philosophy of Medicine*. Elsevier; 2011.
- Copeland DD. Concepts of disease and diagnosis. *Perspect Biol Med.* 1977;20(4):528-538.
- Engelhardt H. The concepts of health and disease. In: Engelhardt HT, ed. *Evaluation and Explanation in the Biomedical Sciences*. Reidel; 1975:125-141.
- Fabrega H Jr. Concepts of disease: logical features and social implications. *Perspect Biol Med.* 1972;15(4):583-616.
- Khushf G. An agenda for future debate on concepts of health and disease. *Med Health Care Philos.* 2007;10(1):19-27; discussion 29-32.
- Taylor FK. A logical analysis of disease concepts. *Compr Psychiatry.* 1983;24(1):35-48.
- D'amico R. Is disease a natural kind? *J Med Philos.* 1995; 20(5):551-569.
- Norman G. Dual processing and diagnostic errors. *Adv Health Sci Educ.* 2009;14(1):37-49.
- Papa FJ, Li F. Evidence of the preferential use of disease prototypes over case exemplars among early year one medical students prior to and following diagnostic training. *Diagnosis.* 2015;2(4):217-225.
- Bishop GD, Briede C, Cavazos L, Grotzinger R, McMahon S. Processing illness information: the role of disease prototypes. *Basic Appl Soc Psychol.* 1987;8(1-2):21-43.
- Lalljee M, Lamb R, Carnibella G. Lay prototypes of illness: their content and use. *Psychol Health.* 1993;8(1):33-49.

25. Sadegh-Zadeh K. The prototype resemblance theory of disease. *J Med Philos.* 2008;33(2):106-139.
26. von Lengerke T. Distinctiveness of disease prototypes in lay illness diagnosis: an exploratory observational study. *Psychol Health Med.* 2005;10(1):108-121.
27. Wakefield JC. Evolutionary versus prototype analyses of the concept of disorder. *J Abnorm Psychol.* 1999;108(3):374-399.
28. Sadegh-Zadeh K. Fuzzy health, illness, and disease. *J Med Philos.* 2000;25(5):605-638.
29. Album D, Johannessen LE, Rasmussen EB. Stability and change in disease prestige: a comparative analysis of three surveys spanning a quarter of a century. *Social Sci Med (1982).* 2017;180:45-51.
30. Album D, Westin S. Do diseases have a prestige hierarchy? A survey among physicians and medical students. *Soc Sci Med.* 2008;66(1):182-188.
31. Rizzi D, Holm S, Bornemann C, Clemmensen T, Pedersen SA. Physicians' grading of disease cases: do disease entities have prototype characteristics? In: Brehmer B, ed. *Models of Human Activities in Work Contexts*. Vol. 2. Grafisk Service; 1992: 155-165.
32. Tikkinen KA, Leinonen JS, Guyatt GH, Ebrahim S, Jarvinen TL. What is a disease? Perspectives of the public, health professionals and legislators. *BMJ Open.* 2012;2(6):e001632.
33. Hofmann B. Do health professionals have a prototype concept of disease? The answer is no. *Philos Ethics Humanit Med.* 2017;12(1):6.
34. Hampton JA. Testing the prototype theory of concepts. *J Mem Lang.* 1995;34(5):686-708.
35. Hampton JA. Concepts and prototypes. *Mind Lang.* 2000; 15(2-3):299-307.
36. Hampton JA. Typicality, graded membership, and vagueness. *Cogn Sci.* 2007;31(3):355-384.
37. Norredam M, Album D. Prestige and its significance for medical specialties and diseases. *Scand J Public Health.* 2007; 35(6):655-661.
38. Haldar M, Engebretsen E, Album D. Legitimizing the illegitimate: How doctors manage their knowledge of the prestige of diseases. *Health.* 2016;20(6):559-577.
39. Hindhede AL, Larsen K. Prestige hierarchies and relations of dominance among healthcare professionals. *Prof Prof.* 2020; 10(1):e3447.
40. Sofer C, Dotsch R, Wigboldus DH, Todorov A. What is typical is good: the influence of face typicality on perceived trustworthiness. *Psychol Sci.* 2015;26(1):39-47.
41. *R: A language and environment for statistical computing* [computer program]. R Foundation for Statistical Computing; 2020.
42. *Ggplot2: Elegant graphics for data analysis* [computer program]. Springer-Verlag; 2016.
43. *sjPlot: Data visualization for statistics in social science* [computer program]. R package version 2.1; 2018.
44. Bynum B. Drapetomania. *Lancet.* 2000;356(9241):1615.
45. WHO. *WHO Methods and Data Sources for Global Burden of Disease Estimates 2000-2011*. Department of Health Statistics and Information Systems; 2013.
46. Global Burden of Disease Collaborative Network. *Global Burden of Disease Study 2017*. Institute for Health Metrics and Evaluation (IHME); 2018.
47. Gelman A, Hill J. *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Vol. 1. Cambridge University Press; 2007.
48. Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects models using lme4. *J Stat Softw.* 2014;67(1):1-48.
49. *Performance: assessment of regression models performance* [computer program]. R package version 0.4.4; 2020.