



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** Pope, J., Olander, E. K. ORCID: 0000-0001-7792-9895, Leitao, S., Meaney, S. and Matvienko-Sikar, K. (2021). Prenatal stress, health, and health behaviours during the COVID-19 pandemic: An international survey.. *Women and Birth*, doi: 10.1016/j.wombi.2021.03.007

This is the published version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/26256/>

**Link to published version:** <https://doi.org/10.1016/j.wombi.2021.03.007>

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

---

---

---

City Research Online:

<http://openaccess.city.ac.uk/>

[publications@city.ac.uk](mailto:publications@city.ac.uk)

---



ELSEVIER

Contents lists available at ScienceDirect

## Women and Birth

journal homepage: [www.elsevier.com/locate/wombi](http://www.elsevier.com/locate/wombi)



# Prenatal stress, health, and health behaviours during the COVID-19 pandemic: An international survey

Johanna Pope<sup>a,\*</sup>, Ellinor K. Olander<sup>b</sup>, Sara Leitao<sup>c</sup>, Sarah Meaney<sup>c</sup>,  
Karen Matvienko-Sikar<sup>a</sup>

<sup>a</sup> School of Public Health, University College Cork, Ireland

<sup>b</sup> Centre for Maternal and Child Health Research, School of Health Sciences, City, University of London, London, UK

<sup>c</sup> National Perinatal Epidemiology Centre, Department of Obstetrics and Gynaecology, University College Cork, Ireland

### ARTICLE INFO

#### Article history:

Received 3 January 2021

Received in revised form 6 March 2021

Accepted 10 March 2021

Available online xxx

#### Keywords:

Pregnancy

Stress

Mental health

Physical health

Health behaviours

COVID-19

### ABSTRACT

**Background:** Pregnant women's stress, mental and physical health, and health behaviours can have important implications for maternal and child health outcomes.

**Aim:** To examine pregnant women's levels of stress, mental and physical health, and health behaviours during the COVID-19 pandemic.

**Methods:** A cross-sectional survey was conducted online, with recruitment and data collection occurring between 16/6/20 and 17/7/20. Participants were pregnant women recruited via online pregnancy/parenting communities. Participants self-reported their levels of general stress, pregnancy-specific stress and COVID-19 related stress, mental and physical health, general health behaviours, and COVID-19 related health behaviours.

**Findings:** 573 pregnant women participated in the survey. Participants were most commonly resident in the United States (42.6%, n = 243), Ireland (41.2%, n = 235) or the United Kingdom (10%, n = 57). The majority (80.0%, n = 457) were married and educated to degree level or above (79.3, n = 453). Pregnant women reported high levels of pregnancy-specific and COVID-19-related stress, and low levels of mental and physical health, during the pandemic. Encouragingly, pregnant women in this study generally reported high levels of adherence to public health advice and pregnancy health behaviours. Stress and general mental health outcomes were best predicted by well-being factors (including stress and social support). Health impairing behaviours (e.g. poor diet) were predicted by both well-being and demographic factors.

**Discussion:** Interventions targeting pregnancy- and pandemic-specific stress at the population level will be essential to support mental health and minimise adverse outcomes for women and children during the pandemic.

© 2021 The Authors. Published by Elsevier Ltd on behalf of Australian College of Midwives. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

### Statement of significance

#### Problem

The COVID-19 pandemic may increase prenatal stress and, in turn, adversely impact both health behaviours and health outcomes.

### What is already known

The pandemic has been associated with increased incidence of mental health symptoms in general and pregnant populations, and poses a threat to protective factors like social support.

### What this paper adds

A multidimensional examination of prenatal general, pregnancy-specific, and COVID-19 related stress, health, and health behaviours during the COVID-19 pandemic. The study found high adherence to both pregnancy- and pandemic-related health advice. Evidence of high levels of pregnancy- and pandemic-related stress provide insight into potentially useful targets for stress intervention.

\* Corresponding author.

E-mail address: [119222524@umail.ucc.ie](mailto:119222524@umail.ucc.ie) (J. Pope).

<http://dx.doi.org/10.1016/j.wombi.2021.03.007>

1871-5192/© 2021 The Authors. Published by Elsevier Ltd on behalf of Australian College of Midwives. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Introduction

Many women experience disruptions to mental health during pregnancy. It is estimated, for example, that between 7% and 30% of women exhibit symptoms of depression during their pregnancies [1]; between 18.2% and 24.6% report anxiety [2]; and up to 84% experience stress [3]. Depression, stress and anxiety are related yet distinct psychological constructs [4,5] which have been associated with a range of poor outcomes including increased risk of premature birth and reduced length of gestation [6] and disruptions to hormonal regulation in children [7]. Prenatal mental health issues may also be associated with poorer physical health and health behaviours, with further direct and indirect effects on maternal and infant outcomes [8]. For instance, low physical activity, poor diet, and substance use during pregnancy are associated with increased risk of adverse prenatal [9], health [9–12], and child developmental [13] outcomes.

Prenatal stress and mental health issues can arise from multiple factors, including from experiencing stressful life events like the COVID-19 pandemic [14,15]. COVID-19, which was declared a pandemic by the World Health Organisation in March 2020 [16], is a significant global public health crisis. The pandemic and pandemic-related restrictions, such as lock-downs and quarantining, are already demonstrating significant negative effects for psychological well-being [17]. In frontline workers and general populations, COVID-19 has been associated with increased risks for depression, anxiety [17–19] vicarious trauma, and sleep disruptions [17,19]. High rates of substance use and suicidal ideation have also been reported among a general population during the pandemic [17].

Pregnant women are potentially at increased risk of adverse outcomes during the pandemic due to compounding pregnancy-specific concerns. For example, women may experience distress due to fear of coronavirus infection during pregnancy and/or in fetuses and children [20]; lack of clarity about pregnancy-specific risks of COVID-19 and which precautions are most effective at reducing infection risks to infants [21]; and changes to maternity services and birth plans due to public health restrictions [20,21]. In addition, COVID-19 restrictions have placed significant limitations on women's levels and types of social support, which has the potential to further strain their coping resources [20,22]. Given the proposed stress-buffering effects of social support, this may also increase women's risk of adverse outcomes both during pregnancy and beyond [23].

To date, some negative outcomes of the COVID-19 pandemic have been demonstrated in pregnant populations. For instance, one Canadian study reported increased incidences of anxiety, depression, post-traumatic stress, and dissociative symptoms among women pregnant during the COVID-19 pandemic, compared to a pre-pandemic cohort [24]. Another study reported increased levels of depressive symptoms in pregnant women in China after COVID-19 was declared an epidemic there in late January, again compared to a pre-pandemic baseline in the same population [25]. Reduced perceived social support has also been found among women pregnant in Ireland during the pandemic [26], and women also self-reported lower physical activity, compared to those pregnant before the pandemic [27]. These early findings suggest that women pregnant during the pandemic may be at risk for impaired prenatal psychological well-being and health behaviours. Examinations of stress have been limited however, with studies tending to focus on depression and/or anxiety symptoms. Similarly there has been little to no examination of prenatal health behaviours. Given the importance of maternal mental health, stress and health behaviours for maternal and child outcomes, examining these factors during the COVID-19 pandemic is essential to inform approaches to supporting pregnant women during this, and potential future, pandemics. The

aim of this study is to examine pregnant women's levels of stress, mental and physical health, and health behaviours during the COVID-19 pandemic.

## Participants, ethics and methods

### Design

A cross-sectional online survey of pregnant women was conducted. This study forms part of a larger project examining prenatal stress, social support and health behaviours [28].

### Ethical considerations

Ethical approval was granted by the [redacted] Research Ethics Committee and the Clinical Research Ethics Committee at [redacted]. Funding organisations were not involved in conceptualising the study, collecting data, analysing/reporting results, or any decisions related to publication.

### Participants

Participants in this study were pregnant women over 18 years of age. No exclusions were made based on nationality or current place of residence. Participants were recruited between June 16th, 2020 and July 17th, 2020. All participants were recruited online via public postings on forums associated with pregnancy websites (e.g. What To Expect, Mumsnet), and on social media pregnancy/parenting communities (e.g. Facebook, Twitter, Instagram, Reddit); a full list of targeted communities is available in Supplement 1. Participants were provided with a brief study description and link to the study survey, containing the full study information leaflet and consent form. Participants provided consent prior to completing the survey.

### Measures

The survey included open and closed questions on demographic factors, stress, mental and physical health, health behaviours, social support, and perceived satisfaction and quality of antenatal care.

Details and findings in relation to social support, quality of antenatal care, stress-reduction strategies, and experiences of distress during the pandemic are presented in our complementary paper [29]. As such they are not reported as outcomes in the current paper but are presented here for completeness.

### Demographic factors

Data on the following demographic factors were collected: age, marital status, education, household income, ethnicity, nationality, country of residence, weeks of pregnancy, and number of previous pregnancies.

### General stress

General stress was measured using the 4-item Perceived Stress Scale Short Form (PSS-4) [30]. This validated scale asks participants to report frequency of experiencing stress on a 5-point Likert scale ranging from 0 = "never" to 4 = "very often". Items include "How often have you felt that you were unable to control the important things in your life?" In the current study the PSS-4 had a reliability coefficient of  $\alpha = .75$ .

### Pregnancy-specific stress

Pregnancy-specific stress was measured using the validated 17-item Revised Prenatal Distress Scale (NuPDQ) [31]. Items ask

participants to report whether they are feeling “bothered, upset, or worried at this point in their pregnancies” about pregnancy-specific issues such as concerns about their pregnancy or baby, and changes to self-image and social relationships. Items are rated on a 3-point Likert scale ranging from 0 = “not at all” to 2 = “very much.” In this study the NuPDQ had a reliability coefficient of  $\alpha = .76$ .

#### COVID-19 related stress

At the time of this study there was no published validated scale available to measure COVID-19 related stress during pregnancy. As such, and guided by available evidence from a recent study by Zhong et al. [32], 6 items were developed to examine women’s concerns about the COVID-19 pandemic in this study. Participants were asked to indicate how worried they were about the coronavirus overall using a single item, and 5 items assessing concerns about coronavirus infection, being able to sleep because of coronavirus concerns, restrictions in place due to the coronavirus, and access to antenatal care due to the coronavirus. Items were measured using a 3-point Likert scale ranging from 0 = “not at all” to 2 = “very much.” The internal validity for this measure was  $\alpha = .77$ .

#### Mental and physical health

Mental and physical health were measured using the mental and physical domains of the validated 12-item Short Form Survey (SF-12) [33]. Participants indicated their general perceived health on a 5-point Likert scale (ranging from 1 = “excellent” to 5 = “poor”). Participants were also asked to indicate health-related limitations to activities using a 3-point Likert scale (ranging from 1 = “Limited a lot” to 3 = “Not at all limited”), and by answering “yes” or “no” to questions about specific limitations. Participants indicated how much pain was interfering with their ability to work using a five-point Likert scale (ranging from 1 = “not at all” to 5 = “extremely”). Participants were also asked about the frequency of emotional experiences, such as “feeling down-hearted or blue” and whether their health interfered with social functioning, on a 6-point Likert scale (ranging from 1 = “all of the time” to 6 = “none of the time”).

#### Prenatal health behaviours

Frequency of engaging in prenatal health behaviours was measured using the validated 20-item Prenatal Health Behaviour (PHBS) scale [34]. Health behaviours were grouped into a health-promoting behaviours subscale ( $\alpha = .72$ ) and a health-impairing behaviours subscale ( $\alpha = .55$ ) as per previous guidance [35]. Health-promoting behaviours included exercise, stretching, getting enough sleep, taking vitamins, and maintaining a good diet. Health-impairing behaviours included skipping meals, consuming snack foods instead of balanced meals, consuming drinks with caffeine, and smoking cigarettes. Items are measured on a Likert scale ranging from 0 = “never” to 4 = “very often”. The internal consistency of the health-promoting subscale was  $\alpha = .72$ . The internal consistency of the health-impairing subscale was  $\alpha = .55$ , which, though lower than desired, is in keeping with previous examinations [34].

#### COVID-19 related health behaviours

Six items were developed to assess women’s adherence to COVID-19 related health behaviours. Women were asked to report whether they were using a mask outside the home and in hospital, avoiding crowded places, leaving the house only when essential, shaking hands or making close contact with people, and disinfecting frequently touched objects, by answering either “yes” or “no.”

#### Social support

Social support was measured using the 12-item Multidimensional Perceived Social Support Scale (MSPSS) [36]. This

validated scale asks women rated their perceived degree of support from significant others, family and friends, by rating statements (for example: “There is a special person with whom I can share my joys and sorrows”) on a 7-point Likert scale ranging from 1 = “very strongly disagree” to 7 = “very strongly agree.” The overall scale reliability was  $\alpha = .96$ ; the reliability of the significant other, family, and friends subscales were  $\alpha = .98$ ,  $\alpha = .94$ , and  $\alpha = .96$  respectively.

#### Open-ended questions

Women were also invited to provide additional qualitative feedback on their prenatal stress experiences including what they found stressful, what stress-reduction strategies they engaged in, and what services and supports they use/want/need during their pregnancies.

#### Analysis

Descriptive statistics were used to summarise information on participant demographics, stress, mental and physical health, social support, health behaviours, and COVID-19 related behaviours. Associations between demographic factors, stress, mental and physical health, social support, and all health behaviours were examined using Spearman’s rank order correlations. Hierarchical multiple regressions were conducted to examine the role of sociodemographic and well-being factors (stress, mental and physical health, and social support) in predicting general stress, pregnancy-specific stress, COVID-19 related stress, mental and physical health, and health behaviours. Sociodemographic factors were entered at Step 1 of the hierarchical regressions; followed by well-being factors at Step 2 to determine the impact of well-being factors when controlling for socio-demographic factors. Dummy variables were created from multilevel categorical variables for inclusion in the regression models. Chi square tests of independence were conducted to examine potential demographic differences in COVID-19 related health behaviours. Due to the use of multiple inferential tests, Bonferroni corrections were applied with a more stringent significance value, calculated by dividing an alpha of .05 by the number of regression tests ( $n = 7$ ), which results in a significance value rounded up to  $p < .01$ . This more stringent significance value was used in all analyses.

#### Results

573 pregnant women took part in the survey. Participants were most commonly resident in the United States (42.6%,  $n = 243$ ), Ireland (41.2%,  $n = 235$ ), or the United Kingdom (10%,  $n = 57$ ). The majority of participants were married (80%,  $n = 457$ ) and educated to degree level or above (79.3%,  $n = 453$ ). The majority (57.2%,  $n = 325$ ) had been pregnant before. See Table 1 for complete participant characteristics.

Details of women’s self-reported general stress, pregnancy-specific stress, COVID-19 related stress, and perceived mental and physical health are presented in Table 2. For COVID-19 related stress, 58% of participants reported feeling “very” concerned about COVID-19 related restrictions; 39.4% of women reported feeling “very” concerned about their babies becoming unwell because of the coronavirus; and only 31.7% of participants felt confident that COVID-19 would be successfully controlled. Participants reported normative levels [37] of general stress ( $M = 6.54$ ,  $SD = 2.84$ ), and high [31] levels pregnancy-specific stress ( $M = 13.89$ ,  $SD = 5.37$ ). Participants also reported low levels of physical ( $M = 44.3$ ,  $SD = 9.2$ ) and mental health ( $M = 42.5$ ,  $SD = 11.3$ ) [38].

Participants reported high frequency of engagement in health-promoting behaviours and low engagement in

**Table 1**  
Participant characteristics.

|                                    | M(SD)        | Range |
|------------------------------------|--------------|-------|
| Age (years)                        | 32.38 (4.6)  | 19–43 |
| No. weeks pregnant                 | 26.22 (9.43) | 4–42  |
| No. children                       | .78 (.993)   | 0–7   |
|                                    | n            | %     |
| Nationality                        |              |       |
| American                           | 220          | 39.2  |
| Other                              | 53           | 9.9   |
| British                            | 60           | 10.7  |
| Canadian                           | 13           | 2.3   |
| Irish                              | 215          | 38.3  |
| Current country of residence       |              |       |
| America                            | 243          | 42.6  |
| Australia                          | 4            | .7    |
| Bahrain                            | 1            | .2    |
| Canada                             | 15           | 2.6   |
| Ireland                            | 235          | 41.2  |
| United Kingdom                     | 57           | 10.0  |
| Germany                            | 2            | .4    |
| Guam                               | 1            | .2    |
| New Zealand                        | 2            | .4    |
| Northern Ireland                   | 3            | .5    |
| Norway                             | 1            | .2    |
| Scotland                           | 1            | .2    |
| South Africa                       | 2            | .4    |
| Spain                              | 1            | .2    |
| Wales                              | 2            | .4    |
| Ethnic/cultural background         |              |       |
| Irish                              | 224          | 39.2  |
| Irish Traveller                    | 1            | .2    |
| Other white background             | 289          | 50.5  |
| African                            | 6            | 1.0   |
| Other Black background             | 3            | .5    |
| Other Asian background             | 7            | 1.2   |
| Hispanic/Latina                    | 13           | 2.3   |
| Mixed Ethnicity                    | 13           | 2.3   |
| Other                              | 10           | 1.7   |
| Not stated                         | 6            | 1.0   |
| Highest level of education         |              |       |
| Secondary school                   | 26           | 4.6   |
| Technical/vocational qualification | 39           | 6.8   |
| Non-degree qualification           | 53           | 9.3   |
| Undergraduate degree               | 213          | 37.3  |
| Postgraduate degree                | 197          | 34.5  |
| Doctorate                          | 43           | 7.5   |
| Relationship status                |              |       |
| Married                            | 457          | 80.0  |
| Cohabiting                         | 76           | 13.3  |
| In a relationship                  | 29           | 5.1   |
| Single                             | 9            | 1.6   |
| First pregnancy                    |              |       |
| No                                 | 325          | 57.2  |
| Yes                                | 243          | 42.8  |
| Prior pregnancy loss               |              |       |
| Never                              | 365          | 63.7  |
| Yes, once                          | 141          | 24.6  |
| Yes, twice                         | 39           | 6.8   |
| Yes, 3 or more times               | 28           | 4.9   |

health-impairing behaviours. For example, most participants reported abstaining entirely from alcohol (87.2%) and cigarettes (96.2%); as such these variables were excluded from further analyses. Participants also reported high levels of adherence to COVID-19 public health advice; for instance, 95.6% of participants reported that they had stopped shaking hands or making close contact with others, while 76.3% reported wearing masks when leaving home (see Table 3).

Findings of associations between stress, mental and physical health, and health behaviours using Spearman's correlations are presented in Supplement 2.

*Predictors of stress, mental and physical health, and health behaviours*

Hierarchical multiple regressions were conducted to examine the role of sociodemographic and well-being factors in predicting stress, mental and physical health, and health behaviours. Preliminary analyses confirmed normality, linearity, multicollinearity and homoscedasticity. A summary of regression outcomes is provided in Table 4 below; full details of each regression are available in Supplements 3–5.

*COVID-19 related stress*

Sociodemographic factors were entered at Step 1 and explained 4.9% of the variance in perceived stress. After entry of the well-being factors (general stress, pregnancy-specific stress, social support, and mental and physical health) at Step 2, the total variance explained by the model was 31.1%,  $F(23, 474) = 9.31$ ,  $p < .001$ . The well-being factors explained an additional 26.2% of the variance in stress after controlling for sociodemographic factors;  $R$  squared change = .26,  $F$  change (5, 474) = 36.07,  $p < .001$ . In the final model, only general stress ( $\beta = -.41$ ,  $p < .001$ ) and pregnancy-specific stress ( $\beta = .17$ ,  $p = .002$ ) predicted COVID-19 related stress. See Supplement 3.

*Pregnancy-specific stress*

Sociodemographic factors entered at Step 1 explained 7.5% of the variance in pregnancy-specific stress. After entry of the COVID-19 related stress, general stress, social support, and mental and physical health at Step 2, the total variance explained by the model was 39.4%,  $F(23, 474) = 13.42$ ,  $p < .001$ . The well-being factors explained an additional 32% of the variance in stress, after controlling for sociodemographic factors;  $R$  squared change = .32,  $F$  change (5, 474) = 50.07,  $p < .001$ . In the final model, the number of children women had ( $\beta = -.21$ ,  $p < .001$ ), COVID-19 related stress ( $\beta = .36$ ,  $p < .001$ ), and mental health ( $\beta = -.23$ ,  $p < .001$ ) predicted pregnancy-specific stress. See Supplement 3.

*General stress*

Sociodemographic factors entered at Step 1 explained 8.1% of the variance in general stress. After entry of the COVID-19 related stress, pregnancy-specific stress, social support, and mental and physical health at Step 2, the total variance explained by the model was 48.9%,  $F(23, 474) = 21.72$ ,  $p < .001$ . The well-being factors explained an additional 43% of the variance in stress, after controlling for sociodemographic factors;  $R$  squared change = .43;  $F$  change (5, 474) = 84.06,  $p < .001$ . In the final model, COVID-19 related stress ( $\beta = .12$ ,  $p = .002$ ), mental health ( $\beta = -.59$ ,  $p < .001$ ) and physical health ( $\beta = -.11$ ,  $p = .003$ ) predicted general stress. See Supplement 3.

*Perceived mental health*

Sociodemographic factors, entered at Step 1, explained 7.9% of the variance in mental health as measured by the SF-12. After entry of the stress measures, social support and physical health at Step 2, the total variance explained by the model was 52.1%,  $F(23, 474) = 24.54$ ,  $p < .001$ . The well-being factors explained an additional 43% of the variance in mental health, after controlling for sociodemographic factors;  $R$  squared change = .43,  $F$  change (5, 474) = 89.60,  $p < .001$ . In the final model, gestational weeks ( $\beta = .09$ ,  $p = .006$ ), pregnancy-specific stress ( $\beta = -.15$ ,  $p < .001$ ), general stress ( $\beta = -.55$ ,  $p < .001$ ), and physical health ( $\beta = -.23$ ,  $p < .001$ ) predicted mental health. See Supplement 4.

*Perceived physical health*

Sociodemographic factors, entered at Step 1, explained 7.6% of the variance in physical health measured by the SF-12. After entry of the stress, social support, and mental health measures at Step 2,

**Table 2**  
Participant stress, mental and physical health, and social support.

|                                          | M(SD)         | Range       |
|------------------------------------------|---------------|-------------|
| General stress (PSS)                     | 6.54 (2.84)   | .00–15.00   |
| Pregnancy-specific stress (PDQ)          | 13.89 (5.37)  | .00–30.00   |
| COVID-related stress                     | 7.14 (2.84)   | .00–12.00   |
| SF-12 Mental Health Component            | 42.49 (11.30) | 10.44–65.55 |
| SF-12 Physical Health Component          | 44.25 (9.17)  | 14.19–64.27 |
| Perceived Social Support (total) (MSPSS) | 67.61 (15.89) | 12.00–84.00 |
| From significant other                   | 24.84 (5.65)  | 4.00–28.00  |
| From family                              | 21.93 (6.21)  | 4.00–28.00  |
| From friends                             | 20.79 (6.42)  | 4.00–28.00  |

  

|                                                                                                          | n   | %    |
|----------------------------------------------------------------------------------------------------------|-----|------|
| Unable to fall or stay asleep due to thinking about the coronavirus                                      |     |      |
| Not at all                                                                                               | 264 | 46.2 |
| Somewhat                                                                                                 | 213 | 37.3 |
| Very much                                                                                                | 94  | 16.5 |
| Worried about being infected by the coronavirus                                                          |     |      |
| Not at all                                                                                               | 111 | 19.4 |
| Somewhat                                                                                                 | 260 | 45.5 |
| Very much                                                                                                | 200 | 35.0 |
| Worried about baby becoming unwell because of the coronavirus                                            |     |      |
| Not at all                                                                                               | 89  | 15.5 |
| Somewhat                                                                                                 | 258 | 45.0 |
| Very much                                                                                                | 226 | 39.4 |
| Worried about restrictions in place due to the coronavirus (e.g. social distancing, travel restrictions) |     |      |
| Not at all                                                                                               | 46  | 8.0  |
| Somewhat                                                                                                 | 194 | 33.9 |
| Very much                                                                                                | 332 | 58.0 |
| Worried about access to antenatal care because of the coronavirus                                        |     |      |
| Not at all                                                                                               | 133 | 23.4 |
| Somewhat                                                                                                 | 206 | 36.2 |
| Very much                                                                                                | 230 | 40.4 |
| Overall, worry about the coronavirus                                                                     |     |      |
| Not at all                                                                                               | 30  | 5.3  |
| Somewhat                                                                                                 | 283 | 49.7 |
| Very much                                                                                                | 256 | 45.0 |
| Belief that COVID-19 will be successfully controlled                                                     |     |      |
| No                                                                                                       | 132 | 23.1 |
| Yes                                                                                                      | 181 | 31.7 |
| I don't know                                                                                             | 258 | 45.2 |

MSPSS = Multidimensional Scale of Perceived Social Support PSS = Perceived Stress Scale PDQ = Prenatal Distress Questionnaire SF-12 = SF-12<sup>®</sup> Health Survey.

the total variance explained by the model was 19.5%,  $F(23, 474) = 6.24, p < .001$ . The well-being factors explained an additional 12.3% of the variance in physical health, after controlling for sociodemographic factors;  $R^2$  change = .123,  $F$  change (5, 474) = 15.20,  $p < .001$ . In the final model, gestational weeks ( $\beta = -.23, p < .001$ ), pregnancy-specific stress ( $\beta = -.29, p < .001$ ), general stress ( $\beta = -.17, p = .003$ ), and mental health ( $\beta = -.38, p < .001$ ) predicted physical health. See Supplement 4.

#### Health-promoting behaviours

Sociodemographic factors entered at Step 1 explained 10.2% of the variance in health-promoting behaviours. After entry of stress measures, social support, and mental and physical health (well-being factors) at Step 2, the total variance explained by the model was 24.8%,  $F(24, 473) = 6.52, p < .001$ . The well-being factors explained an additional 14.7% of the variance in health-promoting behaviours, after controlling for sociodemographic factors;  $R^2$  change = .147,  $F$  change (6, 473) = 15.39,  $p < .001$ . In the final model, having a technical or vocational qualification ( $\beta = .17, p = .008$ ) or a doctorate ( $\beta = -.189, p = .004$ ), physical health ( $\beta = .24, p < .001$ ), mental health ( $\beta = .27, p < .001$ ) and social support ( $\beta = .12, p = .005$ ) predicted health-promoting behaviours. See Supplement 5.

#### Health-impairing behaviours

Sociodemographic factors were entered at Step 1 and explained 12.3% of the variance in health-impairing behaviours. After entry of

stress measures, social support, and mental and physical health at Step 2, the total variance explained by the model was 16.9%,  $F(24, 473) = 3.99, p < .001$ . The well-being factors explained an additional 4.5% of the variance in health-impairing behaviours, after controlling for sociodemographic factors;  $R^2$  change = .045,  $F$  change (6, 473) = 4.31,  $p < .001$ . In the final model, age ( $\beta = -.17, p = .002$ ), number of children ( $\beta = .25, p < .001$ ), being resident in the UK ( $\beta = -.18, p = .007$ ), and pregnancy-specific stress ( $\beta = .19, p < .001$ ) predicted health-impairing behaviours. See Supplement 5.

#### Demographic differences in COVID-19 health behaviours

Chi square tests conducted to examine potential demographic differences in COVID-19 related health behaviours indicated that wearing a mask when leaving home was significantly associated with relationship status ( $\chi^2(3) = 19.49, p < .001$ ): married women were most likely to wear a mask. Wearing a mask was also associated with country of residence ( $\chi^2(3) = 87.30, p < .001$ ), with women in America more likely to report wearing a mask than their counterparts in Ireland or the United Kingdom. Wearing a mask when in hospital/maternity services was also associated with the country of residence ( $\chi^2(3) = 41.08, p < .001$ ): women in America were most likely to do so. Going to crowded places was associated with education ( $\chi^2(5) = 22.92, p < .001$ ), women with a postgraduate degree being least likely to go to crowded places; and with country of residence ( $\chi^2(3) = 24.34, p < .001$ ), women in Ireland being most likely to go to crowded places. Disinfecting and cleaning, leaving

**Table 3**  
General and COVID-19 Specific Health Behaviors.

|                                                               | M (SD)       | Range       |     |      |
|---------------------------------------------------------------|--------------|-------------|-----|------|
| Health-promoting behaviors (PHBS)                             | 24.91 (4.33) | 10.00–32.00 |     |      |
| Health-impairing behaviors (PHBS)                             | 10.22 (3.75) | .00–26.00   |     |      |
|                                                               |              |             | n   | %    |
| Frequency of drinking alcohol <sup>a</sup>                    |              |             |     |      |
| Never                                                         |              |             | 499 | 87.2 |
| Almost never                                                  |              |             | 53  | 9.3  |
| Sometimes                                                     |              |             | 17  | 3.0  |
| Fairly often                                                  |              |             | 2   | .3   |
| Very often                                                    |              |             | 1   | .2   |
| Frequency of smoking cigarettes <sup>a</sup>                  |              |             |     |      |
| Never                                                         |              |             | 551 | 96.2 |
| Almost never                                                  |              |             | 10  | 1.7  |
| Sometimes                                                     |              |             | 6   | 1.0  |
| Fairly often                                                  |              |             | 2   | .3   |
| Very often                                                    |              |             | 4   | .7   |
| Frequency of exercising for at least 15 min <sup>a</sup>      |              |             |     |      |
| Never                                                         |              |             | 39  | 6.8  |
| Almost never                                                  |              |             | 79  | 13.8 |
| Sometimes                                                     |              |             | 171 | 29.8 |
| Fairly often                                                  |              |             | 149 | 26.0 |
| Very often                                                    |              |             | 135 | 23.6 |
| Frequency of getting enough sleep <sup>a</sup>                |              |             |     |      |
| Never                                                         |              |             | 10  | 1.8  |
| Almost never                                                  |              |             | 84  | 14.7 |
| Sometimes                                                     |              |             | 180 | 31.6 |
| Fairly often                                                  |              |             | 198 | 34.7 |
| Very often                                                    |              |             | 98  | 17.2 |
| Going to any crowded place                                    |              |             |     |      |
| No                                                            |              |             | 441 | 77.0 |
| Yes                                                           |              |             | 132 | 23.0 |
| Wearing a mask when leaving home                              |              |             |     |      |
| No                                                            |              |             | 136 | 23.7 |
| Yes                                                           |              |             | 437 | 76.3 |
| Wearing a mask when in hospital/maternity service             |              |             |     |      |
| No                                                            |              |             | 78  | 13.6 |
| Yes                                                           |              |             | 494 | 86.4 |
| Cleaning and disinfecting frequently touched objects/surfaces |              |             |     |      |
| No                                                            |              |             | 98  | 17.1 |
| Yes                                                           |              |             | 475 | 82.9 |
| Leaving the house only when absolutely essential              |              |             |     |      |
| No                                                            |              |             | 152 | 26.6 |
| Yes                                                           |              |             | 420 | 73.4 |
| Shaking hands or making close contact with other people       |              |             |     |      |
| No                                                            |              |             | 24  | 4.2  |
| Yes                                                           |              |             | 548 | 95.8 |

PHBS = Prenatal Health Behavior Scale.

<sup>a</sup> Frequency of prenatal health behaviours is for the preceding month.

home only when essential, and avoiding handshakes/close contact were not significantly associated with any demographic factors. See Supplement 6 for complete details of findings regarding demographic differences in COVID-19 related health behaviours.

## Discussion

This study examined pregnant women’s levels of stress, mental and physical health, and frequency of health behaviours during the COVID-19 pandemic. Building upon existing evidence that pregnant women are experiencing more anxiety, depression, and post-traumatic stress during the pandemic [24,25], this study found high levels of pregnancy-specific stress and COVID-19 related stress, but low levels of mental and physical health and social support. The majority of women in this study engaged in recommended prenatal health behaviours and COVID-19 related behaviours. Psychological outcomes and health-promoting behaviours were mostly predicted by well-being variables including stress and social support; sociodemographic factors had limited impact on outcomes.

Our findings indicated that during the pandemic, pregnant women in our study were experiencing normative levels of general stress [37], but reported high levels of pregnancy-specific stress [31] and COVID-19 related stress. That prenatal and COVID-19 stress were heightened in this sample, but general stress was not, suggests that the experience of being pregnant during the pandemic may function as a contextual stressor. This is supported by our findings that nearly all participants reported feeling somewhat or very concerned about COVID-19/related restrictions, many were concerned about their babies becoming unwell due to the coronavirus, and many reported disruptions to sleep associated with thinking about the coronavirus. That is in line with previous evidence that the coronavirus has prompted pregnancy-specific concerns [20,21]. More generally, our findings of increased COVID-19 related stress are in keeping with previous examinations that indicated high levels of anxiety during the pandemic [24] as stress and anxiety are highly related, though distinct, constructs [5]. These findings provide insight on the specific content of women’s stress during the pandemic, and specifically suggest that stress reduction interventions which target pregnancy and coronavirus-related concerns may be particularly useful.

In addition, although significant differences in stress were not previously observed between women pregnant before or during the pandemic in Ireland [26], the current paper did not find any sociodemographic predictors of any stress outcomes in this international sample. This suggests that while experiences of stress during the pandemic may be contextually specific, they may not be place- or demographic-specific, a finding which suggests that population-level approaches to addressing and supporting prenatal mental health may be effective in the pandemic context. However, given the overall homogeneity of this sample in terms of ethnicity and education, and in light of existing evidence of the relationship between sociodemographic factors and mental health concerns [1], this likely bears further examination.

In this sample, the strongest predictors of stress outcomes were other types of stress and general mental health. Stressors identified by participants included concerns about COVID-19 infection during pregnancy and/or in infants, limitations on travel and social contact imposed by public health restrictions, and limitations on access to prenatal care as a result of public health measures. Additional qualitative feedback from this sample (presented in a complementary paper) suggests that changes to antenatal care, conflict with social support networks, and feelings of uncertainty and isolation were also important sources of stress for this population [29]. This suggests that the development and implementation of prenatal mental health supports and interventions should include a specific focus on pregnancy- and COVID-19-related concerns. Further, though our findings did not suggest an influence of demographic factors, predisposing psychological factors should be considered in the development and implementation of interventions.

In line with previous findings about the relationship between prenatal stress and health outcomes [8], general stress was shown to have a significant impact on both mental and physical health in this sample. Notably, pregnancy-specific stress was also a significant predictor of mental health, suggesting that pregnancy-related concerns may put women at increased risk for mental health impairment and could be useful targets for mental health intervention in pregnant populations. This may be of particular significance as increased mental health symptoms have been reported in pregnant populations since the beginning of the pandemic [24,25]. In addition, our findings suggest a bi-directional relationship between mental and physical health. Previous findings have demonstrated associations between prenatal stress and engagement in health behaviours [8]. However, our findings indicate that despite high levels of pregnancy related and



**Table 4**  
Summary of regression outcomes (full details in Supplementary files 3–5).

|                                                   | R   | R <sup>2</sup> | ΔR <sup>2</sup> | p Values |
|---------------------------------------------------|-----|----------------|-----------------|----------|
| <b>Covid-19 related stress</b>                    |     |                |                 |          |
| Step 1 <sup>a</sup>                               | .22 | .05            | .05             | .14      |
| Step 2 <sup>b</sup>                               | .59 | .31            | .26             | <.005    |
| <b>Pregnancy-specific stress (PDQ)</b>            |     |                |                 |          |
| Step 1 <sup>a</sup>                               | .27 | .08            | .08             | .004     |
| Step 2 <sup>b</sup>                               | .63 | .39            | .32             | <.005    |
| <b>General stress (PSS)</b>                       |     |                |                 |          |
| Step 1 <sup>a</sup>                               | .29 | .08            | .08             | .16      |
| Step 2 <sup>b</sup>                               | .72 | .51            | .43             | .03      |
| <b>Mental health (SF-12 mental component)</b>     |     |                |                 |          |
| Step 1 <sup>a</sup>                               | .34 | .11            | .11             | <.005    |
| Step 2 <sup>c</sup>                               | .74 | .54            | .43             | <.005    |
| <b>Physical health (SF-12 physical component)</b> |     |                |                 |          |
| Step 1 <sup>a</sup>                               | .33 | .11            | .11             | <.005    |
| Step 2 <sup>c</sup>                               | .48 | .23            | .12             | <.005    |
| <b>Health promoting behaviours</b>                |     |                |                 |          |
| Step 1 <sup>a</sup>                               | .32 | .10            | .10             | <.005    |
| Step 2 <sup>c</sup>                               | .49 | .25            | .15             | <.005    |
| <b>Health impairing behaviours</b>                |     |                |                 |          |
| Step 1 <sup>a</sup>                               | .35 | .12            | .12             | <.005    |
| Step 2 <sup>c</sup>                               | .41 | .17            | .05             | <.005    |

<sup>a</sup> Variables entered at Step 1: age, parity, gestational age, country of residence, relationship status, education, previous pregnancy loss.

<sup>b</sup> Variables entered at Step 2: Stress measures, Mental and physical health measures, social support.

<sup>c</sup> Variables entered at Step 2: Stress measures, health measures, social support PSS = Perceived Stress Scale PDQ = Prenatal Distress Questionnaire SF-12 = SF-12<sup>®</sup> Health Survey.

COVID-19 related stress, women in this study reported high adherence to health-promoting behaviours (e.g. eating healthily, exercising, taking vitamins). Similarly, women reported low engagement in health-impairing behaviours (e.g. alcohol consumption and smoking). This finding differs from two recent studies demonstrating lower reported physical activity during the pandemic [26,27]. It is important to note that high levels of self-reported appropriate health behaviours may reflect socially desirable responding in the current study. Unlike stress outcomes, sociodemographic factors including education, number of children, and country of residence, were predictors of health-promoting and health-impairing prenatal behaviours in the current study. The finding that the number of children women have impacts health behaviours is likely related to childcare and caring responsibilities; as there is evidence that changes to schooling and childcare provision during the pandemic have placed significant strain on families with children [39].

Finally, while women generally reported very high levels of adherence to COVID-19 related health behaviours (such as social distancing, wearing masks, and disinfecting surfaces), some sociodemographic differences in adherence were observed. Observed associations between some pandemic-related behaviours (avoiding public places and wearing masks) and country of residence, suggest that national guidance, regulations, and sociocultural contexts may significantly impact on pregnant women's health-related decision-making during the pandemic. Unlike the findings related to stress outcomes, these findings suggest a need for potentially more targeted interventions within countries and groups to improve and/or maintain appropriate prenatal behaviours during a pandemic.

#### Strengths/limitations

The cross-sectional nature of this study is limited as we are unable to track how the pandemic may impact participants' wellbeing- and behaviours over time/as perceived COVID-19 risk and restrictions change. In addition, while assessment of stress during the pandemic was conducted using validated measures of

general stress and pregnancy-specific stress, there was not, at the time of data collection, a published validated scale to measure COVID-19 related stress in pregnant women. While the measure of COVID-19 related stress designed for this study had good internal validity, a validated measure of prenatal stress and anxiety has since been published [40], and this would prove useful for future research in this area. We also did not assess whether women had received a positive COVID-19 test result, which might have influenced their well-being and behaviours.

Furthermore, participants in this study were largely married or in a relationship, educated to degree-level or higher, socioeconomically advantaged, and resident in Ireland, the United Kingdom or America. Thus these results may tend to understate the economic and social drivers of stress and adherence to health advice during the pandemic. However, the international sample enhances generalisability of our findings and interpretations. In addition, our comprehensive approach to examining multiple types of stress and a range of health behaviours provides important insights into as yet under-examined aspects of women's experiences of pregnancy during the pandemic.

#### Conclusion

This study found high levels of pregnancy-specific and COVID-19 related stress, and low levels of mental and physical health during the COVID-19 pandemic. In this sample, these variables were mostly predicted by well-being factors rather than sociodemographic factors. Pregnant women in this study reported good engagement with healthy behaviours and high adherence to COVID-19 public health recommendations during the pandemic.

Our findings highlight the need for prenatal mental health supports that target pregnancy- and coronavirus-specific concerns, and that can be implemented at population levels to support prenatal mental health. While experiences of stress did not appear to be place- or demographic-specific in this study, the association between certain health behaviours (including mask-wearing and avoiding crowded places), and sociodemographic factors, suggest that some elements of prenatal health behaviour may be influenced by national guidance and cultural context, further illustrating the need for structured interventions for women pregnant during the pandemic. These supports are clearly greatly needed and may help to mitigate adverse effects for women and children.

#### Author agreement

The authors hereby certify that the work presented in this submission is the authors' original work, has not been published (and is not under consideration for publication) elsewhere, and has been reviewed and approved by all authors. The authors also agree to abide by the copyright terms and conditions of Elsevier and the Australian College of Midwives.

#### Author contributions

KMS conceptualised and designed the study, collected and analysed the data, wrote and revised the final manuscript.

JP contributed to study design, collected data and assisted with data analysis, and critically reviewed and revised the manuscript.

EKO designed the study, contributed to analyses and critically reviewed and revised the manuscript.

SL conceptualised and designed the study and critically reviewed and revised the manuscript.

SM conceptualised and designed the study, contributed to analyses and critically reviewed and revised the manuscript.

## Funding

This research is supported by an Evidence Synthesis Ireland and Health Research Board Trials Methodology Research Network COVID-19 Emergency Evidence Response Service Summer Scholarship. The research is also supported by a Health Research Board Applying Research into Policy and Practice Fellowship (HRB-ARPP-A-2018-011). The funders had no role in the design, conduct, analysis or interpretation of the research.

## Ethical statement

All procedures were approved by the University College Cork School of Public Health Research Ethics Committee and the Clinical Research Ethics Committee of the Cork University Teaching Hospitals (Ref ECM 4 (zz) 10/01/12).

## Conflict of interest

The authors have no conflict of interest to declare.

## Acknowledgement

The authors would like to thank all the pregnant women who completed the survey.

## Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.wombi.2021.03.007>.

## References

- [1] A. Biaggi, S. Conroy, S. Pawlby, C.M. Pariante, Identifying the women at risk of antenatal anxiety and depression: a systematic review, *J. Affect. Disord.* 191 (February) (2016) 62–77.
- [2] C. Dennis, K. Falah-Hassani, R. Shiri, Prevalence of antenatal and postnatal anxiety: systematic review and meta-analysis, *Br. J. Health Psychol.* 210 (January) (2018) 315–323.
- [3] S.M. Woods, J.L. Melville, Y. Guo, M.Y. Fan, A. Gavin, Psychosocial stress during pregnancy, *Am. J. Obstet.* 202 (September) (2009) 61.e1–61.e617.
- [4] M.W. Eysenck, M. Fajkowska, Anxiety and depression: toward overlapping and distinctive features, *Cogn. Emot.* 32 (November (7)) (2018) 1391–1400.
- [5] V. Glover, Prenatal stress and the origins of psychopathology: an evolutionary perspective, *J. Child Psychol. Psychiatry* 52 (January (4)) (2011) 356–367.
- [6] A. Voellmin, S. Etringer, N. Moog, P.D. Wadhwa, C. Buss, Maternal positive affect over the course of pregnancy is associated with the length of gestation and reduced risk of preterm delivery, *J. Psychosom. Res.* 75 (October) (2013) 336–340.
- [7] A.T. de Bruijn, H.J. van Bakel, H. Wijnen, V.J. Pop, A.L. van Baar, Prenatal maternal emotional complaints are associated with cortisol responses in toddler and preschool aged girls, *Dev. Psychobiol.* 51 (November) (2009) 553–563.
- [8] M. Lobel, D.L. Cannella, J.E. Graham, C. DeVincent, J. Schneider, B.A. Meyer, Pregnancy-specific stress, prenatal health behaviors, and birth outcomes, *Health Psychol.* 27 (5) (2008) 604–615.
- [9] B.A. Bailey, R.J. Sokol, Prenatal alcohol exposure and miscarriage, stillbirth, preterm delivery, and sudden infant death syndrome, *Alcohol Res. Health* 34 (1) (2011) 86–91.
- [10] A. Castles, E.K. Adams, C.L. Melvin, C. Kelsch, M.L. Boulton, Effects of smoking during pregnancy. Five meta-analyses, *Am. J. Prev Med* 16 (April (3)) (1999) 2008–2015.
- [11] M.H. Davenport, S. Ruchat, V.J. Poitras, A.J. Garcia, C.E. Gray, N. Barrowman, Prenatal exercise for the prevention of gestational diabetes mellitus and hypertensive disorders of pregnancy: a systematic review and meta-analysis, *Br. J. Sports Med.* 52 (2008) 1367–1375.
- [12] B.L. Pineles, S. Hsu, E. Park, J. Samet, Systematic review and meta-analyses of perinatal death and maternal exposure to tobacco smoke during pregnancy, *Am. J. Epidemiol.* 84 (July (2)) (2016) 87–97.
- [13] A.J. Malin, S.A. Busgang, A.J. Cantoral, K. Svensson, M.A. Orjuela, I. Pantic, L. Schnaas, et al., Quality of prenatal and childhood diet predicts neurodevelopmental outcomes among children in Mexico city, *Nutrients* 10 (August (8)) (2018) 1093.
- [14] S. King, D. Laplante, Using natural disasters to study prenatal maternal stress in humans, *Adv. Neurobiol.* 10 (2015) 285–313.
- [15] K. Matvienko-Sikar, S. Meedya, C. Ravaldi, Perinatal mental health during the COVID-19 pandemic, *Women Birth* 33 (July (4)) (2020) 309–310.
- [16] World Health Organisation. Coronavirus Disease 2019 (COVID-19) Situation Report – 51. Available at: [https://www.who.int/docs/default-source/coronavirus/situation-reports/20200311-sitrep-51-covid-19.pdf?sfvrsn=1ba62e57\\_10](https://www.who.int/docs/default-source/coronavirus/situation-reports/20200311-sitrep-51-covid-19.pdf?sfvrsn=1ba62e57_10) (11 March 2020, cited 11 December 2020).
- [17] J. Torales, M. O'Higgins, J.M. Castaldelli-Maia, A. Ventriglio, The outbreak of COVID-19 coronavirus and its impact on global mental health, *Int. J. Soc. Psychiatry* 66 (March (4)) (2020) 317–320.
- [18] M.E. Czeisler, R.I. Lane, E. Petrosky, J.F. Wiley, A. Christensen, R. Njai, Mental health, substance use, and suicidal ideation during the COVID-19 pandemic—United States, June 24–30, 2020, *MMWR (of the Center for Disease Control and Prevention)* 69 (2020) 1049–1057.
- [19] R.P. Rajkumar, COVID-19 and mental health: a review of the existing literature, *Asian J. Psychiatr.* 52 (August) (2020) 102066.
- [20] G.A. Corbett, S.J. Milne, M.P. Hehir, S.W. Lindow, M.P. O'Connell, Health anxiety and behavioural changes of pregnant women during the COVID-19 pandemic, *Eur. J. Obstet. Gynecol. Reprod. Biol.* 249 (June) (2020) 96–97.
- [21] M. Yassa, P. Birol, C. Yirmibes, C. Usta, A. Haydar, A. Yassa, et al., Near-term pregnant women's attitude toward, concern about and knowledge of the COVID-19 pandemic, *J. Matern. Fetal Neonatal Med.* 22 (May) (2020) e-publication.
- [22] S.S.B. Thapa, A. Mainali, S.E. Schwank, G. Acharya, Maternal mental health in the time of the COVID-19 pandemic, *Acta Obstet. Gynecol. Scand.* 99 (May (7)) (2020) 817–818.
- [23] S. Cohen, T.A. Wills, Stress, social support, and the buffering hypothesis, *Psychol. Bull.* 98 (2) (1985) 310–357.
- [24] N. Berthelot, R. Lemieux, J. Garon-Bissonnette, C. Drouin-Maziade, É Martel, M. Maziade, Uptrend in distress and psychiatric symptomatology in pregnant women during the coronavirus disease 2019 pandemic, *Acta Obstet. Gynecol. Scand.* 99 (May (7)) (2020) 848–855.
- [25] Y. Wu, C. Zhang, H. Liu, C. Duan, C. Li, J. Fan, et al., Perinatal depressive and anxiety symptoms of pregnant women along with COVID-19 outbreak in China, *Am. J. Obstet. Gynecol.* 223 (August (2)) (2020) 240.e1–240.e9.
- [26] K. Matvienko-Sikar, J. Pope, A. Cremin, H. Carr, S. Leitao, E.K. Colander, S. Meaney, Differences in levels of stress, social support, health behaviours, and stress-reduction strategies for women pregnant before and during the COVID-19 pandemic, and based on phases of pandemic restrictions, in Ireland, *Women Birth* (October) (2020) (in press).
- [27] M.H. Davenport, S. Meyer, V.L. Mesh, M.C. Strynadka, R. Khurana, Moms are not OK: COVID-19 and maternal mental health, *Front. Womens Health* 1 (June (1)) (2020) 1–6.
- [28] K. Matvienko-Sikar, J. Pope, S. Leitao, E.K. Colander, S. Meaney, Prenatal Well-being and Health Behaviours. Available from: <https://doi.org/10.17605/OSF.IO/X357B> (2020 August 23; cited 11 December 2020).
- [29] S. Meaney, J. Pope, S. Leitao, E.K. Colander, K. Matvienko-Sikar, Formal and informal supports for pregnant women during the pandemic: Social support and satisfaction with maternity services, (2020) (under review).
- [30] S. Cohen, T. Kamarck, R. Mermelstein, A global measure of perceived stress, *J. Health Soc. Behav.* 24 (December (4)) (1983) 385–396.
- [31] S. Ibrahim, M. Lobel, Conceptualization, measurement, and effects of pregnancy-specific stress: review of research using the original and revised prenatal distress questionnaire, *J. Behav. Med.* 43 (June) (2019) 16–33.
- [32] B.-L. Zhong, W. Luo, H.-M. Li, Q.-Q. Zhang, X.-G. Liu, W.-T. Li, et al., Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional survey, *Int. J. Biol. Sci.* 16 (10) (2020) 1745–1752.
- [33] J. Ware, M. Kosinski, S.D. Keller, A 12-item short-form health survey: construction of scales and preliminary tests of reliability and validity, *Med. Care* 34 (March (3)) (1996) 220–233.
- [34] M.V. Auerbach, J. Nicoloso-SantaBarbara, L. Rosenthal, C. Kocis, E.R. Weglarz, C. Busso, et al., Psychometric properties of the prenatal health behavior scale in mid- and late pregnancy, *J. Psychosom. Obstet.* 38 (2) (2017) 143–151.
- [35] M.V. Auerbach, M. Lobel, D.T. Cannella, Psychosocial correlates of health-promoting and health-impairing behaviors in pregnancy, *J. Psychosom. Obstet.* 35 (July (4)) (2014) 76–83.
- [36] G.D. Zimet, N.W. Dahlem, S.G. Zimet, G.K. Farley, The multidimensional scale of perceived social support, *J. Pers. Assess.* 51 (1) (1988) 30–41.
- [37] S.L. Wartting, M.J. Forshaw, J. South, A.K. White, New, normative, English-sample data for the short form perceived stress scale (PSS-4), *J. Health Psychol.* 18 (October (2)) (2013) 617–1628.
- [38] B. Gandek, J.E. Ware, N.K. Aaronson, G. Apolone, J.B. Bjorner, J.E. Brazier, M. Bullinger, S. Kaasa, A. Leplege, L. Prieto, M. Sullivan, Cross-validation of item selection and scoring for the SF-12 health survey in nine countries: results from the IQOLA project, *J. Clin. Epidemiol.* 51 (November (11)) (1998) 1171–1178.
- [39] A. Sevilla, SS. Mith. Baby Steps: The Gender Division of Childcare During the COVID19 Pandemic. CEPR Discussion Paper No. DP14804. Available at SSRN: <https://ssrn.com/abstract=3612873>.
- [40] H. Preis, B. Mahaffey, C. Heiselman, M. Lobel, Pandemic-related pregnancy stress and anxiety among women pregnant during the coronavirus disease 2019 pandemic, *Am. J. Obstet. Gynecol. MFM* 2 (August (3)) (2020) 100155.