



City Research Online

City, University of London Institutional Repository

Citation: Lalor, J., Ayers, S., Celleja Agius, J., Downe, S., Gouni, O., Hartmann, K., Nieuwenhuijze, M., Oosterman, M., Turner, J. D., Karlsdottir, S. I. & et al (2021). Balancing restrictions and access to maternity care for women and birthing partners during the COVID-19 pandemic: the psychosocial impact of suboptimal care. *BJOG: An International Journal of Obstetrics and Gynaecology*, 128(11), pp. 1720-1725. doi: 10.1111/1471-0528.16844

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/26461/>

Link to published version: <https://doi.org/10.1111/1471-0528.16844>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Balancing restrictions and access to maternity care for women and birthing partners during the COVID-19 pandemic: the psychosocial impact of suboptimal care

J Lalor,^a S Ayers,^b J Celleja Agius,^c S Downe,^d O Gouni,^e K Hartmann,^f M Nieuwenhuijze,^g M Oosterman,^h JD Turner,ⁱ SI Karlsdottir,^j A Horsch^k

^a School of Nursing and Midwifery, Trinity College Dublin, Dublin 2, Ireland ^b Centre for Maternal and Child Health, City University of London, London, UK ^c Faculty of Medicine and Surgery, University of Malta, Msida, Malta ^d University of Central Lancashire, Preston, UK ^e Cosmoanelixis Prenatal & Life sciences, Athens, Greece ^f Mother Hood e.V., Bundeselterninitiative zum Schutz von Mutter und Kind während Schwangerschaft, Bonn, Germany ^g Research Centre for Midwifery Science Maastricht, Zuyd University, Maastricht, The Netherlands ^h Vrije Universiteit Amsterdam, Amsterdam, The Netherlands ⁱ Department of Infection and Immunity, Luxembourg Institute of Health, Strassen, Luxembourg ^j University of Akureyri Iceland, Akureyri, Iceland ^k University of Lausanne, Lausanne, Switzerland
Correspondence: J Lalor, School of Nursing and Midwifery, Trinity College Dublin, 24 D'Olier Street, Dublin 2, Ireland.
Email: j.lalor@tcd.ie

Accepted 2 July 2021.

Please cite this paper as: Lalor J, Ayers S, Celleja Agius J, Downe S, Gouni O, Hartmann K, Nieuwenhuijze M, Oosterman M, Turner JD, Karlsdottir SI, Horsch A. Balancing restrictions and access to maternity care for women and birthing partners during the COVID-19 pandemic: the psychosocial impact of suboptimal care. BJOG 2021; <https://doi.org/10.1111/1471-0528.16844>.

Key points

- 1 Maternity services across Europe during the pandemic has undergone changes to limit virus transmission; however, many changes are not evidence-based.
- 2 Although these changes were introduced to keep women, babies and healthcare staff safe, the exclusion of companions and the separation of mothers and babies is particularly antithetical to a human rights-based approach to quality care.
- 3 A poll of COST Action 18211 network members showed that inconsistency in the application of restrictions was high, and there were significant deviations from the recommendations of authoritative bodies.
- 4 Concerns have emerged that restrictions in practice may have longer term negative impacts on mothers and their families and, in particular, may impact on the long-term health of babies.
- 5 When practice changes deviate from evidence-based frameworks that underpin quality care, they must be monitored, appraised and evaluated to minimise unintended iatrogenic effects.

Introduction

A woman's right to respectful and dignified care during labour and childbirth is strategically accepted.¹ As management committee members of the EU COST Action CA18211 network ('DEVOTION') focused on traumatic

childbirth (www.ca18211.eu), we are concerned with ensuring a positive birth experience for all. We work on a pan-European level to ensure women's rights to give birth in a clinically and psychologically safe environment including during the current COVID-19 pandemic.

As every country reacted to the COVID-19 pandemic, the swift initial response was based on the basic principles

of infection control, intended to protect all citizens. However, many governments and healthcare workers acted independently, as they waited for emerging evidence and detailed guidance from authoritative organisations and professional bodies to inform appropriate action. The emerging guidance was quickly changing, with fundamental differences in the recommendations of key international bodies, such as the World Health Organization (WHO), Royal College of Obstetricians and Gynaecologists (RCOG), American College of Obstetricians and Gynecologists (ACOG), and Royal College of Midwives (RCM).

While grappling with the public health crisis, many institutional settings imposed significant restrictions on key aspects of maternity services, such as prohibiting a birth companion in labour, placing limitations on breastfeeding, and reducing the contact between a mother and her baby.² Although these interventions were introduced to keep women, babies and healthcare staff safe, excluding companions and separating mothers from their babies are particularly antithetical to a human rights-based approach to quality care. Questions are now being raised about the appropriate balance between infection control and optimal maternity care, particularly in terms of the longer term clinical and psychosocial consequences for the mother, her baby and the family. Women are reporting negative consequences of reduced access to professional care, and of increased interventions, designed to reduce infection risk but associated with increased levels of iatrogenic harm.²

Accounts of restrictions have fuelled fear for some women, especially in the absence of good quality information from official sources, and in the context of alarming social media comments. As a consequence, reports have emerged that substantial minorities of women across Europe have not been accessing publicly provided maternity services, either because they are no longer on offer, or for fear of infection, or because they do not want to be isolated and separated from their accompanying partner. In some cases, this has widened the gap in health equality: where affordable, private consultations were booked and in other cases services have not been accessed at all by some women. Antenatal and childbirth classes were replaced with virtual formats, excluding women without appropriate devices or broadband.³

Women having ultrasound screening had to come alone, facing the possible diagnosis of a fetal anomaly, or even of intra-uterine death, alone. Serious limitations were placed on community services, such as support for breastfeeding. Midwives involved in parentcraft were transferred to public health departments to assist in contact-tracing, implying that their support services to women at this critical time was not essential. Examples of the reorganisation of care from home or birth centres to hospital settings have been seen, as a perception emerged that community care was less safe.

Mapping the European response

In response to these issues, the COST Action CA18211 network undertook a poll of network members, operationalised at a virtual meeting of the network on 25 and 26 November 2021, to explore the situation of maternity care provision in Europe. There were 88 clinicians and researchers from 32 participating countries, representing different disciplines, such as midwifery, obstetrics, nursing, psychology, psychiatry, biology, as well as members of lay advocacy groups. One session focused on the impact of COVID-19 on maternity care: representatives from 11 countries gave presentations and members from 23 countries added information via the chat. Variations in maternity care and restrictions between and within countries were highlighted. Key themes are outlined in Table 1.

These responses illustrate that inconsistency in the application of restrictions was high and that there were significant deviations from the recommendations of authoritative bodies, such as the WHO,³ RCOG⁴ and RCM.⁵ Most consistently, the restrictions excluded birth companions to various degrees, and women were separated from their babies or had significant limitations placed on the level of contact they could have if their baby was in the NICU. Some COST Action CA18211 network respondents were particularly concerned that locally applied restrictions deviated from international guidance (in the absence of evidence to support such restrictions) and also that some services were reporting an increase (without evidence of clinical indication) in interventions, such as induction of labour, and caesarean section rates. Others reported an increase in unplanned out-of-hospital births, as women were delaying coming to hospital. Finally, a recent meta-analysis showed that global maternal and fetal outcomes have worsened since the beginning of the pandemic, e.g. the rate of stillbirths increased by 28%.⁶

What is evident from this network internal poll is that despite the lack of evidence to justify severe restrictions, these were continuing in many maternity services, even though emerging research confirms they are not necessary or helpful to protect mothers, babies and healthcare staff from transmission of the virus. Such restrictions may contribute to an environment in which women may be more at risk of experiencing a traumatic birth, and raise questions about the extent to which women are exposed to human rights violations due to the continued implementation of potentially harmful practices. Data from a systematic review and meta-analysis⁶ show that rates of perinatal mental health disorders such as anxiety and depression have been higher during the pandemic and may be partially attributable to modifications to maternity services. The MBRRACE-UK rapid report⁷ highlighted two instances where women died by suicide after referrals to perinatal

Table 1. Key practice changes in 32 European countries

Key practice change	No. of countries with a practice change	Comments	International Guidance WHO or RCOG/RCM	Countries aligned with International guidance
Companionship	32/32 All countries had some limitations on antenatal visits, attending ultrasound, companionship for birth and postnatally	Cyprus —‘we have good protocols that align with WHO—but no one follows them’ Malta —‘partners are only allowed in (to birthing suite) during established labour and must have had a swab in the last 48 hours’ Estonia —‘no visitors are permitted in the postpartum period’	All women have the right to a safe and positive childbirth experience, whether or not they have C-19- this includes ‘a companion of choice’ (WHO)	Iceland was most closely aligned with WHO guidance—the significant practice change was the exclusion of the partner from the ultrasound examination to protect staff with specialised skills 31/32 countries had an increase in virtual visits and women attending face to face appointments unaccompanied
Breastfeeding	Difficult to ascertain	Cyprus —‘parents need to wear masks but breastfeeding is supported if the baby is positive he/she stays with mother, if the baby is negative they are separated and the midwives feed the baby with breast milk/formula’	Women and their families should be informed that infection with COVID-19 is not a contraindication to breastfeeding. (RCOG/RCM)	It seems that support for breastfeeding is present, but how this is done varies at a local level
Mother/baby separation after birth	Difficult to ascertain	Slovakia —‘mothers are tested for C-19 before birth, mothers and infants are separated after birth’ Czech Republic —‘mothers and babies are separated if mother is C-19 positive, depends on whether the unit can offer a separate quarantine room or not’	Women and their healthy babies should remain together in the immediate postpartum period, if they do not otherwise require maternal critical care or neonatal care. (RCOG/RCM)	It seems all countries are offering skin-to skin contact after birth, and separation is based on the need for maternal or neonatal care
Visiting the postpartum period	30/32	Sweden —‘partners are not allowed to accompany women to the postnatal wards’ Ireland —‘first wave no partners could visit NICU mothers are permitted to visit 15 mins a day’ Poland —‘often parents of premature babies cannot visit them for weeks’	National guidance applies in terms of the models of care in place	Where access to postnatal wards is restricted it seems early discharge home was seen as an alternative for most countries. It seems when the baby has to receive neonatal care visiting restrictions can be particularly severe and prolonged
Mask-wearing in labour	3/32	Luxembourg —‘every woman has to wear a mask during labour even in the second stage’ Estonia —‘the companion must wear a mask at all times (in delivery suite)’ Malta —‘mothers need to have a swab within last 48 hours, otherwise treated	Guidance refers to national policies on wearing face masks. However, overall the guidance is towards the wearing of appropriate PPE by staff and once the woman is in an isolation room she can remove her mask	

Table 1. (Continued)

Key practice change	No. of countries with a practice change	Comments	International Guidance WHO or RCOG/RCM	Countries aligned with International guidance
Impact on interventions	32/32	as C-19 positive and have to labour with a mask' Turkey —'demands for caesarean section increased. More women arrived in active labour and the number of out-of-hospital births increased'		All countries experienced practice changes from antenatal visits, to rules around companionship and access to mother and baby in the postnatal period or to baby in the NICU

Thirty-two countries* were represented and contributed to the chat on practice changes in their country in maternity care in wave 1 of the C-19 pandemic. These findings are not definitive. In addition, the variation between local and regional practices within countries is so wide that ascertaining a definitive description of the changes is not feasible.

*Albania, Austria, Belgium, Bosnia Herzegovina, Croatia, Cyprus, Czech Republic, Denmark, Estonia, France, Germany, Greece, Iceland, Ireland, Israel, Italy, Lithuania, Luxembourg, Malta, Netherlands, North Macedonia, Norway, Poland, Romania, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey, United Kingdom.

mental health teams were refused or delayed because of restrictions related to COVID-19.

Furthermore, the restrictions may lead maternity staff to engage in clinical practices in direct contravention with evidence, professional recommendations or deeply held ethical or moral beliefs and values, as services attempt to control the risk of Covid-19 infection. These changes in clinical practice may result in increasing levels of occupational moral injury, making staff more vulnerable to mental health problems. This may lead to reduced working hours and increased turnover, and may adversely impact service user care.⁸

Getting the balance right

Given the scale and scope of the restrictions that have been imposed across maternity care facilities, it is important now more than ever to ensure that authoritative guidelines are evidence-based and that restrictions in practice are appropriately aligned to evidence-based policy recommendations. To enable this to happen, 'new' approaches to care during a pandemic crisis must be delivered within a quality framework, founded on evidence and analysis of the potential unintended consequences. The current guidance from the WHO^{3,9} continues to emphasise that quality care includes ensuring a woman's right to a safe and positive childbirth experience. When practice changes deviate from evidence-based frameworks that underpin quality care they *must* be monitored, appraised and evaluated to minimise unintended iatrogenic effects.

The COVID-19 pandemic continues with new variants of the virus, resulting in increasing infection rates and

hospital admissions. However, as more evidence has emerged relating to COVID-19 and pregnancy and newborn care, evidence-based principles to ensure equitable, safe, effective, quality maternal and newborn care in a pandemic have been developed by a group of midwifery professors in the UK. These clearly outline that care providers must:

- 1 Continue to provide evidence-based, equitable, safe, compassionate and respectful care for physical and mental health, wherever and whenever care takes place, by remote access if necessary
- 2 Protect the human rights of women and newborn infants, as far as possible
- 3 Ensure strict hygiene measures and social distancing when possible
- 4 Follow national guidance on use of personal protective equipment (PPE)
- 5 Ensure birth companionship
- 6 Prevent unnecessary interventions
- 7 Do not separate a woman from her newborn infant(s) unless absolutely necessary
- 8 Promote and support breastfeeding
- 9 Protect and support staff, including their mental health needs [5, p. 5]

Why getting it right is particularly important for maternity care

Unlike trauma during other life periods, the perinatal period is particularly crucial, as it affects not only the mothers but also their neonates, birth companions and families. Some events during pregnancy, labour, birth and the early

life period appear to have exaggerated lifelong consequences. There is now strong evidence that short, highly stressful exposures that last for weeks are enough to set some individuals on such a negative trajectory and emerging evidence that the COVID-19 pandemic has increased significantly levels of maternal stress for some women during late pregnancy and the immediate postpartum period in a manner reminiscent to the 1998 Quebec Ice Storm; 20 years later, children exposed either in the immediate antenatal period, through chaotic intrapartum maternity care or immediately postpartum, had altered metabolic parameters (body mass index [BMI], insulin resistance) and increased HPA axis reactivity (indicator of increased levels of stress).¹⁰ Furthermore, the mother–infant bond is established in the immediate postpartum period, and any negative psychological or psychosocial event may alter this bond, as well as early interactions and parenting.^{11,12} Evidence is growing that maternal perinatal stress has thus long-term impacts on aspects of child development and health. The importance of this perinatal period for the lifelong health of the infant has been highlighted in a recent retrospective study:¹³ adults aged between 47 and 83 that were breastfed as children had a 12% lower chance of contracting COVID, whereas those exposed to maternal smoking around birth had a 20% higher risk of infection and 24% higher risk of hospitalisation due to COVID-19 after adjustment for later-life socio-economic and environmental factors.¹⁴

Extrapolation of these data to the current maternity care situation suggests that the actions taken to reduce risks due to COVID-19 may negatively impact maternal psychosocial functioning, early parenting and, consequently, child developmental outcomes. It is thus important to document these deviations from best practice, and to reverse them as soon as possible.

Conclusion

Across Europe, commentators on the current pandemic have noted the critical need for health and social care providers to balance reduction of infection risk and loss of life with maintaining compassionate human relationships. The concerns within maternity care echo those in other areas. The difference in maternity care is the potential of ‘just in case’ interventions to have long-term, and even life-course, impacts on mother, baby and the wider family. Variation in maternity care policy or guidelines for practice at a country, regional or facility level cannot be justified. Variation in particular practices for particular women and pregnant people may be justified, but only in relation to their specific values, and clinical and psychological needs. It has been notable that variance from the evidence has disproportionately restricted human contact between pregnant and childbearing women and professionals, partners and

neonates (limiting social, emotional and informational support) and increased unnecessary or unwanted intervention (risking high levels of adverse psychological, physical and/or emotional consequences). This raises serious questions about an underlying ethos of maternity care provision and how it should be reframed when services are rebuilt, once the pandemic is finally over.

Disclosures of interests

None declared. Completed disclosure of interest forms are available to view online as supporting information.

Contribution of authorship

JL and AH are Joint Senior Authors. All other authors contributed to planning and writing equally.

Details of ethics approval

None required.

Funding

Online Open publication funded by COST Action 18211.

Acknowledgements

All authors are members of the management committee of COST Action CA18211 and JL is the Action Chair. This commentary is based on contributions from members of the network. Open access funding provided by IReL.

Data availability

The data that support the findings of this study are available from the corresponding author upon reasonable request. ■

References

- 1 Khosla R, Zampas C, Vogel JP, Bohren MA, Roseman M, Erdman JN. International human rights and the mistreatment of women during childbirth. *Health Hum Rights* 2016;18:131–43.
- 2 Topalidou A, Thomson G, Downe S. COVID-19 and maternal and infant health: are we getting the balance right? A rapid scoping review. *Pract Midwife* 2020;23:36–45.
- 3 Morano S, Calleja-Agius J. Giving birth and dying alone in hospital during the COVID-19 pandemic—a time for shifting paradigm towards continuity of care. *J Perinat Med* 2020;48:551–2.
- 4 WHO. *Corona Virus Disease during Pregnancy and Childbirth*. Geneva: WHO; 2020 [https://www.who.int/news-room/q-a-detail/coronavirus-disease-covid-19-pregnancy-and-childbirth]. Accessed 1 February 2021.
- 5 RCOG guidelines. Coronavirus (COVID-19), pregnancy and women's health. 2020 [https://www.rcog.org.uk/en/guidelines-research-service/s/coronavirus-covid-19-pregnancy-and-womens-health/]. Accessed 26 February 2021.
- 6 RCM. Guidance for provision of midwife-led settings and home birth in the evolving coronavirus (COVID-19) pandemic. 2020. [https://www.rcm.org.uk/media/3893/2020-04-17-guidance-for-provision-of-midwife-led-settings.pdf]. Accessed 5 January 2021.

- 7 Chmielewska B, Barratt I, Townsend R, et al. Effects of the COVID-19 pandemic on maternal and perinatal outcomes: a systematic review and meta-analysis. *Lancet Global Health* 2021;9:e759–72. <https://www.sciencedirect.com/science/article/pii/S2214109X21000796>. Accessed 1 February 2021.
- 8 Yan H, Ding Y, Guo W. Mental health of pregnant and postpartum women during the coronavirus disease 2019 pandemic: a systematic review and meta-analysis. *Front Psychol* 2020;11:617001.
- 9 MBRRACE-UK. *Saving Lives, Improving Mothers' Care Rapid report: Learning from SARS-CoV-2-related and Associated Maternal Deaths in the UK*. MBRRACE-UK; March–May 2020 [<https://www.npeu.ox.ac.uk/mbrrace-uk/reports>].
- 10 Horsch A, Lalor J, Downe S. Moral and mental health challenges faced by maternity staff during the COVID-19 pandemic. *Psychol Trauma Theory Res Pract Policy* 2020;12:S141–2.
- 11 WHO. *Recommendations: Intrapartum Care for a Positive Birth Experience*. Geneva: WHO; 2018 [<https://apps.who.int/iris/bitstream/handle/10665/272447/WHO-RHR-18.12-eng.pdf?ua=1>]. Accessed 26 February 2021.
- 12 Cook N, Ayers S, Horsch A. Maternal posttraumatic stress disorder during the perinatal period and child outcomes: a systematic review. *J Affect Disord* 2018;225:18–31.
- 13 Veru F, Laplante DP, Luheshi G, King S. Prenatal maternal stress exposure and immune function in the offspring. *Stress* 2014;17:133–48.
- 14 Didikoglu A, Maharani A, Pendleton N, Canal MM, Payton A. Early life factors and COVID-19 infection in England: a prospective analysis of UK Biobank participants. *Early Human Dev* 2021;155:105326.