



City Research Online

City St George's, University of London

Citation: Fring, A. & Tenney, R. (2021). Infinite series of time-dependent Dyson maps. *Journal of Physics A: Mathematical and Theoretical*, 54(48), 485201. doi: 10.1088/1751-8121/ac31a0

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/26948/>

Link to published version: <https://doi.org/10.1088/1751-8121/ac31a0>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Infinite series of time-dependent Dyson maps

Andreas Fring and Rebecca Tenney

*Department of Mathematics, City University London,
Northampton Square, London EC1V 0HB, UK
E-mail: a.fring@city.ac.uk, rebecca.tenney@city.ac.uk*

ABSTRACT: We propose and explore a scheme that leads to an infinite series of time-dependent Dyson maps which associate different Hermitian Hamiltonians to a uniquely specified time-dependent non-Hermitian Hamiltonian. We identify the underlying symmetries responsible for this feature respected by various Lewis-Riesenfeld invariants. The latter are used to facilitate the explicit construction of the Dyson maps and metric operators. As a concrete example for which the scheme is worked out in detail we present a two-dimensional system of oscillators that are coupled to each other in a non-Hermitian \mathcal{PT} -symmetrical fashion.

1. Introduction

It is well-known for almost thirty years [1] that a given non-Hermitian Hamiltonian H does not uniquely define a quantum mechanical description of a physical system, in the sense that it always leads to a fixed set of well-defined operators associated to physical observables. Even when this Hamiltonian has a set of real eigenvalues and a complete set of eigenstates one needs to construct a well-defined positive-definite metric operator ρ in order to set up a physical Hilbert space. As dictated by the fundamental axioms of quantum mechanics physical observables are then represented by self-adjoint operators acting in this Hilbert space [2–4]. However, the metric ρ is not uniquely determined when only fixing one of the operators associated to observables, e.g. the Hamiltonian H , but as was argued in [1], one needs to specify at least one further observable to render the metric unique. In [5] it was shown how for time-independent Hamiltonians the ambiguities in the metric may be related to certain symmetries and for many known models explicit solutions, including some of their ambiguities, have been constructed [5–9]. Technically one either solves the time-independent quasi-Hermiticity relation $H^\dagger \rho = \rho H$ for the metric ρ , or uses the fact that the metric factorises into a product of the Dyson map η [10] and its conjugate, $\rho = \eta^\dagger \eta$ and solves instead the time-independent Dyson equation $\eta H \eta^{-1} = h$ for η .

This pseudo-Hermitian approach has also been extended successfully to non-autonomous systems, that is non-Hermitian explicitly time-dependent Hamiltonian systems [11–16]. In this setting also the metric and the Dyson map become explicitly time-dependent, but the ambiguities persist. In fact, the situation worsens as even the Hamiltonian ceases to be related to a physically observable quantity [16]. In the time-independent case it was one of the two natural candidates for operators associated to physical observables that when specified render the metric unique, whereas in the time-dependent case the energy operator has to be defined differently.

Here our main purpose is to identify the symmetries responsible for associating to a uniquely given non-Hermitian Hamiltonian several equivalent Hermitian Hamiltonians corresponding to different types of physical systems. In particular, we propose and explore a scheme that leads to an infinite series of Dyson maps, hence an infinite series of equivalent Hermitian Hamiltonians, albeit different physics as we shall demonstrate. We employ the Lewis-Riesenfeld method of invariants [17] to facilitate the explicit construction of the Dyson maps. We explore the working of our proposal for the concrete example of two-dimensional \mathcal{PT} -symmetrically coupled harmonic oscillators.

Our manuscript is organised as follows: In section 2 we explain the general scheme of how a series of infinite Dyson maps can be constructed iteratively from two known solutions to the time-dependent Dyson equation. Section 3 is devoted to setting up the discussion of this scheme for the concrete example of two \mathcal{PT} -symmetrically coupled oscillators. We point out a previously overlooked relationship between different types of the underlying auxiliary equations and construct various Lewis-Riesenfeld invariants that are crucial in the understanding of the underlying symmetries as well as being a vital aide in the construction of the time-dependent Dyson maps. In section 4 we construct the infinite series for three different purposefully selected sets of two seed Dyson maps, exploring also the limitations of the scheme. Our conclusions and an outlook into open problems are stated in section 5.

2. Infinite symmetries and series of Dyson maps from two seeds

Our starting point is an explicitly non-Hermitian time-dependent Hamiltonian $H \neq H^\dagger$ satisfying the time-dependent Schrödinger equation (TDSE) $H(x, t)\psi(x, t) = i\hbar\partial_t\psi(x, t)$. We further assume that we have two different time-dependent Dyson maps, $\eta(t)$ and $\tilde{\eta}(t)$, satisfying the time-dependent Dyson equations (TDDE)

$$h = \eta H \eta^{-1} + i\hbar\partial_t\eta\eta^{-1}, \quad \text{and} \quad \tilde{h} = \tilde{\eta} H \tilde{\eta}^{-1} + i\hbar\partial_t\tilde{\eta}\tilde{\eta}^{-1}, \quad (2.1)$$

involving two different time-dependent Hermitian Hamiltonians $h = h^\dagger$, $\tilde{h} = \tilde{h}^\dagger$ that also obey their respective TDSEs $h(x, t)\phi(x, t) = i\hbar\partial_t\phi(x, t)$ and $\tilde{h}(x, t)\tilde{\phi}(x, t) = i\hbar\partial_t\tilde{\phi}(x, t)$. The wavefunctions are related as $\phi = \eta\psi$, $\tilde{\phi} = \tilde{\eta}\psi$ and therefore $\tilde{\phi} = A\phi$, where we employed the first of the operators

$$A := \tilde{\eta}\eta^{-1} \quad \text{and} \quad \tilde{A} := \eta^{-1}\tilde{\eta}. \quad (2.2)$$

The operator \tilde{A} is defined for later purposes. Next we eliminate the Hamiltonian H from the two equations in (2.1), such that the two Hermitian Hamiltonians are seen to be related

as

$$\tilde{h} = AhA^{-1} + i\hbar\partial_t AA^{-1}. \quad (2.3)$$

As argued and shown for concrete examples in [18–22], once the Dyson maps are known one may relate the respective Lewis-Riesenfeld invariants $I_{\mathcal{H}}$, with $\mathcal{H} = H, h, \tilde{h}$, satisfying [17]

$$i\hbar\frac{dI_{\mathcal{H}}}{dt} = i\hbar\partial_t I_{\mathcal{H}} + [I_{\mathcal{H}}, \mathcal{H}] = 0, \quad (2.4)$$

simply by means of similarity transformations as

$$I_h = \eta I_H \eta^{-1}, \quad I_{\tilde{h}} = \tilde{\eta} I_H \tilde{\eta}^{-1}, \quad \Rightarrow I_{\tilde{h}} = A I_h A^{-1}. \quad (2.5)$$

Each of the invariants satisfies an eigenvalue equation with time-independent eigenvalues and eigenfunctions that are simply related by a phase factor to the wavefunctions satisfying the respective TDSE. Exploiting the Hermiticity of the invariants I_h and $I_{\tilde{h}}$, the latter relation in (2.5) imply that the operators

$$S := A^\dagger A \quad \text{and} \quad \tilde{S} := A A^\dagger \quad (2.6)$$

are symmetries for the invariants I_h and $I_{\tilde{h}}$, respectively, with

$$[I_h, S] = 0 \quad \text{and} \quad [I_{\tilde{h}}, \tilde{S}] = 0. \quad (2.7)$$

Thus S and \tilde{S} also satisfy the Lewis-Riesenfeld equations for the Hermitian h -Hamiltonian system and the \tilde{h} -Hamiltonian system

$$i\hbar\frac{dS}{dt} = i\hbar\partial_t S + [S, h] = 0, \quad i\hbar\frac{d\tilde{S}}{dt} = i\hbar\partial_t \tilde{S} + [\tilde{S}, \tilde{h}] = 0. \quad (2.8)$$

In turn this means that

$$I'_h = I_h + S, \quad \text{and} \quad I'_{\tilde{h}} = I_{\tilde{h}} + \tilde{S} \quad (2.9)$$

are new invariants for the Hamiltonians h and \tilde{h} , respectively.

Another symmetry with an interesting consequence is the possibility of an \tilde{A} -symmetry, see (2.2), of the non-Hermitian invariant I_H , as it implies that the two invariants related to the Hermitian systems are identical

$$[I_H, \tilde{A}] = 0 \quad \Leftrightarrow \quad I_h = I_{\tilde{h}}, \quad (2.10)$$

and in turn, the equality of two invariants associated to different Hermitian Hamiltonians implies an \tilde{A} -symmetry of the non-Hermitian invariant I_H . This is easily established by making use of the pseudo-Hermiticity relations for the invariants (2.5).

2.1 Iteration of two Dyson maps

While certain symmetries of the invariants imply the presence of two inequivalent Dyson maps and vice versa, we will now construct further time-dependent Dyson maps, say $\check{\eta}$ or $\hat{\eta}$, from two given ones, say η and $\tilde{\eta}$. We start by constructing a third Dyson map making use of either of the following statements:

- (S1) If and only if the adjoint action of A on the invariant $I_{\check{h}}$, $AI_{\check{h}}A^{-1}$, is Hermitian then $\check{h} = A\tilde{h}A^{-1} + i\hbar\partial_t AA^{-1}$ is a new Hamiltonian that is related to the non-Hermitian Hamiltonian H by the time-dependent Dyson equation $\check{h} = \check{\eta}H\check{\eta}^{-1} + i\hbar\partial_t\check{\eta}\check{\eta}^{-1}$ with $\check{\eta} := \tilde{\eta}\eta^{-1}\tilde{\eta}$.
- (S2) If and only if the inverse adjoint action of A on the invariant I_h , $A^{-1}I_hA$ is Hermitian then $\hat{h} = A^{-1}hA - i\hbar A^{-1}\partial_t A$ is a new Hamiltonian that is related to the non-Hermitian Hamiltonian H by the time-dependent Dyson equation $\hat{h} = \hat{\eta}H\hat{\eta}^{-1} + i\hbar\partial_t\hat{\eta}\hat{\eta}^{-1}$ with $\hat{\eta} := \eta\tilde{\eta}^{-1}\eta$.

At first we prove (S1) in reverse: Assuming that $\check{\eta} := \tilde{\eta}\eta^{-1}\tilde{\eta}$ is a new time-dependent Dyson map that maps the non-Hermitian Hamiltonian H to a Hermitian one, the TDDE $\check{h} = \check{\eta}H\check{\eta}^{-1} + i\hbar\partial_t\check{\eta}\check{\eta}^{-1}$ holds by definition. Replacing now H in this equation by means of the first equation in (2.1) and using the definition (2.2) for A , equation $\check{h} = A\tilde{h}A^{-1} + i\hbar\partial_t AA^{-1}$ follows directly. In turn this implies that the adjoint action of A on $I_{\check{h}}$ yields the Lewis-Riesenfeld invariant $I_{\check{h}}$. Since $I_{\check{h}}$ is Hermitian, so is $AI_{\check{h}}A^{-1}$. The direct statement is shown by checking whether $AI_{\check{h}}A^{-1}$ is Hermitian and then reversing the steps in the previous argument. Similarly we may prove (S2).

Thus for practical purposes when given the two time-dependent Dyson maps η , $\tilde{\eta}$ and the invariants $I_{\check{h}}$, I_h we can simply check whether $AI_{\check{h}}A^{-1}$ and/or $A^{-1}I_hA$ are Hermitian and subsequently deduce the form of the new Dyson map. Alternatively one may of course also assume the given forms for $\check{\eta}$ and $\hat{\eta}$ with a subsequent check of whether the right hand sides of the corresponding Dyson equations are Hermitian, thus defining new Hermitian Hamiltonians.

Having now obtained two new time-dependent Dyson maps, we may include them into the set of the two starting Dyson maps to construct yet more and more maps by iteration. We summarize the first step as outlines above, i.e. the construction of $\check{\eta} =: \eta_3$ and $\hat{\eta} =: \eta_4$ from the seed maps η and $\tilde{\eta}$, as

$$\begin{array}{l} \nearrow \\ \eta, \tilde{\eta} \\ \searrow \end{array} \begin{array}{l} \eta_3 = \tilde{\eta}\eta^{-1}\tilde{\eta} = A\tilde{\eta} \\ \eta_4 = \eta\tilde{\eta}^{-1}\eta = A^{-1}\eta \end{array} . \quad (2.11)$$

Replacing now in the next step the seed maps by new maps obtained in the previous step

we obtain, up to the Hermiticity check,

$$\begin{array}{l} \nearrow \eta_5 = \tilde{\eta}\eta^{-1}\tilde{\eta}\eta^{-1}\tilde{\eta}\eta^{-1}\tilde{\eta} = A^3\tilde{\eta} \\ \eta, \eta_3 \searrow \eta_6 = \eta\tilde{\eta}^{-1}\eta\tilde{\eta}^{-1}\eta = A^{-2}\eta \end{array}, \quad (2.12)$$

$$\begin{array}{l} \nearrow \eta_6 = \eta\tilde{\eta}^{-1}\eta\eta^{-1}\eta\tilde{\eta}^{-1}\eta = \eta\tilde{\eta}^{-1}\eta\tilde{\eta}^{-1}\eta \\ \eta, \eta_4 \searrow \tilde{\eta} = \eta\eta^{-1}\tilde{\eta}\eta^{-1}\eta \end{array}, \quad (2.13)$$

$$\begin{array}{l} \nearrow \eta = \tilde{\eta}\tilde{\eta}^{-1}\eta\tilde{\eta}^{-1}\tilde{\eta} \\ \tilde{\eta}, \eta_3 \searrow \eta_7 = \tilde{\eta}\eta^{-1}\tilde{\eta}\tilde{\eta}^{-1}\tilde{\eta}\eta^{-1}\tilde{\eta} = \tilde{\eta}\eta^{-1}\tilde{\eta}\eta^{-1}\tilde{\eta} = A^2\tilde{\eta} \end{array}, \quad (2.14)$$

$$\begin{array}{l} \nearrow \eta_7 = \tilde{\eta}\eta^{-1}\tilde{\eta}\eta^{-1}\tilde{\eta} \\ \tilde{\eta}, \eta_4 \searrow \eta_8 = \eta\tilde{\eta}^{-1}\eta\tilde{\eta}^{-1}\eta\tilde{\eta}^{-1}\eta = A^{-3}\eta \end{array}, \quad (2.15)$$

$$\begin{array}{l} \nearrow \eta_9 = \eta\tilde{\eta}^{-1}\eta\tilde{\eta}^{-1}\eta\tilde{\eta}^{-1}\eta\tilde{\eta}^{-1}\eta = A^{-4}\eta \\ \eta_3, \eta_4 \searrow \eta_{10} = \tilde{\eta}\eta^{-1}\tilde{\eta}\eta^{-1}\tilde{\eta}\eta^{-1}\tilde{\eta}\eta^{-1}\tilde{\eta} = A^4\tilde{\eta} \end{array}. \quad (2.16)$$

Continuing in this manner we obtain a series of Dyson maps of the general form

$$\eta^{(n)} := A^n\eta, \quad \tilde{\eta}^{(n)} := A^n\tilde{\eta}, \quad \text{with } n \in \mathbb{Z}. \quad (2.17)$$

When combined in the way described above we only obtain new maps of the same form

$$\begin{array}{l} \nearrow \tilde{\eta}^{(2m-n)} \\ \tilde{\eta}^{(n)}, \tilde{\eta}^{(m)} \searrow \tilde{\eta}^{(2n-m)} \end{array}, \quad \begin{array}{l} \nearrow \eta^{(2m-n-1)} \\ \tilde{\eta}^{(n)}, \eta^{(m)} \searrow \tilde{\eta}^{(2n-m+1)} \end{array}, \quad (2.18)$$

$$\begin{array}{l} \nearrow \tilde{\eta}^{(2m-n+1)} \\ \eta^{(n)}, \tilde{\eta}^{(m)} \searrow \eta^{(2n-m-1)} \end{array}, \quad \begin{array}{l} \nearrow \eta^{(2m-n)} \\ \eta^{(n)}, \eta^{(m)} \searrow \eta^{(2n-m)} \end{array}. \quad (2.19)$$

As discussed above for the iteration to proceed we need to verify at each step the Hermiticity of the right hand side of the time-dependent Dyson equation or the adjointly mapped invariants. Thus we require the relevant A -operators involving the new maps

$$\tilde{\eta}^{(m)} \left(\tilde{\eta}^{(n)} \right)^{-1} = A^{m-n}, \quad \eta^{(m)} \left(\tilde{\eta}^{(n)} \right)^{-1} = A^{m-n-1}, \quad (2.20)$$

$$\tilde{\eta}^{(m)} \left(\eta^{(n)} \right)^{-1} = A^{m-n+1}, \quad \eta^{(m)} \left(\eta^{(n)} \right)^{-1} = A^{m-n}. \quad (2.21)$$

Naturally we may repeat the symmetry arguments from the previous section using the newly constructed Dyson maps, thus obtaining an infinite set of symmetry operators, provided that the Hermiticity property holds at each of the iterative steps.

3. Two dimensional \mathcal{PT} -symmetrically coupled oscillators

3.1 Six seed Dyson maps

To demonstrate the working of the above scheme we shall be considering two-dimensional time-dependent oscillators with a \mathcal{PT} -symmetric non-Hermitian coupling in space and momenta. This system has been studied extensively recently [20, 23] and is an ideal testing ground for the construction of new Dyson maps as outlined above since six unique solutions have already been identified, that we can build on. The system we will examine is described by the Hamiltonian

$$H(t) = \frac{a(t)}{2} (p_x^2 + p_y^2 + x^2 + y^2) + i \frac{\lambda(t)}{2} (xy + p_x p_y), \quad (3.1)$$

where the time-dependent coefficient functions are taken to be real, $a(t), \lambda(t) \in \mathbb{R}$. This Hamiltonian is symmetric with regards to two realisations of partial \mathcal{PT} -symmetries, i.e. $[\mathcal{PT}_\pm, H] = 0$, where the anti-linear maps are given by $\mathcal{PT}_\pm : x \rightarrow \pm x, y \rightarrow \mp y, p_x \rightarrow \mp p_x, p_y \rightarrow \pm p_y, i \rightarrow -i$. It is convenient to express this Hamiltonian in terms of elements of a Lie algebra

$$H(t) = a(t) (K_1 + K_2) + i\lambda(t)K_3, \quad (3.2)$$

where the Lie algebraic generators are given by

$$K_1 = \frac{1}{2} (p_x^2 + x^2), \quad K_2 = \frac{1}{2} (p_y^2 + y^2), \quad K_3 = \frac{1}{2} (xy + p_x p_y), \quad K_4 = \frac{1}{2} (xp_y - yp_x), \quad (3.3)$$

satisfying the commutation relations

$$\begin{aligned} [K_1, K_2] &= 0, & [K_1, K_3] &= iK_4, & [K_1, K_4] &= -iK_3, \\ [K_2, K_3] &= -iK_4, & [K_2, K_4] &= iK_3, & [K_3, K_4] &= i(K_1 - K_2)/2. \end{aligned} \quad (3.4)$$

The six unique Dyson maps which were constructed using a perturbative approach [23] map the non-Hermitian Hamiltonian (3.2) by means of (2.1) into a general Hermitian Hamiltonian of the form

$$h(t) = f_+(t)K_1 + f_-(t)K_2, \quad (3.5)$$

where the time-dependent coefficient functions $f_\pm(t)$ are specific for each of the maps. The generic form for all of the Dyson maps consists of two factors given by

$$\eta(t) = \exp[\gamma_1(t)q_1] \exp[\gamma_2(t)q_2], \quad (3.6)$$

where the q_1, q_2 are operators taken to be generators of the algebra (3.4).

The time-dependent functions $\gamma_1(t), \gamma_2(t)$ are constrained in each case by two coupled first order differential equations. When suitably parametrising the γ_i , these equation were solved up to some auxiliary equations that may also be solved for a given set of initial conditions. In table 1 we report for a selected set of six Dyson maps, η_i , the corresponding parametrisations for the γ_i , the two operators in the factorisation (3.6) and the resulting time-dependent coefficient functions, $f_\pm(t)$ in (3.5), occurring in the Hermitian counter parts.

η_i	q_1, q_2	γ_1	γ_2	$f_{\pm}(t)$	constraint
η_1	K_4, K_3	*	*	a	*
η_2	K_3, K_4	$\operatorname{arccosh}(\chi)$	$\operatorname{arcsinh}\left(\frac{k}{\chi}\right)$	$a \pm \frac{k\lambda}{2\chi^2}$	$\chi > 1$
η_3	K_4, iK_1	$\operatorname{arcsinh}\left(k_3\sqrt{1+x^2}\right)$	$-\arctan(x)$	$a - \frac{\lambda(\pm 1 + \sqrt{1+(1+x^2)k_3^2})}{2(1+x^2)k_3}$	*
η_4	K_4, iK_2	$\operatorname{arcsinh}\left(k_4\sqrt{1+x^2}\right)$	$\arctan(x)$	$a + \frac{\lambda(\mp 1 + \sqrt{1+(1+x^2)k_4^2})}{2(1+x^2)k_4}$	*
η_5	K_3, iK_1	$\operatorname{arcsinh}\left(k_5\sqrt{1+x^2}\right)$	$-\operatorname{arccot}(x)$	$a + \frac{\lambda(\pm 1 + \sqrt{1+(1+x^2)k_5^2})}{2(1+x^2)k_5}$	*
η_6	K_3, iK_2	$\operatorname{arcsinh}\left(k_6\sqrt{1+x^2}\right)$	$\operatorname{arccot}(x)$	$a - \frac{\lambda(\mp 1 + \sqrt{1+(1+x^2)k_6^2})}{2(1+x^2)k_6}$	*

Table 1: Inequivalent Dyson maps η_i with specific operators q_1, q_2 in the factorisation (3.6), and parametrisations for γ_1, γ_2 in terms of the auxiliary functions χ or x_i together with the time-dependent functions $f_{\pm}(t)$ in $h(t)$.

The auxiliary equations governing the time-dependence were found to be

$$\text{Aux}_1 : \quad \ddot{x}_i - \frac{\dot{\lambda}}{\lambda} \dot{x}_i - \lambda^2 x_i = 0 \quad \text{and} \quad \text{Aux}_2 : \quad \ddot{\chi} - \frac{\dot{\lambda}}{\lambda} \dot{\chi} - \lambda^2 \chi = k^2 \frac{\lambda^2}{\chi^3}, \quad (3.7)$$

where $i = 3, 4, 5, 6$. The second equation in (3.7) is the ubiquitous Ermakov-Pinney equation [24, 25].

3.2 Relation between auxiliary equations

To carry out the discussion as set out in section 2 we have the somewhat unappealing feature that various seed Dyson maps are governed by different types of auxiliary equations. Here we comment briefly on a feature previously not realised, and show that with a different parametrisation also η_2 is in fact constrained by the linear second order equation in (3.7). To demonstrate that this can be achieved we briefly recall how to solve the TDDE for η_2 , but with a different parametrisation.

Assuming for this purpose the Dyson map η_2 to be of the form (3.6) with $\gamma_1(t), \gamma_2(t)$ unknown and $q_1 = K_3, q_2 = K_4$, we substitute η_2 into the TDDE and find that $h_2(t)$ is indeed made to be Hermitian if the following coupled first order differential equations are satisfied

$$\dot{\gamma}_1 = -\lambda \cosh \gamma_2 \quad \text{and} \quad \dot{\gamma}_2 = \lambda \tanh \gamma_1 \sinh \gamma_2. \quad (3.8)$$

To solve these equations for γ_1 and γ_2 we notice first that we can eliminate λ and dt from the above equations to give

$$\gamma_2'(\gamma_1) = -\tanh \gamma_1 \tanh \gamma_2(\gamma_1), \quad (3.9)$$

which we solve to

$$\gamma_2 = \operatorname{arcsinh}(c \operatorname{sech} \gamma_1), \quad (3.10)$$

with c being an integration constant. Previously, [23], we parametrised $\gamma_1 = \operatorname{arccosh} \chi$ which lead to the Ermakov-Pinney equation Aux_2 as auxiliary equation. When instead we

define $\gamma_1 = \operatorname{arccosh} \sqrt{1+x_2^2}$ and let $c = -1/k_2$ we find that the central equation to be satisfied is now also Aux_1 , similarly as for the other cases. We have now a new way of writing the Dyson map η_2 so that all of the maps found are governed by the same central equation, with

$$\eta_2 : \quad \gamma_1 = \operatorname{arccosh} \sqrt{1+x_2^2}, \quad \gamma_2 = \operatorname{arcsinh} \left(-\frac{1}{k_2 \sqrt{1+x_2^2}} \right), \quad f_{\pm} = \mp \frac{\lambda}{2k_2(1+x_2^2)}. \quad (3.11)$$

Thus we have found a way to convert the nonlinear dissipative Ermakov-Pinney equation given by Aux_2 to the linear second order differential equation Aux_1 by the relation

$$\chi = \sqrt{1+x_2^2} \quad \text{with} \quad k = -\frac{1}{k_i}. \quad (3.12)$$

This appears to be somewhat miraculous, but one needs to stress here that this is only possible when employing also the first order equations resulting from (3.8) for the respective variables, i.e.

$$\dot{x}_2 = -\frac{\lambda \sqrt{1+k_2^2(1+x_2^2)}}{k_2}, \quad (3.13)$$

for the new parametrisation. Notice also that the constraint imposed on χ as reported in table 1 is automatically satisfied with the new parametrisation.

3.3 Construction of invariants

As outlined in section 2, in order to construct new Dyson maps we must first calculate invariants for each of the Hermitian Hamiltonians associated to each of the seed Dyson maps $\eta_i, i = 2, \dots, 6$. Using the corresponding expressions for h_i we solve equation (2.4) and find the invariant

$$I_{h_i}(t) = c_1 K_1 + c_2 K_2 + c_3 \cos \left[c_4 - \int^t f_{+-}^i(s) ds \right] K_3 - c_3 \sin \left[c_4 - \int^t f_{+-}^i(s) ds \right] K_4 \quad (3.14)$$

where $f_{+-}^i := f_+^i - f_-^i$ is the difference between the time-dependent functions in (3.5) occurring in the Hermitian Hamiltonian, and the c_1, c_2, c_3, c_4 are real constants. Notice that in all cases the difference takes on the same form

$$f_{+-}^{2,3,4}(t) = -\frac{\lambda}{k_i(1+x_2^2)} = -f_{+-}^{5,6}(t), \quad (3.15)$$

such that the corresponding Hermitian invariants are identical.

While the cases $i = 2, \dots, 6$ have been unified, the case $i = 1$ still stands out as in this case $f_+ = f_-$, so that the invariant in (3.14) is rendered time-independent. We therefore need to construct an additional invariant for $h_1(t)$. To achieve that we need to enlarge the algebra (3.4) by six additional elements. To allow for a more compact notation we temporarily denote $K_1 \rightarrow K_+^x, K_2 \rightarrow K_+^y, K_3 \rightarrow I_+$ and $K_4 \rightarrow J_-$, so that the ten Hermitian generators can be written as

$$K_{\pm}^z = \frac{1}{2}(p_z^2 \pm z^2), \quad K_0^z = \frac{1}{2}\{z, P_z\}, \quad J_{\pm} = \frac{1}{2}(xp_y \pm yp_x), \quad I_{\pm} = \frac{1}{2}(xy \pm p_x p_y), \quad (3.16)$$

where $z = x, y$. As laid out in [20, 26], we then obtain a closed algebra with non-vanishing commutation relations

$$[K_0^x, K_\pm^x] = 2iK_\mp^x, \quad [K_0^y, K_\pm^y] = 2iK_\mp^y, \quad [K_+^x, K_-^x] = 2iK_0^x, \quad [K_+^y, K_-^y] = 2iK_0^y, \quad (3.17)$$

$$[K_0^x, J_\pm] = -iJ_\mp, \quad [K_0^y, J_\pm] = iJ_\mp, \quad [K_0^x, I_\pm] = -iI_\mp, \quad [K_0^y, I_\pm] = -iI_\mp, \quad (3.18)$$

$$[K_\pm^x, J_+] = \pm iI_\mp, \quad [K_\pm^y, J_+] = \pm iI_\mp, \quad [K_\pm^x, J_-] = \mp iI_\pm, \quad [K_\pm^y, J_-] = \pm iI_\pm, \quad (3.19)$$

$$[K_\pm^x, I_+] = \pm iJ_\mp, \quad [K_\pm^y, I_+] = -iJ_\mp, \quad [K_\pm^x, I_-] = \mp iJ_\pm, \quad [K_\pm^y, I_-] = -iJ_\pm, \quad (3.20)$$

$$[J_+, J_-] = \frac{i}{2}(K_0^x - K_0^y), \quad [I_+, I_-] = -\frac{i}{2}(K_0^x + K_0^y), \quad (3.21)$$

$$[J_+, I_\pm] = \pm \frac{i}{2}(K_\mp^x + K_\mp^y), \quad [J_-, I_\pm] = \pm \frac{i}{2}(K_\mp^x - K_\mp^y). \quad (3.22)$$

The \mathcal{PT} -symmetry that leaves this algebra invariant manifests itself as

$$\mathcal{PT}_\pm: \quad K_0^{x,y} \rightarrow -K_0^{x,y}, \quad K_\pm^{x,y} \rightarrow K_\pm^{x,y}, \quad I_\pm \rightarrow -I_\pm, \quad J_\pm \rightarrow J_\pm, \quad i \rightarrow -i. \quad (3.23)$$

Assuming now that the invariant is also spanned by these generators we found as another solution to (2.4) a universal solution for all six cases

$$I_{h_i}(t) = \alpha_+(K_+^x + K_-^x) + \beta_+(K_+^x - K_-^x) + \alpha_-(K_+^y + K_-^y) + \beta_-(K_+^y - K_-^y) + \delta_+K_0^x + \delta_-K_0^y, \quad (3.24)$$

for $i = 1, \dots, 6$. The time-dependent functions are constrained by

$$\alpha_\pm(t) = \rho_\pm(t)^2, \quad \beta_\pm(t) = \frac{1}{\rho_\pm(t)^2} + \frac{\dot{\rho}_\pm(t)^2}{f_\pm(t)^2}, \quad \delta_\pm(t) = -\frac{2\rho_\pm(t)\dot{\rho}_\pm(t)}{f_\pm(t)}, \quad (3.25)$$

where the auxiliary functions ρ_\pm satisfy the dissipative Ermakov-Pinney equation

$$\ddot{\rho}_\pm - \frac{\dot{f}_\pm}{f_\pm}\dot{\rho}_\pm + f_\pm^2\rho_\pm = \frac{f_\pm^2}{\rho_\pm^3}. \quad (3.26)$$

We will exploit these ambiguities and use which ever invariant is most useful in a certain context. Noting that the invariant in (3.14) is much simpler than the one in (3.24), we shall be using it below for the Hermitian Hamiltonians h_i associated with the Dyson maps η_i , $i = 2, \dots, 6$. In turn we shall use the invariant (3.24) only for the Hermitian Hamiltonian h_1 associated with η_1 , for which it simplifies further due to the relation $f_+ = f_-$ that implies $\rho_+ = \rho_-$.

We also construct the non-Hermitian invariant for the non-Hermitian Hamiltonian H in (3.1) by directly solving equation (2.4). We find

$$I_H = C_1(t)K_1 + C_2(t)K_2 + C_3(t)K_3 + iC_4(t)K_4, \quad (3.27)$$

with

$$C_1 = \frac{c_1}{2} + c_3 \cosh\left(c_4 - \int^t \lambda(s)ds\right), \quad C_2 = \frac{c_1}{2} - c_3 \cosh\left(c_4 - \int^t \lambda(s)ds\right), \quad (3.28)$$

$$C_3 = c_2, \quad C_4 = 2c_3 \sinh\left(c_4 - \int^t \lambda(s)ds\right). \quad (3.29)$$

Using the equations in (2.5) we may relate the various invariants up to the stated ambiguities. We have verified that the inverse adjoint actions of η and $\tilde{\eta}$ on I_H in (3.27) are indeed invariants for I_h and $I_{\tilde{h}}$, respectively, albeit different from the invariants in (3.14) and (3.24) up to the aforementioned ambiguities.

4. Infinite series of Dyson maps from two seeds

For our non-Hermitian Hamiltonian $H(t)$ in (3.1) we have now a number of seed Dyson maps η_i at hand together with their associated Hermitian Hamiltonians h_i and their respective Lewis-Riesenfeld invariants I_{h_i} . Thus we can now carry out the scheme laid out in section 2 and construct an infinite series of Dyson maps from two of these seed maps. We will not present here all thirty possibilities that may result from these six maps, but select various examples that exhibit different types of features including an example for which the mechanism breaks down.

4.1 Seed maps $\eta = \eta_3$ and $\tilde{\eta} = \eta_4$ - unitary operator A

The central operator to compute first is A as defined in (2.2). We start with a simple example for which some of its factors commute. Taking $\eta = \eta_3$, $\tilde{\eta} = \eta_4$ as specified in table 1 and setting $k_3 = k_4 = k$, $x_3 = x_4 = x$ we obtain

$$A = \eta_4 \eta_3^{-1} = e^{\operatorname{arcsinh}(k\sqrt{1+x^2})K_4} e^{i \arctan(x)(K_1+K_2)} e^{-\operatorname{arcsinh}(k\sqrt{1+x^2})K_4} \quad (4.1)$$

$$= e^{i \arctan(x)(K_1+K_2)}. \quad (4.2)$$

The last equality results from the fact that $[K_1 + K_2, K_4] = 0$. According to the statement (S1) in section 2.1 we need to guarantee next that $AI_{h_4}A^{-1}$ is Hermitian. For the case at hand this is easily seen to be the case as A is a unitary operator and I_{h_4} is Hermitian. Thus, according to (S1) a new Dyson map is given by

$$\begin{aligned} \eta^{(1)} &= A\eta_4 = e^{i \arctan(x)(K_1+K_2)} e^{\operatorname{arcsinh}(k\sqrt{1+x^2})K_4} e^{i \arctan(x)K_2} \\ &= e^{\operatorname{arcsinh}(k\sqrt{1+x^2})K_4} e^{i \arctan(x)(K_1+2K_2)}, \end{aligned} \quad (4.3)$$

which in turn is not unitary. Next we compute the associated Hermitian Hamiltonian from the TDDE (2.1) simply by substituting into the right hand side all the known quantities

$$h^{(1)} = \left[a + \frac{\lambda \left(3\sqrt{1+k^2(1+x^2)} - 1 \right)}{2k(1+x^2)} \right] K_1 + \left[a + \frac{\lambda \left(3\sqrt{1+k^2(1+x^2)} + 1 \right)}{2k(1+x^2)} \right] K_2. \quad (4.4)$$

Using next the relation (2.17) it is now straightforward to calculate the infinite series of Dyson maps. At each step the Hermiticity of the adjoint action of the higher order A operators, as defined in (2.20), (2.21), on the Hermitian invariants is guaranteed by the fact that also any power of A is a unitary operator. We find

$$\eta^{(n)} = A^n \eta_4 = e^{\operatorname{arcsinh}(k\sqrt{1+x^2})K_4} e^{i \arctan(x)[K_1+(n+1)K_2]}, \quad (4.5)$$

with corresponding infinite series of Hermitian Hamiltonians

$$h^{(n)} = h^{(1)} + \frac{(n-1)\lambda\sqrt{1+k^2(1+x^2)}}{k(1+x^2)}(K_1 + K_2). \quad (4.6)$$

In a similar fashion we use the second relation in (2.17) to obtain the new Dyson maps

$$\tilde{\eta}^{(n)} = A^n \eta_3 = e^{\operatorname{arcsinh}(k\sqrt{1+x^2})K_4} e^{-i \arctan(x)[(n+1)K_1+K_2]}, \quad (4.7)$$

with corresponding Hermitian Hamiltonians

$$\tilde{h}^{(n)} = \left(\frac{\lambda}{2k(1+x^2)} \right) (K_2 - K_1) + \left[a - \frac{(2n+1)\lambda\sqrt{1+k^2(1+x^2)}}{2k(1+x^2)} \right] (K_1 + K_2) \quad (4.8)$$

Since A is a unitary operator the symmetry operator as defined in (2.6) is simply the unit operator, i.e. $S = S^\dagger = \mathbb{I}$. Moreover the unitarity of A also implies that the relation between the two Hermitian Hamiltonians (2.3) simply becomes a non-Abelian gauge symmetry between two Hermitian Hamiltonians. In this case the metric operators do not change in the iteration process

$$\rho^{(n)} = \eta^{(n)\dagger} \eta^{(n)} = \eta_4^\dagger \eta_4 = \rho_4, \quad \text{and} \quad \tilde{\rho}^{(n)} = \tilde{\eta}^{(n)\dagger} \tilde{\eta}^{(n)} = \eta_3^\dagger \eta_3 = \rho_3. \quad (4.9)$$

4.2 Seed maps $\eta = \eta_2$ and $\tilde{\eta} = \eta_3$ - nonunitary operator A

Once again we start with the construction of the operator A

$$A := \eta_3 \eta_2^{-1} = e^{\operatorname{arcsinh}(k\sqrt{1+x^2})K_4} e^{-i \arctan(x)K_1} e^{\operatorname{arcsinh}\left(\frac{1}{k\sqrt{1+x^2}}\right)K_4} e^{-\operatorname{arccosh}(\sqrt{1+x^2})K_3}, \quad (4.10)$$

where we have used $\eta = \eta_2$, $\tilde{\eta} = \eta_3$ as defined in table 1 and set $k_2 = k_3 = k$, such that $x_2 = x_3 = x$. According to (S1) we need to determine again whether the quantity $AI_{h_3}A^{-1}$ is Hermitian in order to proceed. A lengthy computation can be avoided here by noting that the Hermitian invariants I_h for h_2 and h_3 are identical. Thus we have

$$AI_{h_3}A^{-1} = \eta_3 \eta_2^{-1} I_{h_2} \eta_2 \eta_3^{-1} = \eta_3 I_H \eta_3^{-1} = I_{h_3} = I_{h_3}^\dagger = (AI_{h_3}A^{-1})^\dagger, \quad (4.11)$$

Consequently (S1) is implying that

$$\eta^{(1)} = A\eta_3, \quad (4.12)$$

constitutes a new Dyson map. With the help of the TDDE (2.1) we determine the corresponding Hermitian Hamiltonian to

$$h^{(1)} = \left[a - \frac{\lambda \left(1 + 2\sqrt{1+k^2(1+x^2)} \right)}{2k(1+x^2)} \right] K_1 + \left[a - \frac{\lambda \left(2\sqrt{1+k^2(1+x^2)} - 1 \right)}{2k(1+x^2)} \right] K_2. \quad (4.13)$$

As previously, we use the relation (2.17) to calculate the infinite series of Dyson maps. With

$$\eta^{(n)} = A^n \eta_3, \quad \text{and} \quad \tilde{\eta}^{(n)} = A^{-n} \eta_2, \quad (4.14)$$

we can use relation (4.11) repeatedly to ensure that at each level the adjoint action of the higher order A s on the Hermitian invariants is Hermitian. Using the TDDE (2.1) for the new maps we obtain the Hermitian Hamiltonians

$$h^{(n)} = \left(\frac{\lambda}{2k(1+x^2)} \right) (K_2 - K_1) + \left[a - \frac{(n+1)\lambda\sqrt{1+k^2(1+x^2)}}{2k(1+x^2)} \right] (K_1 + K_2) \quad (4.15)$$

from the first map in (4.14) and

$$h^{(n)} = h^{(n)} = \left(\frac{\lambda}{2k(1+x^2)} \right) (K_2 - K_1) + \left[a - \frac{n\lambda\sqrt{1+k^2(1+x^2)}}{2k(1+x^2)} \right] (K_1 + K_2) \quad (4.16)$$

from the second.

We may now also compute the symmetry operators for I_{h_2} and I_{h_3} . The symmetry operator is readily written down as

$$S := A^\dagger A = e^{-\operatorname{arccosh}(\sqrt{1+x^2})K_3} e^{\operatorname{arcsinh}\left(\frac{1}{k\sqrt{1+x^2}}\right)K_4} e^{-i\arctan(x)K_1} e^{\operatorname{arcsinh}(k\sqrt{1+x^2})K_4} \quad (4.17)$$

$$\times e^{\operatorname{arcsinh}(k\sqrt{1+x^2})K_4} e^{-i\arctan(x)K_1} e^{\operatorname{arcsinh}\left(\frac{1}{k\sqrt{1+x^2}}\right)K_4} e^{-\operatorname{arccosh}(\sqrt{1+x^2})K_3}.$$

Thus we may now explicitly verify the symmetry relation (2.17), best calculated in the form $SI_{h_2}S^{-1} = I_{h_2}$. Similarly, the symmetry operator for I_{h_3} should be given by \tilde{S} , which is indeed the case as we verified explicitly.

4.3 Seed maps $\eta = \eta_1$ and $\tilde{\eta} = \eta_2, \eta_3, \eta_4$ - breakdown of the iteration

From the previous two examples one might get the impression that the iteration procedure can always be carried out with any two seed Dyson maps. However, this is not the case when the Hermiticity condition does not hold. To verify this we relied in the previous section on the fact that the invariants for the two Hermitian Hamiltonians resulting from the seed maps were identical. This is not the case when involving η_1 as a seed map and any of the other five maps, as can be seen from (3.24) when comparing the functions f_\pm . Thus in this case the Hermiticity condition needs to be verified more explicitly

Let us now carry out the calculation for the seed Dyson maps chosen to be $\eta = \eta_1$ and $\tilde{\eta} = \eta_2$. We start from the expression for A

$$A := \eta_2\eta_1^{-1} = e^{\operatorname{arccosh}(\sqrt{1+x_2^2})K_3} e^{-\operatorname{arcsinh}\left(\frac{1}{k_2\sqrt{1+x_2^2}}\right)K_4} e^{\int^s \lambda(s)ds K_3} \quad (4.18)$$

where η_2 is defined as in (3.11) and η_1 as in table 1. Next we compute to quantity $AI_{h_2}A^{-1}$ where I_{h_2} is given by (3.24). After a lengthy calculation we find that this quantity is non-

Hermitian and given by

$$\begin{aligned}
 AI_{h_2}A^{-1} = & \left[\frac{1}{2}\Gamma_+^{++} + \frac{\kappa}{2k_2} \right] K_1 \left[\frac{1}{2}\Gamma_+^{++} - \frac{\kappa}{2k_2} \right] K_2 + i \frac{\cosh(g)}{k_2\Delta\sqrt{1+x_2^2}} \Gamma_-^{+-} K_3 \quad (4.19) \\
 & + i \frac{\kappa(1+x_2^2) + \cosh(g)\Delta}{k_2x_2\sqrt{1+x_2^2}} \Gamma_-^{+-} K_4 \\
 & + i \left\{ \frac{(1+k_2^2)(\delta_- + \delta_+)}{k_2^2x_2} \cosh(g) K_5 - \frac{1}{k_2^2x_2} [k_2x_2\Gamma_-^{+-} - \kappa(\delta_- + \delta_+)\Delta] \right\} K_5 \\
 & + i \left\{ \frac{(1+k_2^2)\Gamma_+^{--}}{k_2^2x_2} \cosh(g) + \frac{1}{k_2^2x_2} [k_2x_2(\delta_- - \delta_+) + \kappa\Gamma_+^{--}\Delta] \right\} K_6 \\
 & - \frac{(\delta_- + \delta_+)\Delta - \Gamma_+^{--}((k_2^2+1)x_2 - k_2^2x_2)}{2k_2^2x_2\sqrt{x_2^2+1}} \cosh(g)(K_7 + K_8) \\
 & + \left[\kappa \frac{\delta_- + \delta_+ - x_2\Gamma_+^{--}\Delta}{2k_2^2x_2\sqrt{x_2^2+1}} + \frac{(\delta_+ - \delta_-)x_2 - \Gamma_-^{+-}\Delta}{2k_2\sqrt{x_2^2+1}} \right] K_7 \\
 & + \left[\kappa \frac{-\delta_- - \delta_+ - x_2\Gamma_+^{--}\Delta}{2k_2^2x_2\sqrt{x_2^2+1}} + \frac{\Gamma_-^{+-}\Delta + (\delta_+ - \delta_-)x_2}{2k_2\sqrt{x_2^2+1}} \right] K_8 \\
 & + \frac{\Gamma_+^{--}\Delta - (\delta_- + \delta_+)((k_2^2+1)x_2 - k_2^2x_2)}{2k_2^2x_2\sqrt{x_2^2+1}} \cosh(g)(K_9 + K_{10}) \\
 & + \left[\kappa \frac{\Gamma_+^{--} - (\delta_- + \delta_+)x_2\Delta}{2k_2^2x_2\sqrt{x_2^2+1}} + \frac{(\delta_- - \delta_+)\Delta + x_2\Gamma_-^{+-}}{2k_2\sqrt{x_2^2+1}} \right] K_9 \\
 & + \left[-\kappa \frac{\Gamma_+^{--} + (\delta_- + \delta_+)x_2\Delta}{2k_2^2x_2\sqrt{x_2^2+1}} + \frac{(\delta_+ - \delta_-)\Delta + x_2\Gamma_-^{+-}}{2k_2\sqrt{x_2^2+1}} \right] K_{10},
 \end{aligned}$$

where we introduced the abbreviations

$$\Gamma_{\delta_1}^{\delta_2\delta_3} := \alpha_- + \delta_1\alpha_+ + \delta_2\beta_- + \delta_3\beta_+ \quad \Delta := \sqrt{1+k_2^2(1+x_2^2)} \quad g := \int^t \lambda(s)ds, \quad (4.20)$$

with $\delta_i = \pm 1$, $i = 1, 2, 3$. We simplified here our expressions using the identity

$$\sinh(g) = \frac{\kappa + \cosh(g)\sqrt{1+k_2^2(1+x_2^2)}}{k_2x_2}, \quad (4.21)$$

which is verified using the first order constraint (3.13). As the invariant is non-Hermitian the iteration process breaks down and by (S1) we deduce that $A\eta_2$ is not a Dyson map. We have also carried out the equivalent calculation for the seed map choices $\eta = \eta_1$ and $\tilde{\eta} = \eta_3$ and η_4 , reaching the same conclusion.

5. Conclusions

We proposed a scheme that allows to compute new time-dependent Dyson maps from two seed maps in an iterative fashion for a given non-Hermitian time-dependent Hamiltonian. As argued in general in section 2, in principle the iteration process might continue indefinitely,

thus leading to an infinite series of time-dependent Dyson maps including their associated Hermitian Hamiltonians. The symmetry operators S of the Lewis-Riesenfeld invariants govern this behaviour. Thus when the symmetry is broken also the iteration procedure breaks down. We carried out the procedure in detail for a two-dimensional system of harmonic oscillators that are coupled to each other in a non-Hermitian, but \mathcal{PT} -symmetrical, fashion. We only presented here three examples that exhibit different types of behaviours, but we have verified that similar results are obtained when starting from different sets of seed functions. We have focused in our analysis mainly on the relations between the various Hamiltonians and their corresponding invariants, but having obtained the Dyson maps, and therefore the metric operators, it is straightforward to extend the considerations to the associated wave functions and inner product structures on the physical Hilbert space.

Naturally there are various open issues and extensions possible. As always one may carry out the analysis for more concrete examples. Here we were mainly interested in demonstrating the working of the scheme and selected a model that has a simple two dimensional harmonic oscillator as Hermitian counterpart. It would be interesting to carry out the analysis for an example that has more involved Hermitian counterparts. It is also possible to extend the scheme to start with three or more inequivalent seed Dyson maps leading to new maps of different types, see [27] for details. In addition, one may also try to generalise the seed functions themselves and start from expressions involving more than the two factors as in the Ansatz (3.6).

Acknowledgments: RT is supported by a City, University of London Research Fellowship.

References

- [1] F. G. Scholtz, H. B. Geyer, and F. Hahne, Quasi-Hermitian Operators in Quantum Mechanics and the Variational Principle, *Ann. Phys.* **213**, 74–101 (1992).
- [2] C. M. Bender and S. Boettcher, Real Spectra in Non-Hermitian Hamiltonians Having \mathcal{PT} Symmetry, *Phys. Rev. Lett.* **80**, 5243–5246 (1998).
- [3] A. Mostafazadeh, Pseudo-Hermitian Representation of Quantum Mechanics, *Int. J. Geom. Meth. Mod. Phys.* **7**, 1191–1306 (2010).
- [4] C. M. Bender, P. E. Dorey, C. Dunning, A. Fring, D. W. Hook, H. F. Jones, S. Kuzhel, G. Levai, and R. Tateo, \mathcal{PT} Symmetry: In Quantum and Classical Physics, (World Scientific, Singapore) (2019).
- [5] A. Mostafazadeh, Metric operators for quasi-Hermitian Hamiltonians and symmetries of equivalent Hermitian Hamiltonians, *J. Phys.* **A41**, 055304 (2008).
- [6] H. F. Jones and J. Mateo, An Equivalent Hermitian Hamiltonian for the non-Hermitian $-x^4$ Potential, *Phys. Rev.* **D73**, 085002 (2006).
- [7] P. E. G. Assis and A. Fring, Metrics and isospectral partners for the most generic cubic \mathcal{PT} -symmetric non-Hermitian Hamiltonian, *J. Phys.* **A41**, 244001 (2008).
- [8] P. E. G. Assis and A. Fring, Non-Hermitian Hamiltonians of Lie algebraic type, *J. Phys.* **A42**, 015203 (23p) (2009).

- [9] D. P. Musumbu, H. B. Geyer, and W. D. Heiss, Choice of a metric for the non-Hermitian oscillator, *J. Phys.* **A40**, F75–F80 (2007).
- [10] F. J. Dyson, Thermodynamic Behavior of an Ideal Ferromagnet, *Phys. Rev.* **102**, 1230–1244 (1956).
- [11] C. Figueira de Morisson Faria and A. Fring, Time evolution of non-Hermitian Hamiltonian systems, *J. Phys.* **A39**, 9269–9289 (2006).
- [12] A. Mostafazadeh, Time-dependent pseudo-Hermitian Hamiltonians defining a unitary quantum system and uniqueness of the metric operator, *Physics Letters B* **650**(2), 208–212 (2007).
- [13] M. Znojil, Time-dependent version of crypto-Hermitian quantum theory, *Physical Review D* **78**(8), 085003 (2008).
- [14] H. Břila, Adiabatic time-dependent metrics in PT-symmetric quantum theories, arXiv preprint arXiv:0902.0474 (2009).
- [15] J. Gong and Q.-H. Wang, Time-dependent PT-symmetric quantum mechanics, *J. Phys. A: Math. and Theor.* **46**(48), 485302 (2013).
- [16] A. Fring and M. H. Y. Moussa, Unitary quantum evolution for time-dependent quasi-Hermitian systems with nonobservable Hamiltonians, *Phys. Rev. A* **93**(4), 042114 (2016).
- [17] H. Lewis and W. Riesenfeld, An Exact quantum theory of the time dependent harmonic oscillator and of a charged particle time dependent electromagnetic field, *J. Math. Phys.* **10**, 1458–1473 (1969).
- [18] B. Khantoul, A. Bounames, and M. Maamache, On the invariant method for the time-dependent non-Hermitian Hamiltonians, *The European Physical Journal Plus* **132**(6), 258 (2017).
- [19] M. Maamache, O. K. Djeghiour, N. Mana, and W. Koussa, Pseudo-invariants theory and real phases for systems with non-Hermitian time-dependent Hamiltonians, *The European Physical Journal Plus* **132**(9), 383 (2017).
- [20] A. Fring and T. Frith, Solvable two-dimensional time-dependent non-Hermitian quantum systems with infinite dimensional Hilbert space in the broken PT-regime, *J. of Phys. A: Math. and Theor.* **51**(26), 265301 (2018).
- [21] J. Cen, A. Fring, and T. Frith, Time-dependent Darboux (supersymmetric) transformations for non-Hermitian quantum systems, *J. of Phys. A: Math. and Theor.* **52**(11), 115302 (2019).
- [22] A. Fring and R. Tenney, Exactly solvable time-dependent non-Hermitian quantum systems from point transformations, *Phys. Lett. A* **410**, 127548 (2020).
- [23] A. Fring and R. Tenney, Perturbative approach for strong and weakly coupled time-dependent for non-Hermitian quantum systems, *Physica Scripta* **96**(4), 045211 (2021).
- [24] V. Ermakov, Transformation of differential equations,, *Univ. Izv. Kiev.* **20**, 1–19 (1880).
- [25] E. Pinney, The nonlinear differential equation $y'' + p(x)y + c/y^3 = 0$, *Proc. Amer. Math. Soc.* **1**, 681(1) (1950).
- [26] A. Fring and R. Tenney, Time-independent approximations for time-dependent optical potentials, *The European Physical Journal Plus* **135**(2), 163 (2020).

- [27] R. Tenney, New exact and approximation methods for time-dependent non-Hermitian quantum systems, PhD Thesis, City, University of London (2021).