



# City Research Online

## City St George's, University of London

**Citation:** Durai, R. R., Bharadwaj, S. R., Barbur, J. L., Rodriguez, M. & Rugvedi, A. (2021). Binocular summation of cone- and rod specific signals in flicker-threshold tasks. *Investigative Ophthalmology & Visual Science*, 62(8), ISSN 0146-0404

This is the published version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/27368/>

**Copyright and Reuse:** Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

# Binocular summation of cone- and rod-specific signals in flicker-threshold tasks | IOVS

 [iovs.arvojournals.org/article.aspx](https://iovs.arvojournals.org/article.aspx)

June 2021

Volume 62, Issue 8

ARVO Annual Meeting Abstract | June 2021

Binocular summation of cone- and rod-specific signals in flicker-threshold tasks

Investigative Ophthalmology & Visual Science June 2021, Vol.62, 1442. doi:

- ○ Reena Reena Durai, Shrikant R Bharadwaj, John Barbur, Marisa Rodriguez, Amitha Vikram Rugvedi; Binocular summation of cone- and rod-specific signals in flicker-threshold tasks. *Invest. Ophthalmol. Vis. Sci.* 2021;62(8):1442.

© ARVO (1962-2015); The Authors (2016-present)

x

## Abstract

**Purpose** : Binocular summation of briefly-presented, central and peripheral, spatially-distributed stimuli has been investigated in numerous studies with emphasis on stimulus size, location in the visual field and state of background adaptation. This study examined the binocular threshold summation of cone- and rod-specific, flickering stimuli as a function of stimulus size and retinal location.

**Methods** : Flicker modulation thresholds (FMT's) were measured under monocular and binocular viewing conditions in 13 subjects with normal vision (mean age:23±3yrs, best-corrected visual acuity  $\leq 0.0\log\text{MAR}$ , stereoacuity  $\leq 40\text{arc sec}$  and normal trichromatic colour vision). Rapid flicker thresholds were measured using the Flicker-*Plus* test (City Occupational Ltd., London) for 5 different target sizes (7, 15, 30, 45 and 60arc min) in central vision ( $0^\circ$ ) and at an eccentricity of  $5^\circ$  in each quadrant using a 5-alternative forced-choice psychophysical technique. The temporal frequency, duration, luminance and chromaticity of the flickering stimuli was optimized to favour either cone- or rod-photoreceptors. The ratio of the subject's monocular FMT (measured in the best eye) and the corresponding binocular threshold was used to quantify binocular summation.

**Results** : Monocular FMT's decreased monotonically with target size for both cone- and rod-optimized stimuli ( $p < 0.001$ ). Not surprising, the extreme percentage FMTs corresponded to the smallest (7') and the largest (60') stimuli with  $m \pm s$  values of:  $9.69 \pm 1.06$  and  $2.35 \pm 0.20$  (for cones) and  $20.92 \pm 2.50$  and  $4.45 \pm 0.50$  (for rods). Binocular FMT's followed the same trend for all viewing conditions but were smaller than their

monocular counterparts ( $p < 0.001$ ). The mean summation ratios was  $1.28 \pm 0.49$  for cones and  $1.41 \pm 0.35$  for rods. There was no indication of significant changes in binocular summation with either target size ( $p < 0.12$ ) or retinal location ( $p = 0.67$ ).

**Conclusions** : Detection thresholds for luminance-modulated flicker exhibit large binocular summation with both cone- and rod-enhanced stimuli. Although both monocular and binocular FMTs show large, systematic variation with target size and retinal location, the binocular summation ratio remains largely invariant of these parameters.

This is a 2021 ARVO Annual Meeting abstract.

This work is licensed under a [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License](https://creativecommons.org/licenses/by-nc-nd/4.0/).



Copyright © 2015 Association for Research in Vision and Ophthalmology.

Copyright © 2015 Association for Research in Vision and Ophthalmology.