



City Research Online

City St George's, University of London

Citation: Correa, F., Fring, A. & Taira, T. (2022). Linearly stable and unstable complex soliton solutions with real energies in the Bullough-Dodd model. Nuclear Physics B, 979, 115783. doi: 10.1016/j.nuclphysb.2022.115783

This is the published version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/28100/>

Link to published version: <https://doi.org/10.1016/j.nuclphysb.2022.115783>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).



Linearly stable and unstable complex soliton solutions with real energies in the Bullough-Dodd model

Francisco Correa^b, Andreas Fring^{a,*}, Takanobu Taira^a

^a Department of Mathematics, City, University of London, Northampton Square, London EC1V 0HB, UK

^b Instituto de Ciencias Físicas y Matemáticas, Universidad Austral de Chile, Casilla 567, Valdivia, Chile

Received 24 October 2021; received in revised form 4 April 2022; accepted 7 April 2022

Available online 13 April 2022

Editor: Hubert Saleur

Abstract

We investigate different types of complex soliton solutions with regard to their stability against linear perturbations. In the Bullough-Dodd scalar field theory we find linearly stable complex \mathcal{PT} -symmetric solutions and linearly unstable solutions for which the \mathcal{PT} -symmetry is broken. Both types of solutions have real energies. The auxiliary Sturm-Liouville eigenvalue equation in the stability analysis for the \mathcal{PT} -symmetric solutions can be solved exactly by supersymmetrically mapping it to an isospectral partner system involving a shifted and scaled inverse cosh-squared potential. We identify exactly one shape mode in form of a bound state solution and scattering states which when used as linear perturbations leave the solutions stable. The auxiliary problem for the solutions with broken \mathcal{PT} -symmetry involves a complex shifted and scaled inverse sin-squared potential. The corresponding bound and scattering state solutions have complex eigenvalues, such that when used as linear perturbations for the corresponding soliton solutions lead to their decay or blow up as time evolves.

© 2022 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

* Corresponding author.

E-mail addresses: francisco.correa@uach.cl (F. Correa), a.fring@city.ac.uk (A. Fring), takanobu.taira@city.ac.uk (T. Taira).

1. Introduction

The success of \mathcal{PT} -symmetric quantum mechanics [1] and the applications of the concepts in optical analogues [2] have inspired research in many different areas of physics. One of them is the study of complex \mathcal{PT} -symmetric soliton solutions in nonlinear integrable systems in 1+1 dimensions [3–13] or models that admit complex Skyrmion type solutions in 2+1 dimensions [14]. Despite being complex, these sorts of solutions possess nonetheless real energies, a property that can be attributed to general arguments that make use of the underlying antilinear symmetry, generally referred to as \mathcal{PT} -symmetry [3,7,13,14].

The other aspect, that makes such type of solutions potentially interesting physical objects, is their stability. So far this vital property has been largely ignored in their studies and here the main purpose is to investigate the stability properties of particular complex one-soliton solutions for a specific integrable scalar field theory, the Bullough-Dodd model [15,16]. Besides shedding light on the important question of whether a complex soliton solution will collapse, i.e. grow or decay, when slightly perturbed, it is also well-known that as a by-product of a stability analysis one may obtain the so-called shape mode as a bound state solution for the auxiliary Sturm-Liouville eigenvalue equation. These solutions are instrumental in the understanding of energy transfer effects in the dynamics of multi-soliton solutions that can be unravelled using a collective coordinate or moduli space approach [17,18]. For the model studied here we will investigate these moduli space properties in upcoming work [19].

Our manuscript is organised as follows: In section 2 we assemble some generalities about the stability analysis to be carried out. In addition, we briefly recall the general set up for a supersymmetric formulation of quantum mechanics. While this is not essential for a stability analysis, it greatly facilitates the construction of exact solutions to the auxiliary Sturm-Liouville eigenvalue equations. In section 3 we construct complex one-soliton solutions for the Bullough-Dodd model and analyse in detail their stability.

2. Stability analysis, zero modes, shape modes and supersymmetry

The systems considered here are complex scalar field theories with a Lagrangian density of the general form

$$\mathcal{L} = \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi - V(\varphi), \quad (2.1)$$

with $\varphi(x, t) \in \mathbb{C}$ and potential $V(\varphi) \in \mathbb{C}$. We denote the Lagrangian, i.e. the volume integral over this density, as $L = \int_{-\infty}^{\infty} \mathcal{L} dx$. Our space-time metric is taken to be Lorentzian of the form $\text{diag}(1, -1)$, so that the Euler-Lagrange equation resulting from (2.1) reads

$$\ddot{\varphi} - \varphi'' + \frac{\partial V(\varphi)}{\partial \varphi} = 0. \quad (2.2)$$

We use standard conventions and denote partial derivatives with respect to time t and space x by overdots and dashes, respectively. The static solutions to these equations, i.e. the solutions to (2.2) with $\dot{\varphi} = 0$, are fields $\phi(x, \mathbf{u})$ depending on the space coordinate x and a set of constants that are identified as moduli or collective coordinates $\mathbf{u} = (u^1, \dots, u^n)$. As is well-known, one may construct the time-dependent solutions φ from the time-independent solutions ϕ simply by

a Lorentz boost $\phi[(x - vt)/\sqrt{1 - v^2}] = \varphi(x, t)$ with v denoting the velocity. The energies of particular solutions are computed as

$$E[\varphi] = \int_{-\infty}^{\infty} dx \varepsilon(\varphi), \quad \varepsilon(\varphi) = \left(\frac{1}{2} \dot{\varphi}^2 + \frac{1}{2} (\varphi')^2 + V(\varphi) \right), \quad (2.3)$$

where $\varepsilon(\varphi)$ denotes the energy density. When the Hamiltonian associated to (2.1) admits a modified \mathcal{CPT} -symmetry and φ is mapped by this symmetry to itself or possibly to a second degenerate solution, the energy $E[\varphi]$ is guaranteed to be real, irrespective of whether $\varphi \in \mathbb{R}$, $\varphi \in \mathbb{C}$ or $V^\dagger(\varphi) = V(\varphi)$ [14]. This reasoning may also be extended to ensure the reality of other conserved quantities.

With regard to the main purpose of this paper, we now briefly recall the main argument of a standard stability analysis beginning with the linearisation of the Euler-Lagrange equation (2.2) by replacing

$$\varphi \rightarrow \varphi_s + \varepsilon \delta\varphi \quad \varepsilon \ll 1, \quad (2.4)$$

where φ_s solves (2.2) and $\delta\varphi =: \chi$ is a small perturbation. This converts the Euler-Lagrange equation into

$$\ddot{\chi} - \chi'' + \chi \left(\frac{\partial^2 V(\varphi)}{\partial \varphi^2} \Big|_{\varphi_s} \right) + \mathcal{O}(\varepsilon^2) = 0. \quad (2.5)$$

Evidently the first three terms in (2.5) vanish by construction and when assuming the perturbation to be separable of the form $\chi(x, t) = e^{i\lambda t} \Phi(x)$, the term of first order in ε reduces to a Sturm-Liouville eigenvalue problem of the same form as the time-independent Schrödinger equation

$$-\Phi_{xx} + V_1 \Phi = \lambda^2 \Phi, \quad \text{with } V_1(x) := \frac{\partial^2 V(\varphi)}{\partial \varphi^2} \Big|_{\varphi_s}. \quad (2.6)$$

For the eigenfunction solutions to this equation with $\lambda \in \mathbb{R}$ the linear perturbation will simply introduce an oscillation in time around the solution φ_s , whereas when $\lambda \in \mathbb{C}$ the solutions will grow or decay with time and are therefore unstable.

Let us next see how the energy of the perturbed solution behaves. Expanding up to second order in ε and integrating two terms by parts, we easily derive

$$E[\varphi_s + \chi] = E[\varphi_s] + \int_{-\infty}^{\infty} dx \left[\left(\frac{\partial V(\varphi)}{\partial \varphi} \Big|_{\varphi_s} - \varphi_s'' \right) \chi + \frac{1}{2} \chi \left(\frac{\dot{\chi}^2}{\chi} - \chi'' + \chi \frac{\partial^2 V(\varphi)}{\partial \varphi^2} \Big|_{\varphi_s} \right) \right] + \chi(\chi' + \varphi_s') \Big|_{-\infty}^{\infty} + \mathcal{O}(\varepsilon^3). \quad (2.7)$$

We notice that first bracket of the integrand is simply the Euler-Lagrange equation for the static solution. The second bracket of the integrand vanishes when χ is separable as stated above and $\Phi(x)$ satisfies the Sturm-Liouville equation (2.6). The surface term in the second line of (2.7) is zero when $\Phi(x)$ vanishes asymptotically $\lim_{x \rightarrow \pm\infty} \Phi(x) = 0$, so that the energy of the perturbed solution becomes identical to the one of the static solution φ_s .

The *zero mode*, or translation mode, defined as the solution Φ_0 to (2.6) with $\lambda = 0$, plays a special role for a number of reasons. First of all, it is well-known [20–22] that the dimensionality of the corresponding moduli spaces may be obtained from these modes. Moreover it allows to directly compute the superpotential and to set up a supersymmetric scheme that will be important for the concrete system we consider below.

The zero mode is easily computed: Assuming that the static solution $\phi(x)$ belongs to a class of solutions characterised by a continuous parameter, say a , such that $\phi(x) = \psi(x, a)|_{a=0}$, we may think of $\phi_s + \varepsilon\delta\phi$ in (2.4) as

$$\psi(x, a) = \phi(x) + a \left. \frac{\partial\psi(x, a)}{\partial a} \right|_{a=0} + \mathcal{O}(a^2), \tag{2.8}$$

so that we can identify the zero mode as

$$\Phi_0(x) := \left. \frac{\partial\psi(x, a)}{\partial a} \right|_{a=0}. \tag{2.9}$$

We may now repeat the argument for different continuous parameters $a \rightarrow a_1, \dots, a_n$, so that the number n of independent zero modes constitutes the dimension of the moduli space.

The excited bound states or scattering states for the eigenvalue equation in (2.6) may be constructed in a number of ways. Here it turns out that the potentials we are interested in possess superpartner potentials that have been solved in the literature, so that we may employ the scheme and simply map the known solutions to our problem. Let us therefore briefly recall the general set up of supersymmetric quantum mechanics by following the ideas of Witten and others [23–26]. Assuming that the eigenvalue equation (2.6) has a discrete eigenvalue spectrum, $E_n^{(1)} = \lambda_n^2$, we re-write it as

$$H_1\Phi_n^{(1)} = \left(-\frac{d^2}{dx^2} + V_1\right)\Phi_n^{(1)} = L_+L_-\Phi_n^{(1)} = E_n^{(1)}\Phi_n^{(1)}, \tag{2.10}$$

with $\Phi_0 = \Phi_0^{(1)}$ and $E_0 = 0$. We assumed here that the Hamiltonian factorises into the product of two first order differential operators defined as

$$L_{\pm} := \mp \frac{d}{dx} + W(x), \quad \text{with } W(x) := -\frac{\Phi_0'}{\Phi_0}, \tag{2.11}$$

where $W(x)$ is referred to as the *superpotential* related to the potential in (2.10) as $V_1 = W^2 - W'$. Defining next a new Hamiltonian H_2 with the product of the operators L_{\pm} in reverse order, we have

$$H_2\Phi_m^{(2)} = \left(-\frac{d^2}{dx^2} + V_2\right)\Phi_m^{(2)} = L_-L_+\Phi_m^{(2)} = E_m^{(2)}\Phi_m^{(2)}, \tag{2.12}$$

with a second potential $V_2 = W^2 + W'$. Considering now

$$H_2\left(L_-\Phi_n^{(1)}\right) = L_-L_+L_-\Phi_n^{(1)} = E_n^{(1)}\left(L_-\Phi_n^{(1)}\right), \tag{2.13}$$

$$H_1\left(L_+\Phi_m^{(2)}\right) = L_+L_-L_+\Phi_m^{(2)} = E_m^{(2)}\left(L_+\Phi_m^{(2)}\right), \tag{2.14}$$

and comparing these equations with (2.10), (2.12) we conclude

$$\Phi_{n+1}^{(1)} = N_{n+1}^{(1)}L_+\Phi_n^{(2)}, \quad \Phi_n^{(2)} = N_n^{(2)}L_-\Phi_n^{(1)}, \quad E_n^{(2)} = E_{n+1}^{(1)}, \quad E_0^{(1)} = 0, \quad n \in \mathbb{N}_0. \tag{2.15}$$

Thus apart from $E_0^{(1)}$ the two Hamiltonians H_1 and H_2 are isospectral, hence the referral as *supersymmetric*. When $L_-^\dagger = L_+$ the normalisation constants are simply $N_{n+1}^{(1)} = 1/\sqrt{E_n^{(2)}}$,

$N_n^{(2)} = 1/\sqrt{E_{n+1}^{(1)}}$, but we will not assume this as we allow here for complex superpotentials W . The scheme is directly extended to scattering solutions with continuous eigenvalues spectra.

The crucial feature for our purposes is here the observation from (2.15) that when one of the eigenvalue problems (2.10) or (2.12) has been solved, one may directly deduce the solutions for the other. It is in this sense that we will apply this scheme as the potentials V_1 we construct from the zero modes are not obvious to solve, whereas the solutions for the supersymmetric partner potentials V_2 can be found in the literature.

3. The Bullough-Dodd model

As an example, we are interested here in the stability properties of the complex soliton solutions of the Bullough-Dodd model [15,16], that is a one scalar field integrable field theory already quite well studied in its classical [27] and quantum [28] aspects. It is described by the Lagrangian density of the form

$$\mathcal{L}_{BD} = \frac{1}{2} \partial_\mu \varphi \partial^\mu \varphi - e^\varphi - \frac{1}{2} e^{-2\varphi} + \frac{3}{2} \quad \text{with } \varphi \in \mathbb{C}. \tag{3.1}$$

The resulting classical nonlinear equation of motion

$$\ddot{\varphi} - \varphi'' + e^\varphi - e^{-2\varphi} = 0, \tag{3.2}$$

can be solved by standard techniques from classical theory of integrable systems. Here we exploit the fact that the equation may be solved using Hirota’s direct method [29] when parameterizing the fields as $\varphi = \ln(\tau_0/\tau_1)$ with τ_i denoting the so-called τ -functions. To identify various one-soliton solutions we can therefore make the general Ansatz

$$\varphi(x, t) = \ln \left(\frac{\alpha_0 + \alpha_1 e^{kx+lt} + \alpha_2 e^{2kx+2lt}}{\beta_0 + \beta_1 e^{kx+lt} + \beta_2 e^{2kx+2lt}} \right), \tag{3.3}$$

with unknown constants $\alpha_i, \beta_i, i = 0, 1, 2$, and k, l . Substituting this Ansatz into equation (3.2) and subsequently reading off the coefficients of $e^{jkx+jlt}$ for $j = 0, \dots, 8$ leads to 9 equations, that we do not report here, which may be solved for the unknown constants in (3.3). In this manner we find various types of solutions.

3.1. Complex one-soliton solutions of type I

The first type of solutions we obtain are one-soliton solutions of the form

$$\varphi_I^\pm(x, t) = \ln \left[\frac{\cosh(\beta + \sqrt{k^2 - 3t + kx}) \pm 2}{\cosh(\beta + \sqrt{k^2 - 3t + kx}) \mp 1} \right], \quad \beta \in \mathbb{C}. \tag{3.4}$$

When β is real and $|k| > \sqrt{3}$ the solution φ_I^+ is real but becomes singular for $x_0 = -[\beta + \sqrt{k^2 - 3t}]/k$, whereas φ_I^- is only real for $x < x_l = -[\beta + \text{arccosh}(2) + \sqrt{k^2 - 3t}]/k$ and $x > x_r = [-\beta + \text{arccosh}(2) - \sqrt{k^2 - 3t}]/k$ as the argument of the logarithm becomes negative in the complementary regime. Samples of these types of solutions in the different regimes are depicted in Fig. 1. In the case $|k| < \sqrt{3}$ both solutions φ_I^\pm are always complex, see Fig. 3. The type I solutions found here formally coincide with the solutions constructed in [30].




Fig. 1. Scattering solutions (3.4) of the Bullough-Dodd classical equation of motion (3.2) for $|k| > \sqrt{3}$. Panel (a): Travelling real one-cusp solution φ_I^+ for $k = 2$ with $\beta = 3$ and panel (b) \mathcal{PT} regularized travelling complex solution φ_I^+ for $k = 2$ with $\beta = i3/5$. Solid lines correspond to real and dotted lines to imaginary parts.




Fig. 2. Scattering solutions (3.4) of the Bullough-Dodd classical equation of motion (3.2) for $|k| > \sqrt{3}$. Panel (a): Complex solution φ_I^- for $k = 2$ with $\beta = 3$ and panel (b) \mathcal{PT} regularized complex solution φ_I^- for $k = 2$ with $\beta = i3/5$. Solid lines correspond to real and dotted lines to imaginary parts.

Despite the fact that these solutions are complex, their energies are real governed by an anti-linear (\mathcal{PT} -symmetry), and for N-soliton solutions together with the fact that these theories are integrable [7,13]. The \mathcal{PT} -symmetry is easily identified as

$$\mathcal{PT} : x \rightarrow -x, \quad t \rightarrow -t, \quad i \rightarrow -i, \quad \varphi \rightarrow \varphi. \tag{3.5}$$

The Bullough-Dodd Lagrangian (3.1) is trivially invariant under this symmetry and the solutions $\varphi_I^\pm(x, t)$ evidently respect it for purely imaginary constants β , i.e. $\mathcal{PT} : \varphi_I^\pm \rightarrow \varphi_I^\pm$. Thus to take $\beta \in i\mathbb{R}$ seems to be very suggestive as this choice not only converts the solutions into \mathcal{PT} -symmetric ones, but also regularises the singularities of the solutions at the cost of a discontinuity resulting from the branch cut of the logarithm as depicted in Figs. 1 and 2. The behaviour of these complex shifted solutions is identical to those for the $|k| < \sqrt{3}$ and $\beta \in \mathbb{R}$ already depicted in Fig. 3, when we identify in these solutions $\beta \rightarrow \sqrt{k^2 - 3t}$ and $\sqrt{k^2 - 3t} \rightarrow i\beta$ for fixed time.

Indeed the energies for the solutions (3.4), evaluated according to (2.3), are always real

$$E[\varphi_I^\pm] = -6|k|, \tag{3.6}$$

and independent of the complex parameter β .




Fig. 3. Complex breather solutions (3.4) of the Bullough-Dodd classical equation of motion (3.2) for $|k| < \sqrt{3}$. Panel (a): φ_1^+ for $k = 1 < \sqrt{3}$ with $\beta = 2$ and panel (b) φ_1^- for $k = 1 < \sqrt{3}$ with $\beta = 2$. Solid lines correspond to real and dotted lines to imaginary parts.

3.1.1. Zero modes for type I complex one-soliton solutions

The Sturm-Liouville eigenvalue problem (2.6) of the stability analysis for the Bullough-Dodd model reads

$$-\Phi_{xx} + V(x)\Phi = \lambda^2\Phi \quad \text{with} \quad V(x) = \left(e^{\phi(x)} + 2e^{-2\phi(x)} \right). \tag{3.7}$$

The static solution for the cusp and oscillatory solutions $\phi_I^\pm(x, t)$ acquire the form

$$\phi_I^\pm(x) = \ln \left[\frac{\cosh(\beta + \sqrt{3}x) \pm 2}{\cosh(\beta + \sqrt{3}x) \mp 1} \right], \tag{3.8}$$

which when substituted into (3.7) yield the potentials

$$V_1^+(x) = 1 - \frac{3}{1 - \cosh(\beta + \sqrt{3}x)} + \frac{8 \sinh^4 \left[\frac{1}{2}(\beta + \sqrt{3}x) \right]}{\left[2 + \cosh(\beta + \sqrt{3}x) \right]^2}, \tag{3.9}$$

$$V_1^-(x) = 1 - \frac{3}{1 + \cosh(\beta + \sqrt{3}x)} + \frac{8 \cosh^4 \left[\frac{1}{2}(\beta + \sqrt{3}x) \right]}{\left[2 - \cosh(\beta + \sqrt{3}x) \right]^2}. \tag{3.10}$$

We think now of the larger class of functions depending on the continuous parameter a as $\psi^\pm(x, a) = \phi^\pm(x + a/\sqrt{3})$. By means of (2.9) the zero mode is then easily computed to

$$\Phi_0^\pm(x) = - \frac{3 \left\{ \tanh \left[\frac{1}{2}(\beta + \sqrt{3}x) \right] \right\}^{\mp 1}}{2 \pm \cosh(\beta + \sqrt{3}x)}. \tag{3.11}$$

One verifies that the $\Phi_0^\pm(x)$ indeed satisfy (3.7) for the potentials (3.10) with eigenvalue $\lambda = 0$. Notice that when taking $a \rightarrow ia$, as seems to be natural when $k < \sqrt{3}$, so that we identify $\psi^\pm(x, a) = \phi^\pm(x + ia/\sqrt{3})$, the resulting zero mode is the same as in (3.11) only multiplied by i , which simply corresponds to a different normalization constant.

3.1.2. Bound state solutions for type I complex one-soliton solutions

It is not straightforward or obvious how to solve the eigenvalue equation for the potentials $V_1^\pm(x)$ directly. However, we notice now that the supersymmetric partner Hamiltonians are inverse \cosh^2 -potentials that have been treated extensively in the literature. Since $V_1^+(x, \beta) = V_1^-(x, \beta + i\pi)$, we just focus on the “+”-case from now on, as all quantities related to the “-”-system are computed simply by a shift in β . Using the relation (2.11) we compute the superpotential from the zero modes to

$$W^+(x) = \frac{\sqrt{3} \, 3 + 2 \cosh(\beta + \sqrt{3}x) + \cosh[2(\beta + \sqrt{3}x)]}{2 \left[\cosh(\beta + \sqrt{3}x) + 2 \right] \sinh(\beta + \sqrt{3}x)}, \tag{3.12}$$

which is sufficient to define the corresponding intertwining operators L_\pm in (2.11). Using next the defining relation for the second Hamiltonian H_2 we compute the associated second potential from (2.12) to

$$V_2 = 3 - \frac{3}{2} \operatorname{sech}^2 \left(\frac{\beta}{2} + \frac{\sqrt{3}x}{2} \right). \tag{3.13}$$

When shifting the overall energy by 3, this is an inverse \cosh^2 -potential discussed in the literature. It possesses bound states as treated in section 23 in problem 5 [31] and scattering states for which the potential is entirely reflectionless [32], see also [33]. From [31] we read off the eigenfunction for the energy E with potential $U = -U_0 \operatorname{sech}^2(\alpha x)$, $\alpha, U_0 \in \mathbb{R}$ as

$$\Psi(x) = \left[1 - \tanh^2(\alpha x) \right]^{\epsilon/2} {}_2F_1 \left[\epsilon - s, s + \epsilon + 1; \epsilon + 1; \frac{1}{1 + e^{2\alpha x}} \right], \tag{3.14}$$

where ${}_2F_1(a, b; c; z)$ denotes the hypergeometric function and

$$\epsilon := \frac{\sqrt{-2Em}}{\alpha \hbar}, \quad s := \frac{1}{2} \left(\sqrt{1 + \frac{8mU_0}{\alpha^2 \hbar^2}} - 1 \right). \tag{3.15}$$

The quantization condition for the energy emerges from the requirement $\lim_{x \rightarrow \pm\infty} \Psi(x) = 0$ as $\epsilon - s = -n$ for $n \in \mathbb{N}_0$, which when solved yields

$$E_n = -\frac{\alpha^2 \hbar^2}{8m} \left(\sqrt{\frac{8mU_0}{\alpha^2 \hbar^2} + 1} - 2n - 1 \right)^2. \tag{3.16}$$

As argued in [31], the number of bound states is limited by the constraint $n < s$. When shifting the overall energy by 3 we obtain $V_2 = U$ for the parameter identifications $\hbar = 1, m = 1/2, U_0 = 3/2, \alpha = \sqrt{3}/2$. Since for these values $s = 1$, there is only one bound state for $n = 0$. Thus we obtain the eigensystem solutions

$$\Phi_0^{(2)}(x) = N_0 \operatorname{sech} \left[\frac{1}{2} (\beta + \sqrt{3}x) \right], \quad E_0^{(2)} = \frac{9}{4}, \tag{3.17}$$

$$\Phi_1^{(1)}(x) = L_+ \Phi_0^{(2)} = N_0 \frac{3\sqrt{3} \cosh \left[\frac{1}{2} (\beta + \sqrt{3}x) \right] \coth(\beta + \sqrt{3}x)}{\cosh(\beta + \sqrt{3}x) + 2}, \quad E_1^{(1)} = \frac{9}{4}. \tag{3.18}$$

Notice that the singularity in $\Phi_1^{(1)}(x)$ for $x_0 = -\beta\sqrt{3}$ when $\beta \in \mathbb{R}$, inherited from the superpotential $W(x)$, is eliminated in the \mathcal{PT} -symmetric solution when $\beta \in i\mathbb{R}$. Since we have obtained a well-defined asymptotically vanishing solution, $\lim_{x \rightarrow \pm\infty} \Phi_1^{(1)}(x) = 0$ with real eigenvalues, we conclude that the complex type I soliton solution remains stable when linearly perturbed by the shape mode $\Phi_1^{(1)}$. Let us next consider the case when λ is taken to be a continuous parameter.

3.1.3. Scattering state solutions for type I complex one-soliton solutions

The scattering states for the inverse cosh-squared potential are discussed in detail in [32]. When adjusting the parameters therein to our equation (3.7) for V_2 , we obtain the two scattering solutions

$$\Psi_e^{(2)} = \cosh^2\left(\frac{\hat{x}}{2}\right) {}_2F_1\left[\gamma_-, \gamma_+; \frac{1}{2}; -\zeta^2(\hat{x})\right] = \cos(\tilde{x}) - \frac{\sqrt{3}}{2\kappa} \tanh\left(\frac{\hat{x}}{2}\right) \sin(\tilde{x}), \tag{3.19}$$

$$\begin{aligned} \Psi_o^{(2)} &= \zeta(\hat{x}) \cosh^2\left(\frac{\hat{x}}{2}\right) {}_2F_1\left[\frac{1}{2} + \gamma_-, \frac{1}{2} + \gamma_+; \frac{3}{2}; -\zeta^2(\hat{x})\right] \\ &= \frac{2\sqrt{3}\kappa \sin(\tilde{x}) + 3 \tanh\left(\frac{\hat{x}}{2}\right) \cos(\tilde{x})}{4\kappa^2 + 3}, \end{aligned} \tag{3.20}$$

with $\gamma_{\pm} := 1 \pm i\kappa/\sqrt{3}$, $\zeta(x) := \sinh(x/2)$, $\tilde{x} := \kappa\left(\frac{\beta}{\sqrt{3}} + x\right)$, $\hat{x} := \beta + \sqrt{3}x$ and continuous eigenvalues $\lambda^2 = 3 + \kappa^2$ for κ being a free parameter. As we expect from Abel's identity, the Wronskian for these two solutions is constant, $W(\Psi_e^{(2)}, \Psi_o^{(2)}) = \Psi_e^{(2)} d\Psi_o^{(2)}/dx - d\Psi_e^{(2)}/dx \Psi_o^{(2)} = \sqrt{3}/2$, indicating that they are indeed linearly independent. Thus according to (2.15), the two scattering states for the potential V_1 are obtained as

$$\Psi_e^{(1)} = L_+ \Psi_e^{(2)} = \frac{1}{\kappa} [f_-(\hat{x}) \cos(\tilde{x}) - f_+(\hat{x}) \sin(\tilde{x})], \tag{3.21}$$

$$\Psi_o^{(1)} = L_+ \Psi_o^{(2)} = \frac{2\sqrt{3}}{3 + 4\kappa^2} [f_+(\hat{x}) \cos(\tilde{x}) + f_-(\hat{x}) \sin(\tilde{x})], \tag{3.22}$$

where

$$f_+(x) := \frac{3}{2} - \kappa^2 - \frac{9}{2 \cosh(x) + 4}, \quad f_-(x) := \frac{3\sqrt{3}\kappa \cosh(x) \coth\left(\frac{x}{2}\right)}{2 \cosh(x) + 4}. \tag{3.23}$$

Once again the singularity at $x_0 = -\beta\sqrt{3}$ is eliminated in the \mathcal{PT} -symmetric solution by taking $\beta \in i\mathbb{R}$. Now the Wronskian for these two solutions is computed to $W(\Psi_e^{(1)}, \Psi_o^{(1)}) = (3 + \kappa^2)\sqrt{3}/2$. When $\kappa = \kappa_0 := \pm i\sqrt{3}$ we recover the zero mode Φ_0^+ from $\Psi_e^{(1)}$, whereas $\lim_{x \rightarrow \kappa_0} \Psi_o^{(1)}(x) = 0$.

Since the scattering solutions are well-defined with real continuous eigenvalues one may be tempted to conclude that the complex type I soliton solution remains stable when linearly perturbed by either of the two scattering solutions $\Psi_e^{(1)}$ or $\Psi_o^{(1)}$. However, as the limits for the asymptotic values of x are not well defined, the energies of the perturbed solution (2.7) do not give definite values so that the perturbation by these scattering solutions leads to ill-defined objects. A natural way to overcome this issue would be to restrict the domain of the theory to finite interval.




Fig. 4. Complex one-soliton solutions of type II (3.25) with $\beta = 2, k = 1/2$ for the Bullough-Dodd classical equation of motion (3.2). Solid lines correspond to real and dotted lines to imaginary parts.

An interesting solution arises for $\kappa = 0$

$$\lim_{\kappa \rightarrow 0} \Psi_o^{(1)} = \sqrt{3} \left(1 - \frac{3}{\cosh(\beta + \sqrt{3}x) + 2} \right) =: \Psi_0^{(1)}, \tag{3.24}$$

which has a well-defined finite limit $\sqrt{3}$ for $x \rightarrow \pm\infty$. Since the derivative of this solution is asymptotically vanishing as well as the derivative of the static solution ϕ_I , the second line in (2.7) is vanishing up to order ε^2 so that the energy of the perturbed solution is the same as the original solution. Since we have also a positive eigenvalue for the auxiliary eigenvalue problem we conclude that the solution remains stable when perturbed by $\Psi_0^{(1)}$.

3.2. Complex one-soliton solutions of type II

The second type of one-soliton solutions obtained from (3.3) by fixing the unknown constants from direct substitution into the Bullough-Dodd equation (3.2) is always complex and does not possess a real regime in their range

$$\varphi_{II}^\pm(x, t) = \ln \left\{ \omega^{\mp 2} \left[1 - \frac{6e^{t\sqrt{k^2+3\omega^{\pm 1}}+kx\pm\beta}}{\left(1 + e^{t\sqrt{k^2+3\omega^{\pm 1}}+kx\pm\beta}\right)^2} \right] \right\}. \tag{3.25}$$

Here $\omega = e^{i\pi/3}$ denotes the third root of unity. A sample solution is depicted in Fig. 4, from which we observe that the real part of this solution is an oscillation between a regular shaped soliton solution and a double peakon solution.

The energies when directly computed with (2.3) for the Lagrangian densities \mathcal{L}_{BD} are infinite. However, when shifting the vacuum from $\varphi_0 = 0$ to the complex plane $\varphi_0^\pm = \mp i2\pi/3$ amounts to defining the new Lagrangians densities

$$\mathcal{L}_{BD}^\pm = \mathcal{L}_{BD} + \frac{3}{2} (\omega^{\mp 2} - 1). \tag{3.26}$$

Computing the energies of the solution φ_{II}^\pm with these Lagrangians leads to the same real values as those computed for the type I solution with \mathcal{L}_{BD}

$$E^+[\varphi_{II}^+] = E^-[\varphi_{II}^-] = -6|k|. \tag{3.27}$$

The \mathcal{PT} -symmetry for the individual Lagrangians is now broken, but instead we have $\mathcal{PT} : \mathcal{L}_{\text{BD}}^+ \rightarrow \mathcal{L}_{\text{BD}}^-$. Since the energies are real this suggests that the criteria provided in [14] may still be slightly enlarged. One may easily convince oneself that the energies of two \mathcal{PT} -symmetrically related degenerate solutions are real even when they originate from two different \mathcal{PT} -symmetrically related Hamiltonians. Having obtained real energies suggests that these solutions are potentially well-defined physical objects, but let us investigate whether the solutions are also stable.

3.2.1. Zero modes for complex type II one-soliton solutions

The static solutions corresponding to $\varphi_{II}^\pm(x, t)$ are now obtained as

$$\phi_{II}^\pm(x) = \ln \left[\omega^{\mp 2} \frac{\cos(\sqrt{3}\omega^{\pm 1/2}x \pm \beta) - 2}{\cos(\sqrt{3}\omega^{\pm 1/2}x \pm \beta) + 1} \right]. \tag{3.28}$$

Taking $\beta \in \mathbb{R}$ the two solutions are related to each other by the \mathcal{PT} -symmetry in (3.5) as $\mathcal{PT} : \phi_{II}^+ \rightarrow \phi_{II}^-$. Alternatively, we may also change the sign in β in ϕ_{II}^- and take $\beta \in i\mathbb{R}$ in which case the two solutions are still related by the same symmetry. The potentials for the eigenvalue equation as introduced in (3.7) result to

$$V_1^\pm = \omega^{\mp 2} \left\{ 1 - \frac{3}{\cos(\sqrt{3}\omega^{\pm 1/2}x \pm \beta) + 1} + 2 \frac{[\cos(\sqrt{3}\omega^{\pm 1/2}x \pm \beta) + 1]^2}{[\cos(\sqrt{3}\omega^{\pm 1/2}x \pm \beta) - 2]^2} \right\}. \tag{3.29}$$

Since the two potentials are not \mathcal{PT} -symmetric by themselves one may be tempted to conclude already at this stage that there are no real eigenvalue solutions to (3.7). However, all we can deduce with certainty at this point is that the symmetry as defined in (3.5) is broken and will not ensure the reality of λ . We can not guarantee that there are no other types antilinear symmetries that would explain a real spectrum.

Thus let us show explicitly how the eigensystem may be solved in this case. The function ψ in (2.9) is now taken to $\psi^\pm(x, a) = \phi^\pm(x + a\omega^{\mp 1/2})$, from which we easily calculate the zero mode to

$$\Phi_0^\pm(x) = \mp \frac{3^{3/2} \tan \left[\frac{1}{2}(\sqrt{3}\omega^{\pm 1/2}x \pm \beta) \right]}{\cos(\sqrt{3}\omega^{\pm 1/2}x \pm \beta) - 2}. \tag{3.30}$$

We notice that in these solutions there is no need to take β to be purely imaginary as the singularities in the zero mode are already regularized because x is multiplied by a complex constant.

3.2.2. Bound state solutions for type II complex one-soliton solutions

Next we employ relation (2.11) to compute the superpotentials from the zero modes

$$W^\pm(x) = \pm \frac{\sqrt{3}\omega^{\pm 1/2} [3 + \cos [2(\sqrt{3}\omega^{\pm 1/2}x \pm \beta)] - 2 \cos(\sqrt{3}\omega^{\pm 1/2}x \pm \beta)]}{2 [\cos(\sqrt{3}\omega^{\pm 1/2}x \pm \beta) - 2] \sin(\sqrt{3}\omega^{\pm 1/2}x \pm \beta)}, \tag{3.31}$$

from which we calculate the corresponding intertwining operators L_{\pm} in (2.11). The partner potentials are then obtained by a direct calculation from (2.12) to

$$V_2^{\pm} = \frac{3\omega^{\mp 2}}{1 - \sec\left(\sqrt{3}\omega^{\pm 1/2}x \pm \beta\right)} = \frac{3}{2} \frac{\omega^{\pm 1}}{\sin^2\left[\frac{1}{2}(\pm\beta + \sqrt{3}\omega^{\pm 1/2}x)\right]} - 3\omega^{\pm 1}. \tag{3.32}$$

Again, this is a well studied potential corresponding to a special version of the Pöschl-Teller potential [33–36]. Adapting the shift and scaling to the solutions that can be found in the literature, we obtain the eigensystem solutions

$$\begin{aligned} \Phi_0^{(2)\pm} &= N_0 \frac{1}{\sin\left[\frac{1}{2}(\pm\beta + \sqrt{3}\omega^{\pm 1/2}x)\right]}, & E_0^{(2)} &= -\frac{9}{4}\omega^{\pm 1}, \\ \Phi_1^{(1)\pm} &= L_+ \Phi_0^{(2)} \\ &= \frac{N_0}{2} \frac{3^{3/2}\omega^{\pm 1/2} \cos(\pm\beta + \sqrt{3}\omega^{\pm 1/2}x)}{\left[2 - \cos(\pm\beta + \sqrt{3}\omega^{\pm 1/2}x)\right] \sin\left[\frac{1}{2}(\pm\beta + \sqrt{3}\omega^{\pm 1/2}x)\right]}, & E_1^{(1)} &= E_0^{(2)}. \end{aligned} \tag{3.33}$$

The solutions are asymptotically vanishing, $\lim_{x \rightarrow \pm\infty} \Phi_1^{(1)\pm}(x) = 0$ and come in complex conjugate pairs, which is the signature property of a spontaneously broken \mathcal{PT} -symmetry. Instead we have $\mathcal{PT} : \Phi_1^{(1)+}(x) \rightarrow \Phi_1^{(1)-}(x)$. We conclude that the complex soliton solutions of type II become unstable when perturbed by the bound state solutions $\Phi_1^{(1)\pm}$.

3.2.3. Scattering solutions for type II complex one-soliton solutions

Finally we discuss the scattering solutions for the auxiliary Sturm-Liouville eigenvalue equation obtained from the type II complex one-soliton solutions. By similar means as in the previous section we exploit the fact that the potential becomes a shifted and scaled inverse sin-squared potential. For the V_1^{\pm} -potential we then find the two linearly independent scattering solutions

$$\Psi_e^{(2)\pm} = g^{\pm}(x) \sin(\kappa x) + h^{\pm}(x) \cos(\kappa x), \tag{3.34}$$

$$\Psi_o^{(2)\pm} = g^{\pm}(x) \cos(\kappa x) + h^{\pm}(x) \sin(\kappa x), \tag{3.35}$$

with

$$g^{\pm}(x) = \pm \frac{3^{3/2}\omega^{\pm 1/2}\kappa \cos(\beta + \sqrt{3}\omega^{\pm 1/2}x) \tan\left[\frac{1}{2}(\beta + \sqrt{3}\omega^{\pm 1/2}x)\right]}{2 \cos(\beta + \sqrt{3}\omega^{\pm 1/2}x) - 2} \tag{3.36}$$

$$h^{\pm}(x) = \frac{3}{2}\omega^{\pm 1} \left[\frac{3}{2 - \cos(\beta + \sqrt{3}\omega^{\pm 1/2}x)} - 1 \right] - \kappa^2 \tag{3.37}$$

for the complex continuous eigenvalues $(\lambda^{\pm})^2 = \kappa^2 - 3\omega^{\pm}$. One may be tempted to choose the free parameter κ in such a way that $(\lambda^{\pm})^2$ becomes real. However, when κ acquired a complex part the solutions are easily seen to diverge at $x \rightarrow \pm\infty$. Thus overall we conclude that the complex type II one-solitons solutions become unstable when linearly perturbed with these scattering solutions $\Psi_{e,o}^{(2)\pm}$.

4. Conclusions

We investigated several types of complex one-soliton solutions in the Bullough-Dodd model with regard to their stability when linearly perturbed. The type I solutions were previously known, whereas the type II solutions have been newly constructed. The overall energy for both types of solutions were found to be real when shifting the vacuum of the potential appropriately.

We found that the auxiliary Sturm-Liouville eigenvalue equation of the stability analysis for the type I solutions can be supersymmetrically mapped to an isospectral partner system involving a shifted and scaled inverse cosh-squared potential. Remarkably these potentials also emerge in the stability analysis of the ϕ^4 -theory and sine-Gordon model, but simply with different scalings and overall shifts [20]. We concluded that the type I solutions remain stable when linearly perturbed with bound state and scattering solutions with vanishing continuous parameter.

For the type II solutions the original auxiliary Sturm-Liouville eigenvalue equation can be supersymmetrically mapped to a complex shifted and scaled inverse sin-squared potential. The corresponding bound state and scattering solutions for these equations have complex eigenvalues or are asymptotically divergent, such when used to perturb the type II solutions they become unstable or the energies become infinite.

The underlying \mathcal{PT} -symmetry serves for several purposes: i) When introducing complex rather than real shifts the singularities in the solutions are regularized. ii) As previously pointed out, it governs the reality of the energy of the solutions, but as we found here the set criteria provided in [14] may still be enlarged. The energies of two \mathcal{PT} -symmetrically related degenerate solutions are guaranteed to be real even when they solve the Euler-Lagrange equations of two \mathcal{PT} -symmetrically related Hamiltonians that might even be different. iii) As we have seen here the \mathcal{PT} -symmetry also governs the stability of the solutions. However, in order to obtain stable solutions, that is real eigenvalues in the auxiliary Sturm-Liouville equation, the perturbation must be \mathcal{PT} -symmetric by themselves.

Here we observed that the \mathcal{PT} -symmetry of the soliton solution φ was inherited by the perturbing field χ . The type I solutions and their perturbing fields were found to be \mathcal{PT} -symmetry by themselves, which guaranteed the reality of the energy and their stability. On the other hand the type II solutions were mapped into different degenerate solutions, so that the reality of their energy was still ensured. However, on the level auxiliary of the Sturm-Liouville equation the \mathcal{PT} is broken for a specific solution so that the eigenvalues become complex and the solutions unstable.

While the Bullough-Dodd model appears to unavoidably lead to complex soliton solutions, our finding here also shed more light on theories that have both, complex and real solutions. We refined the criteria under which the measurable physical quantities are guaranteed to be real. In addition, we found here that such complex type of solutions are also physical in the sense of being stable and not artificial artefacts with zero measure. In the corresponding quantum field theories we expect such solutions to play similar roles as complex scalar fields in the description of charged particles, such as Higgs field of the Standard Model or charged pions.

CRedit authorship contribution statement

All authors contributed in equal terms.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

FC was partially supported by FONDECYT grant 1211356. TT is supported by EPSRC grant EP/W522351/1.

References

- [1] C.M. Bender, P.E. Dorey, C. Dunning, A. Fring, D.W. Hook, H.F. Jones, S. Kuzhel, G. Levai, R. Tateo, *PT Symmetry: In Quantum and Classical Physics*, World Scientific, Singapore, 2019.
- [2] R. El-Ganainy, K.G. Makris, M. Khajavikhan, Z.H. Musslimani, S. Rotter, D.N. Christodoulides, Non-Hermitian physics and \mathcal{PT} symmetry, *Nat. Phys.* 14 (2018) 11.
- [3] A. Fring, \mathcal{PT} -symmetric deformations of the Korteweg-de Vries equation, *J. Phys. A, Math. Theor.* 40 (2007) 4215–4224.
- [4] M. Wadati, Construction of parity-time symmetric potential through the soliton theory, *J. Phys. Soc. Jpn.* 70 (2008) 074005.
- [5] P.E.G. Assis, A. Fring, Compactons versus solitons, *Pramana J. Phys.* 74 (2010) 857–865.
- [6] A. Cavaglia, A. Fring, B. Bagchi, \mathcal{PT} -symmetry breaking in complex nonlinear wave equations and their deformations, *J. Phys. A, Math. Theor.* 44 (2011) 325201.
- [7] J. Cen, A. Fring, Complex solitons with real energies, *J. Phys. A, Math. Theor.* 49 (2016) 365202.
- [8] F. Correa, A. Fring, Regularized degenerate multi-solitons, *J. High Energy Phys.* 2016 (9) (2016) 1–16.
- [9] A. Khare, A. Saxena, Novel \mathcal{PT} -invariant solutions for a large number of real nonlinear equations, *Phys. Lett. A* 380 (2016) 856–862.
- [10] M.J. Ablowitz, Z.H. Musslimani, Integrable nonlocal nonlinear Schrödinger equation, *Phys. Rev. Lett.* 110 (2013) 064105.
- [11] J. Cen, F. Correa, A. Fring, Integrable nonlocal Hirota equations, *J. Math. Phys.* 60 (2019) 081508.
- [12] J. Cen, Nonlinear classical and quantum integrable systems with \mathcal{PT} -symmetries, PhD thesis, City, University of London, City Research Online digital archive, 2019.
- [13] A. Fring, T. Taira, Complex BPS solitons with real energies from duality, *J. Phys. A, Math. Theor.* 53 (2020) 455701.
- [14] F. Correa, A. Fring, T. Taira, Complex BPS Skyrmions with real energy, *Nucl. Phys. B* 971 (2021) 115516.
- [15] R. Dodd, R. Bullough, Polynomial conserved densities for the sine-Gordon equations, *Proc. R. Soc. Lond. Ser. A, Math. Phys. Sci.* 352 (1671) (1977) 481–503.
- [16] A.V. Zhiber, A.B. Shabat, The Klein–Gordon equation with nontrivial group, *Dokl. Akad. Nauk* 247 (1979) 1103–1107.
- [17] I. Takyi, H. Weigel, Collective coordinates in one-dimensional soliton models revisited, *Phys. Rev. D* 94 (2016) 085008.
- [18] N.S. Manton, K. Oleś, T. Romańczukiewicz, A. Wereszczyński, Kink moduli spaces: collective coordinates reconsidered, *Phys. Rev. D* 103 (2021) 025024.
- [19] F. Correa, A. Fring, T. Taira, in preparation.
- [20] R. Jackiw, Quantum meaning of classical field theory, *Rev. Mod. Phys.* 49 (1977) 681.
- [21] E.J. Weinberg, Parameter counting for multimonopole solutions, *Phys. Rev. D* 20 (1979) 936.
- [22] S.K. Kim, K.S. Soh, J.H. Yee, Index theory for the nonrelativistic Chern-Simons solitons, *Phys. Rev. D* 42 (1990) 4139.
- [23] E. Witten, Dynamical breaking of supersymmetry, *Nucl. Phys. B* 188 (1981) 513.
- [24] F. Cooper, B. Freedman, Aspects of supersymmetric quantum mechanics, *Ann. Phys.* 146 (1983) 262.
- [25] E. Witten, Constraints on supersymmetry breaking, *Nucl. Phys. B* 202 (1982) 253.
- [26] F. Cooper, A. Khare, U. Sukhatme, Supersymmetry and quantum mechanics, *Phys. Rep.* 251 (1995) 267–385.
- [27] V.A. Andreev, Bäcklund transformations of the Bullough–Dodd–Zhiber–Shabat equation and symmetries of integrable equations, *Theor. Math. Phys.* 79 (1989) 448–450.

- [28] A. Fring, G. Mussardo, P. Simonetti, Form-factors of the elementary field in the Bullough-Dodd model, *Phys. Lett. B* 307 (1993) 83–90.
- [29] R. Hirota, *The Direct Method in Soliton Theory*, vol. 155, Cambridge University Press, 2004.
- [30] P.E.G. Assis, L.A. Ferreira, The Bullough–Dodd model coupled to matter fields, *Nucl. Phys. B* 800 (2008) 409–449.
- [31] L.D. Landau, E.M. Lifschitz, *Quantum Mechanics: Non-relativistic Theory*, second ed., Pergamon Press, Oxford, 1965.
- [32] J. Lekner, Reflectionless eigenstates of the sech^2 potential, *Am. J. Phys.* 75 (2007) 1151–1157.
- [33] S. Flügge, *Practical Quantum Mechanics*, Springer Science & Business Media, 2012.
- [34] G. Pöschl, E. Teller, Bemerkungen zur Quantenmechanik des anharmonischen Oszillators, *Phys. Rev.* 85 (1952) 166–179.
- [35] F. Correa, M.S. Plyushchay, Spectral singularities in \mathcal{PT} -symmetric periodic finite-gap systems, *Phys. Rev. D* 86 (2012) 085028.
- [36] S. Dey, A. Fring, Bohmian quantum trajectories from coherent states, *Phys. Rev. A* 88 (2013) 022116.