



City Research Online

City, University of London Institutional Repository

Citation: Dong, H., Farnaud, S. & Renshaw, D. (2022). Influence of Vitamin D Supplement on Immune Function of the Older Healthy People: A Pilot Randomised Control Trial. *Current Developments in Nutrition*, 6(S1), p. 976. doi: 10.1093/cdn/nzac068.005

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/28462/>

Link to published version: <https://doi.org/10.1093/cdn/nzac068.005>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Influence of Vitamin D Supplement on Immune Function of the Older Healthy People: A Pilot Randomised Control Trial

Honglin Dong,¹ Sebastien Farnaud,² and Derek Renshaw²

¹City, University of London and ²Coventry University

Objectives: The study aims to investigate the influence of vitamin D (VD) supplement on immune function of the healthy older people.

Methods: Designed as a randomized control trial, 21 participants (55–85) years old took part and completed the study (20 White, 1 Indian) during May–November 2018 in Coventry, England. The participants were randomized into two groups, VD and control, stratified by age, gender and body mass index (BMI). The VD group (n = 11) took VD3 of 25 $\mu\text{g}/\text{day}$ for 12 weeks, while the control group (n = 9) were only provided with educational information of VD. At baseline, 6 weeks and 12 weeks, plasma 25(OH)D levels and immune parameters including phagocytic activity of granulocytes and monocytes, tumour necrosis factor (TNF), interleukin 6 (IL-6), lymphocyte subsets (Th, Tc and natural killer cells), and fast blood glucose and lipid were measured. Dietary VD intake was analysed at baseline and week 12. The data were presented as mean \pm standard

error (SE) and analysed by two-way repeated measures ANOVA.

Results: The average age of the participants was (65.8 ± 1.7) years old (67.4 ± 2.7 in VD group and 64.5 ± 2.2 in control group). At baseline, 42.9% of the participants were VD deficiency (25(OH)D < 30 nmol/L), only 10% achieved a level of 25(OH)D > 50 nmol/L. There was a significant increase in plasma 25(OH)D concentration in VD group compared with that in the control group (visit*treatment effect $p = 0.002$). The level of 25(OH)D in the VD group was increased rapidly at 6 weeks (from 38.4 ± 8.7 nmol/L at baseline to 51.0 ± 9.3 nmol/L) and little change has been observed at 12 weeks (51.8 ± 8.6 nmol/L). There was a significant visit*treatment effect ($p = 0.036$) on the plasma creatinine concentration between two groups, indicating a decrease of plasma creatinine concentration in VD group (from 79.8 ± 7.0 $\mu\text{mol}/\text{L}$ at baseline to 75.1 ± 5.4 $\mu\text{mol}/\text{L}$ at week 12). There was a significant correlation between body mass index (BMI) and 25(OH)D concentration ($P < 0.001$). There was no significant effect of VD supplementation on parameters of immune function.

Conclusions: VD deficiency is common among this older population even during the summertime. VD supplementation at 25 $\mu\text{g}/\text{day}$ for 12 weeks showed an anti-inflammatory effect.

Funding Sources: Coventry University.