



City Research Online

City, University of London Institutional Repository

Citation: Madruga, M., Martínez Steele, E., Reynolds, C., Levy, R. B. & Rauber, F. (2022). Trends in food consumption according to the degree of food processing among the UK population over 11 years. *British Journal of Nutrition*, 130(3), pp. 476-483. doi: 10.1017/s0007114522003361

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/29036/>

Link to published version: <https://doi.org/10.1017/s0007114522003361>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

Trends in food consumption according to the degree of food processing among the UK population over 11 years

Mariana Madruga^{1,2}, Eurídice Martínez Steele^{2,3}, Christian Reynolds⁴, Renata Bertazzi Levy^{1,2}, Fernanda Rauber^{1,2,3}

¹Department of Preventive Medicine, School of Medicine, University of São Paulo, São Paulo, Brazil.

²Center for Epidemiological Research in Nutrition and Health, University of São Paulo, São Paulo, Brazil.

³Department of Nutrition, School of Public Health, University of São Paulo, São Paulo, Brazil.

⁴Centre for Food Policy, City, University of London, London, United Kingdom.

Corresponding author: mariana.madruga@gmail.com

Mariana Madruga <https://orcid.org/0000-0002-9141-8079>

Eurídice Martínez Steele <https://orcid.org/0000-0002-2907-3153>

Christian Reynolds <https://orcid.org/0000-0002-1073-7394>

Renata Bertazzi Levy <https://orcid.org/0000-0001-5388-7002>

Fernanda Rauber <https://orcid.org/0000-0001-9693-7954>

Short title: Food consumption trend in the UK population



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114522003361

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

Abstract

Although ultra-processed foods represent more than half of the total energy consumed by the UK population, little is known about the trend in food consumption considering the degree of food processing. We evaluated the trends of the dietary share of foods categorized according to the NOVA classification in a historical series (2008-2019) among the UK population. Data was acquired from the NDNS, a national survey that collects diet information through a 4-day food record. We used adjusted linear regression to estimate the dietary participation of NOVA groups and subgroups and evaluated the linear trends over the years. From 2008-2019, we observed a significant increase in the energy share of culinary ingredients (from 3.7 to 4.9% of the total energy consumed; p -trend<0.001), especially for butter and oils; and reduction of processed foods (from 9.6 to 8.6%; p -trend=0.002), especially for beer and wine. Unprocessed or minimally processed foods (\cong 30%, p -trend=0.505) and ultra-processed foods (\cong 56%, p -trend=0.580) presented no significant change. However, changes in the consumption of some subgroups are noteworthy, such as the reduction in the energy share of red meat, sausages and other reconstituted meat products as well as the increase of fruits, ready meals, breakfast cereals, cookies, pastries, buns and cakes. Regarding the sociodemographic characteristics, no interaction was observed with the trend of the four NOVA groups. From 2008 to 2019 was observed a significant increase in culinary ingredients and a reduction in processed food. Furthermore, it sheds light on the high share of ultra-processed foods in the contemporary British diet.

Keywords: National Diet and Nutrition Survey, food consumption, food processing, United Kingdom

Introduction:

The global increase in obesity and Chronic Non-Communicable Diseases (NCDs) is associated with changes in population dietary habits (1–5). These changes are characterized by the replacement of traditional meals based on fresh foods and culinary ingredients with ultra-processed foods (6–8).

Among various food classification systems, the NOVA classification developed by Monteiro et al. has become the most widely used in research and policy (9). NOVA classifies foods into four groups based on the degree and purpose of their industrial processing: Group 1 - Unprocessed or minimally processed foods, such as whole grains and cereals, legumes, fruits, vegetables, eggs, and meats; Group 2 - Processed culinary ingredients, such as salt, sugar, oils, and fats; Group 3 - Processed foods, such as vegetables and legumes preserved in brine, cheeses, and bread; and Group 4 – Ultra-processed foods, such as soft drinks, ready-to-eat or semi-ready-to-eat meals, prepared sauces, cured meats, breakfast cereals, packaged snacks, and candy (10). Ultra-processed foods have an unfavorable nutritional profile that negatively affects the nutritional quality of foods (11–16), with significant consequences for health (17–20) and the environment (21).

The availability and sale of ultra-processed foods have boomed in high-income countries (1970-2000) (22) and, more recently, in middle-income countries (2000-2013) (23). In a 2019 analysis of global ultra-processed food trends, the United Kingdom had the third-highest sales volume of ultra-processed food per capita (140.7 kg/capita/year) compared with 80 high- and middle-income countries (24). Previous analyzes of the National Diet and Nutrition Survey (NDNS) show that ultra-processed foods already account for more than half of the total food energy consumed by the UK population (25). However, as far as we know, there are no trend studies that have examined the changes in actual food consumption considering industrial processing and the differences between sociodemographic strata in the United Kingdom. In this sense, this study aims to assess the trend of food consumption in the United Kingdom from 2008 to 2019, considering the extent and purpose of industrial food processing, and to analyze how these trends vary according to sociodemographic characteristics.

2.0 Methods

2.1 Data Source and Sample

Data was acquired from the National Diet and Nutrition Survey (NDNS), a national survey conducted in the United Kingdom (England, Wales, Scotland, and Northern Ireland). This annual survey has used a standardized methodology since 2008 enabling combined analysis (26). The 1-11 survey years correspond to the years: 1-2008/09, 2-2009/10, 3-2010/2011, 4-2011/12, 5-2012/2013, 6-2013/2014, 7-2014/2015, 8-2015/2016, 9-2016/2017, 10-2017/2018, and 11-2018/2019. The purpose of the NDNS is to collect detailed information about the food consumption of the UK population. Data collection takes place monthly and considers possible seasonal variations in food consumption (27). The survey samples were randomly drawn from the UK Postcode Address Archive, which contains a list of all addresses in the four constituent countries. One child (from 1.5 to 18 years) or one child together with an adult (19 years or older) were selected from each of the randomly selected addresses (28). Data collection included an interview with the researcher to collect sociodemographic and food consumption data as well as a visit with a nurse (27). Food consumption was assessed using a food diary completed by participants on four consecutive days. Everyone who completed the diary on 3 or 4 days was included in the survey, totaling 15,643 participants.

Data and files used in this study were acquired under license from the UK Data Archive found at <http://www.esds.ac.uk>. The study was approved by the ethics committees of each of the four countries participating in the study.

2.2 Food consumption

Food consumption was assessed using four-day food diaries that included workdays and weekends, thus covering all days of the week. Participants were instructed to record all food and beverages consumed inside and outside the home on that day. Portion sizes were estimated using household measures or weights from packing. After completion, diaries were checked by trained interviewers to improve accuracy. Food consumption data from the complete records were coded and processed using the program DINO (Diet In, Nutrients Out) (29), and energy and nutrient intakes were estimated using the NDNS food nutrient composition table (30).

2.3 Covariates

The sociodemographic variables included in this study were: sex (female and male), age group (1–3 years, 4–10 years, 11–18 years, 19–64 years, 65+ years), region (southern England, central England, northern England, Scotland, Wales, and Northern Ireland), ethnicity (white, mixed ethnic group, black, Asian, and other ethnicities), and occupational social class. Occupational social class was categorized as high (High and lower managerial & professional occupations), intermediate (Intermediate occupations, Small employers and own account workers, Lower supervisory and technical occupations), and low (routine & manual occupations).

2.4 Food classification according to processing

All food items were classified according to NOVA (9), a food classification system based on the nature, extent, and purpose of the industrial processing they undergo before consumption. This classification includes four categories (see box 1).

All foods presented in the NDNS database are coded as food number and grouped into subsidiary food groups (n=155). When possible, the subsidiary food groups were classified directly according to NOVA. When foods within a subsidiary food group belonged to different NOVA groups (n=52), it was the food codes instead of the group, which were classified individually. Hence, it was possible to allocate each underlying ingredient of homemade dishes into the appropriate NOVA group. Food classification details can be found in a previously published article (25).

2.5 Data analysis

For each survey day and age group, we defined values for daily energy intake below 1 percentile and above 99 percentiles as outliers. Based on that, 12 individuals were excluded (n=15,643) and more than 90% completed the 4-day food diary. The mean of all available days of food diary was used for each person.

We estimated the distribution of daily energy intake (% of total energy) according to NOVA groups and subgroups (1. unprocessed or minimally processed=13 subgroups; 2. processed culinary ingredients=4 subgroups, 3. processed foods=6 subgroups, 4. ultra-processed foods=16 subgroups) for each survey year. Linear regression analysis was used to assess how this distribution of NOVA groups and subgroups (% of total energy) varied across the 11 years studied (including year as an ordinal variable). Models were adjusted for the covariates

of sex, age group, region, ethnicity, and occupational social class. In order to estimate the main changes over the period, the difference in the mean energy share (% of total energy) of the extreme years (year 11 (2019) minus year 1 (2008)) of the NOVA subgroups was calculated.

To examine potential differences in energy share trends (% of total energy) by population subgroups, we evaluated an interaction term between the survey year (as an ordinal variable) and each sociodemographic characteristic using the Wald F test.

All analyses were also performed using the contribution of NOVA groups and subgroups to total grams intake (% of total grams).

NDNS study weights were used in all analyses to account for sampling and non-response error. Data analysis was performed using STATA software version 16.1. We considered a p value < 0.05 to test statistical significance.

3.0 Results

Table 1 presents the evolution of the dietary share of NOVA food groups and subgroups (% of total energy) over the 11 years of the NDNS survey (2008-2019). The dietary share of unprocessed or minimally processed foods (group 1) remained similar from 2008 to 2019 ($\cong 30\%$ of total energy, p for linear trend = 0.505). Processed culinary ingredients (group 2) showed an increase from 3.7% in 2008 to 4.9% in 2019 (p for linear trend < 0.001) and processed foods (group 3) a decrease from 9.6% to 8.6%, respectively (p for linear trend = 0.002). No changes were observed in the proportion of ultra-processed foods (group 4), which accounted for more than half of total energy consumed throughout the period ($\cong 56\%$ of total energy, p for linear trend = 0.580).

Regarding unprocessed or minimally processed food subgroups (group 1), there was a significant decrease in the proportion of diet (% of total energy) from 2008 to 2019 for roots and tubers (from 3.9% to 2.7%), for red meat (from 4.0% to 2.2%), and for 100% fruit juice (from 1.2% to 0.8%). In addition, an increase in the dietary contribution was observed for fruits (from 3.4% in 2008 to 3.7% in 2019), for whole grains and cereals (from 2.3% to 3.2%), for pasta (from 1.6% to 1.9%), for eggs (from 1.5% to 1.9%), and for legumes (from 0.5% to 0.9%).

Processed culinary ingredient subgroups (group 2) intake varied from 2008 to 2019. The energy share of table sugar decreased significantly from 1.4% to 1.2%, while butter and plant oil intake increased significantly from 1.4% to 2.2% and from 0.5% to 1.1%, respectively.

Among processed food subgroups (group 3), there was a significant decrease in the energy share of beer and wine from 4.1% to 3.1%, vegetables and legumes preserved in brine from 1.1% to 0.9%, and an increase in other processed foods from 0.3% to 0.5% during the same period.

Among ultra-processed food subgroups (group 4), a significant decrease was observed in the energy share of sausages and other reconstituted meat products (from 4.2% to 3.7%), soft drinks (from 2.5% to 1.1%), and margarine (from 2.4% to 1.6%). On the other hand, significant increase in energy share of packaged pre-prepared meals (from 7.4% to 8.3%), breakfast cereals (from 4.2% to 4.9%), cookies (from 3.2% to 4.2%), pastries, buns and cakes (from 2.9% to 3.7%), packaged salty snacks (from 1.9% to 2.3%) sauces, dressing and gravies (from 2.0% to 2.2%).

The significant differences in the mean energy share of the NOVA subgroups between extreme years, reflecting the entire period studied in the UK population, are shown in Figure 1. For unprocessed or minimally processed foods, the largest differences were observed for red meat (-1.8%), roots and tubers (-1.2%), and whole grains and cereals (0.9%). Among processed culinary ingredients, the differences between butter (0.8%) and plant oil (0.6%) stood out. In processed foods, beer and wine (-1.0%) presented the largest differences. Regarding ultra-processed foods, soft drinks (-1.4%), margarine (-0.8%), sausages and other reconstituted meat products (-0.5%), breakfast cereals (0.7%), pastries, buns and cakes (0.8%), packaged pre-prepared meals (0.9%), and cookies (1.0%) differed the most.

No interaction was observed for the trend of energy share of the four NOVA groups when considering the sociodemographic characteristics.

The trend in the share of NOVA food groups and subgroups (% of total grams) over the 11 years is shown in Supplementary Table 1. The trend in % gram consumption observed in the groups evolved according to the % of energy for processed culinary ingredients (increase from 0.5% in 2008 to 0.6% in 2019, p for linear trend=0.043); processed foods (decreased from 7.7% to 5.5%, p for linear trend < 0.001), with the decrease in consumption of beer and wine being noteworthy (from 6.0% to 3.9%, p for linear trend < 0.001); and ultra-processed foods (remained \cong 29%; p for linear trend=0.090), with the decreased consumption of milk-

based drinks subgroup to be highlighted (from 3.2% to 2.8%, p for linear trend=0.005). There was a significant increase in gram share of foods grouped as unprocessed or minimally processed (from 61.9% in 2008 to 65.2% in 2019, p for linear trend < 0.001), with emphasis on the increase in the water subgroup (from 14.3% to 21.9%, p for linear trend < 0.001) and decreases in the milk and yogurt subgroups (from 7.4% to 6.4%, p for linear trend < 0.001) and fresh fruit juice (from 4.8% to 3.4%, p for linear trend < 0.001).

Discussion

By analyzing the representative data of the UK population from 2008 to 2019, we found that the trend of energy share of culinary ingredients, particularly butter and oils, has increased significantly. Moreover, processed foods, particularly beer and wine and vegetables and legumes preserved in brine, have decreased significantly. The energy share of unprocessed or minimally processed foods and ultra-processed foods did not change significantly during this period. While the energy share of roots and tubers, red meat, and fresh fruit juice in the diet tended to decline, the energy share of fruits, whole grains and cereals, pasta, eggs, and legumes augmented. Among ultra-processed foods, the energy share of packaged pre-prepared meals, breakfast cereals, cookies, pastries, buns and cakes, packaged salty snacks, and sauces, dressing and gravies sauces increased, while the share of sausage and other reconstituted meat products, sweetened beverages, and margarine decreased over the entire period.

The previously described trends in the energy share of Nova groups in the UK population did not vary according to sociodemographic characteristics. A study carried out in a nationally representative sample of US adults, also observed an increase in the energy contribution of processed culinary ingredients from 2001 to 2018 that did not differ by age group, education or income level but increased the most among non-Hispanic blacks, followed by Hispanics and non-Hispanic whites. Among US adults, however, a borderline significant trend toward increased contribution of processed foods that did not differ by age group was observed. A trend toward increased consumption of processed foods was observed only among females, among non-Hispanic whites, and among individuals with a lower household income level. This study observed a decrease in the contribution of unprocessed/minimally processed foods, that was similar among both sexes, among all races/ethnicities, across all education and income levels, decreasing the most among adults aged 60 y and older followed by middle-aged adults and younger adults. Over this period, an increase in the contribution of

ultra-processed foods was observed that was similar across all age groups, education and income levels and that increased to a greater extent among males than among females. This increase was only confirmed among non-Hispanic whites and non-Hispanic blacks (34).

As among US adults, a study carried out in a nationally representative sample of US youths between 1999 and 2018, also observed an increase in the energy contribution of processed culinary ingredients, a non-significant change in the contribution of processed foods, a decrease in the contribution of unprocessed/minimally processed foods and an increase in the contribution of ultra-processed foods. A similar increase in the contribution of ultra-processed foods was observed across all sex, age, education and income level youth population subgroups; however, the increase observed among non-Hispanic Black youths and Mexican American youths was significantly higher than the increase among non-Hispanic White youths (35).

The replacement of unprocessed or minimally processed foods and culinary preparations by ultra-processed foods is occurring worldwide. In the United Kingdom, this substitution can be said to have already taken place and reached a plateau where the overall increase in the ultra-processed food group is no longer observed. Ultra-processed foods have become a significant source, and in some cases the main source, of dietary energy, in high-income countries, including the United Kingdom (31,32). While sales of these products appear to be stagnant in high-income countries (due to market saturation), the sales of ultra-processed foods have grown more rapidly in middle-income countries (22,33).

From 2000 to 2016, an analysis of food sales in 80 countries highlighted the global trend of increasing ultra-processed food sales worldwide, except Western Europe, North America, and Australasia. The United Kingdom had the third-highest volume of ultra-processed food sales (140.7 kg/person), trailing only the Netherlands (143.8 kg/person) and Germany (141.8 kg/person) (24). Our findings are consistent with these studies on global sales trends, highlighting that ultra-processed food consumption in the UK has remained high in recent years (57%).

Studies using household budget surveys conducted in Mexico (from 1984 to 2016) and Canada (from 1938 to 2011) have also shown that there has been a gradual decline in unprocessed or minimally processed foods and culinary ingredients — characterizing culinary preparations — and an increase in ultra-processed food in recent decades (6,8). In the United States, trend analyses over the past two decades (2000 to 2018) have shown an increase in the

consumption of ultra-processed foods and culinary ingredients accompanied by a decrease in the consumption of minimally processed foods among adults and children (34,35). In our study, targeting the recent period 2008-2019, we observed an increase in the energy share of processed culinary ingredients among the UK population due to increased consumption of butter and plant oils substituting margarine, which presented a decline over time. This replacement may result from the fact that margarines which are produced through intensive processing and chemical transformation of vegetable oils, contain trans-fats, which are more harmful than saturated fats (36). We also observed the maintenance of the energy share of unprocessed or minimally processed foods in the British diet, with increases in some subgroups (whole grains and cereals, pasta, eggs, and legumes) and decreases in others (roots and tubers, red meat, and fresh fruit juice), which may indicate a change in the pattern of British culinary preparations during this period.

The decrease in consumption of fresh red and ultra-processed meats, such as sausages and other reconstituted meat products, observed in this study is consistent with current recommendations to reduce consumption of these foods. In the United Kingdom, a review of dietary recommendations by the Scientific Advisory Committee on Nutrition (SACN) in 2011 recommended the reduction of red and processed meat consumption by adults from $\cong 90\text{g/day}$ or more to $\cong 70\text{g/day}$ (26). The report “The Global Syndemic of Obesity, Malnutrition and Climate Change”, published in The Lancet Commission, demonstrates that red meat and ultra-processed foods are among the major contributors to the global syndemic (21).

Our results also showed that despite the reducing energy share of table sugar as a culinary ingredient, there was an increase in the consumption of ultra-processed foods rich in sugars, such as breakfast cereals, biscuits, pastries, buns and cakes. There is already a consensus that high consumption of free sugars contributes to obesity, type 2 diabetes, dyslipidemia, hypertension, and coronary heart disease (37–39) and, consequently, most dietary recommendations limit their intake. However, more focused efforts are needed to put this recommendation into practice as most free sugar is added to foods before they are marketed and sold. A study conducted with data from the national survey showed that ultra-processed foods were the ones that contributed the most to sugar consumption in the UK and the elimination of these products could reduce the prevalence of excessive free sugar intake by 47% (15). Initial stages of the UK government's sugar reduction program, which challenged

the food industry to voluntarily reduce the sugar from some products, produced only slow progress towards proposed targets (40).

Energy share reduction of sweetened beverages observed in the period occurred despite no change in the volume (in grams) of these beverages. That may be due to reductions in the sugar concentration of sweetened beverages associated with the UK sugar reduction program and the taxation initiative of the Soft Drink Industry Levy-SDIL launched in 2015 (41,42). The SDIL was successful in reducing sugar from soft drinks, when compared to that seen for the food categories included in the sugar reduction program (40). However, when targeting sugar, the SDIL has hastily contributed to increases in the consumption of drinks with artificial sweeteners (43), which have also been associated with adverse health effects (44,45). Regarding the contribution of NOVA groups and subgroups to the total grams, an increase of unprocessed or minimally processed foods was observed, unlike the energy share, which is due, at least in part, to the increase in water consumption. The increase in water consumption was accompanied by a decrease in the gram contribution of each one of the remaining beverages, regardless of the degree of processing (milk and plain yogurt, fresh fruit juice, beer and wine and milk-based drinks). This trend of replacing drinks with water in a country like the United Kingdom, with high consumption of ultra-processed foods, is positive taking into account studies assessing an inverse association between the energy share of ultra-processed foods and water intake (46).

Our study pioneers analyzing trends of food consumption considering the degree of industrial processing among the UK population. In addition, we use data sourced from the NDNS, which uses a high-quality dietary assessment method that provides a detailed analysis of different foods consumed, several assessment days, and considers the daily variability of each individual. However, some potential limitations are also noteworthy. The data used was self-reported and may be subject to recording errors. Although food records are recognized as one of the most comprehensive methods for assessing dietary intake, a limitation of this method is the underreporting of some foods, particularly unhealthy foods. Although previous study had shown the mean daily energy intake significant decrease of 164 kJ (39 kcal) between days 1 and 4, the size of the effect was relatively minor (47). The underreporting of ultra-processed foods may have increased over time, considering the growing awareness of the harmful effects of these products on health. However, NDNS data are accurate and validated through painstaking collection methods, with review by trained personnel which helps to minimize misreporting. Although NDNS collects limited information indicative of food processing

(i.e., meal location and product brands), the data is not consistently determined for all food items, which can lead to a food classification error. However, standardized NDNS methods minimize potential error and bias.

Food consumption trends according to the degree of food processing highlight a significant increase in processed culinary ingredients and a reduction in the consumption of processed food in the British diet from 2008 to 2019. Furthermore, it sheds light on the high share of ultra-processed foods in the contemporary British diet.

Conflicts of interest: The authors declare there is no conflict of interest.

Authorship: Planning and analysis were performed by Mariana Ferreira Madruga, Fernanda Rauber, Euri Martínez Steele and Renata Bertazzi Levy. The first draft of the manuscript was written by Mariana Ferreira Madruga and Fernanda Rauber. All authors commented on previous versions of the manuscript. All authors have read and approved the final manuscript.

Financial support: This work was supported by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). CAPES had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References:

1. Rauber F, Martínez Steele E, Louzada ML da C, Millett C, Monteiro CA, Levy RB. Ultra-processed food consumption and indicators of obesity in the United Kingdom population (2008-2016). Meyre D, editor. *PLoS One*. 2020 May 1;15(5):e0232676.
2. Martínez Steele E, Juul F, Neri D, Rauber F, Monteiro CA. Dietary share of ultra-processed foods and metabolic syndrome in the US adult population. *Prev Med (Baltim)* [Internet]. 2019 Aug 1;125(May):40–8. Available from: <https://doi.org/10.1016/j.ypmed.2019.05.004>
3. Hall KD, Ayuketah A, Brychta R, Cai H, Cassimatis T, Chen KY, et al. Ultra-Processed Diets Cause Excess Calorie Intake and Weight Gain: An Inpatient Randomized Controlled Trial of Ad Libitum Food Intake. *Cell Metab*. 2019;30(1):67-77.e3.
4. Levy RB, Rauber F, Chang K, Louzada ML da C, Monteiro CA, Millett C, et al. Ultra-processed food consumption and type 2 diabetes incidence: A prospective cohort study. *Clin Nutr* [Internet]. 2021;40(5):3608–14. Available from: <https://doi.org/10.1016/j.clnu.2020.12.018>
5. Chang K, Khandpur N, Neri D, Touvier M, Huybrechts I, Millett C, et al. Association between Childhood Consumption of Ultraprocessed Food and Adiposity Trajectories in the Avon Longitudinal Study of Parents and Children Birth Cohort. *JAMA Pediatr*. 2021;1–11.
6. Moubarac JC, Batal M, Martins APB, Claro R, Levy RB, Cannon G, et al. Processed and ultra-processed food products: Consumption trends in Canada from 1938 to 2011. *Can J Diet Pract Res*. 2014;75(1):15–21.
7. Martins APB, Levy RB, Claro RM, Moubarac JC, Monteiro CA. Participação crescente de produtos ultraprocessados na dieta brasileira (1987-2009). *Rev Saude Publica*. 2014;47(4):656–65.
8. Marrón-Ponce JA, Tolentino-Mayo L, Hernández-F M, Batis C. Trends in ultra-processed food purchases from 1984 to 2016 in Mexican households. *Nutrients*. 2019 Jan 1;11(1).

9. Monteiro CA, Cannon G, Levy RB, Moubarac JC, Jaime P, Martins AP, Canella D, Louzada ML, Parra D; with Ricardo C, Calixto G, Machado P, Martins C, Martinez E, Baraldi L, Garzillo J SI. Nasce a estrela NOVA. *World Nutr.* 2016;7(7):1–3.
10. Monteiro CA, Cannon G, Levy RB, Moubarac J-C, Louzada ML, Rauber F, et al. Ultra-processed foods: what they are and how to identify them. *Public Health Nutr.* 2019 Apr 12;22(5):936–41.
11. Andrade GC, Julia C, Deschamps V, Srour B, Hercberg S, Kesse-Guyot E, et al. Consumption of ultra-processed food and its association with sociodemographic characteristics and diet quality in a representative sample of French adults. *Nutrients.* 2021;13(2):1–14.
12. Machado PP, Martínez Steele E, Levy RB, Sui Z, Rangan A, Woods J, et al. Ultra-processed foods and recommended intake levels of nutrients linked to non-communicable diseases in Australia: Evidence from a nationally representative cross-sectional study. *BMJ Open.* 2019;9(8):e029544.
13. Martínez Steele E, Baraldi LG, Da Costa Louzada ML, Moubarac JC, Mozaffarian D, Monteiro CA. Ultra-processed foods and added sugars in the US diet: Evidence from a nationally representative cross-sectional study. *BMJ Open.* 2016;6(e009892).
14. Rauber F, Louzada MLDC, Martínez Steele E, De Rezende LFM, Millett C, Monteiro CA, et al. Ultra-processed foods and excessive free sugar intake in the UK: A nationally representative cross-sectional study. *BMJ Open.* 2019;9(10):1–11.
15. Da Costa Louzada ML, Ricardo CZ, Martínez Steele E, Levy RB, Cannon G, Monteiro CA. The share of ultra-processed foods determines the overall nutritional quality of diets in Brazil. *Public Health Nutr.* 2017;21(1):94–102.
16. Juul F, Lin Y, Deierlein AL, Vaidean G, Parekh N. Trends in food consumption by degree of processing and diet quality over 17 years: Results from the Framingham Offspring Study. *Br J Nutr.* 2021;2016(12).

17. Adams J, White M. Characterisation of UK diets according to degree of food processing and associations with socio-demographics and obesity: Cross-sectional analysis of UK National Diet and Nutrition Survey (2008-12). *Int J Behav Nutr Phys Act.* 2015;12:160.
18. Rauber F, Chang K, Vamos EP, da Costa Louzada ML, Monteiro CA, Millett C, et al. Ultra-processed food consumption and risk of obesity: a prospective cohort study of UK Biobank. *Eur J Nutr.* 2021;60(4):2169–80.
19. Lane MM, Davis JA, Beattie S, Gómez-Donoso C, Loughman A, O'Neil A, et al. Ultraprocessed food and chronic noncommunicable diseases: A systematic review and meta-analysis of 43 observational studies. *Obes Rev.* 2021;22(3):1–19.
20. Juul F, Vaidean G, Lin Y, Deierlein AL, Parekh N. Ultra-Processed Foods and Incident Cardiovascular Disease in the Framingham Offspring Study. *J Am Coll Cardiol.* 2021;77(12):1520–31.
21. Swinburn BA, Kraak VI, Allender S, Atkins VJ, Baker PI, Bogard JR, et al. The Global Syndemic of Obesity, Undernutrition, and Climate Change: The Lancet Commission report. *Lancet.* 2019 Feb 23;393(10173):791–846.
22. Monteiro CA, Moubarac JC, Cannon G, Ng SW, Popkin B. Ultra-processed products are becoming dominant in the global food system. Vol. 14, *Obesity Reviews.* 2013. p. 21–8.
23. PAHO. Pan American Health Organization (PAHO). Ultra-processed food and drink products in Latin America: Trends, impact on obesity, policy implications [Internet]. Washington D.C.; 2015. Available from: <http://iris.paho.org/xmlui/handle/123456789/7699>
24. Vandevijvere S, Jaacks LM, Monteiro CA, Moubarac J, Girling-Butcher M, Lee AC, et al. Global trends in ultraprocessed food and drink product sales and their association with adult body mass index trajectories. *Obes Rev.* 2019 May 17;20(Suppl 2):10–9.


25. Rauber F, Louzada ML da C, Martínez Steele E, Millett C, Monteiro CA, Levy RB. Ultra-processed food consumption and chronic non-communicable diseases-related dietary nutrient profile in the UK (2008–2014). *Nutrients*. 2018;10(5):587.
26. NDNS. National Diet and Nutrition Survey. Years 1 to 9 of the Rolling Programme (2008/2009 – 2016/2017): Time trend and income analyses. 2019;1–56. Available from: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/772434/NDNS_UK_Y1-9_report.pdf
27. Public Health England. Appendix A Dietary data collection and editing for Year 9 of the NDNS RP. 2017;1–12. Available from: <https://www.gov.uk/government/statistics/ndns-time-trend-and-income-analyses-for-years-1-to-9>
28. Public Health England. Appendix B : Methodology for Years 9 of the NDNS RP. 2018;1–11.
29. Fitt E, Mak TN, Stephen AM, Prymne C, Roberts C, Swan G, et al. Disaggregating composite food codes in the UK National Diet and Nutrition Survey food composition databank. *Eur J Clin Nutr*. 2010;64:S32–6.
30. Public Health England. McCance and Widdowson's The Composition of Foods Integrated Dataset 2019 User guide. London; 2019.
31. Monteiro CA, Cannon G, Moubarac JC, Levy RB, Louzada MLC, Jaime PC. The UN Decade of Nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health Nutr*. 2018;21(1):5–17.
32. FAO. Ultra-processed foods, diet quality, and health using the NOVA classification system. Prepared by Carlos Augusto Monteiro, Geoffrey Cannon, Mark Lawrence, Maria Laura da Costa Louzada, and Priscila Pereira Machado. [Internet]. Roma: Food and Agriculture Organization of the United Nations; 2019. 48 p. Available from: <http://www.wipo.int/amc/en/mediation/rules>
33. Baker P, Friel S. Food systems transformations, ultra-processed food markets and the nutrition transition in Asia. *Global Health* [Internet]. 2016 Dec 3;12(1). Available from: <http://dx.doi.org/10.1186/s12992-016-0223-3>


34. Juul F, Parekh N, Martinez-Steele E, Monteiro CA, Chang VW. Ultra-processed food consumption among US adults from 2001 to 2018. *Am J Clin Nutr*. 2022;115(1):211–21.
35. Wang L, Martínez Steele E, Du M, Pomeranz JL, O'Connor LE, Herrick KA, et al. Trends in Consumption of Ultraprocessed Foods among US Youths Aged 2-19 Years, 1999-2018. *JAMA - J Am Med Assoc*. 2021;326(6):519–30.
36. Hunter JE. Dietary trans fatty acids: Review of recent human studies and food industry responses. *Lipids*. 2006;41(11):967–92.
37. Morenga L Te, Mallard S, Mann J. Dietary sugars and body weight: Systematic review and meta-analyses of randomised controlled trials and cohort studies. *BMJ*. 2013;345(7891):1–25.
38. Te Morenga LA, Howatson AJ, Jones RM, Mann J. Dietary sugars and cardiometabolic risk: Systematic review and meta-analyses of randomized controlled trials of the effects on blood pressure and lipids. *Am J Clin Nutr*. 2014;100(1):65–79.
39. Scientific Advisory Committee on Nutrition. *Carbohydrates and Health*. 2015.
40. Public Health England. First measure of industry progress to cut sugar unveiled [Internet]. London, UK; 2018. Available from: <https://www.gov.uk/government/news/first-measure-of-industry-progress-to-cut-sugar-unveiled>
41. Alison Tedstone, Victoria Targett, Gabrielle Owtram, Vicki Pyne, Rachel Allen, Kleio Bathrellou, Brittney MacKinlay, Emma Clegg, Kathryn Morgan GS. *Sugar Reduction : Achieving the 20 % A technical report outlining progress to date , guidelines for industry , 2015 baseline levels in key foods and next steps* About Public Health England. Public Health England. London, UK; 2017.

42. Department of HM Government. Childhood Obesity: A Plan For Action. Child obesity a plan action [Internet]. 2018;41(1):59. Available from: www.nationalarchives.gov.uk/doc/open-government-licence/0Awww.nationalarchives.gov.uk/doc/open-government-licence/0Awww.nationalarchives.gov.uk/doc/open-government-licence/0Ahttp://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=0000572
43. Percival R. Ultra-processing is the new frontier in public health policy—reflections on the National Food Strategy [Internet]. The BMJ opinion. 2021 [cited 2021 Apr 12]. Available from: <https://blogs.bmj.com/bmj/2021/07/15/ultra-processing-is-the-new-frontier-in-public-health-policy-reflections-on-the-national-food-strategy/>
44. Suez J, Korem T, Zeevi D, Zilberman-Schapira G, Thaiss CA, Maza O, et al. Artificial sweeteners induce glucose intolerance by altering the gut microbiota. *Nature*. 2014;514(7521):181–6.
45. Pearlman M, Obert J, Casey L. The Association Between Artificial Sweeteners and Obesity. *Curr Gastroenterol Rep*. 2017;19(12):1–8.
46. Baraldi LG, Steele EM, Louzada MLC, Monteiro CA. Associations between ultraprocessed food consumption and total water intake in the US population. *J Acad Nutr Diet* [Internet]. 2021;121(9):1695–703. Available from: <https://doi.org/10.1016/j.jand.2021.02.011>
47. Whybrow S, Horgan GW, MacDiarmid JI. Self-reported food intake decreases over recording period in the National Diet and Nutrition Survey. Vol. 124, *British Journal of Nutrition*. Cambridge University Press; 2020. p. 586–90.

Figure 1 - Changes in food consumption of the UK population according to NOVA classification subgroups between 2008 and 2019.







*Subgroups with statistically significant differences ($p < 0.05$) in dietary participation over 11 years.

Box 1. The NOVA food classification system and its four groups defined according to the extent and purpose of food processing.

Group 1 unprocessed or minimally processed foods	Group 2 processed culinary ingredients	Group 3 processed foods	Group 4 ultra-processed foods
<p>Foods derived directly from plants or animals that have undergone little or no processing after leaving the wild, such as cleaning of inedible parts, grinding, crushing, drying, pasteurizing, and fermenting, and to which no other substances have been added.</p> <p>e.g., beans, grains, fresh or frozen meat, eggs, vegetables, milk</p>	<p>Substances directly obtained from group 1 or from nature by processes such as crushing and pressing, and consumed together with group 1 in culinary preparations.</p> <p>e.g., vegetable oils, butter, salt, and table sugar</p>	<p>Foods produced by industry from the combination of groups 1 and 2.</p> <p>e.g., vegetables in brine, cured meat, cheese and bread made from flour, water and salt</p>	<p>Defined as industrial preparations involving multiple processing steps and techniques and many ingredients, including salt, sugars, oils and fats, substances for industrial use only (soy and milk proteins, hydrogenated fats, and modified starch), and additives added to make the final product more palatable and attractive (flavors, aromas, emulsifiers).</p> <p>e.g., biscuits, confectionary, breakfast cereals, pastries, buns and cakes, packaged salty snacks, sauces, dressing and gravies, soft drinks, milk-based drinks, packaged pre-prepared meals, sausages and other reconstituted meat products</p>

Table 1. Trend of dietary share (% of total energy intake) of NOVA food groups and subgroups over 11 years in the United Kingdom (NDNS 2008 to 2019)

NOVA groups and subgroups	Year 1	Year 2	Year 3	Year 4	Year 5	Year 6	Year 7	Year 8	Year 9	Year 10	Year 11	Adjusted Linear Trend Coeff	p for trend*
	2008/10	2009/11	2010/12	2011/13	2012/14	2013/15	2014/16	2015/17	2016/18	2017/19	2018/20		
Unprocessed or minimally processed foods	30.4 (0.5)	29.7 (0.5)	30.3 (0.5)	30.4 (0.5)	30.3 (0.5)	28.9 (0.5)	30.1 (0.5)	30.2 (0.5)	30.4 (0.6)	30.8 (0.5)	29.8 (0.5)	-0.033	0.505
Milk and plain yoghurt	5.1 (0.2)	5.2 (0.2)	4.7 (0.1)	4.8 (0.1)	5.1 (0.1)	5.0 (0.2)	5.0 (0.2)	5.0 (0.2)	4.9 (0.2)	5.1 (0.1)	4.8 (0.2)	-0.007	0.614
Roots and tubers	3.9 (0.2)	3.5 (0.2)	3.4 (0.1)	3.6 (0.1)	3.4 (0.1)	3.2 (0.1)	2.9 (0.1)	2.8 (0.1)	3.1 (0.1)	3.3 (0.1)	2.7 (0.1)	-0.087	0.000*
Fruit	3.4 (0.2)	3.2 (0.2)	3.2 (0.2)	3.4 (0.2)	3.5 (0.1)	3.2 (0.1)	3.4 (0.1)	3.5 (0.1)	3.6 (0.2)	3.7 (0.1)	3.7 (0.2)	0.042	0.004*
Red meat	4.0 (0.2)	3.1 (0.2)	3.4 (0.2)	3.1 (0.2)	3.2 (0.2)	2.9 (0.2)	3.0 (0.2)	2.7 (0.1)	2.8 (0.2)	2.6 (0.1)	2.2 (0.1)	-0.123	0.000*
Poultry	2.8 (0.2)	2.5 (0.1)	3.0 (0.2)	2.6 (0.1)	2.7 (0.1)	2.7 (0.2)	3.1 (0.2)	3.1 (0.2)	3.0 (0.2)	2.9 (0.2)	2.8 (0.2)	0.030	0.054
Cereal ^a	2.3 (0.1)	2.5 (0.1)	3.0 (0.2)	2.7 (0.2)	2.4 (0.2)	2.5 (0.2)	2.9 (0.2)	2.8 (0.2)	2.7 (0.2)	2.8 (0.2)	3.2 (0.2)	0.052	0.002*
Pastas	1.6 (0.1)	1.7 (0.1)	1.7 (0.1)	1.8 (0.1)	1.9 (0.1)	1.7 (0.1)	1.6 (0.1)	1.8 (0.1)	1.6 (0.1)	2.0 (0.2)	1.9 (0.1)	0.026	0.030*
Eggs	1.5 (0.1)	1.4 (0.1)	1.4 (0.1)	1.5 (0.1)	1.5 (0.1)	1.5 (0.1)	1.6 (0.1)	1.7 (0.1)	1.9 (0.1)	1.9 (0.1)	1.9 (0.1)	0.054	0.000*
Vegetables	1.7 (0.1)	1.5 (0.1)	1.6 (0.1)	1.5 (0.1)	1.4 (0.1)	1.5 (0.1)	1.5 (0.1)	1.6 (0.1)	1.7 (0.1)	1.7 (0.1)	1.6 (0.2)	0.003	0.568
Fresh fruit juice (100% fruit) ^b	1.2 (0.1)	1.3 (0.1)	1.2 (0.1)	1.3 (0.1)	1.3 (0.1)	1.0 (0.1)	0.9 (0.1)	1.0 (0.1)	1.0 (0.1)	0.7 (0.1)	0.8 (0.1)	-0.050	0.000*
Fish	1.3 (0.1)	1.2 (0.1)	1.3 (0.1)	1.2 (0.1)	1.1 (0.1)	1.3 (0.1)	1.2 (0.1)	1.3 (0.1)	1.2 (0.1)	1.4 (0.1)	1.1 (0.8)	-0.005	0.590
Legumes	0.5 (0.0)	0.5 (0.0)	0.7 (0.0)	0.8 (0.1)	0.7 (0.1)	0.6 (0.1)	0.7 (0.1)	0.7 (0.1)	0.9 (0.1)	0.8 (0.1)	0.9 (0.1)	0.028	0.000*
Other unprocessed or minimally processed foods ^c	2.1 (0.1)	2.1 (0.1)	1.9 (0.1)	2.2 (0.2)	2.1 (0.2)	1.9 (0.2)	2.3 (0.2)	2.3 (0.1)	2.1 (0.2)	1.9 (0.2)	2.2 (0.1)	0.003	0.802
Processed culinary ingredients	3.7 (0.2)	4.6 (0.2)	4.3 (0.2)	4.1 (0.2)	4.3 (0.2)	4.4 (0.2)	4.8 (0.2)	4.8 (0.2)	5.2 (0.2)	4.7 (0.2)	4.9 (0.2)	0.101	0.000*
Table sugar	1.4 (0.1)	1.8 (0.1)	1.6 (0.1)	1.7 (0.1)	1.5 (0.1)	1.8 (0.2)	1.7 (0.1)	1.5 (0.1)	1.3 (0.1)	1.3 (0.1)	1.2 (0.1)	-0.040	0.000*
Butter ^d	1.4 (0.1)	1.8 (0.2)	1.5 (0.1)	1.4 (0.1)	1.7 (0.1)	1.5 (0.1)	1.9 (0.1)	1.8 (0.1)	2.4 (0.2)	1.9 (0.1)	2.2 (0.1)	0.071	0.000*
Plant oil	0.5 (0.0)	0.7 (0.1)	0.9 (0.1)	0.7 (0.1)	0.7 (0.0)	0.8 (0.1)	0.9 (0.1)	1.2 (0.1)	1.1 (0.1)	1.1 (0.1)	1.1 (0.1)	0.061	0.000*
Other processed culinary ingredients ^e	0.3 (0.0)	0.4 (0.1)	0.3 (0.1)	0.4 (0.1)	0.4 (0.1)	0.3 (0.0)	0.4 (0.1)	0.4 (0.1)	0.4 (0.1)	0.4 (0.1)	0.4 (0.1)	0.008	0.095
Processed foods	9.6 (0.3)	9.2 (0.4)	8.5 (0.3)	8.9 (0.3)	9.0 (0.4)	8.4 (0.4)	8.4 (0.3)	8.6 (0.3)	7.9 (0.3)	8.5 (0.4)	8.6 (0.4)	-0.101	0.002*
Beer and wine	4.1 (0.3)	3.3 (0.3)	3.4 (0.2)	3.6 (0.2)	3.8 (0.4)	3.3 (0.3)	3.1 (0.3)	3.4 (0.2)	2.8 (0.2)	3.3 (0.3)	3.1 (0.3)	-0.073	0.003*
Cheese	2.8 (0.1)	2.9 (0.2)	2.9 (0.1)	2.8 (0.1)	2.9 (0.1)	2.7 (0.2)	2.9 (0.2)	2.7 (0.1)	2.8 (0.2)	2.9 (0.2)	2.7 (0.2)	-0.003	0.813
Vegetables and other plant foods preserved in brine	1.1 (0.1)	1.0 (0.1)	1.0 (0.1)	0.8 (0.0)	0.9 (0.1)	0.9 (0.1)	0.9 (0.1)	0.9 (0.1)	0.7 (0.0)	0.7 (0.0)	0.9 (0.1)	-0.027	0.000*
Processed breads	0.8 (0.1)	0.9 (0.1)	0.6 (0.1)	0.7 (0.1)	0.6 (0.1)	0.7 (0.1)	0.5 (0.1)	0.6 (0.1)	0.6 (0.1)	0.6 (0.1)	0.7 (0.1)	-0.014	0.171
Salted, dried, or smoked meat or fish	0.5 (0.1)	0.6 (0.1)	0.6 (0.1)	0.6 (0.0)	0.5 (0.1)	0.4 (0.0)	0.6 (0.1)	0.5 (0.0)	0.6 (0.0)	0.5 (0.1)	0.5 (0.0)	-0.003	0.564
Other processed foods ^f	0.3 (0.1)	0.3 (0.1)	0.2 (0.0)	0.4 (0.1)	0.3 (0.1)	0.3 (0.1)	0.3 (0.1)	0.5 (0.1)	0.4 (0.1)	0.4 (0.1)	0.5 (0.1)	0.020	0.003*

Ultra-processed foods	55.3 (0.6)	56.5 (0.6)	56.9 (0.6)	56.5 (0.6)	56.4 (0.7)	58.3 (0.7)	56.7 (0.6)	56.4 (0.6)	56.5 (0.7)	56.0 (0.7)	56.6 (0.7)	0.034	0.580
Ultra-processed breads	10.5 (0.2)	11.1 (0.2)	11.3 (0.3)	11.5 (0.3)	11.4 (0.2)	10.8 (0.3)	10.9 (0.3)	11.1 (0.3)	10.9 (0.3)	10.6 (0.2)	10.3 (0.3)	-0.045	0.074
Packaged pre-prepared meals ^g	7.4 (0.3)	7.5 (0.3)	7.2 (0.3)	7.7 (0.3)	7.6 (0.3)	8.3 (0.4)	7.9 (0.3)	8.6 (0.3)	8.5 (0.3)	8.1 (0.3)	8.3 (0.3)	0.119	0.000*
Breakfast cereals	4.2 (0.2)	4.2 (0.2)	4.3 (0.2)	4.4 (0.2)	4.4 (0.2)	4.9 (0.2)	4.5 (0.2)	4.4 (0.2)	4.7 (0.2)	4.7 (0.2)	4.9 (0.2)	0.064	0.001*
Sausage and other reconstituted meat products	4.2 (0.2)	3.7 (0.2)	4.0 (0.2)	3.8 (0.1)	3.7 (0.2)	3.6 (0.2)	3.7 (0.2)	3.2 (0.1)	3.1 (0.0)	3.2 (0.1)	3.7 (0.2)	-0.076	0.000*
Confectionary	3.5 (0.2)	3.5 (0.2)	3.3 (0.1)	3.4 (0.1)	3.7 (0.2)	3.9 (0.2)	3.5 (0.2)	3.5 (0.2)	3.6 (0.1)	3.5 (0.2)	3.4 (0.2)	0.004	0.796
Biscuits	3.2 (0.1)	3.6 (0.1)	3.3 (0.2)	3.8 (0.2)	3.5 (0.2)	3.5 (0.2)	3.8 (0.2)	3.9 (0.2)	3.9 (0.2)	3.9 (0.2)	4.2 (0.3)	0.076	0.000*
Pastries, buns, and cakes	2.9 (0.1)	3.0 (0.2)	3.2 (0.2)	3.4 (0.2)	3.6 (0.2)	3.5 (0.2)	2.9 (0.2)	3.4 (0.2)	3.3 (0.2)	3.4 (0.2)	3.7 (0.2)	0.049	0.004*
Industrial chips (French fries)	2.9 (0.2)	2.8 (0.2)	2.9 (0.2)	2.4 (0.1)	2.6 (0.2)	2.7 (0.1)	3.0 (0.2)	2.6 (0.2)	2.6 (0.2)	2.5 (0.1)	2.5 (0.2)	-0.027	0.077
Soft drinks, fruit drinks and fruit juices	2.5 (0.1)	2.5 (0.2)	2.7 (0.2)	2.5 (0.1)	2.3 (0.1)	2.5 (0.2)	2.3 (0.2)	2.0 (0.2)	2.0 (0.2)	1.5 (0.1)	1.1 (0.1)	-0.122	0.000*
Milk-based drinks	2.0 (0.1)	2.2 (0.1)	2.2 (0.1)	2.2 (0.1)	2.2 (0.1)	2.4 (0.1)	1.8 (0.1)	2.0 (0.1)	2.0 (0.1)	2.2 (0.1)	2.0 (0.1)	-0.017	0.119
Packaged salty snacks	1.9 (0.1)	2.0 (0.1)	1.9 (0.1)	2.1 (0.1)	2.0 (0.1)	2.1 (0.1)	2.1 (0.1)	2.0 (0.1)	2.3 (0.1)	2.1 (0.1)	2.3 (0.1)	0.030	0.005*
Industrial pizza	1.8 (0.2)	2.0 (0.2)	1.7 (0.2)	1.7 (0.2)	1.6 (0.2)	2.1 (0.2)	2.2 (0.2)	2.0 (0.2)	2.3 (0.2)	2.0 (0.2)	2.0 (0.2)	0.035	0.053
Margarine and other spreads	2.4 (0.1)	2.3 (0.1)	2.3 (0.1)	2.3 (0.1)	2.0 (0.1)	1.9 (0.1)	1.6 (0.1)	1.4 (0.1)	1.4 (0.1)	1.7 (0.1)	1.6 (0.1)	-0.099	0.000*
Sauces, dressing and gravies	2.0 (0.1)	2.1 (0.1)	2.4 (0.1)	2.1 (0.1)	1.9 (0.1)	2.2 (0.1)	2.2 (0.1)	2.3 (0.1)	2.2 (0.1)	2.5 (0.1)	2.2 (0.1)	0.025	0.009*
Industrial desserts	0.7 (0.1)	0.9 (0.1)	0.9 (0.1)	0.8 (0.1)	0.9 (0.1)	1.0 (0.1)	0.9 (0.1)	1.1 (0.1)	0.8 (0.1)	0.9 (0.1)	1.0 (0.1)	0.011	0.168
Other ultra-processed foods ^h	3.3 (0.2)	3.2 (0.2)	3.3 (0.2)	2.6 (0.1)	2.9 (0.2)	3.2 (0.2)	3.3 (0.2)	2.9 (0.2)	3.1 (0.2)	3.2 (0.2)	3.4 (0.2)	0.006	0.746

^aIncluding grains and flours ^bIncluding ultra-high temperature processing (UHT) or pasteurized, and smoothies; ^cIncluding coffee, tea, sea foods, fungi, nuts, and freshly prepared dishes based on one or more unprocessed or minimally processed food; ^dIncluding lard and suet shredded; ^eIncluding starches, coconut and milk cream, gelatin powder, and vinegar; ^fIncluding salted, sweetened, or oil-roasted nuts or seeds, condensed milk, and commercial baby foods; ^gIncluding frozen and shelf-stable dishes and canned soups; ^hIncluding baked beans, meat alternatives, soy and other drinks as milk substitutes, infant formula, and distilled alcoholic drink. SE = standard error.

†Adjusted for age (years), sex, occupational social class and ethnicity (white, mixed ethnic group, black, Asian and other race). *p value<0.05 for linear trend across years.