



## City Research Online

### City, University of London Institutional Repository

---

**Citation:** Spicer-Cain, H., Camilleri, B., Hasson, N. & Botting, N. (2023). Early identification of children at risk of communication disorders: Introducing a novel battery of Dynamic Assessments for infants. *American Journal of Speech-Language Pathology*, 32(2), pp. 523-544. doi: 10.1044/2022\_AJSLP-22-00040

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

---

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/29274/>

**Link to published version:** [https://doi.org/10.1044/2022\\_AJSLP-22-00040](https://doi.org/10.1044/2022_AJSLP-22-00040)

**Copyright:** City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

**Reuse:** Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.



**Early identification of children at risk of communication disorders:**

**Introducing a novel battery of Dynamic Assessments for infants**

Helen Spicer-Cain, Bernard Camilleri, Natalie Hasson, Nicola Botting

Centre for Language and Communication Science Research, City University of London

Corresponding author contact: Nicola.botting.1@city.ac.uk

Conflict of interest: The authors declare no conflict of interest.

Funding: The original study was supported by doctoral funding to the first author from City University of London. No other funding was received.

ACCEPTED INTO AJSLP 8<sup>th</sup> November 2022

## 29 **Abstract**

### 30 *Purpose*

31 Many children with communication disorders (CD) experience lengthy gaps between parental reporting of  
32 concerns and formal identification by professionals. This means that children with CD are denied access to  
33 early interventions that may help to support the development of communication skills and prevent possible  
34 negative sequelae associated with long-term outcomes. This may be due, in part, to the lack of assessment  
35 instruments available for children younger than three years of age. This study therefore reports on  
36 promising preliminary data from a novel set of valid dynamic assessment measures designed for infants.

### 37 *Methods*

38 We recruited 53 low-risk children and two groups of children considered to be at high risk for CD (n=17  
39 social-high-risk and n=22 language high-risk) due to family members with language and social  
40 communication difficulties. Children were between 1 and 2 years of age and were assessed using a battery  
41 of five dynamic assessment (DA) tasks related to receptive vocabulary, motor imitation, response to joint  
42 attention, turn taking and social requesting. A set of standardised measures was also used.

### 43 *Results*

44 The DA tasks showed high levels of inter-rater reliability and relationships with age across a cross-sectional  
45 sample of children from the low-risk group. Three tasks showed moderate to strong correlations with  
46 standardised measures taken at the same age, with particularly strong correlations between the DA of  
47 receptive vocabulary and other receptive language measures. The DA of receptive vocabulary was also the  
48 only task to discriminate between the three risk groups, with the social-high-risk group scoring lower.

### 49 *Conclusions*

50 These results provide preliminary information about early DA tasks, forming the basis for further research  
51 into their utility. DA tasks might eventually facilitate the development of new methods for detecting CD in  
52 very young children, allowing earlier intervention and support.

53

54

## 55 **Introduction**

56 Many children experience communication difficulties that require intervention during development. Autism  
57 and Developmental Language Disorder (DLD) represent two of the most prevalent disorders of childhood.  
58 Roughly 2% (Roman-Urrestarazu et al, 2021) and 8% (Norbury et al, 2016) of all children experience these  
59 disorders respectively, and there is compelling evidence that there are lifelong sequelae including  
60 employment issues (Autism: Harmuth et al, 2018; DLD: Dubois et al, 2020) as well as for mental health  
61 (Autism: Hollocks et al, 2019; DLD: Botting et al, 2016). Yet, for DLD especially, there is relatively low  
62 awareness (Thordardottir et al, 2021) and a paucity of research compared to other developmental disorders  
63 (Bishop, 2010; McGregor, 2020). There is a view that early intervention is optimum for these children, as  
64 language difficulties associate with wider long-term difficulties such as memory impairment (Henry &  
65 Botting, 2017), poorer educational attainment and employment prospects (Conti-Ramsden et al, 2018) and  
66 increased mental health issues (Botting et al, 2016). However, very early diagnosis and associated  
67 intervention services are not yet recommended in many countries including the UK, (e.g. Lindsay et al, 2008;  
68 Boyle, 2011; Wallace et al, 2015; Reilly et al, 2015; Bishop et al, 2017; Law et al, 2020; Jullien et al, 2021), in  
69 part because there are limited reliable assessments which can accurately identify infants with  
70 communication difficulties before the age of 3. In this paper we present preliminary data from a set of novel  
71 assessment tasks as a first step towards developing tools for identifying very early social and communication  
72 difficulties. We have focussed this ‘proof of concept’ study on groups of children at risk of Autism and DLD  
73 because of their combined prevalence and also because these are groups where we expect communication  
74 difficulties to show early signs; however, tools that are applicable to communication difficulties in other risk  
75 groups would be a wider long-term aim.

76

## 77 ***Autism***

78 Autism is a lifelong pervasive developmental disorder which is diagnosed on the basis of impairments of  
79 social communication and social interaction, alongside restricted and repetitive behaviours (American  
80 Psychiatric Association, 2013). Recent prevalence estimates indicate that approximately one in every 68

81 children aged four in the USA has an ASD (Christensen, et al., 2016) and that this figure is approximately 4.5  
82 times as high for boys than for girls (1 in 42 as opposed to 1 in 189 respectively; Christensen, et al., 2016). A  
83 similar estimate of prevalence was derived for the UK by Baron-Cohen et al (2009), although only 60% of  
84 these cases were formally diagnosed before study participation. Because ASD is a spectrum condition, it is  
85 vastly heterogeneous in its presentation. Language abilities can range from minimal use of, or  
86 comprehension of spoken language, to intact structural language skills in the context of difficulties with  
87 pragmatic skills, or language use. ASD can occur both with and without learning disability, and recent  
88 estimates suggest that 44% of children with ASD have average or above average intellectual ability  
89 (Christensen, et al., 2016). Regression of communication and adaptive skills, usually in the second year of  
90 life, is also reported in a subset of cases (Meilleur & Fombonne, 2009).

91

#### 92 ***Developmental Language Disorder (DLD)***

93 DLD is the preferred label for language difficulties of unknown aetiology in children, including conditions that  
94 were previously referred to as Specific Language Impairment or Developmental Dysphasia (Bishop et al,  
95 2017). DLD affects approximately 7-8% of children at school-starting age (Tomblin, et al., 1997; Norbury, et  
96 al., 2016) and typically occurs in the absence of hearing loss, neurological impairment or severe  
97 environmental deprivation which would explain difficulties with language learning. Previous criteria for  
98 Specific Language Impairment required normal non-verbal intellectual ability and/or a discrepancy between  
99 language skills and IQ, but it has recently been established that there are few significant differences between  
100 children with language difficulties in the presence of typical IQ and children who have low abilities in both  
101 language and cognition, and IQ criteria are therefore no longer used to define DLD (Bishop et al, 2017;  
102 Norbury, et al., 2016). The presentation of DLD is also heterogeneous and may involve difficulties at any level  
103 of language processing, including phonology, morphology, syntax, semantics or pragmatics. Individual  
104 children may be affected across one or multiple levels of language and across either receptive or expressive  
105 modalities or both. The relationship of DLD to delayed language acquisition (or “late talking”) in early  
106 childhood is complex, (e.g., Dale et al., 2003; Reilly et al., 2010; Zambrana et al, 2014; Duff et al., 2015;

107 Rudolph & Leonard, 2016), and early language delays are not always predictive of later language impairment  
108 on an individual level. However, at least some children who have DLD are known to have difficulties with  
109 spoken language throughout the lifespan (Botting, 2020).

110

### 111 ***Age of Identification***

112 The age at which children with communication disorders are identified has important implications for early  
113 intervention and support. The potential consequences of not identifying and providing input for children  
114 with communication disorders are large (Hus & Segal, 2021) and may include poorer employment prospects  
115 (Chen et al., 2017; Conti-Ramsden et al., 2017); increased mental health difficulties (Botting et al., 2016);  
116 and economic costs for society (Rogge & Janssen, 2019). For ASD, Zwaigenbaum et al (2009) found that most  
117 parents of children later diagnosed with ASD identified concerns about their children's development  
118 between 12 and 18 months of age, including concerns regarding delayed language development, limited play  
119 skills, decreased social responsiveness, extreme behavioural reactions to external stimuli and difficulties with  
120 sleep or feeding. Some parents report even earlier concerns starting before 12 months of age (Zwaigenbaum  
121 et al., 2005; De Giacomo and Fombonne, 1998; Filipek et al., 1999). However, as noted above, early  
122 identification is not straightforward, and reported age of diagnosis in ASD tends to be much older than  
123 reported age of first concern. Mean age of diagnosis varies between studies and across countries, with a  
124 review by Daniels and Mandell (2014) reporting means ranging from 32 to 120 months of age across  
125 different studies, and a recent meta-analysis reporting a mean age of autism diagnosis of 60.5 months based  
126 on studies from 40 countries (van't Hof et al, 2021). Other recent studies suggest similar age of diagnosis: a  
127 mean of 46 months of age in Australia (Bent et al, 2020), 58 months in France (Rattaz et al, 2022) and 54  
128 months in the USA (Hanley et al, 2021). Indeed, recent prevalence data from the USA suggests that, of 4681  
129 children with an autism diagnosis at the age of 8, only 47% received a diagnosis before the age of 3  
130 (Maenner et al, 2021). Importantly, in the UK context, Crane et al (2016) report an average delay of  
131 approximately 3.5 years between first contact with health professionals and confirmed diagnosis in their  
132 survey of parents of children with ASD, and only 11% of children in their sample were diagnosed before the

age of three. Cohort studies also emphasise the rate of later diagnosis in ASD, with the number of diagnosed children in the Early Language in Victoria Study (ELVS) more than doubling between the ages of four and seven (Veness et al 2014), and percentages of children diagnosed with ASD rising from 0.9% at age five to 1.7% at age seven and 3.5% at age 11 within the Millennium Cohort Study (Dillenburger et al, 2015). Although some children do receive intervention services before they have a diagnosis, these do not appear to start significantly earlier (mean age for first receipt of services = 4.1 years; Hanley et al, 2021).

There is less information on diagnosis/recognition of difficulties among children with DLD, but Rannard et al (2004) reported that a quarter of parents in their sample noticed difficulties in their children's language and communication between 12 and 18 months of age, and a further quarter between 18 and 24 months of age, making around 50% showing concern by the time their child reached their second birthday. Absent or unusual babbling, poor intelligibility and late language onset were the main areas of concern noticed by parents. However, despite roughly half of parents being concerned about their child's communication skills by the age of two, only one third of the children in this study received any input from speech and language therapy services before the age of three, and 21% had no support until they started school (Rannard et al, 2004). Similarly, Tomblin et al (1997) found that only 29% of children who showed evidence of language impairment in Kindergarten had ever been referred to speech and language therapy services. In a more recent study in the UK, Norbury et al (2016) found that, of 9.92% of children who had language difficulties at school entry, only 39% had ever been referred to SLT services, and only 40% received any additional support at school. This figure may be higher in other countries, but a recent study in the USA also found that not all children who have speech and language difficulties are receiving SLT support, with only around 75% ever having received services for these difficulties during their lifetime (Davidson et al, 2022). Thus, there is a need to work towards more sensitive early measures of language and social communication, tapping into pre-verbal behaviours, such as joint attention and turn-taking, which form the foundations of language development (Curtin et al, 2021).



159 UK Referral figures for children of different ages confirm these results from both disorders, with Broomfield  
160 and Dodd (2011) finding that only 6% of referrals made to one speech and language therapy service under  
161 study were for children under two, 67% between the ages of two and five, and 27% for children of five or  
162 older. The Bercow Report (Bercow, 2008) also highlighted that UK parents continue to have difficulties  
163 accessing speech and language therapy services and 28% of those who responded felt they had had to fight  
164 for their child to receive a diagnosis and associated services. As noted earlier, inefficient diagnostic  
165 pathways may lead to poorer outcomes across a variety of areas in later life for children with communication  
166 impairments (Hus & Segal, 2021).

167

#### 168 ***Current assessment issues***

169 There may be many reasons for delayed identification, including limitations in the training of professionals in  
170 early assessment, resource issues and service eligibility criteria (Huerta & Lord, 2012). However, certainly in  
171 the UK and US, there is generally a lack of valid assessments appropriate for infants. Where they are used,  
172 the likelihood of social, cultural and linguistic bias is high (Dockrell & Marshall, 2015), untested adaptations  
173 are sometimes made (Cycyk et al, 2021), and the arbitrary cut-off scores are problematic (Spaulding, Plante  
174 & Farinella, 2006). Furthermore, the appropriateness of a given test is often not considered properly in  
175 practice (Friberg, 2010; Betz et al, 2013). Thus, the addition of appropriate, culturally and linguistically  
176 sensitive infant assessment tools, is one area where there is need for urgent improvement to avoid the  
177 consequences of late- or missed-diagnoses (Hus & Segal, 2021). In particular, the current model relies  
178 heavily on impairment focused assessments that use formal, static approaches – that is measures which tap  
179 into performance at once time point, without considering process (Spaulding et al, 2012; Roulstone et al,  
180 2015; Dockrell & Marshall, 2015). These assessments are primarily designed to identify children scoring  
181 below a particular threshold rather than predicting risk or assessing change over time. (Hasson & Botting,  
182 2010). When considering very young children, especially those at risk of communication difficulties, these  
183 tests may not serve the purpose of assessing possible difficulties because they are not feasible with infants,  
184 and because infants tend to show a wide range of ability at a given age (Law & Roy, 2008).

185

186 Speech and language therapists, teachers, psychologists and others who assess children using static tests  
187 have long known that there are some groups of children who are not well-served by traditional formal  
188 assessment methods (e.g. Spaulding et al, 2012). There are many reasons why a child may fail to perform  
189 well under static testing conditions, including cultural and linguistic diversity, shyness, difficulties with  
190 attention regulation, difficulties with social interaction and lack of familiarity with the formal testing process,  
191 as well as difficulties with the specific knowledge and skills being assessed (Chiat & Roy, 2007; Camilleri &  
192 Law, 2007; Hasson & Joffe, 2007). Because static testing usually seeks to remove the effects of the individual  
193 examiner by making the testing process exactly the same for each child, without environmental support or  
194 examiner feedback, it tells us only how the child performs on a specific measure under those conditions on a  
195 specific day. What it does not tell us is how the same child performs in more natural situations, or where  
196 they are engaged with the examiner in a collective effort to generate correct responses (Peña et al, 2007).  
197 This causes an issue with validity whereby the static assessment only captures a one-point estimate of the  
198 construct, rather than the construct itself (Messick, 1998; Hasson & Joffe, 2007; Camilleri & Law, 2007;  
199 Spaulding et al, 2012) and yet at the same time fails to eliminate all tester input effects (Muskett, Body &  
200 Perkins, 2012). Thus, a different approach is needed. Practitioners often take the approach of very informal  
201 observation or reliance on parent report (for example, by health visitors or doctors; Law et al, 2020) to  
202 counter the lack of formal assessment, but using a 'Dynamic Assessment' to measure emerging skills and  
203 learning potential offers a middle ground providing flexible yet objective measurement (Bamford et al,  
204 2022).

205

### 206 ***Dynamic Assessment (DA)***

207 In contrast to the static formal testing usually used by speech and language therapists, DA is more focused  
208 on the process of learning, and what a child's potential level of performance is, when supported by an adult  
209 who can provide prompting, cueing or teaching to help them improve their performance on the task . DA  
210 arose originally out of the socio-cultural theory of Vygotsky (1978), who described the "zone of proximal

211 development” (ZPD) of the child’s skills in any area of learning. This describes the gap between the child’s  
212 habitual unaided performance and the level that they are able to reach when supported by an adult or more  
213 experienced peer. That is, children’s learning potential can be measured by observing what they can achieve  
214 in a scaffolded paradigm, rather than just their performance in an unaided scenario (Hasson & Joffe, 2007).  
215 This difference in static and dynamic methods has been noted and built on for school age children and now  
216 has widespread awareness (Deutsch & Reynolds, 2000) and some practice among some school psychologists  
217 (Hussein & Woods, 2019). However, to our knowledge no work has been done exploring the use of DA in  
218 preverbal infants.

219

220 Sternberg and Grigorenko (2002) described two main formats into which DA methodologies can be  
221 organised: the “sandwich” and the “cake” (see a recent description by Bamford et al, 2022). The use of test-  
222 teach-retest dynamic assessment procedures may be referred to as a “sandwich”, in which children are  
223 tested using static assessments before and after a brief intervention, to reveal the amount of change, or  
224 ‘gain’ that has taken place. The teaching phase typically involves a metacognitive element, which enables the  
225 child to learn which elements/strategies are required for successful completion of the task in question.  
226 Ratings of the child’s responsiveness during the ‘teach’ phase, together with the gains achieved between the  
227 test and the retest provide an indication of the child’s potential to learn. Within the field of speech and  
228 language assessment, this methodology has been adopted for the diagnostic purpose of distinguishing  
229 between low language ability and typically developing preschool children from specific culturally and  
230 linguistically diverse groups in the United States of America (Kapantzoglou et al., 2012; Peña et al, 2014).  
231 Static, standardised assessments can be biased against these children, leading to low scores for both  
232 typically developing children as well as children with language disorders. Dynamic assessments were found  
233 to reduce this bias, when assessing a range of areas including vocabulary (Kapantzoglou et al., 2012; Pena,  
234 Iglesias & Lidz, 2001), categorization (Ukrainetz, Harpell, Walsh and Coyle, 2000) and narrative (Pena et al.,  
235 2006).

236

237 The “cake” format, which sometimes forms the centre part of the ‘sandwich’ (see below for hybrid  
238 methods), is perhaps more suitable when assessing very young children, below the age of four. This method  
239 usually involves the integration of graduated prompts or feedback into the assessment session, as described  
240 by Campione and Brown (1987) and Carlson and Wiedl (1978) and used more recently by researchers such as  
241 Patterson et al. (2020) in preschool children. The examiner provides support to the child as they are  
242 completing the assessment, typically using a pre-determined cueing hierarchy that provides the child with  
243 increasingly explicit support to reach the correct answer or complete the task. What is measured here and  
244 interpreted as the size of the ZPD is the number of cues given to the child to enable them to complete the  
245 task, with more favourable scores being achieved by children who require less cueing to achieve success  
246 (Campione & Brown, 1987). This methodology was previously adopted with young preschool children (aged  
247 30 to 36 months), who had a specific difficulty with expressive language (Bain & Olswang, 1995; Olswang &  
248 Bain, 1996). This DA targeted the immediate potential for children performing at the one-word stage of  
249 expressive language development, to produce two-term utterances, by using a series of graduated prompts  
250 which facilitated production of the two-term utterance. These prompts included elicitation questions,  
251 sentence completion and direct/indirect modelling. The key findings were that children’s scores on the DA  
252 were highly predictive of change over a nine-week period, both with (Bain & Olswang, 1995) and without  
253 intervention (Olswang & Bain, 1996).

254

255 Some DA research with young children in the United Kingdom and Europe has adopted a hybrid approach,  
256 incorporating both graduated prompts and an element of metacognitive intervention (Hasson et al., 2013;  
257 Camilleri, Hasson & Dodd, 2014). This has included research looking at bilingual and multilingual children in  
258 their first year of schooling (MacLeod & Glaspy, 2022). One big difference between this research and that  
259 from the USA is that children in the UK/Europe derive from a wide range of bilingual backgrounds, whereas  
260 the children in the studies cited above from the USA were recruited from specific linguistic backgrounds  
261 (e.g., Hispanic or Native American). The UK studies compared typically developing bilingual children and  
262 bilingual children with developmental language delays (on the Speech and Language Therapy caseload). The

263 findings were that caseload children required greater assistance and made fewer gains in both vocabulary  
264 and sentence production (Hasson et al., 2013; Camilleri et al., 2013), further extending the evidence base  
265 that DA can be used to distinguish between these two groups. All but one of the children with  
266 developmental language delays were found to experience difficulties with components of the DA  
267 assessment (Camilleri et al., 2013).

268

269 Of the different approaches mentioned above, the graduated prompt approach is particularly suited when  
270 working with very young children, as it does not require the explicit metacognitive element that is crucial to  
271 the 'sandwich' or test-teach-retest approach. Although recent reviews by Hunt et al, (2019) and Orellana et  
272 al (2019) indicate that test-teach-retest methods are mostly chosen, the meta-analysis by Orellana et al  
273 (2019) concluded that modifiability ratings (similar to those used in the graduated prompts approach)  
274 showed more promise as an indicator of typical development vs. language impairment at least in bilingual  
275 children. This is therefore the approach which has been selected for the current study.

276

### 277 ***Present study***

278 As discussed above, there is currently an emerging evidence base for the use of DA in speech and language  
279 therapy. Although the evidence to date is mainly from small-scale studies (see Joffe & Hasson, 2007;  
280 Orellana et al, 2019), there is a growing awareness in the field suggesting that DA can be used successfully  
281 with preschool children to determine the presence or absence of language and communication impairment,  
282 and in order to suggest strategies that may be used to support children in their communication  
283 development, or predict how children will respond to intervention (Hunt et al, 2019). This paper aims to  
284 address some gaps identified in the literature, including exploring the application of DA to children under  
285 two years of age, and the use of DA with high-risk children as a predictor of later language and  
286 communication skills. The areas of focus for these new DA tasks encompass 5 key communicative gestures  
287 and behaviours that have been reported in the literature as predictors of later language or as delayed in  
288 children with later social communication difficulties (Law et al, 2017; Rohlfing, 2019; Ramos-Cabo et al,

289 2019). Namely these areas comprise: Early receptive vocabulary (Markus et al, 2000); Response to joint  
290 attention (Salo et al, 2018); Motor imitation (Hanika & Boyer, 2019); Turn taking (Hendenbro et al, 2014);  
291 and non-verbal requesting behaviour (Ramos-Cabo et al, 2019). We acknowledge that the full development  
292 of a new DA tool for clinical practice will take many iterations. Thus, the objective here is to present work to  
293 establish initial 'proof of concept' and feasibility of an early infant measure.

294 The aims of the study were threefold:

- 295 1. To investigate whether reliable normative scores can be gained from a novel battery of very early DA  
296 procedures for use with infants under two years of age who have no first-degree relatives with  
297 communication or literacy difficulties.
- 298 2. To assess performance on these measures in relation to age, sex and standardised tests of  
299 communication in a low-risk group of children (normative sample).
- 300 3. To explore whether there are early indicators of (known groups) validity, using preliminary  
301 comparisons of infants at high-risk of communication disorders (siblings or parents with ASD, DLD or  
302 Dyslexia) with low-risk infants (siblings or parents with no known difficulties).

303

## 304 **Method**

### 305 ***Recruitment***

306 Recruitment took place via social media, where contacts of the researchers were encouraged to share the  
307 project website on their own feeds. Parents of children in the correct age range could then visit the project  
308 website, view the project information sheet, and contact the research team if they agreed to take part.  
309 Children with bilingual exposure were not excluded from this sample, as long as they were exposed to  
310 English as one of the main languages of the home and could be assessed in English<sup>1</sup> Informed consent was  
311 taken from parents of all infants at the start of the research visit. The infants who participated were too  
312 young to give formal assent, but willingness to interact with the researcher and participate in activities was

---

<sup>1</sup> For the purposes of developing this task, English was the only language assessed. However, we acknowledge that in clinical practice, it is preferable to assess all home languages.

taken to indicate assent. The study was granted ethical approval from City University of London, Language and Communication Science Research Ethics Committee.

### **Participants**

Two groups of children participated in the study: those at low-risk ( $n=51$ ) and high-risk ( $n=41$ ) of communication difficulties based on family history. The novel tasks were assessed for feasibility, reliability age relations and preliminary validity using the low-risk group only, to establish how the test performs for a normative sample (Pena, Spaulding & Plante, 2006). The high-risk groups were then used to compare scores to explore clinical usefulness and preliminary known-group validity.

#### *Low-risk children (with typical siblings and parents)*

Participants in the first part of this study were 51 low-risk children (25 female and 26 male) and had a mean age of 12.2m ( $SD = 3.0$ ) at the time of assessment (see Table 1 for demographics). The majority of this group were white British ( $n = 37$ ) with 4 who had mixed ethnicity, 2 who were Asian and 2 reported as being of 'other' ethnicity (6 children had no ethnicity data recorded). The inclusion criteria for this group were that children had no known developmental, physical or sensory difficulties at the time of recruitment, and their parents and elder siblings showed no evidence of language, communication or literacy difficulties. This sample included five children who were exposed to other European languages within their home in addition to English (Swedish ( $n=1$ ), Finnish ( $n=1$ ), German ( $n=2$ ) and Italian ( $n=1$ ), with exposure to their additional language varying between 20 to 40 hours per week ( $M = 24.8$ ;  $SD = 8.6$ ) as reported by parents on the UKCDI demographic questionnaire (Alcock et al, 2020). Children were largely recruited from Greater London (72.5%) although some were from other parts of England.

In total 31 of the children were first-born, and had parents with no reported history of difficulties with language, social communication or literacy development. Twenty children had older siblings ( $n=26$  siblings), who were reported by parents to be developing typically. Parents completed the Children's Communication

339 Checklist (CCC-2) (Bishop, 2003) for 18 elder siblings of children in the sample who were aged 4;0 and above.  
340 All 18 elder siblings scored within the average range for the General Communication Composite score on the  
341 CCC-2, indicating no communication impairments, and none scored within the range of clinical concern on  
342 the Social Interaction Deviance Composite score. Of the remaining 8 siblings, 4 were older than 4;0 but did  
343 not have a CCC-2 completed by parents, and 4 were younger than 4;0 and therefore the CCC-2 could not be  
344 completed. However, in all cases, parents reported no concerns about their development. Additionally, 3 out  
345 of 4 siblings under the age of 4;0 were present during the assessment of the infant in the study, and were  
346 judged by the first researcher, who is an experienced speech and language therapist, to have language and  
347 communication skills within the typical range for their age. All elder siblings were therefore assumed to be  
348 typically developing. In addition, none of these 21 infants had parents who reported a history of difficulties  
349 with language, literacy or social communication.

350

351 Demographic data showed that 79.0% of mothers and 81.8% of fathers of these infants were aged 31 or  
352 older. The sample had high levels of parental education, with 95.5% of mothers and 86.4% of fathers  
353 reporting an undergraduate or postgraduate degree, and no parents reporting no formal educational  
354 qualifications. Overall, 76.7% of the sample reported family annual income of £42,000 or more. It is  
355 therefore acknowledged in the data that follow that these infants may not be representative across a  
356 broader range of socioeconomic status. See statistical group comparisons below.

357

### 358 *High-risk children*

359 For the final research question, a further 41 children were recruited who were considered at high risk of  
360 communication difficulty on account of their siblings or parents having existing developmental disorders.  
361 These children fell into 2 groups with the following inclusion criteria: i) those with siblings who had a  
362 diagnosis of autism or social communication disorder or were being assessed for this diagnosis, or whose  
363 siblings fell below the clinical threshold for the Social Interaction Deviance Composite score on the Children's  
364 Communication Checklist, 2<sup>nd</sup> Edition (CCC-2; Bishop, 2003). We refer to this group as the Social-High-Risk



365 (SHR) group; ii) infants with siblings and/or parents who had a diagnosis of Developmental Language  
366 Disorder, Dyslexia or other Speech Language and Communication Needs, or who were late to speak (defined  
367 as fewer than 50 single words at the age of two). We refer to these children as the Language-High-Risk (LHR)  
368 group. Children were not excluded on the basis of siblings with other genetic syndromes but infants with  
369 genetic syndromes, physical disabilities or sensory impairments were excluded. High-risk children were  
370 recruited from across England and are detailed below.

371

372 In the SHR group there were 18 children, 10 female and 8 male, with a mean age of 15.4 months ( $SD = 3.9$ ).  
373 Of these 11 were white British and 1 was of mixed ethnicity (6 children with missing ethnicity data). Ten of  
374 these children had elder siblings or half-siblings with a confirmed ASD diagnosis. For the other 8 elder  
375 siblings, concerns were raised by parents about their social interaction skills. Where the elder sibling was  
376 aged 4;0 or above, parents completed the CCC-2, and a conservative Social Interaction Deviance Composite  
377 (SIDC) score of -10 or less was taken to indicate the presence of a social communication impairment in elder  
378 siblings who did not have an ASD diagnosis. Where the elder sibling was aged less than 4;0, the younger child  
379 was considered to fall into the SHR group if the elder sibling was under assessment for an ASD diagnosis.  
380 Two of the eight elder siblings without a formal autism diagnosis had been given a diagnosis of Social  
381 Communication Difficulties by a Speech and Language Therapist; a further child had previously been  
382 assessed for ASD and not given the diagnosis (although traits that could be consistent with mild ASD were  
383 identified), and two were currently undergoing ASD assessment. For three undiagnosed elder siblings,  
384 parents did not refer specifically to ASD when describing their elder child but made reference to difficulties  
385 interacting with others. All of these children showed low SIDC scores on the CCC-2 whilst no sibling of any  
386 child in the low-risk or LHR group had SIDC scores that would indicate significant social impairment. Two  
387 elder siblings in this group had additional diagnoses: one of Attention Deficit Disorder and one of Cri-Du-Chat  
388 Syndrome. One SHR child was exposed to another language (French) for 24 hours a week. In total 16.7% of  
389 SHR children were from Greater London.

390 In the LHR group there were 23 children, 15 female and 8 male, with a mean age of 13.2 months ( $SD = 3.0$ ).  
391 Sixteen were white British and 5 were of mixed ethnicity (2 children had missing ethnicity data). Eight  
392 children in this group had elder siblings or half-siblings with concurrent speech and language difficulties, and  
393 three had parents, elder siblings or half-siblings with a history of late language emergence (no single words  
394 before the age of two). The remaining 12 children had parents and/or siblings/half-siblings with a diagnosis  
395 of dyslexia. In two out of eight cases of concurrent LI, the elder siblings also had learning difficulties and  
396 global developmental delays. In cases where elder siblings were 4;0 or older and speaking in sentences, the  
397 CCC-2 was used to confirm that they had impairments of language but did not have social communication  
398 difficulties. Where the elder sibling was younger than 4;0, their parents reported in all cases that they were  
399 receiving support from Speech and Language Therapy services for language or speech and that no concerns  
400 had been raised about social communication. In total, 47.8% of LHR children were from the Greater London  
401 area and one child was exposed to Spanish for 15 hours a week.

402

#### 403 *Demographic group comparisons*

404 The parental age profile of the high-risk group was similar to that of the low-risk group, with 91.7% of  
405 mothers and 83.4% of fathers in the SHR group, and 86.3% of mothers and 91.0% of fathers in the LHR group  
406 aged 31 or older. Chi squared analysis using three age categories (30 or younger, 31 to 35 and 36 or older)  
407 showed a similar pattern of maternal ( $\chi^2 (4) = 0.893$ ,  $p=0.926$ ) and paternal age ( $\chi^2 (4) = 0.911$ ,  $p=0.923$ )  
408 across groups.

409 Parental education levels were lower for both mothers and fathers in the SHR group, and for fathers in the  
410 LHR group, than the low-risk group (66.7% of mothers and 58.3% of fathers in the SHR group and 95.5% of  
411 mothers and 68.2% of fathers in the LHR group reported having an undergraduate or postgraduate degree).  
412 A significant group difference was found for maternal education level (with categories up to and including  
413 Level 3 qualifications collapsed<sup>2</sup> ;  $\chi^2 (4) = 12.376$ ,  $p=0.015$ ), such that the SHR group contained more

---

<sup>2</sup> Level 3 qualifications are end of high-school qualifications such as A-Levels usually taken at 18 years of age

mothers educated to Level 3 or lower, and fewer mothers educated at degree or postgraduate level than the other groups. Analysis of paternal education level using the same categories revealed that group differences were not statistically significant ( $\chi^2 (4) = 8.049$ ,  $p=0.090$ ). Family income across three categories (£24,000 or less, £24,001 to £42,000, and £42,000 or more) also did not differ significantly across groups ( $\chi^2 (4) = 3.055$ ,  $p=0.549$ ). 50% of SHR families and 71.4% of LHR families reported family annual income of £42,000 or more. Groups showed a significant difference in birth order distribution ( $\chi^2 (4) = 20.557$ ,  $p<0.001$ ), with the low-risk group containing higher numbers of first-born children, and the SHR and LHR groups containing more third or fourth children. In part, this was a function of how the groups were defined, as first-borns could not occur in the SHR group, but could be classified as control/LHR depending on parental dyslexia status. The children from bilingual households do not appear different in terms of SES status or background from the main sample, but the number of children was too small to statistically analyse.

[Table 1 about here]

## Measures

**Demographic data** were collected via questionnaire to parents for all groups, and which were completed for 45/51 low-risk infants, 20/23 LHR infants, 12/18 SHR infants. This questionnaire was the one included on the front of the published UK-CDI measure (please see Alcock et al, 2020 for more details), and asked about parental age, education and family income, as well as hours of exposure to additional languages. These variables were not used for categorisation into groups, which was done solely on family risk factors as described for each group below. An additional question about family history was also included. This question asked parents to say whether any family member had any of the following difficulties: Hearing Impairment; visual impairment; physical disability; autism spectrum disorder; Asperger Syndrome; speech and language difficulties; dyslexia or other problems with reading and spelling; learning difficulties; other developmental difficulties. Family history was also discussed with all parents by the first author who is a qualified and experienced SLT.

A set of **standardised measures** was administered for validation of novel assessment tasks. These included:

- The UK Communicative Development Inventory (UK-CDI) (Alcock et al, 2020). This is a parent-report measure, adapted from the MacArthur Bates Communicative Development Inventory (Fenson et al, 2007). Parents are given a list of 395 words across 19 categories, and asked to indicate whether their child understands and/or says these words. There is also a checklist of 63 gestures and pretend play actions, which parents are asked to indicate whether their child ever performs. The UK-CDI was normed on 1210 children from the UK, who were selected to match the demographic composition of the UK population, and may therefore represent children with a broader range of parental education levels than those included in this sample. However, the UKCDI demonstrates high internal validity for all scales (receptive vocabulary:  $\alpha=0.99$ ; expressive vocabulary  $\alpha=0.99$ ; gesture scale  $\alpha=0.99$ ). Strong correlations were also observed in the standardisation sample with scores on standardised measures of language and an object selection task that measured comprehension directly (Alcock et al, 2020). Parents were asked to complete the CDI for their children in English, as this was one of the main languages for all families.
- The Infant Toddler Checklist (ITC) from the Communication and Symbolic Behaviour Scales – Developmental Profile (CSBS-DP) (Wetherby & Prizant, 2002). This is a 24-item questionnaire, completed by parents, which generates three subscale scores for Social, Speech and Symbolic aspects of communication. The ITC was initially standardised on more than 2000 children in the USA, many of whom were recruited from the same geographic area. However, the standardisation sample matches that included in the present study in terms of having high levels of infants whose parents have completed degree-level or postgraduate education. The ITC shows good levels of internal consistency ( $\alpha=0.93$ ) and test-retest reliability ( $r=0.88$ ), as well as strong correlation in the standardisation sample with other aspects of the CSBS-DP that involve more detailed parent questionnaires and examiner assessment (Wetherby & Prizant, 2002). Additionally, a large cohort study in Australia found the ITC to be a valid clinical tool for measuring early communication skills (Eadie et al, 2010).

- The Modified Checklist for Autism in Toddlers (M-CHAT) (Robins et al, 2001). This is a 23-item checklist, where parents are asked to answer “yes” or “no” to each item, based on their child’s typical behaviour. A subset of 6 items of this questionnaire (the “Core 6 items”) is considered to be particularly indicative of risk for a later diagnosis of ASD (Robins et al, 2001). The M-CHAT shows a high level of internal consistency ( $\alpha=0.85$ ) and also has high levels of sensitivity (0.97) and specificity (0.95) (Robins et al, 2001). Although the M-CHAT is designed for use from 18 months of age, it was included in this study due to its clear format and its potential for indicating emergent difficulties that are linked to ASD. Scoring for this checklist is according to the number of items failed, and higher scores therefore indicate more symptoms related to ASD.
- The Pre-school Language Scales, 4<sup>th</sup> Edition (PLS-4) (Zimmerman, Steiner and Pond, 2002). This is a standardised language assessment, providing scores for receptive and expressive language for children aged from birth to 6 years 11 months. In infancy, scores are mainly given from observation of infant communication during natural interaction, although some older children in the sample were administered receptive language items using toys or picture material. The PLS-4 was originally standardised in the USA, on a sample of 2400 children selected to match the demographic characteristics of the US population. The assessment then received additional UK standardisation with a sample of 800 children matching the UK demographic profile, who were similar in ethnicity to the children in this study, but had a broader range of parental education levels. Test-retest reliability ( $r=0.82-0.95$ ) and internal consistency of this measure were high ( $\alpha=0.72-0.95$ ) in the standardisation sample, and standardisation of the measure showed a good ability to distinguish typically developing children from those with language disorders (Zimmerman, Steiner & Pond, 2002).

A set of novel **dynamic assessment** measures, designed and piloted by the authors for use in this study, was also administered to the children. These measured skills in five areas found in previous studies, to be associated with early communication skills including:

- Receptive vocabulary
- Motor imitation
- Response to joint attention
- Turn taking
- Requesting

These areas of development were chosen as representing core elements of early communication derived from a number of sources including existing reviews (e.g., Ramos-Cabo et al, 2019) and a review of the early communication literature (Spicer-Cain, 2019). These tasks were then tested in a feasibility phase involving 8 children aged 9-17 months (all monolingual; 6 white and 2 mixed ethnicity), and were judged to be engaging for the children, that children were able to complete the assessment and that parents found them acceptable. We concluded that the tasks formed an appropriate assessment for this age range and were likely to be predictors of later language (Spicer-Cain, 2019). This feasibility pilot also helped to guide scoring and number of trials on each task. Note that although early expressive language may be an important predictor, because of the very young target age of the children (12 months), a dynamic cueing hierarchy for this skill was not considered feasible. We therefore acknowledge that this set of DA tasks is preliminary and serves as a 'proof of concept' battery to determine whether initial reliability and feasibility can be achieved.

Based on the principles of DA, graded cueing hierarchies were devised to support children to achieve each of the tasks (Orellana et al, 2019). These are detailed for each task in more detail below, but overall were designed to provide three prompts if the child could not achieve the task independently. Generally, the first of these prompts was a repetition of the instruction, designed to draw the child's attention to the task and give the child more processing time. The second prompt was more specific, and aimed to reduce the difficulty of the target task. The third prompt provided full support for the child to achieve the task. Administration of all DA tasks was videorecorded for reliability checking.

516 *DA of receptive vocabulary*

517 For the dynamic receptive vocabulary task, children were shown a series of five common items, which were  
518 taken out of a bag and placed in front of the child, without naming them. The items (cup, car, duck, ball and  
519 spoon) were chosen to represent words a child would typically acquire as part of their early vocabulary. For  
520 each of the five items, the child's attention was drawn using their name, and pointing to the array of items.  
521 The child was then asked to give one of the items to the researcher, accompanied by an open-hand gestural  
522 prompt. The cueing hierarchy in Appendix 1 was then used for each item. Items were returned to the array  
523 after each had been tested, so that the child was always looking at a choice of five items.

524

525 *DA of motor imitation*

526 Motor imitation was tested via imitation of actions on objects using a toy cup and spoon. The list of gestures  
527 included in the Actions and Gestures section of the UCKDI (Alcock et al, 2020) was reviewed, and used to  
528 choose these objects for use in the motor imitation task, considering previous research showing that young  
529 children are more likely to imitation actions involving objects (Kim et al, 2015). Actions were then chosen  
530 that could be performed with these objects, but which were mostly unrelated to their typical use, to enable  
531 the experimenter to be sure whether the infant was truly imitating the action, as opposed to just showing  
532 understanding of object function. A cup and spoon were given to the child at the start of the activity, and  
533 the experimenter then demonstrated the action using their own set of objects, and encouraged the child to  
534 copy using the phrases "X do it" and "your turn". Animated sound effects were also used by the  
535 experimenter to maintain the child's attention, although the child was not required to copy the sound, and  
536 most did not attempt to do so. The actions used were:

- 537 • Pretending to eat from the cup using the spoon
- 538 • Banging the spoon on the bottom of the cup
- 539 • Touching the spoon to the experimenter's nose
- 540 • Placing the cup upside down on the experimenter's head
- 541 • Stroking the spoon on the experimenter's arm

542 Allowances were made for the children's level of motor development, and any clear attempt to perform the  
543 target action was considered as correct, with no requirement for completely correct execution. The child  
544 was also credited for using either their own set of items or those of the experimenter, or for performing the  
545 actions on their own body or the experimenter's. For each of the five items, the cueing hierarchy in  
546 Appendix 1 was used.

547

#### 548 *DA of response to joint attention (point following)*

549 Response to joint attention (RJA) was assessed based on the child's ability to follow adult pointing, during a  
550 picture-book reading task. A first words picture book containing large colourful photographs of everyday  
551 objects was used, with several objects pictured on each page. Unlike the other subtests, ten trials were run  
552 for this task, because the pilot study suggested both that infants at this age were more difficult to score on  
553 this item; and that increased items on this task were better tolerated than for other DA items (ideally all  
554 elements would have run with ten trials). For each RJA trial, the experimenter pointed at an item on the  
555 page, saying "Look! A (name of item)". To aid the scoring of the task, the items used for each child were  
556 chosen so that the child would have to make an obvious gaze shift from where they were currently looking  
557 to look at the item to which the adult was pointing. The sequence of cueing in Appendix 1 was used. If the  
558 child pointed to items in the book, the experimenter named these, and the child was allowed to look at each  
559 page until they lost interest, although only one trial was made on each page.

560

#### 561 *DA of turn-taking*

562 Turn-taking skills were assessed using a ball-run toy designed for infants, where a ball is put into a hole and  
563 then runs down a spiral track. The experimenter first demonstrated the toy for the child by taking a turn, and  
564 then encouraged the child to take a turn using the phrase "X's turn" / "you do it". Once the child was  
565 engaged with the toy, the experimenter initiated a turn-taking sequence by taking a turn themselves (see  
566 Appendix 1). Five turn sequences were then scored according to the procedure in Appendix 1. The



567 experimenter and child then continued to play with the toy until the child lost interest, although only the  
568 first five turns were scored.

569

#### 570 *DA of social requesting*

571 Requesting was measured using a disco ball, which spun and displayed colourful lights when it was switched  
572 on. The child was shown the toy, and once they were engaged with it, the toy was then switched off.

573 Appendix 1 shows the cueing sequence which was then used to support the child to make a request to have  
574 the toy turned back on. Requests did not have to be verbal, and could be made using gesture, touch or  
575 vocalisation, as long as this was considered to be socially referenced (accompanied by eye contact to the  
576 experimenter or parent). Five trials were scored, and then the experimenter and the child continued to play  
577 with the toy until the child lost interest in it.

578

#### 579 ***Procedure***

580 Children were assessed by the first author who is a qualified SLT, in their home with a parent present. For  
581 the first fifteen minutes of the session, parents and children were video recorded playing with a standard set  
582 of toys. During this time, aspects of the PLS-4 which could be rated from observation were completed. The  
583 remainder of the appropriate items from the PLS-4 were administered, depending on the age and abilities of  
584 the child. The dynamic assessment measures were then administered and scored live during task  
585 completion. However, all tasks were videorecorded for later reliability checking. The total duration of the  
586 session was around 60 minutes for each child. This included DA administration and scoring of between 10 to  
587 25 minutes. Parents were then given a set of questionnaires to complete and return to the research team,  
588 including the three standardised questionnaire measures listed above.

589

590

591

592

593 ***Analysis***

594 Results were analysed using SPSS version 23.

595 For research question 1, intraclass correlations were used to assess reliability. Cronbach's alpha was used to  
596 report internal consistency.

597

598 For research question 2, due to the non-normal distribution of some variables, Spearman correlations were  
599 used to investigate the relationship between age and scores on each of the dynamic assessment measures.

600 As age was significantly related to most scores, partial correlations were used to establish relationships  
601 between dynamic assessment scores and scores on other measures taken concurrently. Mann-Whitney U  
602 tests were used to compare scores across biological sex.

603

604 For research question 3, ANCOVAs were used to compare all 3 groups on the DA tasks, controlling for age.

605 Assumptions of ANCOVA were checked: Homoscedasticity was verified via scatterplots of predicted against  
606 standardised residuals, and there were no outliers for any task. However, Shapiro Wilks tests showed that  
607 standardised residuals were significantly non-normally distributed in at least one group for all tasks.

608 Transformation of data did not normalise the distributions. No difference in the pattern of results was  
609 observed when combining both high-risk groups and comparing to low-risk groups, thus this analysis is not  
610 reported. There was not enough variability in maternal education scores to consider this as a covariate.

611

612 No results changed substantively on any analysis when children from bilingual families (n=7) were removed,  
613 therefore all children are retained in the analyses that follow.

614

615

616

617

618

619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644

Descriptive statistics for the five DA measures for the children in the low-risk control group only are reported in Table 2. This group of children act as a normative sample and thus are the basis for the development of the tool. Adjusted means from all groups are reported later when we compare scores across known groups. Scores on all measures were not normally distributed, with floor effects present on the measures of receptive language, motor imitation and turn taking, and ceiling effects present on the measures of response to joint attention and social requesting. Interquartile ranges are reasonably wide across all tasks.

### ***Aim1: Reliability of the DA tasks***

In order to ensure the reliability of the above scores, a random subset of 27 videos, selected using stratified sampling to represent 25% of each risk group, was scored by a second rater to investigate inter-rater reliability for the dynamic assessment measures. The independent rater was given the DA scoring hierarchies, training in the scoring methods used, and the basic information about the project was explained, but they were not told any other information about the children, and so were blind to group status or other scores. Intra-class correlation coefficients for the dynamic assessment measures represent 'good' ( $>0.75$ ) or 'excellent' ( $>0.90$ ) agreement for all dynamic assessment measures except Turn Taking which was moderate at 0.70 (Koo & Li, 2016) when the whole sample was considered. All values were good or excellent for our normative (low-risk) and SHR samples. All values except turn-taking were in this range for the LHR group (see Table 3).

645 *Inter-task correlations and internal consistency*

646 With the effect of age statistically controlled, the five DA measures did not show significant partial  
647 correlations with one another, suggesting that they should not be combined into a single scale (see Table 4).  
648 Unsurprisingly, internal consistency of the battery was therefore low at  $\alpha = 0.594$ .

649

650 [Table 4 about here]

651

652 ***Aim 2: Relationship of DA tasks with age and standardised measures of communication***

653 *Relationship with Age*

654 There were significant positive relationships between age and all but one of the DA tasks. For Receptive  
655 Language ( $r = .553, p < .001$ ), Motor Imitation ( $r = .640, p < .001$ ) and Turn-taking ( $r = .777, p < .001$ ) these  
656 associations were all strong, whilst for Response to Joint Attention the relationship was moderate ( $r = .495, p$   
657  $< .001$ ). There was no age relationship with the DA of Requesting ( $r = -.072, p = .620$ ), with high variability of  
658 scores present at all ages. Fig 1 illustrates the findings.

659

660 [Fig 1 about here]

661

662 *Relationship with Sex*

663 Due to the non-normal distribution of scores on the DA measures, Mann-Whitney U tests were used to  
664 compare the scores of boys and girls from the low-risk control group on the five tasks. None of the  
665 comparisons showed significant differences, although there was a marginal difference on the motor  
666 imitation task in favour of girls (see Table 5).

667

668 [Table 5 about here]

669

670

671 *Relationships with Other Measures*

672 Scores on the DA tasks were compared with scores on other parent-reported and experimenter-  
673 administered standardised measures of communication ability. Three DA tasks showed moderate to large  
674 associations with at least one other measure taken concurrently. For the DA receptive language task,  
675 significant correlations were found with parent-reported receptive vocabulary on the UKCDI, and with  
676 receptive and expressive language scores on the PLS-4. The ITC Symbolic and Social subscale scores showed  
677 a significant association with the DA turn taking task, which was also significantly correlated with Total  
678 Gestures scores on the UKCDI. For the DA social requesting task, significant correlations were found with  
679 parent-reported expressive vocabulary on the UKCDI, and the ITC Social subscale (see Table 6).

680

681 After correcting for multiple comparisons using the Bonferroni method, only the association between the DA  
682 measure of receptive language and the PLS-4 Auditory Comprehension score remained significant.

683

684 [Table 6 about here]

685

686 ***Aim 3: Comparison of DA tasks across low-risk and high-risk groups*** One-way between-groups ANCOVAs  
687 with age entered as a covariate were run to evaluate group differences on the DA measures. Adjusted mean  
688 scores for each group on all tasks can be seen in Table 7.

689

690 [Table 7 about here]

691

692 *Receptive Language*

693 Age was significantly related to scores on the DA of receptive language within the ANCOVA model ( $F(1, 88) =$   
694  $73.669, p < .001$ ). Significant group differences were found on the dynamic receptive language task ( $F(2, 88)$

695 = 5.218,  $p = .007$ ,  $\eta^2_p = 0.106$ )<sup>3</sup>. Scores in the SHR group were significantly lower than those of the low-risk  
696 group ( $p = .002$ ) and the LHR group ( $p = .016$ ). Scores in the LHR group did not differ from the low-risk group  
697 ( $p = .558$ ). See Table 7 for adjusted means and SEs.

698

#### 699 *Motor Imitation*

700 The association between age and motor imitation scores was significant within the ANCOVA model ( $F(1, 88)$   
701 = 53.021,  $p < .001$ ). Scores on the DA of motor imitation did not differ significantly between groups ( $F(2, 88)$   
702 = 1.212,  $p = .302$ ,  $\eta^2_p = 0.027$ ), although the mean score of the SHR group was lower than those of the other  
703 two groups.

704

#### 705 *Response to Joint Attention*

706 The covariate age was significantly related to scores on the DA of response to joint attention within the  
707 ANCOVA model ( $F(1, 88) = 15.997$ ,  $p < .001$ ). The ANCOVA did not indicate significant differences between  
708 risk groups for response to joint attention scores ( $F(2, 88) = 0.511$ ,  $p = .602$ ,  $\eta^2_p = 0.011$ ), although the mean  
709 score of the SHR group was lower than for the low-risk and LHR groups.

710

#### 711 *Turn-taking*

712 Age was significantly related to score on the DA of turn taking within the ANCOVA model ( $F(1, 87) = 42.582$ ,  
713  $p < .001$ ). Turn taking scores showed no significant differences among risk groups ( $F(2, 87) = 0.461$ ,  $p = .633$ ,  
714  $\eta^2_p = 0.010$ ), although the mean scores in both high-risk groups were lower than for the low-risk group.

715

716

717

718

---

<sup>3</sup> Notably the UK-CDI receptive vocabulary scale (Alcock et al, 2017) was not sensitive enough to detect differences across these groups ( $F(2, 74) = 2.114$ ,  $p = .128$ ,  $\eta^2_p = 0.054$ ).

## 719 *Social requesting*

720 Age did not show significant relationship to scores on the DA of social requesting within the ANCOVA model  
721 ( $F(1, 86) = 2.879, p = .093$ ) and neither did the ANCOVA show significant group differences in scores on the  
722 dynamic requesting task ( $F(2, 86) = 1.028, p = .362, \eta^2_p = 0.023$ ).

723

724 This pattern of results was identical when combining the two high risk groups in comparison to the low-risk  
725 infants.

726

## 727 **Discussion**

728 The current study aimed to present ‘proof of concept’ findings from a set of new dynamic tasks of early  
729 communication for infants, with the long term aim of developing a tool that can be used reliably and easily  
730 within family homes. This study is unique in using DA to investigate the skills of infants at high risk of  
731 communication difficulties. The tasks presented here are a first step towards more reliable communication  
732 assessment of children at very early ages before most static measures are appropriate.

733

## 734 ***Task characteristics***

735 While the five tasks showed only weak inter-correlations and therefore appear to be measuring different  
736 constructs, each measure had good inter-rater reliability (across all groups). In addition, four of the five DA  
737 tasks showed a significant correlation with age, indicating sensitivity to developing abilities in children within  
738 the age range studied here. However, the fact that there were some floor and ceiling effects, indicates that  
739 the tasks may need to be refined to capture a fuller range of language and communicative potential in both  
740 clinical and typically developing populations. This would eventually enable the creation of norms so that  
741 individual child scores on each task can be interpreted appropriately on a clinical basis according to age. No  
742 sex differences were observed in our normative sample, suggesting that DA might in the future serve as a  
743 useful tool for equality of diagnosis across boys and girls.

744

745 The tasks also showed significant correlations with standardised measures for our normative, low-risk  
746 sample, suggesting that they are valid and are tapping important constructs relevant to communication. Our  
747 DA tasks were also sensitive enough to detect some early differences between risk groups, especially for  
748 receptive language, which was the only task out of the five to show statistically significant differences. The  
749 SHR group also received lower mean scores than low-risk infants on the motor imitation, point following and  
750 turn taking tasks, but these did not constitute significant differences. No differences were found between  
751 LHR children and the children in the low-risk group. These findings are now discussed in more detail below.

752

### 753 ***Receptive language and other group comparisons***

754 Receptive language was the only task to show significant differences across groups. Nevertheless, at this  
755 early stage of test development, the trend towards lower scores for children in the SHR group for turn  
756 taking, RJA and motor imitation (which all relate to the development of receptive language), is also worth  
757 noting and taking forward to the next iteration. Receptive language was also the only DA task to correlate  
758 with standardised tasks. This result in itself does not entirely limit the usefulness of the other DA tasks, since  
759 it may be that they are more sensitive than standardised tests at this age, or that they are measuring slightly  
760 different aspects of communication. However, taken together, our results highlight receptive language as  
761 the most promising early assessment domain, especially for children at risk of autism. This is particularly  
762 interesting given that receptive language has been found to be a strong predictor of language outcome both  
763 for children with autism and for late talkers (Brignall et al, 2019; Fisher, 2017).

764

765 We were somewhat surprised that the LHR group showed no differences compared to the low-risk group.  
766 This may be due to the group criteria including family members with dyslexia, which is diagnosed later, or  
767 because the pathway of difficulties for those with language disorders is more gradual and less identifiable in  
768 infancy. Notably receptive language was not different for the LHR group whereas this was already showing  
769 signs of impact for the SHR group. Several research studies have demonstrated the instability between early  
770 language delay and later language impairment (e.g. Dale et al, 2003; Reilly et al, 2010; Zambrana et al, 2014;



771 Duff et al, 2015; Rudolph et al, 2016). It may be the case that group differences based on *family history* were  
772 not evident here, but that individual children who later receive diagnoses of communication difficulties will  
773 show differences on the DA tasks as infants, indicating their predictive validity. The crucial aspect in  
774 validating these DA tasks is whether they can be used to identify children who require support early on.  
775 Work is ongoing to follow up the current cohort at school age to investigate this very question.

776

### 777 ***Strengths and limitations of the present study***

778 The present study addressed a number of key gaps in the literature. Firstly, studies of language high-risk  
779 children are few, while numerous studies of social-high-risk children exist. Although few differences were  
780 evident between LHR and low-risk children, it may be that these will manifest later in childhood, particularly  
781 in terms of literacy outcomes (e.g., Zambrana et al. 2014). Secondly, this study is one of only a few studies to  
782 use DA methods to assess infants, particularly infants at high risk of communication disorders.

783

### 784 ***Age of the children***

785 In order to recruit a sufficient sample, the age range of the children was wider than ideal. Our key aim for  
786 developing the tool was to keep the age range of the *low-risk children* reasonably tight and this was  
787 achieved, with only 3 of this group older than 16 months. However, the age range of the social high-risk  
788 group was wider. As with most clinical measures, we anticipate that a tool of this kind may be useful for  
789 identifying older children who are at risk of language and communication difficulties, and who are  
790 functioning at a lower level than expected for age. Attempts were made to control for age effects in  
791 statistical analyses, but it is acknowledged that results would be clearer and easier to interpret in a cohort  
792 that had a narrower age range at each assessment time point. There is also a suggestion in existing research  
793 that the profiles of children with language difficulties change with age, such that social communication  
794 difficulties and features relevant to ASD diagnosis become more prominent over time in children whose  
795 language difficulties appeared more specific earlier in childhood (e.g., Conti-Ramsden et al, 2006 Chiat &  
796 Roy, 2013; Charman et al, 2015). Replication of similar results in a sample with higher proportions of high-

797 risk children, and with follow-up of the sample at later ages, would lend weight to the conclusions of this  
798 study, and would allow analysis of some trends that did not reach significance in the current sample, but  
799 appeared to have large effect size.

800

#### 801 *Sample diversity*

802 In addition, the self-selecting nature of the sample means that results are not necessarily generalisable to  
803 the total population of young children. The parents who responded to recruitment advertising were typically  
804 educated at degree level or higher, were older than 30, and had higher levels of family income than the  
805 population as a whole. Their children may therefore not accurately represent children from a broader range  
806 of socio-economic status. It would therefore be useful to recruit a sample of participants more diverse in  
807 socio-economic factors such as family income and parental education, to assess whether this affects the  
808 pattern of results. Although 7 children came from bilingual families, removing them from analyses did not  
809 affect results. However, it was not possible to statistically compare children exposed to other languages to  
810 those in monolingual households due to small numbers. Increasing the diversity of the sample would be an  
811 important step for future research as this would help to establish the utility of culturally-sensitive  
812 assessments. For the purposes of this study the only language tested was English, but we acknowledge that  
813 in clinical practice, all home languages should be assessed. One of the potential strengths of DA is that tests  
814 may be more easily adapted to work across several languages.

815

816 It is also the case that parents who already had concerns about the development of their child may have  
817 been more likely to enrol them to participate in this research project, so that their communication  
818 development could be evaluated. It is therefore possible that the groups of high-risk children who  
819 participated in the study contain higher numbers of children with developmental communication difficulties  
820 than would be the case in an unselected sample, although our results suggest this is not the case. Indeed,  
821 the opposite may also be true, in that parents with concerns about their second child might have avoided a  
822 study where issues could have been revealed.

823 *Measurement issues*

824 Another potential limitation was that some of the siblings were too young to complete the CCC-2 (although  
825 we introduced alternative criteria for these few) and some parents did not return sibling questionnaires.  
826 Furthermore, there was also no direct assessment of sibling/parent probands with communication or  
827 literacy difficulties and classification of children was dependent on parental reports. Together, these  
828 limitations reveal that establishing sibling status is not a straightforward process. In future research, more  
829 objective classification using direct measures of parent and elder sibling language, social communication and  
830 literacy skills could be of benefit.

831

832 The range of measures taken in this study allowed evaluation across many areas of development, and also  
833 allowed the comparison of novel tasks with established measures for a normative sample. However, it is  
834 important to note that the assessments were not blinded, as the first author and assessor had knowledge of  
835 the children's risk status. Nevertheless, good inter-rater reliability was achieved for novel tasks after coding  
836 by researchers who were blind to the children's group status. We did not include a DA of expressive  
837 vocabulary. This is because we felt it would be difficult to define a cuing hierarchy with children who were  
838 mostly non-verbal, but assessment of utterances or vocalisations could be explored in further studies.

839

840 Finally, the fact that there was only one task per skill may also serve to limit the assessment battery. It  
841 should be noted however, that infants are very restricted in terms of the time and attention they can apply  
842 to formal tasks. Indeed, we originally intended all DA items to have 10 trials, but our feasibility pilot  
843 suggested that only RJA was tolerated sufficiently for this number. The DA tasks presented here are being  
844 developed with a view to offering a quick yet reliable addition to tools used by health visitors and other  
845 infant-care professionals.

846

847

848

### ***Future directions and Dynamic Assessment in practice***

Some adaptations could be useful in future versions of our tasks. In particular, it may be beneficial to combine some of the tasks into one composite subtask tapping into early non-verbal communication skills. The receptive vocabulary task could also be modified to incorporate novel word learning opportunities; an expressive task could be included for older children; and the scoring of tasks could be standardised across areas. Other aspects of communication which are emerging, and may be more easily measurable than vocabulary (such as vocalizations), might also be useful target behaviours to include as part of a DA of early communication. Thus, further development of the DA tasks should be a focus of future research, including data from large diverse samples, trials for additional items and more detailed investigations into reliability and validity of the measures especially at an individual level. Ultimately, if key tasks differentiating children at high risk of later language delay and/or social communication difficulties could be identified through further studies, it would be possible to trial intervention programmes for children who show early signs of these difficulties and evaluate the effect on outcomes. It may be that the children in our high-risk groups do not go on to develop communication disorders, despite their family histories and conversely, some children in the low-risk group may develop a communication disorder. Therefore, the utility and inter-rater reliability established for the DA tasks in the current study warrant future work on developing and formalising these tasks in order to improve the prediction of future difficulties.

In spite of the need for further development of these procedures, we concur with Hasson & Joffe (2007), in believing that dynamic approaches are a promising way forward for providing practitioners with a reliable user-friendly screening tool for identifying infants at risk. DA tasks have the advantage of being a quick to administer direct assessment that is infant-friendly and ecologically valid in contrast to existing standardised tools that are often used inappropriately (Dockrell & Marshall, 2015; Cych et al, 2021; Spaulding, Plante & Farinella, 2006; Betz et al, 2013). We have shown that they can also be carried out with reliability. However, further work is needed to establish clinical discrimination between individual children who will need language and social support, and those who do not (Szatmari et al. 2016).

875

876 Lastly, for DA to be used in practice, the issue of training would need developing and evaluating at an  
877 individual case level. In the present study, all assessments were completed by the same Speech and  
878 Language Therapist (SLT; first author), but in practice any infant assessments would ideally be available to a  
879 wider group of professionals such as health visitors following full test development. Continuing work  
880 suggests that this training would be much easier and quicker than for most DAs, but rigorous further  
881 development of the tool is needed before any clinical implementation.

882

### 883 ***Conclusions and implications***

884 This study suggests that Dynamic Assessment for infants may be feasible and useful, especially in the domain  
885 of receptive language. Many children with communication disorders are still being identified too late to  
886 access the critical early intervention they need, and appropriate standardised tests are not currently  
887 available. Although we have focussed here on children at risk of Autism and DLD, our hope for the long term  
888 is that tools can be developed to identify communication issues in a wide range of children.

889 We acknowledge that there are numerous barriers to early identification, especially in the UK. However, if  
890 parents and professionals were to have a reliable screening tool to identify the key markers of  
891 communication disorders in early life, this may increase the number of children identified before the age of  
892 two or three. This would in turn allow more children to access intervention designed to improve outcomes.  
893 This work represents just the first steps towards such a tool. Further longitudinal work will play a key part in  
894 determining which skills in infancy are predictive of communication problems not just at preschool age, but  
895 into the school years and adolescence. Only then can children receive earlier intervention and thereby attain  
896 more positive academic and psychosocial outcomes later in life.

897

898

899

900

901 **Acknowledgements**

902 The authors would like to thank all the families who took part in this study, City University of London for  
903 doctoral funding to the first author, and the Heather van der Lely Trust for supporting follow up work  
904 resulting from this project.

905

906 **Data Availability Statement**

907 The data produced by this study is preliminary and therefore is not hosted on an open access platform.  
908 However, data requests can be made to the first or corresponding author.

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926 **References**

927

928 Alcock, K., Meints, K., & Rowland, C. (2020). *The UK Communicative Development Inventories: Words and*  
929 *Gestures*. Havant: J&R Press.

930

931 American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders, 5th Edition*.  
932 Washington DC: American Psychiatric Association.

933

934 Bain, B. A. & Olswang, L. B. (1995). Examining readiness for learning two-word utterances by children with  
935 specific expressive language impairment: Dynamic assessment validation. *American Journal of Speech-*  
936 *Language Pathology*, 4(1), 81-91

937

938 Baron-Cohen, S., Scott, F., Allison, C., Williams, J., Bolton, P., Matthews, F., & Brayne, C. (2009). Prevalence of  
939 autism spectrum conditions: UK school-based population study. *The British Journal of Psychiatry*, 194, 500-  
940 509.

941

942 Bent, C.A., Barbaro, J. & Dissanayke, C. (2020). Parents' experiences of the service pathway to an autism  
943 diagnosis for their child: what predicts an early diagnosis in Australia? *Research in Developmental*  
944 *Disabilities*, 103, epub 103689.

945

946 Bercow, J. (2008). *The Bercow Report: A Review of Services for Children and Young People (0-19) with Speech,*  
947 *Language and Communication Needs*. Nottingham: DCSF.

948

949 Betz, S.K., Eickhoff, J.R., & Sullivan, S.F. (2013). Factors influencing the selection of standardized tests for the  
950 diagnosis of specific language impairment. *Language Speech and Hearing Services in Schools*, 44, 133–146.

951

952 Bishop, D.V.M. (2003). *The Children's Communication Checklist, 2nd Edition*. London: Pearson.

953

954 Bishop D.V.M. (2010). Which neurodevelopmental disorders get researched and why? *PLoS ONE* 5(11):

955 e15112.

956

957 Bishop, D., Snowling, M., Thompson, P., Greenhalgh, T., & Consortium (2017). Phase 2 of CATALISE: a multi-

958 national and multi-disciplinary Delphi consensus study of problems with language development:

959 terminology. *Journal of Child Psychology and Psychiatry*, 58(10), 1068-1080.

960

961 Botting, N. (2020). Language, literacy and cognitive skills of young adults with developmental language

962 disorder (DLD). *International Journal of Language & Communication Disorders*, 55(2), 255-265.

963

964 Botting N., Durkin K., Toseeb U., Pickles A. & Conti-Ramsden G. (2016). Emotional health, support, and self-

965 efficacy in young adults with a history of language impairment, *British Journal of Developmental Psychology*,

966 34, 538–554.

967

968 Boyle, J. (2011). Speech and language delays in preschool children. *British Medical Journal*, 343, d5181.

969

970 Brignell, A., May, T., Morgan, A. T., & Williams, K. (2019). Predictors and growth in receptive vocabulary from

971 4 to 8 years in children with and without autism spectrum disorder: A population-based study. *Autism*, 23(5),

972 1322-1334.

973

974 Broomfield, J., & Dodd, B. (2011). Children with speech and language disability: caseload characteristics.

975 *International Journal of Language and Communication Disorders*, 39(3), 303-324.

976



977 Camilleri, B. & Botting, N. (2013). Beyond static assessment of children's receptive vocabulary: the dynamic  
 978 assessment of word learning (DAWL). *International Journal of Language and Communication Disorders*,  
 979 48(5), 565-581.

980

981 Camilleri, B., Hasson N., & Dodd, B. (2014). Dynamic assessment of bilingual children's language at the point  
 982 of referral. *Educational & Child Psychology*, 31(2) 58-73.

983

984 Camilleri, B., & Law, J. (2007). Assessing children referred to speech and language therapy: Static and  
 985 dynamic assessment of receptive vocabulary. *Advances in Speech Language Pathology*, 9(4), 312-322.

986

987 Campione, J. C., & Brown, A. L. (1987). *Linking dynamic assessment with school achievement*. In C. S. Lidz  
 988 (Ed.), *Dynamic assessment: An interactional approach to evaluating learning potential*. New York: Guilford.

989

990 Charman, T., Ricketts, J., Dockrell, J., Lindsay, G., & Palikara, O. (2015). Emotional and behavioural problems  
 991 in children with language impairments and children with autism spectrum disorders. *International Journal of*  
 992 *Language and Communication Disorders*, 50(1), 84-93.

993

994 Chen, J. L., Leader, G., Sung, C., & Leahy, M. (2015). Trends in employment for individuals with autism spectrum  
 995 disorder: A review of the research literature. *Review Journal of Autism and Developmental Disorders*, 2(2), 115-127.

996

997 Chiat, S. & Roy, P. (2007). The preschool repetition test: an evaluation of performance in typically developing and  
 998 clinically referred children. *Journal of Speech, Language and Hearing Research*, 50, 429-443.

999

1000 Chiat, S., & Roy, P. (2013). Early predictors of language and social communication impairments at aged 9-11  
 1001 years: a follow-up study of early-referred children. *Journal of Speech, Language and Hearing Research*, 56(6),  
 1002 1824-1836.

1003

1004 Christensen, D., Bilder, D., Zahorodny, W., Pettygrove, S., Durkin, M., Fitzgerald, R., Rice, C., Kurzius-Spencer,  
1005 M., Baio, J. & Yeargin-Allsop, M. (2016). Prevalence and characteristics of autism spectrum disorder among 4  
1006 year old children in the Autism and Developmental Disabilities Monitoring Network. *Journal of*  
1007 *Developmental Behavioural Paediatrics*, 37(1), 1-8.

1008

1009 Conti-Ramsden, G., Simkin, Z., & Botting, N. (2006). The prevalence of autism spectrum disorders in  
1010 adolescents with a history of specific language impairment (SLI). *Journal of Child Psychology and Psychiatry*,  
1011 47(6), 621-628.

1012

1013 Conti-Ramsden, G., Durkin, K., Toseeb, U., Botting, N. and Pickles, A. (2018). Education and employment  
1014 outcomes of young adults with a history of developmental language disorder. *International Journal of*  
1015 *Language and Communication Disorders*, 53(2), pp. 237–255.

1016

1017 Crane, L., Chester, J., Goddard, L., Henry, L., & Hill, E. (2016). Experiences of autism diagnosis: a survey of  
1018 over 1000 parents in the United Kingdom. *The International Journal of Research and Practice*, 20(2), 153-162.

1019

1020 Cocyk, L. M., De Anda, S., Moore, H., & Huerta, L. (2021). Cultural and linguistic adaptations of early language  
1021 interventions: Recommendations for advancing research and practice. *American Journal of Speech-Language*  
1022 *Pathology*, 30(3), 1224-1246.

1023

1024 Dale, P., Price, T., Bishop, D., & Plomin, R. (2003). Outcomes of early language delay: I. Predicting persistent  
1025 and transient language difficulties at 3 and 4 years. *Journal of Speech, Language and Hearing Research*,  
1026 46(3), 544-560.

1027

1028 Daniels, A., & Mandel, D. (2014). Explaining differences in age at autism spectrum disorder diagnosis: a  
1029 critical review. *Autism*, 18(5), 583-597.

1030

1031 Davidson, M., Alonzo, C. & Stransky, M. (2022). Access to speech and language services and service providers  
 1032 for children with speech and language disorders. *American Journal of Speech-Language Pathology*, online  
 1033 ahead of print.

1034

1035 De Giacomo, A., & Fombonne, E. (1998). Parental recognition of developmental abnormalities in autism.  
 1036 *European Child and Adolescent Psychiatry*, 7(3), 131-136.

1037

1038 Deutsch, R., & Reynolds, Y. (2000). The use of dynamic assessment by educational psychologists in the UK.  
 1039 *Educational Psychology in Practice*, 16(3), 311-331.

1040

1041 Dillenburger, K., Jordan, J., McKerr, L., & Keenan, M. (2015). The millennium child with autism: early  
 1042 childhood trajectories for health, education and economic wellbeing. *Developmental Neurorehabilitation*,  
 1043 18(1), 37-46.

1044

1045 Dockrell, J. E., & Marshall, C. R. (2015). Measurement issues: assessing language skills in young children.  
 1046 *Child and Adolescent Mental Health*, 20(2), 116-125.

1047

1048 Dubois, P., St-Pierre, M. C., Desmarais, C., & Guay, F. (2020). Young adults with developmental language  
 1049 disorder: a systematic review of education, employment, and independent living outcomes. *Journal of*  
 1050 *Speech, Language, and Hearing Research*, 63(11), 3786-3800.

1051

1052 Duff, F., Reen, G., Plunkett, K., & Nation, K. (2015). Do infant vocabulary skills predict school-age language  
 1053 and literacy outcomes? *Journal of Child Psychology and Psychiatry*, 56(8), 848-856.

1054

1055 Filipek, P.A., Accardo, P.J., Baranek, G.T., Cook, E.H., Dawson, G., Gordon, B., Gravel, J.S., Johnson, C.P.,  
 1056 Kallen, R.J., Levy, S.E., Minshew, N.J., Prizant, B.M., Rapin, I. Rogers, S.J., Stone, W.L., Teplin, S., Tuchman,  
 1057 R.F. & Volkmar, F. (1999). The screening and diagnosis of autism spectrum disorders. *Journal of Autism and*  
 1058 *Developmental Disorders*, 29(6), 439-484.  
 1059  
 1060 Fisher, E. L. (2017). A systematic review and meta-analysis of predictors of expressive-language outcomes  
 1061 among late talkers. *Journal of Speech, Language, and Hearing Research*, 60(10), 2935-2948.  
 1062  
 1063 Friberg, J. (2010). Considerations for test selection: how do validity and reliability impact diagnostic  
 1064 decisions. *Child Language Teaching and Therapy*, 26(1), 77-92.  
 1065  
 1066 Hanika, L., & Boyer, W. (2019). Imitation and social communication in infants. *Early Childhood Education*  
 1067 *Journal*, 47(5), 615-626.  
 1068  
 1069 Hanley, A., Nguyen, Q., Badawi, D., Chen, J., Ma, T. & Slopen, N. (2021). The diagnostic odyssey of autism: a  
 1070 cross-sectional study of 3 age cohorts of children from the 2016-2018 National Survey of Children's Health.  
 1071 *Child and Adolescent Psychiatry and Mental Health*, 15(58).  
 1072  
 1073 Harmuth, E., Silletta, E., Bailey, A., Adams, T., Beck, C., & Barbic, S. P. (2018). Barriers and facilitators to  
 1074 employment for adults with autism: A scoping review. *Annals of International Occupational Therapy*, 1(1),  
 1075 31-40.  
 1076  
 1077 Hasson, N. & Joffe, V. (2007). The case for dynamic assessment in speech and language therapy. *Child*  
 1078 *Language Teaching and Therapy*, 23(1), 9-25.  
 1079

1080 Hasson, N. & Botting, N. (2010). Dynamic assessment of children with language impairments: a pilot study.  
1081 *Child Language Teaching and Therapy*, 29, 52-70.

1082

1083 Hasson, N., Camilleri, B., Jones, C., Smith, J., & Dodd, B. (2013). Discriminating disorder from difference:  
1084 using dynamic assessment with bilingual children. *Child Language Teaching and Therapy*, 29(1), 57-75.

1085

1086 Hedenbro, M., & Rydelius, P. A. (2014). Early interaction between infants and their parents predicts social  
1087 competence at the age of four. *Acta Paediatrica*, 103(3), 268-274

1088

1089 Henry, L.A. and Botting, N. (2017). Working memory and developmental language impairments. *Child*  
1090 *Language Teaching and Therapy*, 33(1), 19–32.

1091

1092 Hollocks, M. J., Lerh, J. W., Magiati, I., Meiser-Stedman, R., & Brugha, T. S. (2019). Anxiety and depression in  
1093 adults with autism spectrum disorder: a systematic review and meta-analysis. *Psychological Medicine*, 49(4),  
1094 559-572.

1095

1096 Huerta, M., & Lord, C. (2012). Diagnostic evaluation of autism spectrum disorders. *Pediatric Clinics*, 59(1),  
1097 103-111.

1098

1099 Hunt, E., Nang, C., Meldrum, S. & Armstrong, E. (2019). Can dynamic assessment identify language disorder  
1100 in multilingual children? Clinical applications from a systematic review. *Language, Speech and Hearing*  
1101 *Services in Schools*, 53, 598-625.

1102

1103 Hussain, S., & Woods, K. (2019). The use of dynamic assessment by educational psychologists in the early  
1104 years foundation stage. *Educational Psychology in Practice*, 35(4), 424-439.

1105

1106 ICAN & RCSLT (2018). *Bercow: Ten years on*. London: ICAN & RCSLT.

1107

1108 Jullien, S. Secreening for language and speech delay in children under five years. *BMC Pediatrics*, 21 (Suppl

1109 1), 362.

1110

1111 Kapantzoglou, M., Restrepo, M. A., & Thompson, M. S. (2012). Dynamic assessment of word learning skills:

1112 Identifying language impairment in bilingual children. *Language, Speech and Hearing Services in Schools*, 43

1113 (1), 81-96.

1114

1115 Kim, Z., Oturai G., Kiraly, I. & Knopf, M. (2015). The role of object and effects in action imitation: comparing

1116 the imitation of object-related actions vs gestures in 18-month-old infants. *Infant Behaviour and*

1117 *Development*, 41, 43-51.

1118

1119 Koo, T.K. & Li, M.Y. (2016). A guideline of selecting and reporting intraclass correlation coefficients for

1120 reliability research. *Journal of Chiropractic Medicine*, 15(2), 155–163.

1121

1122 Law, J. & Roy, P. (2008). Parental report of infant language skills: a review of the development and

1123 application of the communicative development inventories. *Child and Adolescent Mental Health*, 13(4), 198-

1124 206.

1125

1126 Law, J., Charlton, J., Dockrell, J., Gascoigne, M., McKean, C., & Theakston, A. (2017). *Early Language*

1127 *Development: Needs, provision and intervention for pre-school children from socio-economically*

1128 *disadvantaged backgrounds*. London: Education Endowment Foundation.

1129

1130 Law, J., Charlton, J., McKean, C., Watson, R., Roulstone, S., Holme, C., Gilroy, V., Wilson, P. & Rush, R. (2020).  
 1131 *Identifying and supporting children's early language needs*. London: Department of Education and Public  
 1132 Health England.

1133

1134 Lindsay, G., Desforges, M., Dockrell, J., Law, J., Peacey, N. & Beecham, J. (2008). *Effective and efficient use of*  
 1135 *resources in services for children and young people with speech, language and communication needs*.  
 1136 Warwick: DCSF.

1137

1138 MacLeod, A. A., & Glaspey, A. M. (2022). Dynamic assessment of multilingual children's word learning.  
 1139 *International Journal of Language & Communication Disorders*, Early Online.

1140

1141 Maenner, M.J., Shaw, K.A., Bakian, A.V., Bilder, D.A., Durkin, M.S., Esler, A., Furnier, S.M., Hallas, L., Hall-  
 1142 Lande, J., Hudson, A., Hughes, M.M., Patrick, M., Pierce, K., Poynter, J.N., Salinas, A., Shenouda, J., Vehorn,  
 1143 A., Warren, Z., Constantino, J.N., DiRienzo, M., Fitzgerald, R.T., Grzybowski, A., Spivey, M.H., Pettygrove, S.,  
 1144 Zahorodny, W., Ali, A., Andrews, J.G., Baroud, T., Gutierrez, J., Hewitt, A., Lee, L.C., Lopez, M., Mancilla, K.C.,  
 1145 McArthur, D., Schwenk, Y.D., Washington, A., Williams, S. & Cogswell, M.E. Prevalence and characteristics of  
 1146 Autism Spectrum Disorder among children aged 8 years - Autism and Developmental Disabilities Monitoring  
 1147 Network, 11 Sites, United States, 2018. *MMWR Surveillance Summaries*, 70(11), 1-16.

1148

1149 MacGregor, K. (2020). How we fail children with developmental language disorder. *Language, Speech and*  
 1150 *Hearing Services in Schools*, 51(4), 981-992.

1151

1152 Markus, J., Mundy, P., Morales, M., Delgado, C. E., & Yale, M. (2000). Individual differences in infant skills as  
 1153 predictors of child-caregiver joint attention and language. *Social Development*, 9(3), 302-315.

1154

1155 Meilleur, A., & Fombonne, E. (2009). Regression of language and non-language skills in pervasive  
 1156 developmental disorders. *Journal of Intellectual Disability Research*, 53(2), 115-124.  
 1157

1158 Messick, S. (1998). Test validity: A matter of consequence. *Social Indicators Research*, 45(1), 35-44.  
 1159

1160 Muskett, T., Body, R., & Perkins, M. (2012). Uncovering the dynamic in static assessment interaction. *Child*  
 1161 *Language Teaching and Therapy*, 28(1), 87-99.  
 1162

1163 Norbury, C., Gooch, D., Wray, C., Baird, G., Charman, T., Simonoff, E., Vamvakas, G. & Pickles, A. (2016). The  
 1164 impact of non-verbal ability on prevalence and clinical presentation of language disorder: evidence from a  
 1165 population study. *Journal of Child Psychology and Psychiatry*, 57(11), 1247-1257.  
 1166

1167 Olswang, L.B. & Bain, B. A. (1996). Assessment information for predicting upcoming change in language  
 1168 production. *Journal of Speech and Hearing Research*, 39, 414-423.  
 1169

1170 Orellana, C. I., Wada, R., & Gillam, R. B. (2019). The use of dynamic assessment for the diagnosis of language  
 1171 disorders in bilingual children: A meta-analysis. *American Journal of Speech-Language Pathology*, 28(3),  
 1172 1298-1317.  
 1173

1174 Patterson, J. L., Rodríguez, B. L., & Dale, P. S. (2020). Dynamic assessment language tasks and the prediction  
 1175 of performance on year-end language skills in preschool dual language learners. *American Journal of Speech-*  
 1176 *Language Pathology*, 29(3), 1226-1240.  
 1177

1178 Peña, E.D., Iglesias, A., & Lidz, C.S. (2001). Reducing test bias through dynamic assessment of children's word  
 1179 learning ability. *American Journal of Speech-Language Pathology*, 10, 138-154.  
 1180



1181 Peña, E.D., Gillam, R.B., Malek, M., Felter, R., Resendiz, M. & Fiestas, C. (2006). Dynamic assessment of  
 1182 children from culturally diverse backgrounds: Application to narrative assessment. *Journal of Speech,*  
 1183 *Language and Hearing Research, 49*, 1037-1057.

1184

1185 Peña, E., Resendiz, M. & Gillam, R. (2007). The role of clinical judgements of modifiability in the diagnosis of  
 1186 language impairment. *Advances in Speech-Language Pathology, 9*(4), 332-345.

1187

1188 Peña, E. D., Gillam, R. B., & Bedore, L. M. (2014). Dynamic assessment of narrative ability in English  
 1189 accurately identifies language impairment in English language learners. *Journal of Speech, Language, and*  
 1190 *Hearing Research, 57*(6), 2208-2220.

1191

1192 Peña, E. D., Spaulding, T. J., & Plante, E. (2006). The composition of normative groups and diagnostic decision  
 1193 making: Shooting ourselves in the foot. *American Journal of Speech-Language Pathology, 15*, 247–254.

1194

1195 Ramos-Cabo, S., Vulchanov, V., & Vulchanova, M. (2019). Gesture and language trajectories in early  
 1196 development: An overview from the autism spectrum disorder perspective. *Frontiers in Psychology, 10*,  
 1197 1211.

1198

1199 Rannard, A., Lyons, C., & Glenn, S. (2004). Children with specific language impairment: parental accounts of  
 1200 the early years. *Journal of Child Health Care, 8*(2), 165-176.

1201

1202 Rattaz, C., Loubersac, J., Michelon, C., Geoffray, M.M., Picot, M.C., Munir, K. & Baghdadli, A. (2022). Factors  
 1203 associated with age of diagnosis in children with autism spectrum disorders: report from a French cohort.  
 1204 *Autism*, Online ahead of print.

1205

1206 Reilly, S., Wake, M., Ukomunne, O., Bavin, E., Prior, M., Cini, E., Conway, L., Eadie, P. & Bretherton, L. (2010).  
1207 Predicting language outcomes at 4 years of age: findings from the Early Language in Victoria Study.  
1208 *Pediatrics*, 126(6), e1530-e1537.  
1209  
1210 Reilly, S., McKean, C., Morgan, A. & Wake, M. (2015). Identifying and managing common childhood language  
1211 and speech impairments. *British Medical Journal*, 350, h2318.  
1212  
1213 Robins, D., Fein, D., Barton, M., & Green, J. (2001). The Modified Checklist for Autism in Toddlers: an initial  
1214 study investigating the early detection of autism and pervasive developmental disorders. *Journal of Autism*  
1215 *and Developmental Disorders*, 31(2), 131-144.  
1216  
1217 Rogge, N., & Janssen, J. (2019). The economic costs of autism spectrum disorder: A literature review. *Journal*  
1218 *of Autism and Developmental Disorders*, 49(7), 2873-2900.  
1219  
1220 Rohlfsing, K. J. (2019). Learning language from the use of gestures. In J. S. Horst & J. von Koss Torkildsen  
1221 (Eds.), *International Handbook of Language Acquisition* (pp. 213–233). London: Routledge.  
1222  
1223 Roman-Urrestarazu, A., van Kessel, R., Allison, C., Matthews, F.E., Brayne, C. & Baron-Cohen, S. Association  
1224 of race/ethnicity and social disadvantage with autism prevalence in 7 million school children in England.  
1225 *JAMA Pediatrics*, 175(6), e210054.  
1226  
1227 Roulstone, S., Marshall, J., Powell, G., Goldbart, J., Wren, Y., Coad, J., Daykin, N., Powell, J., Lascelles, L.,  
1228 Hollingworth, W., Emond, A., Peters, T., Pollock, J., Fernandes, C., Moultrie, J., Harding, S., Morgan, L.,  
1229 Hambly, H., Parker, N. & Coad, R. (2015). Evidence-based intervention for preschool children with primary  
1230 speech and language impairments: Child Talk – an exploratory mixed methods study. *Programme Grants for*  
1231 *Applied Research*, 3(5).

1232

1233 Rudolph, J., & Leonard, L. (2016). Early language milestones and specific language impairment. *Journal of*

1234 *Early Intervention*, 38(1), 41-58.

1235

1236 Salo, V. C., Rowe, M. L., & Reeb-Sutherland, B. C. (2018). Exploring infant gesture and joint attention as

1237 related constructs and as predictors of later language. *Infancy*, 23(3), 432-452.

1238

1239 Spaulding, T. J., Plante, E., & Farinella, K. A. (2006). Eligibility criteria for language impairment. *Language,*

1240 *Speech and Hearing Services in Schools*. 37, 61-72.

1241

1242 Spaulding, T., Szulga, M. & Figueroa, C. (2012). Using norm-referenced tests to determine severity of

1243 language impairment in children: disconnect between US policy makers and test developers. *Language,*

1244 *Speech and Hearing Services in Schools*, 43, 176-190.

1245

1246 Spicer-Cain, H. (2019). *Early markers of communication difficulties in children at high risk of Autism Spectrum*

1247 *Disorder and Language Impairment*. [Unpublished doctoral thesis]. City, University of London.

1248

1249 Sternberg, R. J., & Grigorenko, E. L. (2002). *Dynamic testing: The nature and measurement of learning*

1250 *potential*. New York: Cambridge University Press.

1251

1252 Szatmari, P., Chawarska, K., Dawson, G., Georgiades, S., Landa, R., Lord, C., ... & Halladay, A. (2016).

1253 Prospective longitudinal studies of infant siblings of children with autism: lessons learned and future

1254 directions. *Journal of the American Academy of Child & Adolescent Psychiatry*, 55(3), 179-187.

1255

1256 Thordardottir, E., Tobpas, S. & Working Group 3 of COST Action IS1406. (2021). How aware is the public of  
 1257 the existence, characteristics and causes of language impairment in childhood and where have they heard  
 1258 about it? A European survey. *Journal of Communication Disorders*, 89, 106057.

1259

1260 Tomblin, J., Records, N., Buckwalter, P., Zhang, X., Smith, E., & O'Brien, M. (1997). Prevalence of specific  
 1261 language impairment in kindergarten children. *Journal of Speech, Language and Hearing Research*, 40(6),  
 1262 1345-1260.

1263

1264 Ukrainetz, T.A., Harpell, S., Walsh, C., & Coyle, C. (2000). A preliminary investigation of dynamic assessment  
 1265 with Native American kindergartners. *Language, Speech and Hearing Services in Schools*, 31, 142-154.

1266

1267 van 't Hof, M., Tisseur, C., van Berckeleer-Onnes, I., van Nieuwenhuyzen, A., Daniels, A.M., Deen, M., Hoek,  
 1268 H.W. & Ester, W.A. (2021). Age at autism spectrum disorder diagnosis: a systematic review and meta-  
 1269 analysis from 2012 to 2019. *Autism*, 25(4), 862-873.

1270

1271 Veness, C., Prior, M., Eadie, P., Bavin, E., & Reilly, S. (2014). Predicting autism diagnosis by 7 years of age  
 1272 using parent report of infant social communication skills. *Journal of Paediatrics and Child Health*, 50(9), 693-  
 1273 700.

1274

1275 Vygotsky, L. S. (1978). *Mind in society: The development of higher psychological processes* Cambridge, MA.:  
 1276 Harvard University Press.

1277

1278 Wetherby, A., & Prizant, B. (2002). *Communication and Symbolic Behaviour Scales - Developmental Profile*.  
 1279 Baltimore: Brookes.

1280

1281 Zambrana, I., Pons, F., Eadie, P., & Ystrom, E. (2014). Trajectories of language delay from age 3 to 5:  
 1282 persistence, recovery and late onset. *International Journal of Language and Communication Disorders*, 49(3),  
 1283 304-316.

1284

1285 Zimmerman, I., Steiner, V., & Pond, R. (2002). *Preschool Language Scales, 4th Edition*. San Antonio: The  
 1286 Psychological Corporation.

1287

1288 Zwaigenbaum, L., Bryson, S., Rogers, T., Roberts, W., Brian, J., & Szatari, P. (2005). Behavioural  
 1289 manifestations of autism in the first year of life. *International Journal of Developmental Neuroscience*, 23,  
 1290 143-152.

1291

1292 Zwaigenbaum, L., Bryson, S., Lord, C., Rogers, S., Carter, A., Carver, L. C., Chawarska, K., Constantino, J.,  
 1293 Dawson, G., Dobkins, K., Fein, D., Iverson, J., Klin, A., Landa, R., Messinger, D., Ozonoff, S., Sigman, M., Stone,  
 1294 W., Tager-Flusberg, H. & Yirmiya, N. (2009). Clinical assessment and management of toddlers with suspected  
 1295 autism spectrum disorder: insights from studies of high-risk infants. *Pediatrics*, 123(5), 1383-1391.

1296

1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307 *Table 1: Demographic information for the whole sample by group*

	Whole Sample (n=92)	Low-risk Group (n=51)	SHR Group (n=18)	LHR Group (n=23)
Child Age (mean/SD)	13.1(3.4)	12.2 (3.0)	15.4 (3.9)	13.2(3.0)
Child Sex				
Male	42 (45.6%)	26 (51.0%)	8 (44.4%)	8 (34.8%)
Female	50 (54.4%)	25 (49.0%)	10 (55.6%)	15 (65.2%)
Birth order				
1	40 (43.5%)	30 (58.8%)	0 (0%)	10 (43.5%)
2	36 (39.1%)	16 (31.4%)	12 (66.7%)	8 (34.8%)
3 or higher	16 (17.4%)	5 (9.8%)	6 (33.3%)	5 (21.7%)
Maternal Age				
25 or younger	2 (2.6%)	2 (4.7%)	0 (0%)	0 (0%)
26-30	11 (14.3%)	7 (16.3%)	1 (8.3%)	3 (13.6%)
31-35	39 (50.6%)	21 (48.8%)	6 (50.0%)	12 (54.5%)
36 or older	25 (32.5%)	13 (30.2%)	5 (41.7%)	7 (31.8%)
Paternal Age				
25 or younger	1 (1.3%)	1 (2.3%)	0 (0%)	0 (0%)
26-30	11 (14.1%)	7 (15.9%)	2 (16.7%)	2 (9.1%)
31-35	26 (33.3%)	14 (31.8%)	5 (41.7%)	7 (31.9%)
36 or older	40 (51.3%)	22 (50.0%)	5 (41.7%)	13 (59.1%)
Maternal Education				
No formal qualifications	1 (1.3%)	0 (0%)	0 (0%)	1 (4.5%)
Level 2 (GCSE or equivalent)	2 (2.6%)	0 (0%)	2 (16.7%)	0 (0%)
Level 3 (A-Level or equivalent)	4 (5.1%)	2 (4.5%)	2 (16.7%)	0 (0%)
Degree or equivalent	30 (38.5%)	20 (45.5%)	2 (16.7%)	8 (36.4%)
Postgraduate Qualification	41 (52.6%)	22 (50.0%)	6 (50.0%)	13 (59.1%)
Paternal Education				
No formal qualifications	1 (1.3%)	0 (0%)	0 (0%)	1 (4.5%)
Level 2 (GCSE or equivalent)	8 (10.3%)	3 (6.8%)	2 (16.7%)	3 (13.6%)
Level 3 (A-Level or equivalent)	9 (11.5%)	3 (6.8%)	3 (25.0%)	3 (13.6%)
Degree or equivalent	29 (37.2%)	20 (45.5%)	4 (33.3%)	5 (22.7%)
Postgraduate Qualification	31 (39.7%)	18 (40.9%)	3 (25.0%)	10 (45.5%)
Family Income				
£14,000 or less	3 (3.9%)	1 (2.3%)	1 (8.3%)	1 (4.8%)
£14,001 to £24,000	4 (5.3%)	1 (2.3%)	2 (16.7%)	1(4.8%)
£24,001 to £42,000	15 (19.7%)	8 (18.6%)	3 (25.0%)	4 (19.0%)
£42,001 or more	54 (71.1%)	33 (76.7%)	6 (50.0%)	15 (71.4%)

1308  
1309

1310 *Table 2: Descriptive information on DA tasks for low-risk infants*

Measure	Range of Scores	Mean (SD)	Median (IQR)
Receptive Language	0-15	4.02 (4.54)	2.0 (1-7)
Motor Imitation	0-15	4.38 (4.11)	3.5 (1-6.5)
Response to Joint Attention	0-30	19.54 (9.44)	23.0 (11.75-27)
Turn Taking	0-15	2.86 (3.79)	1.0 (0-5.25)
Social Requesting	0-15	10.0 (4.59)	12.0 (76.75-14)

1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332

1333 Table 3: Inter-rater reliability for each DA task for the whole sample and by group

Measure	Whole Sample ICC	Low-Risk ICC	SHR ICC	LHR ICC
Receptive Language	.902	.886	.966	.820
Motor Imitation	.869	.799	.918	.895
Response to Joint Attention	.958	.929	.977	.972
Turn Taking	.702	.815	.828	.386
Social Requesting	.852	.750	.971	.838

1334 \*ICC of 0.75-0.9 is considered ‘good’ whilst 0.9 and above is considered excellent.

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348

1349

1350

1351

1352

1353



1354 Table 4: Correlations between DA subtests for the low-risk group

	DA Receptive Language	DA Motor Imitation	DA Response to Joint attention	DA Turn Taking
DA Motor Imitation	$r = .210$ $p = .182$			
DA Response to Joint Attention	$r = .240$ $p = .126$	$r = -.025$ $p = .875$		
DA Turn Taking	$r = .009$ $p = .954$	$r = -.108$ $p = .496$	$r = .140$ $p = .377$	
DA Social Requesting	$r = .115$ $p = .467$	$r = .053$ $p = .739$	$r = -.003$ $p = .986$	$r = -.090$ $p = .570$

1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371

1372     *Table 5: Mann Whitney statistics for sex comparisons on DA tasks in the low-risk group*

	Mann-Whitney <i>U</i>	<i>P</i> value	1373
	statistic		1374
Receptive	282.5	.449	1375
Language			1376
Motor	231.5	.085	1377
Imitation			1378
Response to Joint	307.0	.776	1379
Attention			1380
Turn-Taking	250.5	.155	1381
			1382
Social Requesting	292.5	.761	1383
			1384

1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404

1405 **Table 6: Relationships between DA tasks and standardised tests for the low-risk group**

	DA Receptive Language	DA Motor Imitation	DA Response to Joint Attention	DA Turn Taking	DA Social Requesting
UKCDI	<b><i>r</i> = .389</b>	<i>r</i> = .182	<i>r</i> = .258	<i>r</i> = .071	<i>r</i> = .144
Receptive Vocabulary	<b><i>p</i> = .011</b>	<i>p</i> = .249	<i>p</i> = .100	<i>p</i> = .653	<i>p</i> = .362
UKCDI	<i>r</i> = .154	<i>r</i> = .169	<i>r</i> = -.080,	<i>r</i> = -.285,	<b><i>r</i> = .404</b>
Expressive Vocabulary	<i>p</i> = .331	<i>p</i> = .285	<i>p</i> = .617	<i>p</i> = .067	<b><i>p</i> = .008</b>
UKCDI Total	<i>r</i> = .209	<i>r</i> = -.164	<i>r</i> = .154	<b><i>r</i> = .315</b>	<i>r</i> = -.272
Gestures	<i>p</i> = .183	<i>p</i> = .301	<i>p</i> = .332	<b><i>p</i> = .042</b>	<i>p</i> = .082
ITC Social Subscale	<i>r</i> = .157 <i>p</i> = .321	<i>r</i> = .021 <i>p</i> = .895	<i>r</i> = .081 <i>p</i> = .610	<b><i>r</i> = .389 <i>p</i> = .011</b>	<b><i>r</i> = -.307, <i>p</i> = .048</b>
ITC Speech Subscale	<i>r</i> = .223 <i>p</i> = .156	<i>r</i> = -.066 <i>p</i> = .677	<i>r</i> = -.083 <i>p</i> = .601	<i>r</i> = .083 <i>p</i> = .599	<i>r</i> = -.040 <i>p</i> = .802
ITC Symbolic Subscale	<i>r</i> = .102 <i>p</i> = .522	<i>r</i> = -.098 <i>p</i> = .536	<i>r</i> = .255, <i>p</i> = .102	<b><i>r</i> = .355 <i>p</i> = .021</b>	<i>r</i> = -.139 <i>p</i> = .380
M-CHAT Core-6 Items Failed	<i>r</i> = -.074 <i>p</i> = .641	<i>r</i> = -.125 <i>p</i> = .431	<i>r</i> = .050 <i>p</i> = .754	<i>r</i> = -.257 <i>p</i> = .101	<i>r</i> = .272 <i>p</i> = .081
PLS-4 Auditory Comprehension Score	<b><i>r</i> = .477 <i>p</i> = .001**</b>	<i>r</i> = .184 <i>p</i> = .244	<i>r</i> = .221 <i>p</i> = .159	<i>r</i> = .252 <i>p</i> = .107	<i>r</i> = -.069 <i>p</i> = .665
PLS-4 Expressive Communication Score	<b><i>r</i> = .353 <i>p</i> = .022</b>	<i>r</i> = .246 <i>p</i> = .117	<i>r</i> = -.056 <i>p</i> = .723	<i>r</i> = .039 <i>p</i> = .806	<i>r</i> = -.133 <i>p</i> = .400

1406 **Bold type indicates significant relations**

1407 **\*\*This relationship remains significant after adjusting for multiple comparisons using the Bonferroni method.**

1408 *Table 7: Adjusted mean scores and standard errors for each group on DA tasks*

Group	Adjusted Mean Score	Standard Error	Significant differences
Receptive Language			
Low-risk	4.87	0.49	SHR<LHR=LR
SHR	1.62	0.86	
LHR	4.36	0.72	
Motor Imitation			
Low-risk	5.13	0.46	No differences across groups
SHR	3.65	0.80	
LHR	4.86	0.67	
Response to joint attention			
Low-risk	20.36	1.19	No differences across groups
SHR	17.94	2.07	
LHR	20.17	1.74	
Turn taking			
Low-risk	3.50	0.45	No differences across groups
SHR	2.72	0.78	
LHR	2.94	0.67	
Social requesting			
Low-risk	9.80	0.64	No differences across groups
SHR	10.42	1.10	
LHR	11.43	0.94	

1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418

1419     *Fig 1: DA task correlations with age for the low-risk group (separate file)*

1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452

1453 **Appendix 1: Cueing and Scoring for the DA tasks**

1454

1455 *Receptive Language*

The array of items was indicated and the child asked “Give me the....”	Correct response – score 3
If the child made no response, or selected the wrong item, the wrong item was returned to the array and the instruction repeated.	Correct response after repetition – score 2
If the child made no response, or selected the wrong item, three of the items were removed from the array, leaving a choice of two, and the instruction was repeated.	Correct response from choice of 2 – score 1
If the child still made no response or selected the wrong item, the correct item was indicated and the child told “Here is the....”. The child was then praised.	Incorrect response after prompting – score 0

1456

1457

1458 *Motor Imitation*

Modelling of action, preceded by “Look what I can do!” and followed by the command “X do it...”	Copying of action following model – score 3
If no response was made, the action and the command “X do it... was repeated”	Copying of action with extra verbal prompt – score 2
If the action was not copied, or the child performed another action, the action was repeated, exaggerating the action.	Copying of action after repetition – score 1
If the action was not copied following this, the child was praised and the next item presented.	No attempt to copy or another action produced – score 0

1459

1460

1461

1462

1463

1464

1465

1466 *Joint Attention*

The experimenter pointed at an item on the page and said "Look! A...."	Gaze shift to focus of pointing – score 3
If the child did not response to this, the instruction was repeated using the child's name.	Gaze shift following repetition – score 2
If the child still did not shift their gaze, a light touch was used to gain their attention and direct them to the page, with the instruction "look!"	Gaze shift following physical prompt – score 1
If the child still did not attend to the focus of pointing, the next item was presented.	No gaze shift after cueing – score 0

1467

1468

1469

1470 *Turn Taking*

Experimenter handed ball to child following child's turn	Child spontaneously stops and waits for experimenter to take a turn - score 3
If the child moved their ball towards the run without waiting for the experimenter to take a turn, the verbal prompt "My turn" was used	Child stops and waits for experimenter to take a turn following verbal prompt - score 2
If the child did not respond to the verbal prompt, a physical prompt of blocking the child's entry to the run was used, along with a repetition of "My turn"	Child stops and waits for experimenter to take a turn following a physical prompt - score 1
If the child persisted in taking their turn, the experimenter then allowed this, and began the procedure again for the next trial	Child does not stop and wait for experimenter's turn despite physical prompting - score 0

1471

1472

1473

1474

1475

1476

The experimented turned off the toy and waited for a response.	Child spontaneously bids for the toy to be turned back on – score 3
If the child made no response, the verbal prompt “Would you like more?” was used.	Child bids for toy to be turned back on after verbal prompt – score 2
If the child still made no response, a light touch was used to call their attention to the toy and the verbal prompt was repeated.	Child bids for toy to be turned back on after verbal and physical prompt – score 1
If the child still made no response, the toy was turned back on, with the word “More” and the procedure was repeated.	No bid for help – score 0

1478

1479

1480

1481

1482

1483

1484