



City Research Online

City, University of London Institutional Repository

Citation: Giuntella, O., McManus, S., Mujcic, R., Oswald, A. J., Powdthavee, N. & Tohamy, A. (2023). The Midlife Crisis. *Economica*, 90(357), pp. 65-110. doi: 10.1111/ecca.12452

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/29373/>

Link to published version: <https://doi.org/10.1111/ecca.12452>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

The Midlife Crisis

By OSEA GIUNTELLA*, SALLY MCMANUS†, REDZO MUJIC‡,
ANDREW J. OSWALD§, NATTAVUDH POWDTHAVEE¶ and AHMED TOHAMY**

*University of Pittsburgh †National Centre for Social Research ‡University of Warwick
§University of Warwick and IZA ¶Nanyang Technological University and IZA
**Oxford University

This paper documents a longitudinal crisis of midlife among the inhabitants of rich nations. Yet middle-aged citizens in our datasets are close to their peak earnings, have typically experienced little or no illness, reside in some of the safest countries in the world, and live in the most prosperous era in human history. This is paradoxical and troubling. The finding is consistent, however, with the prediction—one little-known to economists—of Elliott Jaques (1965). Our analysis does not rest on elementary cross-sectional analysis. Instead, the paper uses panel and through-time data on, in total, approximately 500,000 individuals. It checks that the key results are not due to cohort effects. Nor do we rely on simple life satisfaction measures. The paper shows that there are approximately quadratic hill-shaped patterns in data on midlife suicide, sleeping problems, alcohol dependence, concentration difficulties, memory problems, intense job strain, disabling headaches, suicidal feelings, and extreme depression. We believe that the seriousness of this societal problem has not been grasped by the affluent world's policy-makers.

I. INTRODUCTION

The myth of the midlife crisis ... (Tergesen 2014)

Epidemiological study of psychological distress in adulthood does not suggest that midlife is a time of out-of-the-ordinary distress. (Wethington 2000)

Worried about a midlife crisis? Don't. There's no such thing. (Whitbourne 2015)

Residents of today's affluent nations are citizens of the richest societies in human history. By the midlife point, these men and women have normally encountered no significant illness or disability. They are also close to their peak lifetime earnings (which occurs in the late forties for those with low levels of education, and in the early fifties for those with high levels of education; see Bhuller *et al.* 2017). It would be expected, therefore, that middle-aged adults in the industrialized nations would have extraordinarily cushioned and enjoyable lives.

We demonstrate in this paper that measures of extreme distress paint a different picture. Something elemental appears to be going wrong in the middle of many of our citizens' lives. The paper's main finding is consistent—in ways that remain to be fully understood—with the ideas of Jaques (1965), and with the notion of a paradox of economic progress, and perhaps with the problem that humans are not influenced merely by absolute prosperity, as discussed in different ways by researchers such as Richard Easterlin, Richard Layard, Fredrik Carlsson and colleagues, and Carol Graham and colleagues (Layard 1980; Easterlin 2003; Carlsson *et al.* 2007; Graham 2017; Graham *et al.* 2017).

The paper offers new evidence that midlife is a time when people disproportionately take their own lives, have trouble sleeping, are clinically depressed, spend time thinking about suicide, feel life is not worth living, find it hard to concentrate, forget things, feel overwhelmed in their workplace, suffer from disabling headaches, and become dependent

on alcohol. As shown below, markers of distress routinely follow the approximate shape of a quadratic equation that is concave from below (although, unsurprisingly, a second-order equation should not be thought of as holding in an exact way close to the end of the human lifespan). It should be emphasized that in the later analysis, a quadratic is not forced onto the data. Instead, a set of age dummy variables is used in order to allow more general non-parametric estimation.

The paper's central result does not rest on cross-sectional correlations. Nor is it dependent on data from a single nation; nor is it the result of young children in the household; nor is it driven merely by high effort among workers; nor is the observed pattern a temporary one or the result of cohort effects. The latter possibility is especially important to consider (see, for example, Gunnell *et al.* 2003). In the analysis below, we attempt to ensure that the later evidence adjusts directly for cohort effects, and/or examines the key conclusion for widely different time periods, and/or uses within-person fixed effects analysis where the same individuals are followed longitudinally over many years. Nor does the paper's analysis rely merely on data on subjective wellbeing scores, although, as will be seen, one special strand of the previous cross-sectional happiness literature is potentially relevant. Nor do there appear to be especially large differences between male and female patterns, so it is not straightforward to believe that female menopause, for example, plays any major explanatory role.

Later analysis focuses primarily on *ceteris paribus* patterns. The age trajectory of human distress will be studied here, where feasible, by using conventional regression equations. Adjustment will thus generally be made for a set of other influences (measured contemporaneously in the equations), such as marriage, employment and having children. The study is therefore an analytical inquiry into the other-factors-constant consequences of ageing. It is not designed as a description, or empirical summary, of the simple mean values for different age groups in society.¹ The aim is instead to try to evaluate the tendency to—or not to—a crisis of middle age. Moreover, the paper's purpose is not to imply that all midlife individuals exhibit high levels of distress. The later analysis will attempt, as normal in applied research, to understand broad statistical influences among representative individuals.

II. BACKGROUND

The analysis builds on three literatures. They appear to have started independently of one another. In each case, it was initially rare to mention or cross-reference writings from the other two literatures, although today these literatures are beginning, if tentatively, to connect.

All three literatures are currently disputatious in tone. They are:

- an economics and behavioural science literature on the idea that subjective wellbeing may be 'U-shaped' through life;
- a social science and medical literature on the idea that white low-education American men in midlife are currently experiencing new levels of psychological despair;
- a psychotherapy and psychology literature on the idea that human beings have a 'midlife crisis'.

More generally, we examine data on extreme-distress measures rather than simpler life-satisfaction kinds of scores or experienced-utility measures (such as Winkelmann and Winkelmann 1998; Dolan and Kahneman 2007; Dolan *et al.* 2008).

Within the first literature, there has been much debate. This style of work, which is predominantly cross-sectional, appears particularly in economics and certain kinds of social science journals (e.g. Blanchflower and Oswald 2008; Graham and Pozuelo 2017). It explores the claim that the mean level of life satisfaction and contentment may follow a convex

quadratic equation over the bulk of the lifespan. Consistent with some aspects of this putative U-shape in subjective wellbeing, there is also growing evidence that old age seems to help human beings to engage in fewer regretful emotions (Charles *et al.* 2003). One cross-sectional study (Blanchflower and Oswald 2016) finds that middle-aged Europeans consume the largest numbers of antidepressants. One of the few pieces of genuine longitudinal evidence examines the mean levels of life satisfaction rather than extreme distress (Cheng *et al.* 2017; see also Kunzmann *et al.* (2013). Another identifies one aspect of the U-shape (Van Landeghem 2012). Non-technical books have also appeared on this topic (Setiya 2017; Rauch 2018).

The notion of a midlife nadir is not generally accepted. An early review paper (Diener *et al.* 1999) concludes that wellbeing is approximately flat through the lifespan. Such scholars argue that midlife is a time of high satisfaction (Mroczek and Spiro 2005); that, at best, only low-income individuals experience a U-shape (Lang *et al.* 2011); others conclude that the cross-sectional U-shape is simply illusory (Galambos *et al.* 2020; Frijters and Beaton 2012). Some researchers are also reluctant to place any weight on subjective wellbeing scores because of their subjectivity (although Oswald and Wu (2010) offer a robustness check on such data).

A second, still-expanding literature, stimulated particularly by Case and Deaton (2015), has sprung up. Researchers have documented evidence of rising levels of mental distress among particular groups of US citizens, especially white midlife Americans with few educational qualifications (Graham 2017; Gaydosch *et al.* 2019; Blanchflower and Oswald 2020; Daly 2022). There has been debate about this important hypothesis (Goldman *et al.* 2018). To our knowledge, the literature generally views the American trend as recent, distinct and unusual.

In a third and earlier literature, there is discussion about, and a general rejection of, the psychological concept of a ‘midlife crisis’ in humans. That term was adopted by Jaques (1965) in an article published in a psychotherapy journal. The argument made by Jaques is that in midlife, a human being is forced to come to terms, painfully, with the certainty of his or her own eventual mortality.

Most psychologists since then have treated the notion with extreme scepticism (Wethington 2000; Freund and Ritter 2009; Galambos *et al.* 2020)—and even, occasionally, derision. Lachman (2015), however, calls persuasively for more research on these issues.

III. DATA AND METHODS

In this section, we describe the different datasets and methods used to analyse eight intuitively natural ‘distress’ indicators. We test both objective and subjective markers of extreme distress over the human lifespan. Depending on the nature of the data analysed, our empirical approach adjusts for cohort effects and/or examines widely different time periods and/or uses within-person fixed effects analysis where the same individuals are followed longitudinally through time. The paper offers various forms of evidence. Regression equations and estimated coefficients are discussed in Section III.

The data types are discussed briefly below.

Cross-national data on suicide

Suicide data are collected from the mortality deaths database provided by the World Health Organization (WHO).² The data were classified into population 5-year bands to generate suicide rates per 100,000 individuals per nation. We focus our analysis on affluent countries and especially on English-speaking countries. The suicide rates are averaged over 5-year periods to be able to adjust for cohort effects and period effects. The unit of

observation in the formal analysis is by country by 5-year-period-band by 5-year-age-band by gender.

Cohorts are constructed as the difference between periods and age bands. The standard difficulty faced in this type of analysis is the famous ‘age–period–cohort’ (APC) problem. This occurs because as $period = age + cohort$, the three influences are, together, perfectly collinear. Therefore in order to estimate separate coefficients on each of the age bands, further assumptions need to be made. In this paper, this is dealt with using the so-called intrinsic estimator approach due to Fu, Yang and others (Fu 2000; Yang *et al.* 2004, 2008). It is known that the numbers of age groups and time periods (the so-called design matrix) in the APC accounting model may affect the estimates obtained from certain kinds of estimators. One way to think about the rationale for the Yang intrinsic estimator (IE), therefore, is that it removes the influence of the design matrix on the coefficient estimates. The IE approach produces an estimator that has attractive statistical properties. Although its application to suicide rates appears to be rare, our later analysis is similar in spirit to a perhaps little-known paper on Canadian cohort data by Thibodeau (2015). Our results, in international data, are consistent with her findings for certain provinces of Canada. The IE approach also can be seen as a particular kind of principal components regression estimator. For this analysis, we have also checked—results not presented but available on request—the method of equality constraints as a robustness test.

Here, and throughout the paper, we concentrate on annual data because we lack data of higher frequency. We omit independent variables on childhood experience because the datasets do not offer such information.

UK data on clinical depression and generalized anxiety disorder

UK data on serious psychological illness are available to us in three separate years of the Adult Psychiatric Morbidity Survey (APMS); see, example, McManus *et al.* (2020). Here, we use the long so-called Clinical Interview Schedule—Revised (CIS-R) to assess a range of different depressive and anxiety disorders. The schedule was administered face to face by an interviewer using computer-assisted interviewing. The CIS-R is a structured interview schedule on the presence of symptoms in the week prior to interview. It comprises over 130 questions, spanning 14 types of symptoms (e.g. fatigue, sleep problems, worry). These items were used to assess for different types of clinical disorders (e.g. depression, generalized anxiety disorder). The 14 sections of the CIS-R cover somatic symptoms, fatigue, concentration and forgetfulness problems, sleep problems, irritability, worry about physical health, depressive symptoms, depressive ideas, worry, anxiety, phobias, panic, and compulsions and obsessions. Each section starts with two filter questions to establish the presence of the specific symptom in the past month. A positive response leads to further questions about the symptom in the past week, including frequency, duration, severity and time since onset.

Our approach is standard in this branch of psychological medicine. The participants’ answers were used to generate what are known as 10th International Classification of Disease (ICD-10) diagnoses of anxiety and depressive disorders by applying an algorithm to operationalize the ICD-10 diagnostic criteria (WHO 1992). These ICD-10 diagnoses were then amalgamated to produce categories of disorder. The two most prevalent disorders classified were generalized anxiety disorder (GAD) and depression (combining mild, moderate and severe). The ICD-10 criteria for so-called code F41.1 (symptoms lasting for at least six months, general anxiety that is not restricted to any one environment, symptoms that include persistent nervousness, trembling, muscle tension, sweating, light-headedness, palpitations

and dizziness, and an overall anxiety symptom score of at least 2) were used to identify GAD.

Clinical depression is measured in an equivalent way. The ICD-10 depressive episode criteria include: symptoms lasting at least two weeks, some of depressed mood, loss of interest and fatigue; some of reduced concentration, reduced self-esteem, ideas of guilt, pessimism about the future, suicidal ideas or acts, disturbed sleep, diminished appetite; perceived social impairment; and some of lack of normal pleasure /interest, loss of normal emotional reactivity, morning waking more than 2 hours early, loss of libido, diurnal variation in mood, diminished appetite, loss of more than 5% of body weight, psychomotor agitation and psychomotor retardation. These were scored, within the dataset rather than by us, to generate so-called ICD-10 codes F32.00, F32.01, F32.10, F32.11 and F32.2, which were combined to produce the final standardized technical category of 'depression'.

Cross-national data on sleeping problems

Time-use data for Austria, Canada, Finland, France, the Netherlands, Spain, the UK and the USA were collected using the multinational time-use study extract builder (MTUS-X; Fisher *et al.* 2015).³ The studied data cover the 1965–2012 period. However, as these data were collected from multiple sources, the samples are not balanced, and in each country, individuals were interviewed in different years. For this reason, we have checked the pattern separately by country of residence of the respondents. The US sample is the largest (160,445 citizens) and covers the entire period (1965–2012), although it should be noted that the early 1965 sample is extremely small. The Dutch sample covers the period 1975–2005, including 84,028 observations. We have data from the UK time-use survey for the period 1974–2005, covering 66,959 individuals. Data for France are available for the period 1985–1999, covering 31,488 individuals. For the other countries, only one wave of data, in each case, was available: the 1992 wave for Austria (25,233 individuals), the 2009 wave for Spain (19,295 individuals), the 2010 wave for Canada (15,390 individuals), and the 1979 wave for Finland (12,038). Data for Germany were drawn from the 2012–13 German Time-use Survey.

Sleep duration is defined as the total time assigned to sleep and naps. We also included 'imputed sleep', defined as short gaps in the early hours at the beginning or end of the diary, where the diarist is at home or in the same location where they report sleeping on the diary day, and being asleep before the gap at the end of the diary, or they were asleep following the gap at the beginning of the diary.

Data on suicidal feelings, concentration problems and forgetfulness, alcohol dependence

For this part of the study, we drew again on the APMS series.⁴ The latest survey is the fourth in the series and was conducted by NatCen Social Research, in collaboration with the University of Leicester, for National Health Service (NHS) Digital. A previous survey was conducted in 2000 (for 16–74-year-olds) by the Office for National Statistics, which covered England, Scotland and Wales. Another, the 2007 APMS, included people aged over 16, and covered England only. The survey uses a robust stratified, multi-stage probability sample of households and assesses psychiatric morbidity using diagnostic criteria for several disorders.

The 'Not worth living' measure is based on responses to the following questions:

Have you ever felt that life was not worth living?

1 Yes

2 No

- If yes, was this ...
- 1 in the last week?
 - 2 in the last year?
 - 3 or at some other time?

The 'Concentration and forgetfulness' symptom score is constructed by scoring 1 for each of the criteria below that is met:

- Noticed problems with concentration/memory for four days or more in the past week.
- Could not always concentrate on a TV programme, read a newspaper article or talk to someone without mind wandering in past week.
- Problems with concentration actually stopped you from getting on with things you used to do or would like to do.
- Forgot something important in past seven days.

We also use APMS data to create a measure of 'Alcohol dependence'. Participants were given the Alcohol Use Disorders Identification Test (AUDIT; Saunders *et al.* 1993), which takes the year before the interview as a reference period. In broad outline, our statistical analysis draws upon a so-called AUDIT score, and takes the cut-point of an AUDIT score of greater than or equal to 16 to indicate dependence.

The measure of 'Suicidal thoughts', which is clearly related to, but perhaps even more sharply focused than, the earlier 'Not worth living' score, is derived from the question:

There may be times in everyone's life when they become very miserable and depressed and may feel like taking drastic action because of these feelings. Have you ever thought of taking your life, even if you would not really do it?

Those responding 'Yes' were asked when this last occurred. A derived variable then identified those reporting suicidal thoughts in the past year. See also Spiers *et al.* (2012).

Longitudinal data on job stress

Longitudinal data on job stress come from the Household, Income and Labour Dynamics in Australia (HILDA) Survey.⁵ This is a nationally representative longitudinal survey that was first conducted in 2001 (Wooden and Watson 2007). It collects annual information from members of Australian households who are at least 15 years old. It began providing information on a total of 13,969 individuals from 7682 different households interviewed since the first survey wave. In interviews, information is collected on a variety of topics, including labour market dynamics, income and education levels, family composition and lifestyle choices, as well as the physical and psychological wellbeing of individuals.

Longitudinal data on migraine

The analysed data on migraine problems come from the British Household Panel Survey (BHPS).⁶ The dataset is a nationally representative survey of British households, with over 10,000 adult respondents, and was first conducted between September and Christmas in 1991.

IV. RESULTS

This section summarizes a set of mutually complementary findings. It draws upon the different markers of extreme psychological problems and mental crisis described above.

Presence of clinical depression, generalized anxiety disorder and life not worth living

First, as shown in Figure 1, there is a repeated hump-shaped age pattern through time of clinical depression and GAD in England. The data are once more from the APMS. Both indicators of distress use the internationally agreed ICD-10 diagnostic definitions. The subfigures in Figure 1 are for three separate years (2000, 2007, 2014); at the time of writing, the 2021 data have not been released. The underlying regression equations represented in the Appendix tables control for gender, educational level, marital status, number of children, type of housing in which the respondent lives, and employment status. On balance, throughout the paper, we think it best to control for standard kinds of socioeconomic covariates, even though it is technically possible to remove all potentially endogenous variables and still find




FIGURE 1. The age profile of distress, using three different measures, in England (2000, 2007 and 2014).

Notes: These diagrams plot the age dummy coefficients from linear probability regression equations controlling for gender, education, marriage, dependent children, housing type and unemployment, using data from direct survey questions. The data come from the APMS series (about 7500 individuals) of the general population in England (for the years 2000, 2007, 2014—2021 data not yet available) using a detailed clinical schedule (the Clinical Interview Schedule—Revised), which applies ICD-10 diagnostic criteria. Age bands are depicted on the x -axis. For clarity, standard error bands are omitted, but underlying regression equations are given in Appendix Tables A1–A9. Further information on the construction of depression and GAD scores is available online at <https://www.ncbi.nlm.nih.gov/books/NBK262332> and http://doc.ukdataservice.ac.uk/doc/6379/mrdoc/pdf/6379_apms_2007_interviewer_instructions.pdf (both accessed 30 October 2022).

evidence of the paper's main pattern—as in, for example, the paper's later results on suicide. Appendix Table A1 gives raw numbers. Appendix Tables A2–A9 provide the background regression equations behind Figure 1.

Because the APMS dataset is not a panel, it is infeasible to estimate within-person estimates of the ageing effect. That means that it is less easy to rule out cohort effects that could lead to fallacious inferences about the role of human ageing. Therefore Figure 1 deliberately shows the age curve for each of the years rather than pooled together. The approximate stability across the subfigures suggests that the hill-shaped pattern is not merely the result of simple cohort effects. The hump shape is apparently a consistent feature.

How large is the age effect? Figure 1 reveals that being in midlife approximately doubles the probability of depression, *ceteris paribus*, compared to being aged 65–74 or under the age of 25. Further details are in Model (2) of Appendix Table A4.

Figure 1(c) examines a 'Not worth living' measure, using the patterns in people's answers to the question: 'Have you ever felt that life was not worth living? Yes/No.' We code these as 1 for those citizens who answered 'Yes, in the last year'. Again, in Figure 1(c), an equivalent middle-age peak is evident.

The Appendix gives miscellaneous APMS results for related markers of distress, including concentration and memory problems.

Suicide in cross-national data

Human suicide is the starkest measure of mental crisis. Once again, the data seem to exhibit a hump shape that exists across the age groups. Figure 2 reports cross-national findings on suicide rates. It corrects for cohort effects in a now conventional way described in work by Yang *et al.* (2008). The supplementary information in the Appendix provides details on the so-called intrinsic estimator. For the statistical analysis, we created a dataset that contains a non-balanced panel of 9 affluent non-English countries and 8 English-speaking countries. The data run through the years 1950 to 2015. The dataset has annual data and gives rise to 2389 and 2320 observations, respectively, in each of the two subsamples. The suicide rate is in logarithms in order to focus on percentage deviations.




FIGURE 2. The age profile of suicide in the English-speaking nations since the 1950s.

Notes: Suicide rates by age group after adjusting for cohort + time effects for seven decades of data, 1950–2015 (2020 standardized data not yet available). This figure uses WHO data from 1950 to 2015. Age is on the horizontal axis. The studied nations are Australia, Canada, Ireland, New Zealand, England and Wales, Northern Ireland, Scotland, USA. In some nations, the suicide rate spikes up again at the end of life, but that is not the focus of this study. The plots in the above use the intrinsic estimator due to Yang *et al.* (2004, 2008). This adjusts for country fixed effects, cohort effects and period effects. Moreover, similar results are obtained for cohort adjustment using the previously traditional method of equality constraints. Those are available on request. For clarity, standard error bands are omitted, but underlying regression equations are given in Appendix Tables A1–A9.

A midlife maximum is evident in Figure 2 after correction for cohort effects, time dummies and country dummies. Age coefficients are plotted. When averaging males and females in the English-speaking countries, for example, the peak risk is estimated to be approximately in people's early fifties. These hump shapes are again consistent with a psychological low in the middle of people's lives in affluent countries. The estimated female peak here may occur slightly earlier than the male peak. Appendix Table A10 gives the formal regression estimates that lie behind these curves. Appendix Table A11 is for a set of rich countries where English is not the first language.

It could be argued that suicide data are special, in the sense that they capture an exceptional tail of the distress distribution in a society. We therefore turn to a range of further indicators to check for signs of extreme distress in midlife.

Sleeping problems and hospital admissions in cross-national data through time

Sleep is important to health (Spiegel *et al.* 1999), and it has been known since at least the time of William Shakespeare that sleep and the level of mental stress are closely connected (Furman *et al.* 1997). Worry keeps people awake at night; sleeping problems are a marker of anxiety; they also depress the strength of the immune system, raise the risk of personal accident, and have negative effects on people's productivity. There is also evidence linking greater sleep hours to reduced risk of depression (Furman *et al.* 1997; Spiegel *et al.* 1999; Taylor *et al.* 2005; Roenneberg 2013; Giuntella *et al.* 2017; Hafner *et al.* 2017). Official US medical guidelines, such as those issued by the Centers for Disease Control (CDC), state that healthy human adults need 8 hours of sleep per night.

What happens in the middle of people's lives? The answer is that adults in rich countries tend to have sleeping difficulties. These are sometimes of such worrying severity that they are akin to physical problems.

One simple piece of evidence is available from data from the National Health Service of the UK. Figure 3, which draws upon official NHS records, reveals a marked hill-shaped pattern. We are not sure exactly how ill a person has to be in order to have to go to a hospital because of a sleeping problem, but it can be presumed that he or she would have to have really extreme symptoms of distress. To the best of our knowledge, this kind of diagram (Figure 3) is not known to many social scientists. It is perhaps worth emphasizing that, unlike the wellbeing and happiness literature, such evidence is not based on reported feelings.

Figure 4, for the USA, is related, but measures different variables. These plot the mean hours of sleep for individuals for different periods spanning half a century. Further countries' results are in the Appendix. In each case, in a robust way, the low point in hours of sleep is reached in midlife, including for those individuals without young children.

Figure 4(c) is of a complementary kind to Figure 3. Here, again, the information is not on subjective sleeping levels; instead, it is objective and comes from hospitals in the UK. Figures 4(c)–4(d) are for the UK and Greater London area, on sleeping problems so acute that they led to admission to hospital.

Finally, in genuine panel data, from the BHPS (explained more fully later in the paper), following the same individuals as they grow older through time, Figure 4(f) shows that a sleep hours U-shape pattern in age is present even in 'within-person' longitudinal data from Great Britain.

See Appendix Tables A12 and A13 for further details.

The midlife sleep problem phenomenon in humans appears to be little known to most researchers. A search of the literature using the Web of Science, however, shows that the




FIGURE 3. Evidence of a midlife sleep ‘crisis’, UK 2012–13.

Notes: Hospital admissions for sleep disorders, official data from the NHS of the UK. Finished admissions episodes for sleep disorders by age, September 2012 to August 2013. The data come from the Health and Social Care Information Centre (see <https://digital.nhs.uk/data-and-information/find-data-and-publications/supplementary-information/2018-supplementary-information-files/patients-treated-for-sleep-disorders-by-age-and-gender-2012-13-to-2017-18>, accessed 31 October 2022). There were approximately 35,000 hospital admissions for sleep disorders in the year 2013. A finished admission episode (FAE) is the first period of inpatient care under one consultant within one healthcare provider. FAEs are counted against the year or month in which the admission episode finishes. Admissions do not represent the number of inpatients, as a person may have more than one admission within the period. Sleep disorders include insomnias, hypersomnias, disorders of sleep–wake schedule, sleep apnoea, narcolepsy and cataplexy, other sleep disorders, non-organic insomnia, non-organic hypersomnia, non-organic disorder of the sleep–wake schedule, sleep terrors, nightmares, other non-organic sleep disorders. This is the population, hence no standard error bands are given.

results on sleep hours are consistent with one published cross-sectional US study (Krueger and Friedman 2009), although age was not the authors’ focus.

Disabling headaches in longitudinal data

Headaches of a disabling kind—migraine headaches—are known to affect a significant proportion of the citizens of affluent countries. Their cause is not fully understood, but it is believed that migraine is a correlate of anxiety and depression (Ratcliffe *et al.* 2008; Spiers *et al.* 2012; Lampl *et al.* 2016). A longitudinal study of 17,600 Canadians found that migraine headaches were, prospectively, one of the strongest predictors of who would be (newly) diagnosed with major depression within the ensuing 24 months (Patten 2001).

We therefore use fixed effects methods to examine this indicator—extreme headache attacks—as an additional potential marker to assess the possible incidence of mental strain in midlife. For the exercise, data are drawn from waves 1–18 of the BHPS. The migraine variable itself is derived from the BHPS health questionnaire: ‘Do you have any of the listed health problems: ... migraine?’

We focus on individuals aged 16–75 years old. This produces 217,645 observations. Of those observations, 18,058 people–interviews (or 8.3%) listed migraine as a health problem. Fixed effects logit regressions are estimated here and control for marital status, employment status, highest qualification level, homeownership status, number of days spent in the hospital




FIGURE 4. Age profiles of sleep issues in the USA and the UK since the 1960s.

Notes: Parts (a) and (b) document U-shaped sleep hours versus age in a sample of half a million individuals over two periods of time in the USA. Age is plotted on the x -axis. The U-shape pattern continues to hold in regression equations that include standard demographic controls, including for children in the household.

Parts (c) and (d) document hospital admissions for sleep disorders in the UK, from the NHS. Finished admissions episodes for sleep disorders by age, September 2012 to August 2018. The data come from the Health and Social Care Information Centre (see Notes for Figure 3). There are typically approximately 35,000 hospital admissions for sleep disorders per year in the UK. (See Figure 3 for further information.) This is the population, hence no standard error bands are given.

Part (e) gives hospital admissions for sleep disorders in the Greater London area. These are official data from the NHS, but for Greater London, which is the largest urban area in the UK. Finished admission episodes for sleep disorders by age, covering years 2017–2018. See <https://digital.nhs.uk/data-and-information/find-data-and-publications/supplementary-information/2018-supplementary-information-files/hospital-admissions-for-sleep-disorders-covering-the-london-commissioning-region> (accessed 30 October 2022).

The plot in part (f) uses longitudinal data drawn from the BHPS for the period 1994–2004. The estimates are from fixed effects equations, hence the pattern is derived solely from within-person, not cross-sectional, variation. For clarity, standard error bands are omitted, but underlying regression equations are given in Appendix Tables A1–A9.

last year, self-rated health, number of children under the age of 16 living in the same household, year fixed effects and regional fixed effects.

Figure 5(a) plots the outcome. A hill shape in disabling headaches, with the maximum reached in midlife, emerges once again. These are within-person patterns obtained by following the same randomly selected group of British people as they age through time. See Appendix Table A14 for more information.

Extreme job stress in longitudinal data

Work is a major part of life, so it is of interest to consider also what happens during that section of people's lives. It is known, for example, that job strain is a predictor of




FIGURE 5. The age profile of disabling headaches, job stress, alcohol dependence and suicidal thoughts in the UK and Australia since 2002.

Notes: Parts (a) and (b) illustrate within-person longitudinal changes in migraine and within-person longitudinal severe job stress by age group, respectively. These are based on coefficients from fixed effects regression equations with six banded dummy variables for age groups. The plots use within-person longitudinal data from the BHPS (waves 1–18) and the HILDA Survey (year waves 2002–18). The regression equations also control for other socioeconomic variables, including income and number of young children. Total samples contain 213,011 and 127,199 person–year observations, respectively.

The data in parts (c) and (d), on two other distress measures, come from the APMS series (7500 individuals) for 2014. Age bands are depicted on the x -axis. These diagrams plot the age dummy coefficients from linear probability regression equations controlling for gender, education, marriage, dependent children, housing type and unemployment. For clarity, standard error bands are omitted, but underlying regression equations are given in Appendix Tables A1–A9.

elevated blood pressure (Ramirez *et al.* 1996). Research evidence suggests that stress at work is a longitudinal predictor of depression and poor mental health (Stansfeld *et al.* 1999; Choi 2018).

The next form of evidence applies a little-used measure of employees' strain in the workplace. Our analysis draws upon 17 waves (years 2002–18) of the HILDA Survey.⁷ We focus on extreme stress evaluations. These are made by respondents relating to their current job. The survey respondents were asked to assign an integer value between 1, 'Strongly disagree', and 7, 'Strongly agree', to each of the following statements, among others:

- I fear the amount of stress in my job will make me physically ill.
- My job is complex and difficult.
- My job is more stressful than I had ever imagined.

We average responses to these three statements. This creates an overall job stress measure, which is then used as the dependent variable in fixed effects regression equations. The grouping was also checked, and supported, by a principal components factor analysis (not reported).

A hill shape is again traced out. Figure 5(b), based on Appendix Table A15, illustrates our longitudinal evidence following the same 20,648 working individuals (aged 15–75) through time (127,199 person–year observations). The 'within-person' regression equations adjust for income, total hours of work, job or industry type, education, marital status, number of children, alcohol consumption, and a range of variables for exercise and healthy diet.

The maximum level of work stress is reached at approximately age 45. On these estimates, the size of the implied age effect is substantial. It is 0.1–0.2 in size, which can be compared to the mean value of stress in the sample of 3.22, but that would make the estimated effect look smaller than is truly correct. A more appropriate way to understand the size of the effect is to compare with other coefficients in a stress regression equation. For instance, in the underlying regression equation in the Appendix, approximately 0.15 would then be the implied rise in stress associated with a full extra 8 hours of work per week.

Other results

Information on additional markers of distress is available from the earlier discussed APMS for England. These, it should be emphasized, are not longitudinal results, but we present them here for completeness.

The data in Figures 5(c) and 5(d), on two distress measures, namely dependence on alcohol and having had suicidal thoughts, come from the APMS series (7500 individuals) for 2014. Age bands are once more depicted on the x -axis. These diagrams plot the age dummy coefficients from linear probability regression equations controlling for gender, education, marriage, dependent children, housing type and unemployment. The by-now-usual hill shape seems still to be in evidence.

Tables A16–A21 contain further results and details.

V. CONCLUSIONS

The human midlife crisis seems to be an important and under-recognized phenomenon. We document longitudinal evidence of extreme distress among middle-aged adults in affluent countries. These individuals are close to their peak lifetime earnings and in general have experienced no serious illness. Our findings therefore appear to point to a disturbing paradox within modern society.

Using eight different measures, an approximate hump shape in severe distress over the lifecycle emerges in data from industrialized nations such as the UK, Australia and the USA. This paper's methods go beyond cross-sectional analyses based on simple measures of subjective wellbeing (for example, Graham and Pozuelo 2017). As far as we know, our recurring longitudinal patterns—to be thought of as a collection of complementary types of evidence—are not widely known by policymakers.

The late Elliott Jaques (1965) is believed to have coined the term 'midlife crisis' in the year 1965. He offered anecdotal evidence, and psychoanalytic arguments, for it. Using modern datasets and conventional statistical methods, this paper explores, and provides empirical support consistent with, the hypothesis advanced by Jaques. The paper's analysis finds hill-shaped patterns in data on:

- suicide
- sleeping problems
- extreme depression
- intense job strain
- disabling headaches
- suicidal feelings
- concentration and memory problems
- alcohol dependence.

In some cases, a particular mental distress marker is available in many nations; in other cases, it is available for only a few nations.

The explanation for the midlife shape currently remains open. Could the paper's empirical result be the product of the stresses of having dependent children, or a country-specific or new phenomenon, or something to do with selection effects, or an illusion caused by cohort effects? These are natural and important possibilities. Nevertheless, the balance of our evidence appears to suggest not. It also does not seem that envy of others causes the midlife shape. (Mujcic and Oswald (2018) test for that possibility, although not with extreme distress measures as the dependent variable.) The notion of unmet aspirations as part of the explanation does, however, have intuitive appeal, in our judgement (see particularly Schwandt 2016). Perhaps so also, more speculatively, does some role for rising 'wisdom' seem possible (Jeste and Oswald 2014) in the observed reduction in distress levels later in life.

There is some published evidence for a midlife psychological low in data on chimpanzees and orangutans (Weiss *et al.* 2012). So sheer ageing biology in primates may play some kind of role. That would take the ultimate explanation out of the social sciences and into the natural sciences. Much is still to be understood.

Scientific caution remains appropriate. The evidence described here is based on a particular, if large, set of indicators. It is possible to think of objections to those indicators. A caveat on that, however, should arguably also be entered. It would be incumbent upon a critic of our chosen extreme distress measures to suggest what would count instead as a set of better markers of human crisis. Most especially, it would not seem scientifically acceptable to suggest something like 'indicator X is less than perfect so I reject the repeated pattern of these multiple indicators'.

Finally, we believe it is not currently clear whether (i) there is a timeless and innate form of human middle-aged crisis, or (ii) the midlife pattern documented here is some kind of perplexing, and perhaps temporary, byproduct of today's affluent world.

Whichever of these turns out to be true, the hill-shaped pattern of extreme distress over the human life course in rich countries appears to constitute a foundational puzzle for economists, behavioural scientists and perhaps other kinds of scientific researchers.

APPENDIX

Supplementary tables

TABLE A1
APMS DATA FOR ENGLAND: DESCRIPTIVE STATISTICS—PREVALENCE OF DEPRESSION, GAD AND FELT LIFE NOT WORTH LIVING, 2000, 2007 AND 2014

	Age group						Total
	16–24	25–34	35–44	45–54	55–64	65–74	
<i>Year 2000</i>							
Depression	2.0%	2.3%	3.3%	3.4%	2.9%	0.8%	2.6%
SE	0.5%	0.4%	0.4%	0.5%	0.5%	0.2%	0.2%
Lower CI	1.2%	1.6%	2.5%	2.6%	2.1%	0.5%	2.2%
Upper CI	3.1%	3.1%	4.3%	4.5%	4.0%	1.4%	3.0%
Generalized anxiety disorder	1.4%	4.2%	5.7%	6.8%	4.6%	2.5%	4.4%
SE	0.4%	0.5%	0.6%	0.7%	0.6%	0.4%	0.2%
Lower CI	0.8%	3.4%	4.6%	5.6%	3.6%	1.7%	4.0%
Upper CI	2.4%	5.3%	6.9%	8.3%	5.8%	3.5%	4.9%
Felt life not worth living	18.3%	21.1%	21.6%	22.0%	20.5%	14.2%	20.1%
SE	1.5%	1.1%	1.0%	1.1%	1.1%	1.0%	0.5%
Lower CI	15.6%	19.1%	19.6%	19.9%	18.4%	12.3%	19.2%
Upper CI	21.4%	23.3%	23.6%	24.3%	22.7%	16.3%	21.0%
Base	794	1683	1848	1545	1442	1268	8580
<i>Year 2007</i>							
Depression	3.2%	2.5%	3.3%	4.8%	2.5%	1.6%	3.1%
SE	0.8%	0.4%	0.5%	0.6%	0.4%	0.4%	0.2%
Lower CI	2.0%	1.8%	2.5%	3.7%	1.8%	1.0%	2.7%
Upper CI	5.1%	3.6%	4.4%	6.2%	3.5%	2.5%	3.6%
Generalized anxiety disorder	3.6%	4.2%	5.3%	6.1%	4.1%	3.3%	4.6%
SE	0.7%	0.6%	0.6%	0.8%	0.6%	0.6%	0.3%
Lower CI	2.4%	3.1%	4.3%	4.7%	3.2%	2.3%	4.0%
Upper CI	5.3%	5.6%	6.6%	7.8%	5.4%	4.6%	5.1%
Felt life not worth living	20.0%	17.2%	20.0%	21.3%	17.1%	16.1%	18.8%
SE	1.8%	1.2%	1.2%	1.2%	1.1%	1.1%	0.6%
Lower CI	16.6%	14.9%	17.7%	19.1%	15.0%	13.9%	17.7%
Upper CI	23.9%	19.7%	22.4%	23.8%	19.3%	18.4%	19.9%
Base	568	1035	1413	1130	1279	1028	6453
<i>Year 2014</i>							
Depression	2.3%	3.5%	4.1%	4.5%	4.3%	2.1%	3.5%
SE	0.6%	0.6%	0.6%	0.6%	0.7%	0.5%	0.3%
Lower CI	1.4%	2.4%	3.1%	3.4%	3.2%	1.4%	3.1%
Upper CI	3.7%	5.0%	5.5%	5.9%	5.8%	3.2%	4.1%
Generalized anxiety disorder	6.3%	6.1%	6.9%	7.3%	6.4%	4.0%	6.3%
SE	1.0%	0.9%	0.8%	0.8%	0.8%	0.5%	0.3%
Lower CI	4.6%	4.7%	5.4%	5.8%	5.1%	3.0%	5.7%
Upper CI	8.7%	8.1%	8.8%	9.0%	8.1%	5.1%	7.0%
Felt life not worth living	21.4%	19.8%	20.7%	23.5%	25.4%	13.5%	21.0%
SE	1.7%	1.4%	1.3%	1.3%	1.3%	1.0%	0.6%
Lower CI	18.3%	17.2%	18.3%	20.9%	22.8%	11.7%	19.8%
Upper CI	24.9%	22.8%	23.4%	26.2%	28.1%	15.6%	22.1%
Base	559	1034	1178	1293	1226	1187	6477

TABLE A2
 LINEAR REGRESSION MODELS OF DEPRESSIVE DISORDER: APMS, 2000

Dependent variable:	Model (1)			Model (2)		
Depressive disorder	b	SE	95% CI	b	SE	95% CI
<i>Age group</i>						
25–34	0.004	0.006	[−0.008, 0.015]	0.011	0.007	[−0.002, 0.025]
35–44	0.012	0.006	[0.000, 0.025]	0.022	0.008	[0.007, 0.037]
45–54	0.014	0.007	[0.001, 0.027]	0.023	0.008	[0.007, 0.039]
55–64	0.011	0.006	[−0.002, 0.023]	0.005	0.008	[−0.011, 0.021]
65–74	−0.011	0.005	[−0.021, 0.000]	−0.035	0.009	[−0.052, −0.017]
<i>Gender</i>						
Female				−0.002	0.004	[−0.010, 0.005]
<i>Marital status</i>						
Separated				0.016	0.012	[−0.008, 0.040]
Single				0.008	0.005	[−0.003, 0.018]
Divorced				0.030	0.008	[0.014, 0.047]
Widowed				0.019	0.011	[−0.002, 0.040]
<i>Children</i>						
Children in household				−0.003	0.005	[−0.012, 0.006]
<i>Employment status</i>						
Unemployed				0.013	0.014	[−0.013, 0.040]
Economic inactivity				0.039	0.006	[0.028, 0.051]
<i>Educational qualification</i>						
Teaching/HND/nursing				0.009	0.008	[−0.007, 0.024]
A level				0.005	0.006	[−0.007, 0.017]
GCSE/equivalent				0.007	0.005	[−0.002, 0.016]
None				0.012	0.006	[0.000, 0.025]
<i>Tenure</i>						
Social renter				0.005	0.006	[−0.007, 0.017]
Private or other renter				0.007	0.005	[−0.002, 0.016]
Constant	0.019	0.005	[0.010, 0.028]	−0.009	0.010	[−0.028, 0.010]
Overall R ²		0.002			0.028	
Number of individuals		8580			8495	

Notes

Assessment using the Revised Clinical Interview Schedule (CIS-R) to identify presence of current depressive disorder according to diagnostic criteria.

TABLE A3
LINEAR REGRESSION MODELS OF DEPRESSIVE DISORDER: APMS, 2007

Dependent variable:	Model (1)			Model (2)		
Depressive disorder	b	SE	95% CI	b	SE	95% CI
<i>Age group</i>						
25–34	−0.006	0.009	[−0.024, 0.012]	0.015	0.010	[−0.004, 0.035]
35–44	0.002	0.009	[−0.016, 0.019]	0.021	0.011	[0.001, 0.042]
45–54	0.016	0.010	[−0.004, 0.036]	0.033	0.012	[0.009, 0.057]
55–64	−0.007	0.009	[−0.025, 0.011]	−0.002	0.011	[−0.024, 0.019]
65–74	−0.016	0.009	[−0.033, 0.001]	−0.029	0.011	[−0.052, −0.007]
<i>Gender</i>						
Female				0.004	0.004	[−0.005, 0.013]
<i>Marital status</i>						
Cohabiting				−0.010	0.006	[−0.021, 0.002]
Single				0.018	0.007	[0.003, 0.032]
Widowed				0.020	0.008	[0.005, 0.035]
Divorced				0.032	0.010	[0.012, 0.052]
Separated				0.037	0.018	[0.002, 0.071]
<i>Children</i>						
Children in household				−0.008	0.006	[−0.021, 0.004]
<i>Employment status</i>						
Unemployed				0.041	0.026	[−0.011, 0.092]
Economic inactivity				0.034	0.007	[0.021, 0.048]
<i>Educational qualification</i>						
Teaching/HND/nursing				0.011	0.009	[−0.006, 0.028]
A level				0.004	0.006	[−0.008, 0.016]
GCSE/equivalent				0.010	0.005	[−0.001, 0.020]
Other				0.003	0.009	[−0.014, 0.020]
None				0.015	0.006	[0.003, 0.026]
<i>Tenure</i>						
Social renter				0.032	0.008	[0.017, 0.048]
Private or other renter				−0.007	0.006	[−0.020, 0.005]
Constant	0.032	0.008	[−0.040, 0.014]	−0.013	0.014	[−0.040, 0.014]
Overall R ²		0.003			0.030	
Number of individuals		7403			7212	

Notes

Assessment using the Revised Clinical Interview Schedule (CIS-R) to identify presence of current depressive disorder according to diagnostic criteria.

TABLE A4
 LINEAR REGRESSION MODELS OF DEPRESSIVE DISORDER: APMS, 2014

Dependent variable:	Model (1)			Model (2)		
Depressive disorder	b	SE	95% CI	b	SE	95% CI
<i>Age group</i>						
25–34	0.011	0.008	[−0.005, 0.028]	0.034	0.010	[0.014, 0.054]
35–44	0.018	0.009	[0.001, 0.035]	0.048	0.011	[0.026, 0.070]
45–54	0.022	0.009	[0.005, 0.039]	0.051	0.011	[0.029, 0.072]
55–64	0.020	0.008	[0.003, 0.036]	0.034	0.011	[0.014, 0.055]
65–74	−0.002	0.007	[−0.016, 0.012]	−0.012	0.012	[−0.035, 0.010]
75+	−0.011	0.007	[−0.024, 0.003]	−0.034	0.012	[−0.058, −0.009]
<i>Gender</i>						
Female				0.004	0.005	[−0.005, 0.013]
<i>Marital status</i>						
Single				0.020	0.008	[0.003, 0.036]
Divorced/Separated/Widowed				0.004	0.006	[−0.008, 0.016]
<i>Children</i>						
Children in household				−0.006	0.007	[−0.019, 0.006]
<i>Employment status</i>						
Unemployed				0.034	0.017	[0.000, 0.069]
Economic inactivity				0.058	0.008	[0.042, 0.074]
<i>Educational qualification</i>						
Teaching/HND/nursing				0.000	0.008	[−0.015, 0.015]
A level				0.003	0.007	[−0.010, 0.016]
GCSE/equivalent				0.008	0.007	[−0.005, 0.021]
Other/foreign				0.010	0.011	[−0.011, 0.031]
None				0.015	0.008	[−0.001, 0.031]
<i>Tenure</i>						
Social renter				0.050	0.009	[0.032, 0.068]
Private or other renter				0.009	0.006	[−0.003, 0.021]
Constant	0.023	0.006	[0.012, 0.034]	−0.037	0.012	[−0.060, −0.014]
Overall R ²		0.004			0.043	
Number of individuals		7546			7438	

Notes

Assessment using the Revised Clinical Interview Schedule (CIS-R) to identify presence of current depressive disorder according to diagnostic criteria.

TABLE A5
 LINEAR REGRESSION MODELS OF GENERALIZED ANXIETY DISORDER: APMS, 2000

Dependent variable:	Model (1)			Model (2)		
GAD	b	SE	95% CI	b	SE	95% CI
<i>Age group</i>						
25–34	0.029	0.006	[0.016, 0.041]	0.032	0.008	[0.017, 0.047]
35–44	0.042	0.007	[0.028, 0.055]	0.041	0.009	[0.023, 0.058]
45–54	0.054	0.008	[0.038, 0.070]	0.048	0.010	[0.028, 0.068]
55–64	0.034	0.007	[0.020, 0.047]	0.010	0.010	[−0.010, 0.030]
65–74	0.012	0.006	[0.000, 0.024]	−0.034	0.012	[−0.057, −0.010]
<i>Gender</i>						
Female				−0.003	0.005	[−0.013, 0.007]
<i>Marital status</i>						
Separated				0.036	0.015	[0.006, 0.066]
Single				0.000	0.007	[−0.014, 0.014]
Divorced				0.048	0.011	[0.027, 0.069]
Widowed				0.002	0.011	[−0.019, 0.024]
<i>Children</i>						
Children in household				−0.007	0.006	[−0.019, 0.005]
<i>Employment status</i>						
Unemployed				0.011	0.013	[−0.014, 0.036]
Economic inactivity				0.051	0.008	[0.035, 0.067]
<i>Educational qualification</i>						
Teaching/HND/nursing				0.000	0.010	[−0.020, 0.021]
A level				0.003	0.008	[−0.014, 0.019]
GCSE/equivalent				0.002	0.007	[−0.012, 0.017]
None				0.017	0.009	[0.000, 0.034]
<i>Tenure</i>						
Social renter				0.015	0.008	[−0.019, 0.059]
Private or other renter				0.000	0.008	[0.020, 0.983]
Constant	0.014	0.004	[0.006, 0.022]	0.000	0.012	[−0.023, 0.024]
Overall R ²		0.007			0.026	
Number of individuals		8580			8495	

Notes

Assessment using the Revised Clinical Interview Schedule (CIS-R) to identify presence of current GAD according to diagnostic criteria. GAD is the most common type of mental disorder in the population and is characterized by feelings of fear and worry severe enough to impact on day-to-day living.

TABLE A6

LINEAR REGRESSION MODELS OF GENERALIZED ANXIETY DISORDER: APMS, 2007

Dependent variable:	Model (1)			Model (2)		
GAD	b	SE	95% CI	b	SE	95% CI
<i>Age group</i>						
25–34	0.006	0.009	[−0.012, 0.025]	0.016	0.010	[−0.005, 0.037]
35–44	0.018	0.009	[0.000, 0.036]	0.030	0.012	[0.007, 0.054]
45–54	0.025	0.011	[0.003, 0.047]	0.034	0.014	[0.007, 0.061]
55–64	0.006	0.009	[−0.012, 0.024]	0.005	0.013	[−0.020, 0.030]
65–74	−0.003	0.009	[−0.021, 0.015]	−0.015	0.014	[−0.043, 0.012]
<i>Gender</i>						
Female				0.015	0.006	[0.004, 0.026]
<i>Marital status</i>						
Cohabiting				0.013	0.010	[−0.007, 0.034]
Single				−0.001	0.009	[−0.018, 0.017]
Widowed				0.011	0.010	[−0.009, 0.031]
Divorced				0.028	0.013	[0.002, 0.053]
Separated				0.026	0.019	[−0.010, 0.063]
<i>Children</i>						
Children in household				−0.011	0.007	[−0.026, 0.003]
<i>Employment status</i>						
Unemployed				0.071	0.028	[0.015, 0.127]
Economic inactivity				0.027	0.007	[0.014, 0.041]
<i>Educational qualification</i>						
Teaching/HND/nursing				0.020	0.011	[−0.001, 0.040]
A level				0.016	0.009	[−0.001, 0.034]
GCSE/equivalent				0.004	0.007	[−0.010, 0.019]
Other				0.005	0.012	[−0.018, 0.029]
None				0.012	0.008	[−0.004, 0.027]
<i>Tenure</i>						
Social renter				0.042	0.009	[0.024, 0.060]
Private or other renter				0.011	0.009	[−0.007, 0.029]
Constant	0.035	0.007	[0.021, 0.050]	−0.020	0.017	[−0.053, 0.013]
Overall R ²		0.003			0.023	
Number of individuals		7403			7212	

Notes

Assessment using the Revised Clinical Interview Schedule (CIS-R) to identify presence of current GAD according to diagnostic criteria. GAD is the most common type of mental disorder in the population and is characterized by feelings of fear and worry severe enough to impact on day-to-day living.

TABLE A7
 LINEAR REGRESSION MODELS OF GENERALIZED ANXIETY DISORDER: APMS, 2014

Dependent variable:	Model (1)			Model (2)		
GAD	b	SE	95% CI	b	SE	95% CI
<i>Age group</i>						
25–34	–0.002	0.014	[–0.029, 0.025]	0.023	0.016	[–0.008, 0.055]
35–44	0.006	0.013	[–0.020, 0.032]	0.042	0.015	[0.012, 0.073]
45–54	0.009	0.013	[–0.017, 0.035]	0.040	0.016	[0.008, 0.071]
55–64	0.001	0.013	[–0.025, 0.027]	0.019	0.016	[–0.012, 0.051]
65–74	–0.024	0.012	[–0.046, –0.001]	–0.028	0.016	[–0.060, 0.004]
75+	–0.038	0.012	[–0.061, –0.016]	–0.059	0.017	[–0.093, –0.024]
<i>Gender</i>						
Female				0.016	0.007	[0.002, 0.029]
<i>Marital status</i>						
Single				0.034	0.010	[0.014, 0.054]
Divorced/Separated/Widowed				0.018	0.008	[0.002, 0.034]
<i>Children</i>						
Children in household				–0.014	0.009	[–0.032, 0.004]
<i>Employment status</i>						
Unemployed				–0.008	0.016	[–0.040, 0.024]
Economic inactivity				0.043	0.009	[0.024, 0.061]
<i>Educational qualification</i>						
Teaching/HND/nursing				0.010	0.012	[–0.013, 0.033]
A level				–0.002	0.009	[–0.020, 0.016]
GCSE/equivalent				0.008	0.009	[–0.009, 0.025]
Other/foreign				–0.002	0.013	[–0.028, 0.024]
None				0.026	0.011	[0.006, 0.047]
<i>Tenure</i>						
Social renter				0.043	0.011	[0.022, 0.065]
Private or other renter				0.011	0.009	[–0.008, 0.029]
Constant	0.063	0.010	[0.043, 0.084]	–0.016	0.018	[–0.050, 0.019]
Overall R ²		0.004			0.025	
Number of individuals		7546			7438	

Notes

Assessment using the Revised Clinical Interview Schedule (CIS-R) to identify presence of current GAD according to diagnostic criteria. GAD is the most common type of mental disorder in the population and is characterized by feelings of fear and worry severe enough to impact on day-to-day living.

TABLE A8

LINEAR REGRESSION MODELS OF REPORTING 'LIFE NOT WORTH LIVING': APMS, 2000

Dependent variable:	Model (1)			Model (2)		
Felt life not worth living	b	SE	95% CI	b	SE	95% CI
<i>Age group</i>						
25–34	0.034	0.018	[−0.001, 0.069]	0.068	0.020	[0.029, 0.106]
35–44	0.040	0.018	[0.005, 0.075]	0.090	0.021	[0.050, 0.130]
45–54	0.040	0.018	[0.004, 0.076]	0.095	0.022	[0.051, 0.139]
55–64	0.027	0.019	[−0.010, 0.064]	0.055	0.023	[0.009, 0.101]
65–74	−0.040	0.018	[−0.075, −0.004]	−0.053	0.024	[−0.101, −0.005]
<i>Gender</i>						
Female				0.066	0.010	[0.047, 0.085]
<i>Marital status</i>						
Separated				0.144	0.030	[0.085, 0.203]
Single				0.072	0.015	[0.043, 0.102]
Divorced				0.152	0.018	[0.116, 0.188]
Widowed				0.119	0.021	[0.078, 0.161]
<i>Children</i>						
Children in household				−0.014	0.013	[−0.039, 0.011]
<i>Employment status</i>						
Unemployed				0.049	0.030	[−0.011, 0.109]
Economic inactivity				0.082	0.013	[0.056, 0.108]
<i>Educational qualification</i>						
Teaching/HND/nursing				−0.007	0.020	[−0.046, 0.031]
A level				−0.005	0.018	[−0.041, 0.031]
GCSE/equivalent				−0.010	0.015	[−0.039, 0.019]
None				−0.015	0.016	[−0.047, 0.017]
<i>Tenure</i>						
Social renter				0.077	0.014	[0.000, 0.153]
Private or other renter				0.035	0.018	[−0.009, 0.053]
Constant	0.181	0.015	[0.152, 0.209]	−0.022	0.027	[−0.075, 0.031]
Overall R ²		0.004			0.050	
Number of individuals		8574			8491	

Notes

Dependent variable equals 1 if responding 'Yes' to a question on whether the person has felt that life is not worth living, 0 otherwise.

TABLE A9

LINEAR REGRESSION MODELS OF REPORTING 'LIFE NOT WORTH LIVING': APMS, 2014

Dependent variable:	Model (1)			Model (2)		
Felt life not worth living	b	SE	95% CI	b	SE	95% CI
<i>Age group</i>						
25–34	–0.016	0.023	[–0.062, 0.030]	0.046	0.027	[–0.008, 0.099]
35–44	–0.007	0.022	[–0.049, 0.036]	0.086	0.026	[0.034, 0.137]
45–54	0.021	0.022	[–0.022, 0.063]	0.100	0.026	[0.049, 0.151]
55–64	0.040	0.022	[–0.004, 0.083]	0.091	0.028	[0.037, 0.145]
65–74	–0.079	0.020	[–0.118, –0.040]	–0.061	0.029	[–0.118, –0.005]
75+	–0.067	0.020	[–0.106, –0.027]	–0.092	0.030	[–0.152, –0.033]
<i>Gender</i>						
Female				0.046	0.011	[0.025, 0.066]
<i>Marital status</i>						
Single				0.074	0.018	[0.039, 0.108]
Divorced/Separated/Widowed				0.085	0.013	[0.060, 0.110]
<i>Children</i>						
Children in household				–0.049	0.015	[–0.078, –0.020]
<i>Employment status</i>						
Unemployed				0.058	0.036	[–0.013, 0.129]
Economic inactivity				0.072	0.015	[0.043, 0.100]
<i>Educational qualification</i>						
Teaching/HND/nursing				0.007	0.020	[–0.033, 0.047]
A level				0.001	0.017	[–0.033, 0.035]
GCSE/equivalent				0.010	0.015	[–0.019, 0.039]
Other/foreign				–0.007	0.027	[–0.060, 0.046]
None				0.033	0.017	[0.000, 0.065]
<i>Tenure</i>						
Social renter				0.132	0.015	[0.102, 0.162]
Private or other renter				0.062	0.015	[0.033, 0.091]
Constant	0.214	0.017	[0.181, 0.247]	0.011	0.029	[–0.047, 0.068]
Overall R ²		0.009			0.056	
Number of individuals		7537			7429	

Notes

Dependent variable equals 1 if responding 'Yes' to a question on whether the person has felt that life is not worth living, 0 otherwise.

Suicide in cross-national data

In the analysis, we have omitted the very highest ages, because the study's focus is on whether there is midlife distress. In some countries, it should be emphasized, including in the USA in some years where there is widespread access to guns, there is evidence of a turn-up in suicidality towards the end of life (though this is less true of females). That is consistent with the intuitive idea that very old people, with major illnesses, are statistically more prone to take their own lives. The current study does not focus on that segment of the lifespan, and some might wish to argue that self-inflicted deaths at the very end of life, by those in pain or with extreme illness, may not be a major public policy concern.

Figure A1 presents a simple illustrative pattern. Table A10 gives more details on the cohort-adjusted estimates. The dependent variable in that table is the natural logarithm of the suicide rate (expressed per 100,000 citizens). Each data point is a 5-year average. The countries covered in Table A10 are the English-speaking ones of Canada, Australia, Ireland, New Zealand, England and Wales (necessarily combined in our dataset), Northern Ireland, Scotland, and the USA. The base category in Table A10 is Canada, so the coefficients on the country dummy variables are level effects relative to Canada.

Of primary interest is the pattern of the age dummies. For English-speaking females in Table A10, the age dummy variables have negative coefficients through youth, and then turn positive at ages 35–39. From there they rise steadily to ages 50–54, with the coefficients altering, across the age blocks, from -0.014 at ages 30–34, to 0.13 at ages 35–39, 0.29 at ages 40–44, 0.41 at ages 45–49, and 0.47 at ages 50–54. Then the coefficients decline through the numbers, respectively, 0.37 at ages 55–59, 0.23 at ages 60–64, 0.06 at ages 65–69, and -0.17 at ages 70–74. This evidence of a marked hill shape in suicide risk by age is also found for English-speaking males. However, in the case of the second column of Table A10, which is for the male subsample, it would certainly be fair to conclude that the hill shape reaches its literal peak somewhat beyond what would usually be called 'midlife'. It does so at ages 55–59. The coefficients suggest a slightly smoother hill-top, where the pattern is rising from 0.09 at ages 35–39, to 0.22 at ages 40–44, to 0.27 at ages 45–49, to 0.28 at ages 50–54, to 0.29 at ages 55–59, and then down to 0.20 at ages 60–64, 0.08 at ages 65–69, and -0.016 at ages 70–74. These coefficients depict in a visual way the age profile of suicide in the principal English-speaking nations of the world. The period dummies in Table A10 reveal a significant amount of variation, with the highest values occurring in the late 1970s, throughout the 1980s, and early in the 1990s. These were the periods of high unemployment rates in the industrialized world; but in this analysis it is not possible to give an explanation for the observed pattern. There are also strong patterns in the cohort dummies. Low suicide-risk values occur among birth cohorts that were born in the 1920s to the 1940s.




FIGURE A1. (Unadjusted) suicide data for the English-speaking nations in 2015/16.

Notes: These are the raw data (that is, not regression-adjusted).

TABLE A10

SUICIDE REGRESSION EQUATIONS FOR THE ENGLISH-SPEAKING NATIONS SINCE 1950

Dependent variable: Logged 5-year average suicide rate per 100,000 citizens

	Female	Male	Both
Female			−1.06 (−64.9)***
<i>Country dummies</i>			
Australia	0.19 (4.05)***	0.052 (1.31)	0.12 (3.66)***
Ireland	−0.78 (−16.3)***	−0.72 (−18.3)***	−0.75 (−22.4)***
New Zealand	0.11 (2.27)**	−0.059 (−1.53)	0.023 (0.71)
England and Wales	−0.19 (−4.00)***	−0.52 (−13.5)***	−0.36 (−10.8)***
Northern Ireland	−0.38 (−8.14)***	−0.67 (−17.4)***	−0.53 (−16.0)***
Scotland	0.012 (0.26)	−0.35 (−9.13)***	−0.17 (−5.19)***
USA	−0.018 (−0.38)	0.034 (0.86)	0.0078 (0.23)
<i>Age dummies</i>			
15–19	−1.09 (−26.9)***	−1.08 (−32.3)***	−1.08 (−38.2)***
20–24	−0.49 (−12.7)***	−0.24 (−7.52)***	−0.37 (−13.5)***
25–29	−0.19 (−4.84)***	−0.094 (−2.89)***	−0.14 (−5.15)***
30–34	−0.014 (−0.35)	−0.017 (−0.52)	−0.015 (−0.56)
35–39	0.13 (3.35)***	0.094 (2.90)***	0.11 (4.10)***
40–44	0.29 (7.35)***	0.22 (6.87)***	0.26 (9.29)***
45–49	0.41 (10.4)***	0.27 (8.42)***	0.34 (12.4)***
50–54	0.47 (11.9)***	0.28 (8.53)***	0.37 (13.5)***
55–59	0.37 (9.41)***	0.29 (8.94)***	0.33 (12.0)***
60–64	0.23 (5.90)***	0.20 (6.28)***	0.22 (7.90)***
65–69	0.062 (1.58)	0.081 (2.50)**	0.072 (2.60)***
70–74	−0.17 (−4.38)***	−0.016 (−0.50)	−0.095 (−3.41)***
<i>Period dummies</i>			
1950–1954	−0.55 (−13.7)***	−0.49 (−14.9)***	−0.52 (−18.5)***
1955–1959	−0.30 (−7.32)***	−0.29 (−8.65)***	−0.29 (−10.3)***
1960–1964	0.012 (0.30)	−0.15 (−4.49)***	−0.069 (−2.41)**

TABLE A10
Continued

Dependent variable: Logged 5-year average suicide rate per 100,000 citizens			
	Female	Male	Both
1965–1969	0.18 (4.42)***	−0.085 (−2.54)**	0.047 (1.66)*
1970–1974	0.21 (5.12)***	−0.063 (−1.89)*	0.072 (2.54)**
1975–1979	0.36 (8.84)***	0.081 (2.42)**	0.22 (7.73)***
1980–1984	0.32 (8.00)***	0.24 (7.13)***	0.28 (9.90)***
1985–1989	0.23 (5.55)***	0.30 (9.02)***	0.26 (9.27)***
1990–1994	0.086 (2.13)**	0.28 (8.46)***	0.18 (6.50)***
1995–1999	0.013 (0.32)	0.20 (6.13)***	0.11 (3.84)***
2000–2004	−0.090 (−2.23)**	0.073 (2.19)**	−0.008 (−0.30)
2005–2009	−0.095 (−2.23)**	0.030 (0.84)	−0.033 (−1.10)
2010–2014	−0.38 (−7.07)***	−0.13 (−2.98)***	−0.25 (−6.79)***
<i>Cohort dummies</i>			
1878	0.69 (5.17)***	0.75 (6.81)***	0.72 (7.70)***
1883	0.48 (5.01)***	0.53 (6.64)***	0.50 (7.48)***
1888	0.44 (5.54)***	0.35 (5.33)***	0.40 (7.09)***
1893	0.21 (2.97)***	0.20 (3.46)***	0.21 (4.15)***
1898	0.089 (1.38)	0.069 (1.30)	0.079 (1.74)*
1903	−0.044 (−0.74)	−0.033 (−0.66)	−0.039 (−0.92)
1908	−0.14 (−2.51)**	−0.19 (−4.01)***	−0.16 (−4.15)***
1913	−0.16 (−3.07)***	−0.30 (−6.78)***	−0.23 (−6.18)***
1918	−0.25 (−4.85)***	−0.41 (−9.74)***	−0.33 (−9.19)***
1923	−0.31 (−6.48)***	−0.44 (−11.0)***	−0.38 (−11.1)***
1928	−0.45 (−9.74)***	−0.49 (−13.1)***	−0.47 (−14.6)***
1933	−0.48 (−10.9)***	−0.52 (−14.4)***	−0.50 (−16.3)***
1938	−0.53 (−11.9)***	−0.57 (−15.4)***	−0.55 (−17.6)***
1943	−0.49 (−10.5)***	−0.52 (−13.5)***	−0.50 (−15.4)***
1948	−0.42 (−8.70)***	−0.41 (−10.1)***	−0.41 (−12.2)***

TABLE A10
Continued

Dependent variable: Logged 5-year average suicide rate per 100,000 citizens			
	Female	Male	Both
1953	−0.35 (−6.96)***	−0.23 (−5.50)***	−0.29 (−8.20)***
1958	−0.29 (−5.44)***	−0.12 (−2.72)***	−0.21 (−5.48)***
1963	−0.20 (−3.56)***	0.018 (0.39)	−0.092 (−2.30)**
1968	−0.11 (−1.76)*	0.18 (3.58)***	0.036 (0.86)
1973	0.046 (0.69)	0.27 (4.97)***	0.16 (3.42)***
1978	0.24 (3.29)***	0.33 (5.42)***	0.29 (5.53)***
1983	0.40 (4.64)***	0.36 (5.17)***	0.38 (6.35)***
1988	0.67 (6.20)***	0.50 (5.59)***	0.59 (7.71)***
1993	0.96 (5.19)***	0.65 (4.28)***	0.81 (6.22)***
Constant	2.09 (58.1)***	3.32 (111)***	3.23 (122)***
Observations	1198	1199	2397

Notes

Intrinsic estimator approach to solve the APC problem of identification with person overlaps, using 5-period (years) averages for the English-speaking subsample. Cohort dummies overlap in this model just as in Thibodeau's work on Canada. z -statistics in parentheses.

***, **, * indicate $p < 0.01$, $p < 0.05$, $p < 0.1$, respectively.

TABLE A11
SUICIDE REGRESSION EQUATIONS FOR NON-ENGLISH-SPEAKING NATIONS SINCE 1950

Dependent variable: Logged 5-year average suicide rate per 100,000 citizens			
	Females	Males	Both
Female			−0.98 (−83.4)***
<i>Country dummies</i>			
Belgium	−0.078 (−2.17)**	−0.29 (−10.1)***	−0.19 (−7.44)***
Denmark	0.065 (1.79)*	−0.25 (−8.69)***	−0.094 (−3.75)***
Finland	−0.000091 (−0.0025)	0.20 (7.02)***	0.10 (4.07)***
France	−0.19 (−5.33)***	−0.25 (−8.79)***	−0.22 (−8.95)***
Netherlands	−0.52 (−14.5)***	−1.02 (−35.4)***	−0.77 (−31.0)***

TABLE A11

Continued

Dependent variable: Logged 5-year average suicide rate per 100,000 citizens			
	Females	Males	Both
Norway	−0.62 (−17.0)***	−0.64 (−21.9)***	−0.63 (−25.0)***
Sweden	−0.097 (−2.68)***	−0.31 (−10.6)***	−0.20 (−8.07)***
Switzerland	0.015 (0.40)	−0.12 (−4.09)***	−0.052 (−2.09)**
<i>Age dummies</i>			
15–19	−1.20 (−38.1)***	−1.20 (−47.2)***	−1.20 (−54.9)***
20–24	−0.58 (−19.6)***	−0.40 (−16.9)***	−0.49 (−24.0)***
25–29	−0.38 (−13.1)***	−0.25 (−10.5)***	−0.32 (−15.6)***
30–34	−0.18 (−6.35)***	−0.15 (−6.29)***	−0.16 (−8.24)***
35–39	0.005 (0.16)	−0.003 (−0.13)	0.001 (0.040)
40–44	0.16 (5.61)***	0.16 (6.78)***	0.16 (7.98)***
45–49	0.32 (11.2)***	0.28 (12.0)***	0.30 (15.0)***
50–54	0.45 (15.5)***	0.33 (14.4)***	0.39 (19.6)***
55–59	0.42 (14.6)***	0.35 (14.8)***	0.39 (19.1)***
60–64	0.36 (12.4)***	0.29 (12.2)***	0.33 (16.1)***
65–69	0.34 (11.6)***	0.28 (11.8)***	0.31 (15.2)***
70–74	0.29 (9.67)***	0.32 (13.3)***	0.30 (14.7)***
<i>Period dummies</i>			
1950–1954	−0.44 (−7.47)***	−0.33 (−6.90)***	−0.39 (−9.40)***
1955–1959	−0.12 (−3.79)***	−0.072 (−2.86)***	−0.095 (−4.40)***
1960–1964	−0.092 (−2.98)***	−0.061 (−2.46)**	−0.077 (−3.58)***
1965–1969	−0.0092 (−0.30)	0.0042 (0.17)	−0.0025 (−0.12)
1970–1974	0.15 (5.09)***	0.11 (4.35)***	0.13 (6.20)***
1975–1979	0.28 (9.36)***	0.20 (8.50)***	0.24 (11.7)***
1980–1984	0.35 (11.8)***	0.31 (13.0)***	0.33 (16.0)***
1985–1989	0.34 (11.4)***	0.28 (11.9)***	0.31 (15.2)***

TABLE A11
Continued

Dependent variable: Logged 5-year average suicide rate per 100,000 citizens			
	Females	Males	Both
1990–1994	0.16 (5.57)***	0.17 (7.23)***	0.17 (8.22)***
1995–1999	0.016 (0.55)	0.049 (2.05)**	0.032 (1.59)
2000–2004	−0.095 (−3.20)***	−0.088 (−3.68)***	−0.091 (−4.45)***
2005–2009	−0.20 (−6.79)***	−0.23 (−9.58)***	−0.22 (−10.5)***
2010–2014	−0.33 (−10.2)***	−0.34 (−13.0)***	−0.34 (−14.9)***
<i>Cohort dummies</i>			
1878	0.55 (2.80)***	0.69 (4.41)***	0.62 (4.58)***
1883	0.33 (3.69)***	0.31 (4.29)***	0.32 (5.16)***
1888	0.18 (2.51)**	0.22 (3.91)***	0.20 (4.08)***
1893	0.12 (1.95)*	0.15 (3.03)***	0.13 (3.16)***
1898	0.040 (0.72)	0.11 (2.56)**	0.076 (2.01)**
1903	−0.0040 (−0.079)	0.043 (1.06)	0.019 (0.56)
1908	−0.013 (−0.28)	−0.051 (−1.35)	−0.032 (−0.99)
1913	−0.076 (−1.75)*	−0.15 (−4.21)***	−0.11 (−3.71)***
1918	−0.11 (−2.69)***	−0.22 (−6.67)***	−0.16 (−5.81)***
1923	−0.16 (−4.20)***	−0.25 (−7.96)***	−0.20 (−7.66)***
1928	−0.18 (−4.97)***	−0.24 (−8.39)***	−0.21 (−8.46)***
1933	−0.19 (−5.56)***	−0.27 (−9.96)***	−0.23 (−9.80)***
1938	−0.15 (−4.67)***	−0.29 (−11.2)***	−0.22 (−9.85)***
1943	−0.17 (−5.31)***	−0.28 (−10.6)***	−0.23 (−9.99)***
1948	−0.14 (−4.00)***	−0.18 (−6.67)***	−0.16 (−6.76)***
1953	−0.091 (−2.60)***	−0.084 (−2.98)***	−0.088 (−3.60)***
1958	−0.095 (−2.57)**	−0.051 (−1.73)*	−0.073 (−2.86)***
1963	−0.14 (−3.63)***	−0.047 (−1.49)	−0.094 (−3.48)***

TABLE A11

Continued

Dependent variable: Logged 5-year average suicide rate per 100,000 citizens			
	Females	Males	Both
1968	−0.18 (−4.45)***	−0.020 (−0.60)	−0.10 (−3.55)***
1973	−0.15 (−3.22)***	−0.018 (−0.49)	−0.081 (−2.61)***
1978	−0.063 (−1.26)	0.063 (1.58)	0.00025 (0.0073)
1983	0.094 (1.65)*	0.12 (2.54)**	0.11 (2.66)***
1988	0.20 (2.83)***	0.14 (2.44)**	0.17 (3.46)***
1993	0.40 (3.74)***	0.29 (3.39)***	0.35 (4.66)***
Constant	2.56 (90.5)***	3.71 (163)***	3.63 (177)***
Observations	1320	1320	2640

Notes

Intrinsic estimator approach to solve the APC problem of identification with person overlaps, using 5-period (years) averages for affluent non-English-speaking countries. z-statistics in parentheses.


***, **, * indicate $p < 0.01$, $p < 0.05$, $p < 0.1$, respectively.

Table A11 gives equivalent results for non-English affluent countries in Europe. The countries covered in Table A11 are Belgium, Denmark, Finland, France, the Netherlands, Norway, Sweden and Switzerland, and the base category is Austria.


Sleeping problems in cross-national data

Figures A2 and A3 use information only on work-week days. However, we have found that a U-shaped midlife pattern is observed even during weekends; all calculations can be redone with 7-day data. Furthermore, we excluded from the analysis any children under 10 years of age, and did not include naps—defined as any sleep taking place between 11am and 9pm—in our measure of sleep duration. However, including sleep naps and focusing on overall sleep duration throughout the day yields similar results.

The U-shaped pattern uses self-reported sleeping hours, and thus might be some form of statistical illusion or error of measurement. This is a natural, and scientifically appropriate, concern. Yet the UK hospital admissions data cast doubt on that concern. Lauderdale *et al.* (2008) also offers more general support for the similarity of subjective and objective sleep information. We additionally explored the patterns in data from the Cleveland Family Study; that suggests again, in objectively measured sleeping hours data, an approximate U-shape in age. Also, using panel data from Germany, and graphical and Granger-causality methods, we checked that the sleep–age profile mirrors an age U-shape in life satisfaction.




(a) Midlife and sleep hours, by country




(b) Midlife and sleep hours, by demographic group (pooling 9 countries)

FIGURE A2. Sleep hours, by nation and type of person.

Notes: This figure documents U-shaped sleep hours versus age in a sample of half a million individuals. Time-use data for Austria, Canada, Finland, France, Germany, the Netherlands, Spain, the UK and the USA, 1965–2012. Age is plotted on the x-axis. The U-shape pattern continues to hold in regression equations that include standard demographic controls, including for children in the household. In part (b), low-skilled are defined as individuals with completed secondary education or below. High-skilled are defined as individuals with greater than secondary education. For clarity, standard error bands are omitted, but underlying regression equations are given in Appendix Tables A1–A9.



(a) Midlife and sleep hours, by time period (pooling 9 countries)



(b) Within-person longitudinal data on sleep hours and age (Great Britain)

FIGURE A3. Sleep hours, by time period and in intrapersonal data.
Notes: This figure documents U-shaped sleep hours versus age in a sample of half a million individuals. Time-use data for Austria, Canada, Finland, France, Germany, the Netherlands, Spain, the UK and the USA, 1965–2012. Age is plotted on the x -axis. The U-shape pattern continues to hold in regression equations that include standard demographic controls, including for children in the household. The plot in part (b) uses data drawn from the BHPS for the period 1994–2004. This is based on fixed effects equations, so the pattern is derived solely from within-person, not cross-sectional, variation. For clarity, standard error bands are omitted, but underlying regression equations are given in Appendix Tables A1–A9.

TABLE A12
REGRESSION EQUATIONS FOR SLEEP

	Sleep hours			Sleep less than 6 hours		
	(1)	(2)	(3)	(4)	(5)	(6)
<i>Age group</i>						
20–24	−0.491*** (0.089)	−0.496*** (0.090)	−0.236* (0.118)	0.036*** (0.007)	0.035*** (0.007)	0.030*** (0.004)
25–30	−0.795*** (0.057)	−0.800*** (0.057)	−0.441*** (0.101)	0.040*** (0.009)	0.039*** (0.009)	0.036*** (0.005)
31–34	−0.959*** (0.064)	−0.965*** (0.065)	−0.556*** (0.110)	0.041*** (0.009)	0.040*** (0.010)	0.038*** (0.006)
35–40	−1.053*** (0.070)	−1.058*** (0.071)	−0.620*** (0.117)	0.042*** (0.010)	0.042*** (0.010)	0.039*** (0.006)
41–44	−1.082*** (0.074)	−1.089*** (0.074)	−0.662*** (0.114)	0.043*** (0.011)	0.043*** (0.011)	0.040*** (0.007)
45–50	−1.047*** (0.098)	−1.053*** (0.098)	−0.681*** (0.108)	0.046*** (0.011)	0.045*** (0.011)	0.043*** (0.006)
51–54	−0.999*** (0.123)	−1.003*** (0.123)	−0.702*** (0.108)	0.044*** (0.011)	0.044*** (0.011)	0.042*** (0.006)
55–60	−0.836*** (0.145)	−0.840*** (0.144)	−0.655*** (0.110)	0.028*** (0.006)	0.027*** (0.006)	0.030*** (0.004)
61–64	−0.555** (0.184)	−0.559** (0.183)	−0.556*** (0.144)	0.011* (0.005)	0.010* (0.005)	0.021*** (0.006)
65–70	−0.287 (0.153)	−0.287 (0.153)	−0.442** (0.141)	−0.002 (0.004)	−0.003 (0.004)	0.014* (0.006)
71–74	−0.083 (0.184)	−0.086 (0.182)	−0.316 (0.185)	−0.010* (0.005)	−0.011* (0.005)	0.007 (0.005)
75–80	0.089 (0.223)	0.086 (0.219)	−0.187 (0.241)	−0.013*** (0.003)	−0.014*** (0.002)	0.004 (0.007)
81–84	0.436* (0.217)	0.433* (0.214)	0.114 (0.250)	−0.018*** (0.002)	−0.020*** (0.002)	−0.002 (0.006)
85–90	1.570*** (0.142)	1.560*** (0.128)	1.267*** (0.146)	−0.061** (0.020)	−0.057** (0.021)	−0.043* (0.020)
90+	2.654*** (0.269)	2.642*** (0.272)	2.352*** (0.283)	−0.059** (0.020)	−0.054** (0.022)	−0.041* (0.020)
Year FE	Yes	Yes	Yes	Yes	Yes	Yes
Country FE	No	Yes	Yes	No	Yes	Yes
Controls	No	No	Yes	No	No	Yes
Observations	256,776	256,776	256,776	256,776	256,776	256,776
R-squared	0.094	0.095	0.126	0.033	0.034	0.040

Notes

Time-use data for Austria, Canada, Finland, France, the Netherlands, Spain, the UK and the USA (1965–2012). Dependent variables are the amount of ‘Sleep hours’ and a binary indicator for individuals who ‘Sleep less than 6 hours’. Age is in banded 5-year intervals. Controls include income, education, employment status, number of children, marital status, gender and self-reported health. All estimates include year, month and day-of-the-week dummies. Standard errors (in parentheses) are clustered at the country level. All estimates used the survey proposed weights.

***, **, * indicate $p < 0.01$, $p < 0.05$, $p < 0.1$, respectively.

TABLE A13

WITHIN-PERSON REGRESSION EQUATIONS FOR SLEEP (GREAT BRITAIN)

	Sleep hours and personal care			Short sleep hours and personal care		
	(1)	(2)	(3)	(4)	(5)	(6)
<i>Age group</i>						
20–24	−0.319*** (0.017)	−0.142*** (0.011)	−0.132*** (0.012)	0.099*** (0.009)	0.035*** (0.008)	0.034*** (0.010)
25–30	−0.697*** (0.020)	−0.309*** (0.014)	−0.281*** (0.016)	0.273*** (0.012)	0.129*** (0.011)	0.129*** (0.016)
31–34	−0.811*** (0.020)	−0.366*** (0.015)	−0.329*** (0.019)	0.349*** (0.012)	0.173*** (0.012)	0.165*** (0.019)
35–40	−0.858*** (0.020)	−0.395*** (0.015)	−0.363*** (0.021)	0.387*** (0.012)	0.194*** (0.013)	0.188*** (0.021)
41–44	−0.810*** (0.020)	−0.375*** (0.015)	−0.353*** (0.023)	0.370*** (0.012)	0.184*** (0.013)	0.179*** (0.023)
45–50	−0.637*** (0.021)	−0.295*** (0.016)	−0.297*** (0.024)	0.266*** (0.011)	0.117*** (0.013)	0.145*** (0.025)
51–54	−0.468*** (0.021)	−0.230*** (0.016)	−0.248*** (0.026)	0.182*** (0.011)	0.073*** (0.013)	0.111*** (0.027)
55–60	−0.226*** (0.022)	−0.110*** (0.017)	−0.142*** (0.027)	0.082*** (0.010)	0.005 (0.013)	0.058** (0.028)
61–64	0.164*** (0.022)	0.060*** (0.018)	−0.001 (0.029)	−0.022** (0.009)	−0.050*** (0.013)	0.013 (0.029)
65–70	0.550*** (0.020)	0.291*** (0.020)	0.166*** (0.031)	−0.071*** (0.008)	−0.073*** (0.014)	−0.003 (0.030)
71–74	0.811*** (0.021)	0.476*** (0.021)	0.311*** (0.033)	−0.079*** (0.008)	−0.067*** (0.014)	−0.002 (0.030)
75–80	1.090*** (0.021)	0.698*** (0.022)	0.491*** (0.034)	−0.080*** (0.008)	−0.062*** (0.014)	−0.002 (0.030)
81–84	1.397*** (0.023)	0.963*** (0.023)	0.735*** (0.037)	−0.082*** (0.008)	−0.059*** (0.015)	−0.004 (0.030)
85–90	1.734*** (0.029)	1.269*** (0.026)	1.017*** (0.040)	−0.082*** (0.008)	−0.062*** (0.016)	−0.004 (0.031)
90+	2.145*** (0.045)	1.659*** (0.035)	1.326*** (0.048)	−0.081*** (0.008)	−0.064*** (0.020)	−0.005 (0.031)
Year FE	Yes	Yes	Yes	Yes	Yes	Yes
Controls	No	Yes	Yes	No	Yes	Yes
Person FE	No	No	Yes	No	No	Yes
Observations	91,686	91,686	91,686	91,686	91,686	91,686
R-squared	0.608	0.805	0.931	0.164	0.349	0.692

Notes

Data are drawn from the BHPS calibrated time-use data (1994–2004). Time-use variables were measured using evidence derived from a smaller-scale panel survey that collected time-use information by both the survey and diary methods (Home OnLine, 1998–2001). Dependent variables are the amount of ‘Sleep hours and personal care’, and a binary indicator for individuals who spent less than 9.18 hours on ‘Sleep and personal care’, equivalent to the bottom quartile of the distribution. Age is in banded 5-year intervals. Controls include gender, marital status, household type, number of children, employment status and retirement status. All estimates include year, month and day-of-the-week dummies. Standard errors (in parentheses) are clustered at the individual level. All estimates used the survey proposed weights.

***, **, * indicate $p < 0.01$, $p < 0.05$, $p < 0.1$, respectively.

Disabling headaches and job stress

TABLE A14
MIGRAINE AND AGE: FIXED EFFECT LOGIT REGRESSIONS

Dependent variable: Migraine	Model (1)	Model (2)
Age	0.076*** (0.008)	
Age-squared	−0.001*** (0.000)	
Age 26–35		0.194*** (0.051)
Age 36–45		0.244*** (0.062)
Age 46–55		0.219*** (0.069)
Age 56–65		−0.084 (0.080)
Age 66–75		−0.496*** (0.101)
Log of real equivalent income	−0.075*** (0.021)	−0.071*** (0.021)
Disabled	0.766*** (0.065)	0.765*** (0.065)
Unemployed	−0.116** (0.056)	−0.127** (0.056)
Self-employed	−0.422*** (0.080)	−0.425*** (0.080)
Retired	0.154** (0.071)	0.072 (0.068)
Not in the labour market	0.317*** (0.044)	0.277*** (0.043)
Married	0.193*** (0.064)	0.238*** (0.064)
Cohabiting	0.297*** (0.058)	0.342*** (0.058)
Divorced	0.257*** (0.088)	0.310*** (0.087)
Widowed/widower	0.289** (0.112)	0.293*** (0.112)
Separated	0.211** (0.104)	0.268*** (0.104)
Completed higher degree	−0.079 (0.142)	−0.060 (0.142)
Completed first degree	−0.300*** (0.084)	−0.277*** (0.084)
HND/HNC/teaching qualification	−0.042 (0.090)	−0.028 (0.090)
A level qualifications	−0.119* (0.062)	−0.108* (0.062)
O level qualifications	0.042 (0.054)	0.045 (0.054)
CSE qualifications	0.096 (0.091)	0.103 (0.090)
Homeowner	−0.173*** (0.041)	−0.177*** (0.041)

TABLE A14
Continued

Dependent variable: Migraine	Model (1)	Model (2)
Number of days spent in hospital last year	0.002 (0.001)	0.002 (0.001)
Number of children aged under 16	0.015 (0.019)	0.035* (0.020)
Outer London	0.197 (0.151)	0.189 (0.151)
Rest of South-East	0.208 (0.133)	0.196 (0.133)
South-West	0.195 (0.144)	0.185 (0.144)
East Anglia	0.096 (0.162)	0.083 (0.162)
East Midlands	0.007 (0.144)	−0.003 (0.144)
West Midlands conurbation	0.190 (0.157)	0.177 (0.157)
Rest of West Midlands	0.079 (0.158)	0.066 (0.158)
Greater Manchester	0.079 (0.186)	0.072 (0.186)
Merseyside	0.058 (0.198)	0.051 (0.198)
Rest of North-West	0.034 (0.167)	0.022 (0.167)
South Yorkshire	0.326* (0.196)	0.321 (0.196)
West Yorkshire	0.226 (0.165)	0.217 (0.165)
Rest of York and Humberside	0.000 (0.177)	−0.011 (0.177)
Tyne and Wear	−0.091 (0.222)	−0.097 (0.222)
Rest of North	0.173 (0.172)	0.168 (0.172)
Wales	0.141 (0.131)	0.131 (0.131)
Scotland	0.112 (0.130)	0.104 (0.131)
Northern Ireland	−0.177 (0.134)	−0.182 (0.134)
Constant	−3.228*** (0.274)	−2.089*** (0.235)
Year dummies	Yes	Yes
Observations	213,011	213,011
Log-likelihood	−59,641.145	−59,700.759

Notes

Longitudinal data from the BHPS, waves 1–18. These are within-person, not cross-sectional, estimates. Dependent variable is derived from the BHPS health questionnaire: ‘Do you have any of the listed health problems: ... migraine or frequent headaches?’ Migraine variable equals 1 if respondent answered ‘Yes’, 0 otherwise. The youngest age group (16–25) is the base reference category. Standard errors are given in parentheses.

***, **, * indicate $p < 0.01$, $p < 0.05$, $p < 0.1$, respectively.

TABLE A15
JOB STRESS EQUATIONS FOR AUSTRALIAN DATA

Dependent variable:	Model (1)			Model (2)			Model (3)		
Severe job stress	β	95% CI	<i>p</i>	β	95% CI	<i>p</i>	β	95% CI	<i>p</i>
<i>Age group</i>									
25–34	0.52	[0.48, 0.56]	0.00	0.23	[0.19, 0.26]	0.00	0.10	[0.06, 0.14]	0.00
35–44	0.67	[0.62, 0.72]	0.00	0.35	[0.31, 0.40]	0.00	0.14	[0.09, 0.20]	0.00
45–54	0.77	[0.71, 0.83]	0.00	0.40	[0.34, 0.45]	0.00	0.13	[0.06, 0.20]	0.00
55–64	0.72	[0.65, 0.78]	0.00	0.38	[0.32, 0.45]	0.00	0.05	[−0.03, 0.13]	0.25
65–75	0.48	[0.39, 0.56]	0.00	0.32	[0.23, 0.41]	0.00	−0.08	[−0.19, 0.03]	0.14
<i>Income and job-related variables</i>									
Log of household income				0.09	[0.07, 0.10]	0.00	0.06	[0.05, 0.08]	0.00
Work hours per week				0.02	[0.02, 0.03]	0.00	0.02	[0.02, 0.03]	0.00
Recently promoted				0.13	[0.11, 0.15]	0.00	0.13	[0.11, 0.15]	0.00
Recently changed jobs				−0.17	[−0.18, −0.15]	0.00	−0.17	[−0.19, −0.15]	0.00
Recently bankrupt				0.09	[0.03, 0.14]	0.00	0.09	[0.04, 0.14]	0.00
Recently received major financial gain				−0.02	[−0.05, 0.02]	0.36	−0.02	[−0.05, 0.02]	0.32
<i>Education level</i>									
Masters or doctorate							0.25	[0.13, 0.37]	0.00
Bachelor or honours							0.32	[0.21, 0.43]	0.00
Graduate diploma/certificate							0.24	[0.16, 0.32]	0.00
Advanced diploma							0.10	[0.01, 0.19]	0.23
Professional qualification							0.02	[−0.04, 0.08]	0.50
Completed high school							−0.08	[−0.13, −0.02]	0.00
Currently full-time student							−0.05	[−0.09, −0.01]	0.02
<i>Marital status</i>									
Married							0.06	[0.02, 0.11]	0.01
De facto							0.03	[−0.01, 0.07]	0.14
Separated							0.01	[−0.07, 0.08]	0.84
Divorced							−0.01	[−0.09, 0.07]	0.78
Widowed							0.01	[−0.14, 0.15]	0.91
<i>Number of dependent children</i>									
No. children under age 4							0.00	[−0.02, 0.02]	0.74
No. children aged 5–14							0.01	[−0.00, 0.03]	0.16
<i>Lifestyle variables</i>									
Long-term health issues							0.03	[0.01, 0.05]	0.00
Non-smoker							−0.02	[−0.05, 0.01]	0.19
Drink alcohol every day							0.06	[0.01, 0.10]	0.00
Constant	2.67	[2.63, 2.71]	0.00	0.92	[0.77, 1.08]	0.00	1.21	[1.05, 1.38]	0.00
Year dummies	No			No			Yes		
Industry dummies	No			Yes			Yes		
Overall R ²	0.03			0.15			0.17		
Number of individuals	20,648			20,648			20,648		
Number of observations	127,199			127,199			127,199		

Notes

Longitudinal (fixed effects) regression models of severe job stress on age, HILDA Survey (2002–18). Respondents assigned an integer value between 1, 'Strongly disagree', and 7, 'Strongly agree', to each statement: 'I fear the amount of stress in my job will make me physically ill'; 'My job is complex and difficult'; 'My job is more stressful than I had ever imagined'. Averaged responses to the three statements form a combined 'Job stress' measure. The youngest age group (15–24) is the base reference category. The analysed sample is restricted to employed individuals; aged between 15 and 75; working between 5 and 90 hours per week. 'Year dummies' control for each of the 17 survey waves. 'Industry dummies' control for 19 different job–industry categories (e.g. manufacturing, construction, mining, financial services, healthcare, education, hospitality).

Miscellaneous

TABLE A16

LINEAR REGRESSION MODELS OF CONCENTRATION PROBLEMS AND FORGETFULNESS: APMS, 2014

Dependent variable:	Model (1)			Model (2)		
Concentration problems and forgetfulness	b	SE	95% CI	b	SE	95% CI
<i>Age group</i>						
25–34	–0.004	0.017	[–0.037, 0.029]	0.028	0.018	[–0.008, 0.063]
35–44	0.016	0.017	[–0.017, 0.049]	0.058	0.019	[0.020, 0.096]
45–54	0.017	0.016	[–0.016, 0.049]	0.055	0.019	[0.017, 0.093]
55–64	–0.002	0.016	[–0.034, 0.030]	0.015	0.020	[–0.023, 0.054]
65–74	–0.055	0.015	[–0.085, –0.025]	–0.084	0.021	[–0.125, –0.042]
75+	–0.047	0.015	[–0.077, –0.017]	–0.101	0.022	[–0.145, –0.057]
<i>Gender</i>						
Female				0.028	0.008	[0.012, 0.043]
<i>Marital status</i>						
Single				0.021	0.013	[–0.004, 0.046]
Divorced/Separated/Widowed				0.030	0.010	[0.010, 0.051]
<i>Children</i>						
Children in household				–0.006	0.011	[–0.027, 0.016]
<i>Employment status</i>						
Unemployed				0.059	0.029	[0.002, 0.116]
Economic inactivity				0.103	0.012	[0.079, 0.126]
<i>Educational qualification</i>						
Teaching/HND/nursing				0.004	0.013	[–0.023, 0.030]
A level				0.009	0.012	[–0.014, 0.033]
GCSE/equivalent				0.031	0.011	[0.009, 0.054]
Other/foreign				0.020	0.019	[–0.018, 0.058]
None				0.023	0.011	[0.001, 0.046]
<i>Tenure</i>						
Social renter				0.071	0.014	[0.043, 0.098]
Private or other renter				0.031	0.011	[0.009, 0.052]
Constant	0.104	0.013	[0.078, 0.131]	–0.030	0.023	[–0.076, 0.015]
Overall R ²		0.007			0.050	
Number of individuals		7546			7438	

Notes

As indicated on the symptom scores administered as part of the Clinical Interview Schedule—Revised (CIS-R).

TABLE A17
 LINEAR REGRESSION MODELS OF CONCENTRATION PROBLEMS AND FORGETFULNESS: APMS, 2000

Dependent variable: Concentration problems and forgetfulness	Model (1)			Model (2)		
	b	SE	95% CI	b	SE	95% CI
<i>Age group</i>						
25–34	0.024	0.013	[−0.001, 0.049]	0.037	0.013	[0.010, 0.063]
35–44	0.030	0.013	[0.005, 0.056]	0.046	0.015	[0.018, 0.075]
45–54	0.040	0.015	[0.011, 0.068]	0.056	0.017	[0.022, 0.089]
55–64	0.008	(0.013	[−0.018, 0.034]	−0.003	0.017	[−0.037, 0.030]
65–74	−0.020	(0.013	[−0.045, 0.005]	−0.068	0.018	[−0.104, −0.032]
<i>Gender</i>						
Female				0.004	0.007	[−0.011, 0.018]
<i>Marital status</i>						
Separated				0.062	0.022	[0.019, 0.105]
Single				0.013	0.011	[−0.009, 0.035]
Divorced				0.030	0.013	[0.005, 0.055]
Widowed				0.032	0.015	[0.001, 0.062]
<i>Children</i>						
Children in household				−0.002	0.010	[−0.021, 0.018]
<i>Employment status</i>						
Unemployed				0.010	0.019	[−0.028, 0.048]
Economic inactivity				0.078	0.010	[0.058, 0.098]
<i>Educational qualification</i>						
Teaching/HND/nursing				0.019	0.015	[−0.010, 0.047]
A level				0.023	0.014	[−0.003, 0.050]
GCSE/equivalent				0.017	0.011	[−0.004, 0.038]
None				0.030	0.012	[0.006, 0.053]
<i>Tenure</i>						
Social renter				0.020	0.011	[1.810, 0.070]
Private or other renter				0.031	0.013	[2.410, 0.017]
Constant	0.080	0.010	[0.060, 0.101]	0.015	0.020	[−0.024, 0.054]
Overall R ²		0.004			0.023	
Number of individuals		8580			8495	

Notes

As indicated on the symptom scores administered as part of the Clinical Interview Schedule—Revised (CIS-R).

TABLE A18

LINEAR REGRESSION MODELS OF ALCOHOL DEPENDENCE: APMS, 2014

Dependent variable:	Model (1)			Model (2)		
	b	SE	95% CI	b	SE	95% CI
<i>Alcohol dependence</i>						
<i>Age group</i>						
25–34	0.001	0.012	[−0.022, 0.025]	0.019	0.013	[−0.007, 0.045]
35–44	0.000	0.011	[−0.020, 0.021]	0.028	0.013	[0.003, 0.054]
45–54	−0.014	0.011	[−0.036, 0.008]	0.011	0.013	[−0.014, 0.035]
55–64	−0.014	0.011	[−0.035, 0.007]	0.006	0.013	[−0.019, 0.031]
65–74	−0.031	0.010	[−0.051, −0.011]	−0.010	0.012	[−0.034, 0.014]
75+	−0.040	0.010	[−0.059, −0.021]	−0.019	0.013	[−0.043, 0.006]
<i>Gender</i>						
Female				−0.023	0.005	[−0.033, −0.014]
<i>Marital status</i>						
Single				0.028	0.008	[0.013, 0.044]
Divorced/Separated/Widowed				0.005	0.005	[−0.005, 0.016]
<i>Children</i>						
Children in household				−0.022	0.006	[−0.034, −0.010]
<i>Employment status</i>						
Unemployed				0.015	0.019	[−0.022, 0.051]
Economic inactivity				−0.001	0.006	[−0.014, 0.012]
<i>Educational qualification</i>						
Teaching/HND/nursing				−0.004	0.010	[−0.023, 0.016]
A level				−0.003	0.008	[−0.020, 0.013]
GCSE/equivalent				0.006	0.008	[−0.010, 0.022]
Other/foreign				0.008	0.016	[−0.023, 0.038]
None				0.002	0.007	[−0.012, 0.017]
<i>Tenure</i>						
Social renter				0.008	0.008	[−0.007, 0.023]
Private or other renter				0.011	0.008	[−0.005, 0.027]
Constant	0.042	0.010	[0.024, 0.061]	0.052	0.017	[0.019, 0.085]
Overall R ²		0.006			0.020	
Number of individuals		7264			7164	

Notes

Alcohol dependence indicated by an AUDIT score of 16 or more.

TABLE A19
 LINEAR REGRESSION MODELS OF ALCOHOL DEPENDENCE: APMS, 2000

Dependent variable:	Model (1)			Model (2)		
	b	SE	95% CI	b	SE	95% CI
Alcohol dependence						
<i>Age group</i>						
25–34	–0.035	0.012	[–0.058, –0.012]	–0.008	0.013	[–0.034, 0.017]
35–44	–0.051	0.012	[–0.074, –0.028]	–0.012	0.013	[–0.037, 0.014]
45–54	–0.064	0.011	[–0.086, –0.041]	–0.031	0.013	[–0.057, –0.005]
55–64	–0.071	0.011	[–0.093, –0.048]	–0.042	0.013	[–0.068, –0.015]
65–74	–0.074	0.011	[–0.096, –0.052]	–0.046	0.014	[–0.074, –0.018]
<i>Gender</i>						
Female				–0.036	0.004	[–0.045, –0.028]
<i>Marital status</i>						
Separated				0.037	0.013	[0.011, 0.063]
Single				0.035	0.007	[0.021, 0.049]
Divorced				0.028	0.008	[0.013, 0.043]
Widowed				0.011	0.005	[0.001, 0.021]
<i>Children</i>						
Children in household				–0.020	0.006	[–0.033, –0.008]
<i>Employment status</i>						
Unemployed				0.042	0.020	[0.002, 0.082]
Economic inactivity				0.002	0.005	[–0.008, 0.013]
<i>Educational qualification</i>						
Teaching/HND/nursing				0.014	0.010	[–0.006, 0.034]
A level				0.012	0.008	[–0.004, 0.029]
GCSE/equivalent				0.009	0.007	[–0.004, 0.022]
None				0.009	0.007	[–0.005, 0.023]
<i>Tenure</i>						
Social renter				0.008	0.006	[–0.330, 0.185]
Private or other renter				0.012	0.010	[–0.013, 0.259]
Constant	0.082	0.011	[0.061, 0.103]	0.091	0.016	[0.059, 0.123]
Overall R ²		0.017			0.042	
Number of individuals		8538			8467	

Notes

Alcohol dependence indicated by an AUDIT score of 16 or more. Note that this table has large standard error bands.

TABLE A20

LINEAR REGRESSION MODELS OF SUICIDAL THOUGHTS IN THE PAST YEAR: APMS, 2014

Dependent variable:	Model (1)			Model (2)		
Suicidal thoughts in past year	b	SE	95% CI	b	SE	95% CI
<i>Age group</i>						
25–34	–0.029	0.015	[–0.058, 0.000]	0.003	0.017	[–0.030, 0.036]
35–44	–0.032	0.014	[–0.061, –0.004]	0.011	0.017	[–0.022, 0.044]
45–54	–0.031	0.014	[–0.058, –0.003]	0.008	0.016	[–0.024, 0.039]
55–64	–0.035	0.014	[–0.062, –0.008]	–0.011	0.016	[–0.043, 0.021]
65–74	–0.066	0.013	[–0.092, –0.041]	–0.061	0.018	[–0.096, –0.026]
75+	–0.065	0.013	[–0.090, –0.041]	–0.075	0.018	[–0.110, –0.039]
<i>Gender</i>						
Female				–0.004	0.006	[–0.017, 0.008]
<i>Marital status</i>						
Single				0.036	0.011	[0.014, 0.058]
Divorced/Separated/Widowed				0.034	0.007	[0.019, 0.048]
<i>Children</i>						
Children in household				–0.017	0.009	[–0.034, 0.000]
<i>Employment status</i>						
Unemployed				0.062	0.027	[0.009, 0.115]
Economic inactivity				0.043	0.009	[0.026, 0.060]
<i>Educational qualification</i>						
Teaching/HND/nursing				–0.012	0.009	[–0.030, 0.006]
A level				–0.007	0.010	[–0.027, 0.014]
GCSE/equivalent				0.011	0.009	[–0.007, 0.029]
Other/foreign				–0.024	0.009	[–0.043, –0.005]
None				0.010	0.009	[–0.008, 0.027]
<i>Tenure</i>						
Social renter				0.027	0.011	[0.006, 0.048]
Private or other renter				0.011	0.009	[–0.006, 0.028]
Constant	0.084	0.012	[0.060, 0.108]	0.033	0.019	[–0.005, 0.071]
Overall R ²		0.008			0.031	
Number of individuals		7546			7438	

Notes

The question is: ‘There may be times in everyone’s life when they become very miserable and depressed and may feel like taking drastic action because of these feelings. Have you ever thought of taking your life, even if you would not really do it?’ Note that this table has large standard error bands.

TABLE A21

LINEAR REGRESSION MODELS OF SUICIDAL THOUGHTS IN THE PAST YEAR: APMS, 2000

Dependent variable:	Model (1)			Model (2)		
Suicidal thoughts in past year	b	SE	95% CI	b	SE	95% CI
<i>Age group</i>						
25–34	−0.026	0.011	[−0.048, −0.003]	−0.015	0.012	[−0.039, 0.008]
35–44	−0.025	0.011	[−0.047, −0.003]	−0.010	0.013	[−0.036, 0.015]
45–54	−0.033	0.012	[−0.055, −0.010]	−0.019	0.014	[−0.046, 0.008]
55–64	−0.049	0.010	[−0.070, −0.029]	−0.048	0.012	[−0.072, −0.023]
65–74	−0.059	0.010	[−0.079, −0.039]	−0.074	0.013	[−0.100, −0.049]
<i>Gender</i>						
Female				−0.002	0.005	[−0.011, 0.008]
<i>Marital status</i>						
Separated				0.051	0.015	[0.021, 0.081]
Single				0.018	0.007	[0.004, 0.032]
Divorced				0.020	0.007	[0.006, 0.035]
Widowed				0.038	0.012	[0.015, 0.061]
<i>Children</i>						
Children in household				−0.003	0.006	[−0.016, 0.010]
<i>Employment status</i>						
Unemployed				0.016	0.017	[−0.017, 0.050]
Economic inactivity				0.031	0.007	[0.017, 0.045]
<i>Educational qualification</i>						
Teaching/HND/nursing				−0.015	0.008	[−0.031, 0.001]
A level				−0.003	0.009	[−0.021, 0.015]
GCSE/equivalent				−0.004	0.007	[−0.019, 0.010]
None				0.004	0.009	[−0.013, 0.021]
<i>Tenure</i>						
Social renter				0.018	0.008	[0.002, 0.026]
Private or other renter				0.008	0.010	[−0.079, 0.430]
Constant	0.069	0.010	[0.049, 0.088]	0.045	0.015	[0.015, 0.074]
Overall R ²		0.008			0.022	
Number of individuals		8572			8489	

Notes

The question is: ‘There may be times in everyone’s life when they become very miserable and depressed and may feel like taking drastic action because of these feelings. Have you ever thought of taking your life, even if you would not really do it?’ Note that this table has large standard error bands.

ACKNOWLEDGMENTS

For valuable comments, we thank the editor and three referees, plus Danny Dorling, Amanda Goodall, Daniel Hamermesh, and attendees in presentations at a large number of research seminars, including Oxford, IZA Bonn, NBER Cambridge, UCLA, Carnegie Mellon, Copenhagen, Bristol, Mannheim, BRIQ Bonn, UCL-Harvard, Monash, and the University of British Columbia. This work started in late 2018 and has had a long gestation. Blanchflower (2020a), which described and cited an early draft of our work while our paper was at a confidential stage, examines cross-sectional data on ill-being. The published version of that paper, Blanchflower (2020b), does not cite our work.

NOTES

1. This study's approach is likely to seem natural to most economists and epidemiologists, but some psychologists prefer the tradition of non-adjusted studies of age and ageing. Because of different intellectual traditions, occasionally the debate in related literatures has been at cross-purposes and perhaps needlessly confrontational.
2. The data are available from <https://platform.who.int/> mortality (accessed 29 October 2022).
3. The data are available from www.mtusdata.org (accessed 29 October 2022).
4. The data are available from www.ukdataservice.ac.uk (accessed 29 October 2022).
5. The data and other access details are available from the HILDA website: <https://melbourneinstitute.unimelb.edu.au/hilda> (accessed 29 October 2022).
6. The data are available from: <https://www.understandingsociety.ac.uk> (accessed 29 October 2022).
7. The HILDA survey data were extracted using PanelWhiz (Haisken-DeNew and Hahn 2006).

REFERENCES

- BHULLER, M., MOGSTAD, M. and SALVANES, K. G. (2017). Life-cycle earnings, education premiums, and internal rates of return. *Journal of Labor Economics*, **35**, 993–1030.
- BLANCHFLOWER, D. G. (2020a). Unhappiness and age. NBER Working Paper no. 26642.
- (2020b). Unhappiness and age. *Journal of Economic Behavior and Organization*, **176**, 461–88.
- and OSWALD, A. J. (2008). Is well-being U-shaped over the life cycle? *Social Science & Medicine*, **66**, 1733–49.
- and ——— (2016). Antidepressants and age: a new form of evidence for U-shaped well-being through life. *Journal of Economic Behavior & Organization*, **127**, 46–58.
- and ——— (2020). Trends in extreme distress in the United States, 1993–2019. *American Journal of Public Health*, **110**(10), 1538–44.
- CARLSSON, F., JOHANSSON-STENMAN, O. and MARTINSSON, P. (2007). Do you enjoy having more than others? Survey evidence of positional goods. *Economica*, **74**, 586–98.
- CASE, A. and DEATON, A. (2015). Rising morbidity and mortality in midlife among white non-Hispanic Americans in the 21st century. *Proceedings of the National Academy of Sciences of the USA*, **112**(49), 15078–83.
- CHARLES, S. T., MATHER, M. and CARSTENSEN, L. L. (2003). Aging and emotional memory: the forgettable nature of negative images for older adults. *Journal of Experimental Psychology—General*, **132**(2), 310–24.
- CHENG, T., POWDTHAVEE, N. and OSWALD, A. J. (2017). Longitudinal evidence for a midlife nadir in human well-being: results from four data sets. *Economic Journal*, **127**(599), 126–42.
- CHOI, B. (2018). Job strain, long work-hours, and suicide ideation in US workers: a longitudinal study. *International Archives of Occupational and Environmental Health*, **91**, 865–75.
- DALY, M. (2022). Prevalence of psychological distress among working-age adults in the United States, 1999–2018. *American Journal of Public Health*, **112**, 1045–9.
- DIENER, E., SUH, E. M., LUCAS, R. E. and SMITH, H. L. (1999). Subjective well-being: three decades of progress. *Psychological Bulletin*, **125**(2), 276–302.
- DOLAN, P. and KAHNEMAN, D. (2007). Interpretations of utility and their implications for the valuation of health. *Economic Journal*, **118**, 215–34.
- , PEASGOOD, T. and WHITE, M. (2008). Do we really know what makes us happy? A review of the economic literature on the factors associated with subjective wellbeing. *Journal of Economic Psychology*, **29**, 94–122.
- EASTERLIN, R. A. (2003). Explaining happiness. *Proceedings of the National Academy of Sciences of the USA*, **100**(19), 11176–83.
- FISHER, K., HOFFERTH, S., FLOOD, S., ROMAN, J. G., LEE, Y. and GERSHUNY, J. (2015). Introducing the American Heritage Time-use Study Data Extract Builder (AHTUS-X). *Electronic International Journal of Time-Use Research*, **12**(1), 169–73.
- FREUND, A. M. and RITTER, J. O. (2009). Midlife crisis: a debate. *Gerontology*, **55**, 582–91.
- FRUTERS, P. and BEATON, T. (2012). The mystery of the U-shaped relationship between happiness and age. *Journal of Economic Behavior & Organization*, **82**, 525–42.
- FU, W. J. (2000). Ridge estimator in singular design with application to age-period-cohort analysis of disease rates. *Communications in Statistics—Theory and Method*, **29**, 263–78.
- FURMAN, Y., WOLF, S. M. and ROSENFELD, D. S. (1997). Shakespeare and sleep disorders. *Neurology*, **49**(4), 1171–2.
- GALAMBOS, N. L., KRAHN, H. J., JOHNSON, M. D. and LACHMAN, M. E. (2020). The U shape of happiness across the life course: expanding the discussion. *Perspectives on Psychological Science*, **15**, 898–912.

- GAYDOS, L., HUMMER, R. A., HARGROVE, T. W., HALPERN, C. T., HUSSEY, J. M., WHITSEL, E. A., DOLE, N. and HARRIS, K. M. (2019). The depths of despair among US adults entering midlife. *American Journal of Public Health*, **109**(5), 774–80.
- GIUNTELLA, O., HAN, W. and MAZZONNA, F. (2017). Circadian rhythms, sleep and cognitive skills: evidence from an unsleeping giant. *Demography*, **54**, 1715–42.
- GOLDMAN, N., GLEI, D. A. and WEINSTEIN, M. (2018). Declining mental health among disadvantaged Americans. *Proceedings of the National Academy of Sciences of the USA*, **115**, 7290–5.
- GRAHAM, C. (2017). *Happiness for All? Unequal Hopes and Lives in Pursuit of the American Dream*. Princeton, NJ: Princeton University Press.
- and POZUELO, J. R. (2017). Happiness, stress, and age: how the U curve varies across people and places. *Journal of Population Economics*, **30**(1), 225–64.
- , ZHOU, S. and ZHANG, J. (2017). Happiness and health in China: the paradox of progress. *World Development*, **96**, 231–44.
- GUNNELL, D., MIDDLETON, N., WHITLEY, E., DORLING, D. and FRANKEL, S. (2003). Influence of cohort effects on patterns of suicide in England and Wales, 1950–1999. *British Journal of Psychiatry*, **182**, 164–70.
- HAFNER, M., TROXEL, W. M., STEPANEK, M., TAYLOR, J. and VAN STOLK, C. (2017). Why sleep matters: the macroeconomic costs of insufficient sleep. *Journal of Sleep and Sleep Disorders Research*, **40**(Suppl. 1), A297.
- HAISKEN-DENEW, J. P. and HAHN, M. (2006). PanelWhiz: a flexible modularized Stata interface for accessing large scale panel data sets; available online at http://panelwhiz.de/docs/PanelWhiz_Introduction.pdf (accessed 31 October 2022).
- JAKES, E. (1965). Death and the mid-life crisis. *International Journal of Psychoanalysis*, **46**, 502–14.
- JESTE, D. V. and OSWALD, A. J. (2014). Individual and societal wisdom: explaining the paradox of human aging and high well-being. *Psychiatry: Interpersonal and Biological Processes*, **77**, 317–30.
- KRUEGER, P. M. and FRIEDMAN, E. M. (2009). Sleep duration in the United States: a cross-sectional population-based study. *American Journal of Epidemiology*, **169**(9), 1052–63.
- KUNZMANN, U., RICHTER, D. and SCHMUKLE, S. C. (2013). Stability and change in affective experience across the adult life span: analyses with a national sample from Germany. *Emotion*, **13**(6), 1086–95.
- LACHMAN, M. E. (2015). Mind the gap in the middle: a call to study midlife. *Research in Human Development*, **12**, 327–34.
- LAMPL, C., THOMAS, H., TASSORELLI, C., *et al.* (2016). Headache, depression and anxiety: associations in the Eurolight project. *Journal of Headache and Pain*, **17**, article 59.
- LANG, I. A., LLEWELLYN, D. J., HUBBARD, R. E., LANGA, K. M. and MELZER, D. (2011). Income and the midlife peak in common mental disorder prevalence. *Psychological Medicine*, **41**, 1365–72.
- LAUDERDALE, D. S., KNOTSON, K. L., YAN, L. L., LIU, K. and RATHOUZ, P. J. (2008). Self-reported and measured sleep duration: how similar are they? *Epidemiology*, **19**(6), 838–84.
- LAYARD, R. (1980). Human satisfactions and public policy. *Economic Journal*, **90**, 737–50.
- MCMANUS, S., BEBBINGTON, P. E., JENKINS, R., MORGAN, Z., BROWN, L., COLLINSON, D. and BRUGHA, T. (2020). Data resource profile: Adult Psychiatric Morbidity Survey (APMS). *International Journal of Epidemiology*, **49**(2), 361–2.
- MROCEK, D. K. and SPIRO III, A. (2005). Change in life satisfaction during adulthood: findings from the Veterans Affairs normative aging study. *Journal of Personality and Social Psychology*, **88**(1), 189–202.
- MUJIC, R. and OSWALD, A. J. (2018). Is envy harmful to a society's psychological health and wellbeing? A longitudinal study of 18,000 adults. *Social Science & Medicine*, **198**, 103–11.
- OSWALD, A. J. and WU, S. (2010). Objective confirmation of subjective measures of human well-being: evidence from the USA. *Science*, **327**, 576–9.
- PATTEN, S. B. (2001). Long-term medical conditions and major depression in a Canadian population study at waves 1 and 2. *Journal of Affective Disorders*, **63**, 35–41.
- RAMIREZ, A. J., GRAHAM, J., RICHARDS, M. A., CULL, A. and GREGORY, W. M. (1996). Mental health of hospital consultants: the effects of stress and satisfaction at work. *Lancet*, **347**, 724–8.
- RATCLIFFE, G. E., ENNS, M. W., BELIK, S. L. and SAREEN, J. (2008). Chronic pain conditions and suicidal ideation and suicide attempts: an epidemiologic perspective. *Clinical Journal of Pain*, **24**, 204–10.
- RAUCH, J. (2018). *The Happiness Curve: Why Life gets Better after Midlife*. London: Green Tree.
- ROENNEBERG, T. (2013). Chronobiology: the human sleep project. *Nature*, **498**(7455), 427–8.
- SAUNDERS, J. B., AASLAND, O. G., BABOR, T. F., DELA FUENTE, J. R. and GRANT, M. (1993). Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption, Part II. *Addiction*, **88**, 791–804.
- SCHWANDT, H. (2016). Unmet aspirations as an explanation for the age U-shape in wellbeing. *Journal of Economic Behavior & Organization*, **122**(1), 75–87.

- SETIYA, K. (2017). *Midlife: A Philosophical Guide*. Princeton, NJ: Princeton University Press.
- SPIEGEL, K., LEPROULT, R. and VAN CAUTER, E. (1999). Impact of sleep debt on metabolic and endocrine function. *Lancet*, **354**(9188), 1435–9.
- SPIERS, N., BRUGHA, T. S., BEBBINGTON, P., MCMANUS, S., JENKINS, R. and MELTZER, H. (2012). Age and birth cohort differences in depression in repeated cross-sectional surveys in England: the National Psychiatric Morbidity Surveys, 1993 to 2007. *Psychological Medicine*, **42**, 2047–55.
- STANSFELD, S. A., FUHRER, R., SHIPLEY, M. J. and MARMOT, M. G. (1999). Work characteristics predict psychiatric disorder: prospective results from the Whitehall II study. *Occupational and Environmental Medicine*, **56**, 302–7.
- TAYLOR, D. J., LICHSTEIN, K. L., DURRENCE, H. H., REIDEL, B. W. and BUSH, A. J. (2005). Epidemiology of insomnia, depression, and anxiety. *Sleep*, **28**(11), 1457–64.
- TERGESEN, A. (2014). The myth of the midlife crisis. *Wall Street Journal*, 12 October; available online at <https://www.wsj.com/articles/the-myth-of-the-midlife-crisis-1413147918> (accessed 29 October 2022).
- THIBODEAU, L. (2015). Suicide mortality in Canada and Quebec, 1926–2008: an age–period–cohort analysis. *Canadian Studies in Population*, **42**, 1–23.
- VAN LANDEGHEM, B. (2012). A test for the convexity of human well-being over the life cycle: longitudinal evidence from a 20-year panel. *Journal of Economic Behavior & Organization*, **81**(2), 571–82.
- WEISS, A., KING, J. E., INOUE-MURAYAMA, M., MATSUZAWA, T. and OSWALD, A. J. (2012). Evidence for a midlife crisis in great apes consistent with the U-shape in human well-being. *Proceedings of the National Academy of Sciences of the USA*, **109**(49), 19949–52.
- WETHINGTON, E. (2000). Expecting stress: Americans and the midlife crisis. *Motivation and Emotion*, **24**, 85–103.
- WHITBOURNE, S. K. (2015). Worried about a midlife crisis? Don't. There's no such thing. *Psychology Today*, 11 July; available online at <https://www.psychologytoday.com/gb/blog/fulfillment-any-age/201507/worried-about-midlife-crisis-dont-theres-no-such-thing> (accessed 29 October 2022).
- WORLD HEALTH ORGANIZATION (WHO) (1992). The ICD-10 classification of mental and behavioural disorders: clinical descriptions and diagnostic guidelines; available online at <https://apps.who.int/iris/handle/10665/37958> (accessed 31 October 2022).
- WINKELMANN, L. and WINKELMANN, R. (1998). Why are the unemployed so unhappy? Evidence from panel data. *Economica*, **65**, 1–15.
- WOODEN, M. and WATSON, N. (2007). The HILDA survey and its contribution to economic and social research (so far). *Economic Record*, **83**(261), 208–31.
- YANG, Y., FU, W. J. and LAND, K. C. (2004). A methodological comparison of age–period–cohort models: the intrinsic estimator and conventional generalized linear models. *Sociological Methodology*, **34**(1), 75–110.
- , SCHULHOFER-WOHL, S., FU, W. J. and LAND, K. C. (2008). The intrinsic estimator for age–period–cohort analysis: what it is and how to use it. *American Journal of Sociology*, **113**(6), 1697–736.