



# City Research Online

## City St George's, University of London

**Citation:** Goldsmith, L., Anderson, K., Clarke, G., Crowe, C., Jarman, H., Johnson, S., Lomani, J., McDaid, D., Park, A. L., Smith, J. G. & et al (2023). Service use preceding and following first referral for psychiatric emergency care at a short-stay crisis unit: A cohort study across three cities and one rural area in England. *International Journal of Social Psychiatry*, 69(4), pp. 928-941. doi: 10.1177/00207640221142530

This is the published version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

**Permanent repository link:** <https://openaccess.city.ac.uk/id/eprint/29577/>

**Link to published version:** <https://doi.org/10.1177/00207640221142530>

**Copyright and Reuse:** Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

# Service use preceding and following first referral for psychiatric emergency care at a short-stay crisis unit: A cohort study across three cities and one rural area in England

International Journal of  
Social Psychiatry  
1–14

© The Author(s) 2022



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/00207640221142530

journals.sagepub.com/home/isp



Lucy Pollyanna Goldsmith<sup>1</sup> , Katie Anderson<sup>2</sup>,  
Geraldine Clarke<sup>3</sup>, Chloe Crowe<sup>4</sup>, Heather Jarman<sup>1,5</sup>,  
Sonia Johnson<sup>6</sup>, Jo Lomani<sup>7</sup>, David McDaid<sup>8</sup>,  
A-La Park<sup>8</sup>, Jared G Smith<sup>1</sup> and Steven Gillard<sup>9</sup>

## Abstract

**Background:** Internationally, hospital-based short-stay crisis units have been introduced to provide a safe space for stabilisation and further assessment for those in psychiatric crisis. The units typically aim to reduce inpatient admissions and psychiatric presentations to emergency departments.

**Aims:** To assess changes to service use following a service user's first visit to a unit, characterise the population accessing these units and examine equality of access to the units.

**Methods:** A prospective cohort study design (ISCTRN registered; 53431343) compared service use for the 9 months preceding and following a first visit to a short-stay crisis unit at three cities and one rural area in England. Included individuals first visited a unit in the 6 months between 01/September/2020 and 28/February/2021.

**Results:** The prospective cohort included 1189 individuals aged 36 years on average, significantly younger (by 5–13 years) than the population of local service users (<.001). Seventy percent were White British and most were without a psychiatric diagnosis (55%–82% across sites). The emergency department provided the largest single source of referrals to the unit (42%), followed by the Crisis and Home Treatment Team (20%). The use of most mental health services, including all types of admission and community mental health services was increased post discharge. Social-distancing measures due to the COVID-19 pandemic were in place for slightly over 50% of the follow-up period. Comparison to a pre-COVID cohort of 934 individuals suggested that the pandemic had no effect on the majority of service use variables.

**Conclusions:** Short-stay crisis units are typically accessed by a young population, including those who previously were unknown to mental health services, who proceed to access a broader range of mental health services following discharge.

## Keywords

Mental health crisis, psychiatric admission, emergency department, experience of care, mental health nursing, short-stay crisis care

<sup>1</sup>Population Health Research Institute, St George's, University of London, UK

<sup>2</sup>Middlesex University London, UK

<sup>3</sup>The Health Foundation, London, UK

<sup>4</sup>North East London NHS Foundation Trust, Goodmayes Hospital, Ilford, UK

<sup>5</sup>St George's University Hospitals NHS Foundation Trust, London, UK

<sup>6</sup>NIHR Mental Health Policy Research Unit, Division of Psychiatry, University College London – Bloomsbury, UK

<sup>7</sup>NHS England and NHS Improvement, London, UK

<sup>8</sup>Care Policy and Evaluation Centre, Department of Health Policy, London School of Economics and Political Science, UK

<sup>9</sup>School of Health and Psychological Sciences, City, University of London, London, UK

## Corresponding author:

Lucy Pollyanna Goldsmith, Population Health Research Institute, St George's, University of London, Cranmer Terrace, Tooting, London SW17 0RE, UK.

Email: lgoldsmi@sgul.ac.uk

## Introduction

Internationally, psychiatric crisis care services are under intense pressure (The Strategy Unit, 2019). An increasing number of visits to emergency departments (ED) for mental health crisis care and decreasing numbers of inpatient beds (Fleury et al., 2019) has result in long waits in ED (Nicks & Manthey, 2012). Concomitantly it is unclear whether short stays on psychiatric wards followed by early discharge are beneficial (Clibbens et al., 2018). Inpatient stays can be unnecessary (Stulz et al., 2015), costly (McCrone et al., 2009) and detrimental (Thibaut et al., 2019). In crisis services, service users' needs are the most acute, yet their voices can often be ignored (Srebnik & Russo, 2008). For service users, shared decision-making, feeling respected and having basic comforts are priorities when in crisis (Thomas et al., 2018). Within this difficult context, a new type of service is emerging to reduce the pressure on ED and psychiatric wards. Short-stay crisis units, also known as Psychiatric Decision Units (PDU), Mental Health Decision Units, Crisis Stabilisation Units, or Psychiatric Emergency Services have been introduced in Australia (Braitberg et al., 2018), Belgium (Spooren et al., 1997), Canada (Mok & Walker, 1995), The Netherlands (Van Der Sande et al., 1997), the US (Stamy et al., 2021) and England (Trethewey et al., 2019).

In England, psychiatric ward occupancy rates in the publicly funded National Health Service (NHS) have long tended to exceed recommended levels (Crisp et al., 2016), sometimes resulting in long delays for a hospital bed for those in crisis. People presenting with psychological distress at a general hospital ED are over six times more likely than those with physical presentations to breach the 4-hour waiting time target (NHS England et al., 2014), resulting in difficult experiences in noisy environments. PDUs, which are typically designed to alleviate pressure on ED and reduce inpatient admissions can provide an alternative (Goldsmith et al., 2021). A national survey in England identified six units – all were open 24/7 and had maximum stays of 12 to 72 hours. All accepted voluntary (informal) service users only and provided recliner chairs for sleeping rather than beds. Units are staffed by senior mental health nurses and healthcare assistants, with input/oversight from psychiatrists. The staff: patient ratio is high, approximately one member of staff per two service users in daytime hours (Goldsmith et al., 2021). As in other countries, PDUs in England emerged to address pressures on the crisis care pathway, leaving formal evaluation of the units to follow their development, in response to policy initiatives (Department of Health and Concordat Signatories, 2014; NHS England et al., 2014).

A subsequent, comprehensive systematic literature review on hospital-based short-stay crisis units included 12 studies from six countries with a total of 67,505 participants (Anderson et al., 2022). A wide range of comparison groups were used in the included studies including four

pre/post studies comparing service use data before and after the unit opened (Braitberg et al., 2018; Browne et al., 2011; Lester et al., 2018; Stamy et al., 2021). A single interrupted time-series compared pre-post data (Parwani et al., 2018). Five studies compared the PDU to other units (Gillig et al., 1989; Kealy-Bateman et al., 2019; Mok & Walker, 1995), one compared referral routes (Trethewey et al., 2019) and one compared to an inpatient service (Schneider & Ross, 1996). Additionally there was a case-control study (Spooren et al., 1997) and a randomised controlled trial (Van Der Sande et al., 1997). Pooled estimates indicated a reduction in ED length of stay by about 2 hours and 40 minutes ( $-164.24$  minutes, 95% CI  $[-261.24$  to  $-67.23]$  minutes), and that the odds ratio of an ED presentation for psychiatric crisis leading to an inpatient admission were approximately halved ( $OR=0.55$ , 95% CI  $[0.43-0.68]$ ) (Anderson et al., 2022). The units were found to have a wide range of effects which seem likely to be connected to staffing levels, how the units are oriented towards their stated aims and the surrounding crisis care services. None of these prior studies examined equality of access to the service, changes to service use following a visit to a unit at an individual patient level, or examined the effects of the COVID-19 pandemic on service use following discharge from a unit. The present study aims to (1) understand who uses PDUs; (2) examine changes to service use following a service user's first visit to a unit, (3) assess the effects of the COVID-19 pandemic on service use post-discharge and (4) assess equality of access to the units.

## Methods

### Setting

This prospective cohort study was conducted in four participating mental health service providers (NHS mental health trusts) in England – London, Birmingham, Sheffield (city sites) and Lincolnshire (a rural site), each with an operational PDU before the study commenced. Key characteristics of study sites, including configuration of the units and other local crisis care services, are in Table 1. Changes to the unit or other crisis care provision during the timeframe of the research, including those resulting from the COVID-19 pandemic are indicated in parentheses.

### Design

#### Primary cohort study

The prospective, primary cohort study included all individuals at the four study sites experiencing their first visit to a PDU over a 6-month period between 01/September/2020 and 28/February/2021. The study compared routine service use data for the 9 months preceding this visit (the 'pre'

**Table 1.** Key characteristics of study sites.

	South West London	Sheffield	Lincolnshire	Birmingham
Date opened	November-16	March-19	January-18	November-14
Location	Psychiatric hospital	Psychiatric unit on a general hospital site	Psychiatric hospital	Psychiatric hospital
Maximum length of stay	48 hours	48 hours (decreased to 24 hours during the COVID-19 pandemic)	24 hours	Target 24 hours (initially 72 hours maximum, reduced to 12 hours in 2019)
Unit capacity	5 (increased to 7 during the COVID-19 pandemic)	5	6	8 (decreased to 5 during the COVID-19 pandemic)
Referral routes	ED, CRHT, Street Triage	ED, CRHT, Street Triage, CMHT	ED, CRHT, Street Triage (16 months after short-stay crisis unit opens), AMHPs	ED, CRHT, Street Triage
Staffing	Mental Health Nurse, HCA, Psychiatrist	Mental Health Nurse, Support Worker, Psychiatrist	Mental Health Nurse, HCA, Psychiatrist (plus CMHT during the COVID-19 pandemic)	Mental Health Nurse, HCA, Psychiatrist
Staff:patient ratio	1:1 (decreased to 5:7 during the COVID-19 pandemic)	4:5	1:2	1:4
Setting: other local trust-run crisis services	Crisis house, crisis café (×2), (mental health ED opened during the COVID-19 pandemic)	Crisis house	Triage ward, Crisis house	Triage ward (16 months only), Acute day unit, Crisis café, Crisis house

Note. ED=Emergency Department, also known as Accident and Emergency (A&E) in the UK; CRHT=Crisis Resolution & Home Treatment Team – a team which provides intensive short-term support to service users in crisis at home including telephone calls and visits; CMHT=Community Mental Health Team – a secondary mental health care service composed of a multidisciplinary team operating in the community; AMHP=Approved Mental Health Professional – these professionals can make formal legal decisions regarding whether to detain a person under the Mental Health Act; HCA=Health Care Assistant; Street Triage – an emergency response service in which mental health professionals provide telephone support or accompany police and paramedics attending disturbances.

period) to the 9 months following (the ‘post’ period). The follow-up period for the study closed on 30/November/2021, allowing complete follow-up for service users who joined the study at the end of the ‘pre’ period. The study is prospective, and was registered with the ISCTRN (53431343) (Goldsmith, 2020) on 11/February/2020.

To characterise the population accessing PDUs, we summarised socio-demographic data including age, ethnicity, gender, diagnosis, marital and employment status, housing, sexual orientation and referral source. The English indices of deprivation 2019 (McLennan et al., 2019) is a measure of relative deprivation, based on seven different, weighted, domains of deprivation: income, employment, crime, living environment, barriers to housing and services, education, skills and development deprivation and health deprivation and disability. We conducted the equalities impact assessment by comparing the demographic characteristics of people in the primary cohort accessing the crisis unit to the general population of service users (calculated over a recent 1-year period) at the corresponding NHS mental health trust.

To address the third aim, to identify whether there were changes to service use following a visit to a unit, we examined two primary outcomes as PDUs are designed to reduce

the pressure on both general hospital ED and inpatient wards. First, we examined whether there is a change in informal (voluntary) admissions for the cohort between the pre- and post- periods. Second, we examined whether there were changes to the number of ED presentations, measured as liaison psychiatry episodes (referral to the services providing psychiatry support to ED). We also checked for changes in inpatient admissions, short-stay (0–5 days inpatient admissions), average length of inpatient stays, compulsory admissions, use of community mental health team (CMHT) and other mental health trust specialist services to identify the wider effects of a PDU visit.

### *Adaptation due to COVID-19: an additional retrospective cohort*

Widespread restrictions of movement and social interaction were introduced in England in response to the COVID-19 pandemic on 23/03/2020, during the follow-up period for the primary cohort. The effects of the pandemic on access to care and help-seeking behaviours could affect the generalisability of the results to other periods. Hence the study was adapted to include a retrospective cohort of people who visited a unit 1 year prior to the prospectively designed

cohort (eligible first visits to the units were between 01/September/2019 and 28/February/2020). Choosing dates exactly 1 year prior controls for any effect of seasonality on the results. The retrospective cohort includes three of the sites as the fourth site opened their unit during the 'pre' period, meaning that complete data for the retrospective cohort could not be obtained from this site.

### Coproduction

The present study is part of a larger project, 'Evaluating mental health decision units on acute crisis care pathways' (National Institute for Health Research Funding and Awards, Award ID 17/49/70, 2019). The project was coproduced with a national expert panel of people with lived experience of mental health crisis services, a local service user research advisory group and researchers who also have lived experience of psychological distress. This lived experience informed decisions about which outcomes we would collect, which data we included in our equalities impact assessment, and interpretation of results in a data synthesis workshop.

### Data extraction and statistical analysis

Data were extracted from electronic patient records by Business Intelligence teams at each site, pseudonymised, securely transferred to the study team, cleaned and analysed. Each cohort population was summarised using descriptive statistics. We tested for differences in service use in the pre- and post-periods using McNemar's  $\chi^2$  tests (Fagerland et al., 2013) for binary paired data and paired  $t$ -tests bootstrapped with 2,000 replications for continuous data (Efron & Tibshirani, 1994). Unpaired  $t$ -tests were used for the average length of inpatient stay – many in the cohort were not admitted to a ward in both the pre- and post-periods – an unpaired test enabled all data to be included in the analysis. The additional, retrospective cohort facilitated examination of whether there were changes in service use due to the COVID-19 pandemic. The number of service users in each cohort was used as a denominator and tests for significant differences in service use across the two post-periods at each site were conducted using odds ratios and unpaired  $t$ -tests.

For the equalities impact assessment, summary statistics describing the demographics of those using mental health services in the catchment area of each unit during a recent 1-year period were compared to the demographics of the primary cohort.  $Z$ -tests were used to compare the proportions of each demographic (where numbers were sufficient) – this is a valid way of comparing a subgroup drawn from a population to a wider population. Correction for the overlap was not needed as each subgroup represented less than 10% of the whole population (Hayes & Berry, 2006).

## Results

### Population accessing hospital-based short-stay crisis units

In total, 1,176 people were included in the primary cohort, 277 (23.55%) in South West London, 308 (26.19%) in Lincolnshire, 387 (32.91%) in Birmingham and 204 (17.35%) in Sheffield, as shown in Table 2. Six hundred and nine participants (51.79%) were female, with an average age of 36 years. The predominant ethnicity was White British, which describes 821 (69.81%) of the cohort. Between 55% and 82% of service users at each site had no primary diagnosis. Most people were living in mainstream housing, but not with a partner, and had an occupation. There were no data recorded about sexuality for the majority of the cohort (842 people; 71.60% of participants). A sizeable portion of people were relatively new to mental health services; 372 (38.27%) had their first contact with mental health services in the 9 months preceding their first crisis unit visit (based on data from 3 sites; 972 participants). The English indices of deprivation 2019 (McLennan et al., 2019) mean scores ranged from 3.45 (2.58) to 5.78 (2.34) at each site (lower scores indicate higher deprivation). The large standard deviations indicate substantial variation within sites in the disadvantage of the people in the relatively small local areas accessing the unit. Comprehensive data on participant characteristics by site are provided in Table 2.

The ED was the largest single source of referrals (488; 41.50%), followed by the Crisis Resolution and Home Treatment Team, who provide intensive home-based therapies as an alternative to inpatient admission (246; 20.92%). At one site, the majority of participants (157; 76.96%) accessed the crisis unit via single point of access – a single route into all crisis services at that Trust.

### Changes to service use following the first visit to a short-stay crisis unit

Typically, there was a marked increase in the use of mental health services following a first visit to a PDU. There were more informal (voluntary) admissions in the 'post' period, a difference which reached significance at all but one site (see Table 3). ED presentations were significantly reduced at two sites and unchanged at a further two sites. There were significant increases in both compulsory admissions and inpatient admissions at all sites. At three of the four sites the number of individuals experiencing a short (0–5 days) admission increased, this was significant at two sites. At all four sites there was a very substantial increase of at least 8 days in the average length of an inpatient stay. Due to relatively small numbers admitted to wards combined with a huge variation in the length of these stays (indicated by the very large  $SD$ ), this increase was

**Table 2.** Primary cohort – participant characteristics by site.

	South West London	Lincolnshire	Birmingham	Sheffield
	<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)
Sample size	277	308	387	204
N female (%)	162 (58.48)	167 (54.25)	171 (44.19)	109 (54.43)
Age (mean, SD)	36.74 (13.77)	37.43 (15.22)	34.64 (13.53)	37.63 (13.46)
Ethnicity				
Asian/British Asian	19 (6.86)	0	36 (9.30)	10 (4.90)
Black/Black British	24 (8.66)	3 (0.97)	27 (6.98)	11 (5.39)
White British	163 (58.84)	265 (86.04)	257 (66.41)	136 (66.67)
White Irish/White other	40 (14.44)	9 (2.92)	15 (3.88)	5 (2.45)
Other	26 (9.39)	0	8 (2.07)	7 (3.43)
Mixed	1 (0.36)	1 (0.32)	18 (4.65)	2 (0.98)
Missing	4 (1.44)	30 (9.74)	26 (6.72)	33 (16.18)
Primary diagnosis				
F1: Psychoactive substance use	12 (4.33)	10 (3.25)	4 (1.04)	0
F2: Schizophrenia	8 (2.89)	16 (5.19)	14 (3.62)	15 (7.35)
F3: Mood disorders	39 (14.08)	13 (4.22)	21 (5.43)	2 (0.98)
F4: Neurotic disorders	22 (7.94)	13 (4.22)	11 (2.84)	1 (0.49)
F6: Personality disorders	21 (7.58)	14 (4.55)	10 (2.58)	15 (7.35)
F8: Disorders of psych development	3 (1.08)	0	0	0
F9: Behavioural and emotional disorders	1 (0.36)	0	3 (0.78)	0
Other	19 (6.86)	10 (3.25)	5 (1.29)	3 (1.47)
Not stated/no diagnosis	152 (54.87)	232 (75.32)	319 (82.43)	168 (82.35)
Secondary diagnosis				
Mental health	81 (29.24)	–	11 (2.84)	–
Alcohol or substance abuse	17 (6.14)	–	8 (2.07)	–
Physical health	7 (2.53)	–	4 (1.03)	–
Other	12 (4.33)	–	1 (0.26)	–
No secondary diagnosis	160 (57.76)	–	363 (93.80)	–
Marital status				
Single/separated/divorced/widowed	224 (80.9)	163 (52.92)	143 (36.96)	116 (56.86)
Married/civil partnership/cohabiting	39 (14.1)	39 (12.66)	26 (6.72)	36 (17.65)
Missing	14 (5.1)	106 (42)	218 (56.33)	52 (25.49)
Employment status				
Working/student/volunteer/homemaker	90 (32.5)	92 (29.87)	32 (8.27)	65 (31.86)
Unemployed	114 (41.2)	133 (43.18)	60 (15.50)	85 (41.67)
Retired	5 (1.8)	9 (2.92)	1 (0.26)	5 (2.45)
Missing	68 (24.6)	74 (24.03)	294 (75.97)	49 (24.02)
Housing				
Mainstream housing	192 (69.3)	200 (64.94)	74 (19.12)	130 (63.73)
Supported housing	6 (2.2)	12 (3.90)	9 (2.33)	8 (3.92)
Insecure housing/homeless	12 (4.3)	23 (7.47)	8 (2.07)	9 (4.41)
Missing	67 (24.2)	73 (23.70)	296 (76.49)	57 (27.94)
Sexual orientation				
Gay/lesbian/bi	4 (1.4)	13 (4.22)	3 (0.78)	11 (5.39)
Heterosexual	39 (14.1)	143 (46.43)	58 (14.99)	63 (30.88)
Missing	234 (84.5)	152 (49.35)	326 (84.24)	130 (63.73)
Referral source				
ED	123 (44.4)	113 (36.69)	237 (61.24)	15 (7.35)
CMHT	31 (11.2)	27 (8.77)	0	–
Home treatment team	107 (38.6)	135 (43.83)	4 (1.03)	–
Local authority and other provider	–	8 (2.60)	0	–
Primary care and other medical	12 (4.3)	6 (1.95)	137 (35.40)	30 (14.71)
Section 136 and Police	4 (1.4)	19 (6.17)	6 (1.55)	1 (0.49)

(Continued)

**Table 2.** (Continued)

	South West London	Lincolnshire	Birmingham	Sheffield
	<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)
Other source within the trust	0	0	0	157 (76.96)
Missing	0	0	3 (0.78)	1 (0.49)
Admission to crisis unit within 7 days of first contact with trust	63 (22.74)	62 (20.13)	110 (28.42)	–
First contact with trust during the ‘pre’ period	114 (41.16)	97 (31.49)	161 (41.60)	–
English indices of deprivation 2019 mean (SD) score	18,360 (7,282)	12,990 (8,475)	7,240 (7,679)	10,245 (8,450)
Category of social deprivation; mean (SD)	5.78 (2.34)	4.40 (2.60)	2.65 (2.34)	3.45 (2.58)
Category of social deprivation range	2–10	1–10	1–10	1–10

Note. – = data not available.

significant at only one site – where the mean increase in duration was over 30 days.

At every site the number of contacts the cohort had with community services was significantly increased in the ‘post’ period. Broken down into face-to-face and remote (telephone/video) appointments, the number of remote appointments significantly increased at all sites – and despite the ‘post’ period overlapping with the COVID-19 pandemic, the number of face-to-face appointments also significantly increased at two sites. Contacts with the community mental health team (CMHT) approximately doubled at three sites, and significantly increased at all sites. Contact with the crisis resolution/home treatment team was similarly increased. Use of the single point of access operating at a single trust also significantly increased. Use of specialised services, which typically increased, is summarised towards the lower part of Table 3. Tests for significance were not used on these data due to the relatively low counts. Following their visit to the unit, participants were typically discharged to their usual place of residence (540; 45.92% participants). A smaller proportion were discharged into the care of the Crisis and Home Treatment Team (156; 16.05%), and 12.76% (124 participants) were discharged to a psychiatric inpatient ward. Across all four sites, social-distancing measures as a result of the COVID-19 pandemic were in place for just over 50% of the follow-up period.

### Effect of the COVID-19 pandemic on service use

Data describing the post-period of the primary and pre-COVID cohorts were compared to examine whether the COVID-19 pandemic had an effect on service use (see Table 4 for comparisons). The pre-COVID cohort is summarised in the Supplemental Materials. In general, service use was unchanged during the pandemic, with a few exceptions. In Lincolnshire a significantly higher proportion of people experienced an involuntary inpatient admission in the pre-COVID cohort ( $OR = 1.49, p < .05$ ); this is

also reflected as a higher proportion of people experiencing any type of admission in the pre-COVID cohort at this site ( $OR = 1.77, p < .01$ ). The proportion of face-to-face appointments at two sites (London and Birmingham) were significantly reduced in the primary cohort ( $t = 2.51, p = .01$  and  $t = 3.62, p < 0.001$  at each site, respectively). Similarly, appointments with the CRHT decreased significantly in both London ( $t = 4.15, p < .001$ ) and Birmingham ( $t = 50.03, p < .001$ ). There was no evidence of difference for any of the other outcomes.

### Equalities impact assessment

To assess equality of access to the units, demographic characteristics of the primary cohort and a recent sample of all people using mental health services at the local mental health Trust were compared (Table 5). Across all sites there is a consistent trend that the population accessing the units was significantly younger (by, on average, 5–13 years) than the population of service users at the Trust. At one city site, first-time visitors to the PDU were significantly less likely to be female compared to the Trust-wide population (171/387 [44.19%] compared to 37,669/69,513 [54.19%];  $z = 2.62, p < .01$ ) and significantly more likely to be White British (257 [66.41%] compared to the Trust population 41,019 [59.01%],  $z = 2.41, p = .02$ ). At the rural site the population was predominately White British, both at the Trust and cohort levels. At a second city site, service users accessing the crisis unit were more likely to be Black than the Trust population (11/204 [5.39%] compared to 828/31,190 [2.65%];  $z = 2.06, p = .04$ ), although the small number of Black individuals in the cohort (11) make it difficult to draw statistically reliable conclusions.

### Discussion

This study aimed to understand who uses short-stay psychiatric units, assess equality of access to the units and examine change in service use following a service user’s first visit to a crisis unit (termed PDU in the UK). We also

**Table 3. Pre-post comparisons (primary cohort).**

	South West London			Lincolnshire			Birmingham			Sheffield		
	Pre	Post	Pre-post comparison	Pre	Post	Pre-post comparison	Pre	Post	Pre-post comparison	Pre	Post	Pre-post comparison
<b>Primary outcomes</b>												
Mental health: <i>n</i> experiencing an informal admission	2	54	$\chi^2=48.29, p<.001$	11	51	$\chi^2=30.77, p<.001$	13	64	$\chi^2=36.63, p<.001$	9	15	$\chi^2=1.5, p=.31$
Acute trust ED presentations (mean, SD)	1.33 (1.30)	0.67 (1.86)	$t=5.98, p<.001$	0.68 (0.85)	0.45 (1.00)	$t=3.30, p<.001$	1.64 (1.45)	1.73 (2.20)	$t=0.98, p=.33$	1.82 (2.11)	1.70 (5.42)	$t=0.32, p=.40$
<b>Secondary outcomes: binary events data (number experiencing one or more)</b>												
Inpatient admissions	6	64	$\chi^2=49.47, p<.001$	17	78	$\chi^2=52.41, p<.001$	26	95	$\chi^2=50.12, p<.001$	24	41	$\chi^2=6.42, p=.01$
0–5 days inpatient admissions	1	26	$\chi^2=23.15, p<.001$	4	11	$\chi^2=3.77, p=.05$	3	11	$\chi^2=4.57, p=.06$	8	4	$\chi^2=1.33, p=.39$
Compulsory admissions	4	21	$\chi^2=11.56, p<.001$	8	31	$\chi^2=19.59, p<.001$	13	37	$\chi^2=15.16, p<.001$	20	37	$\chi^2=7.41, p<.01$
<b>Continuous outcomes, <i>n</i> (SD)</b>												
Length of non-null inpatient stays (days; mean, SD) <sup>a</sup>	15.00 (16.10), 6	23.35 (26.71), 64	$t=0.75, p=.46$	26.11 (21.65), 17	35.77 (41.31), 78	$t=-0.93, p=.35$	30.04 (39.96), 26	41.75 (60.88), 95	$t=0.93, p=.36$	19.63 (14.11), 24	54.61 (57.73), 41	$t=4.03, p<.001$
<b>Use of CMHT and other trust services</b>												
Community contacts with services (mean, SD)	12.47 (19.75)	23.31 (27.42)	$t=7.78, p<.001$	12.03 (18.29)	27.38 (31.37)	$t=9.76, p<.001$	3.09 (7.51)	5.52 (11.82)	$t=5.38, p<.001$	25.45 (38.28)	36.60 (48.38)	$t=3.71, p<.001$
Contact type: face to face (mean, SD)	8.79 (13.42)	13.62 (16.61)	$t=5.27, p<.001$	7.71 (12.25)	12.07 (12.45)	$t=5.98, p<.001$	2.58 (6.64)	2.45 (6.91)	$t=0.39, p=.70$	16.37 (26.06)	19.43 (25.46)	$t=1.59, p=.11$
Contact type: telephone/video (mean, SD)	3.69 (8.10)	9.69 (13.68)	$t=9.23, p<.001$	4.32 (8.46)	15.31 (22.49)	$t=9.77, p<.001$	0.51 (1.63)	3.06 (7.91)	$t=6.90, p<.001$	9.08 (15.05)	17.17 (26.71)	$t=5.16, p<.001$
<b>Use of CMHT and other trust services by category; continuous data; mean (SD)</b>												
CMHT	3.21 (8.58)	7.50 (13.30)	$t=6.36, p<.001$	4.52 (13.12)	9.75 (22.91)	$t=5.41, p<.001$	3.16 (7.72)	5.76 (12.29)	$t=5.70, p<.001$	15.65 (29.50)	21.00 (36.97)	$t=2.56, p<.01$
Crisis resolution or home treatment team	4.34 (9.82)	9.18 (12.44)	$t=5.74, p<.001$	5.13 (8.75)	13.84 (17.64)	$t=8.29, p<.001$	5.22 (14.79)	8.64 (18.38)	$t=5.51, p<.005$	8.62 (18.87)	13.73 (23.17)	$t=2.62, p<.01$
Mental health telephone support line	0.42 (3.35)	0.72 (5.90)	$t=1.78, p=.76$	–	–	–	–	–	–	–	–	–
Single point of access	–	–	–	–	–	–	–	–	–	–	–	–
<b>Number of contacts (for small numbers), <i>n</i></b>												
Mental health support line (also summarised above)	117	199	–	–	–	–	–	–	–	–	–	–
Dialectic behaviour therapy service	0	52	–	–	–	4	4	4	–	78	32	–
Service contracted by trust	32	80	–	–	–	0	0	0	–	–	–	–
Art therapies	8	52	–	–	–	0	40	–	–	4	79	–
Other	71	97	–	4	26	–	–	–	–	35	47	–
Outpatient eating disorder service	67	54	–	81	72	6	6	3	–	–	–	–
Exercise therapy team	11	59	–	–	–	–	–	–	–	–	–	–
Personality disorder (PD)/PD intensive team	0	8	–	–	–	–	–	–	–	–	–	–
Complex needs service	221	258	–	0	1	109	109	91	–	–	–	–
Family therapy	7	21	–	–	–	–	–	–	–	–	–	–
Early intervention in psychosis team	153	407	–	168	369	93	93	41	–	311	754	–
Learning disability service	2	80	–	3	14	–	–	–	–	0	0	–
Perinatal community team	9	35	–	61	108	252	252	166	–	29	0	–
Traumatic stress service	45	63	–	–	–	–	–	–	–	0	0	–
Older peoples service (including but not limited to dementia services)	3	44	–	55	245	–	–	–	–	122	316	–

(Continued)

Table 3. (Continued)

	South West London			Lincolnshire			Birmingham			Sheffield		
	Pre	Post	Pre-post comparison	Pre	Post	Pre-post comparison	Pre	Post	Pre-post comparison	Pre	Post	Pre-post comparison
Forensic/criminal justice liaison service	-	-	-	53	3	-	142	85	-	-	-	-
Individual placement and support (for work)	-	-	-	38	57	-	-	-	-	-	-	-
SARC (sexual assault referral centre)	-	-	-	10	38	-	-	-	-	-	-	-
Short-stay crisis unit visits (people experiencing one or more): n	0	31	-	-	54	-	-	54	-	-	20	-
Days in the pandemic; mean (SD)	-	163.15 (54.60)	-	-	157.25 (53.92)	-	-	170.99 (52.72)	-	-	158.62 (52.48)	-
Summary of discharge destinations												
Inpatient – at trust or out of area		7 (2.5)			-			-			28 (13.73)	
Inpatient at trust		-			31 (10.06)			53 (13.70)			-	
Inpatient out of area/non-NHS		-			19 (6.17)			14 (3.62)			-	
Home treatment team		47 (17.0)			109 (35.59)			-			-	
Community mental health team		9 (3.3)			1 (0.32)			-			-	
Primary care		23 (8.3)			0			-			-	
Temporary place of residence		36 (13.0)			0			31 (8.01)			28 (13.73)	
Usual place of residence		142 (51.3)			34 (11.04)			228 (89.91)			136 (66.67)	
Other		-			28 (9.09)			20 (5.17)			4 (1.96)	
Ongoing care – not specified		-			57 (18.51)			29 (7.42)			-	
Missing		13 (4.7)			29 (9.42)			12 (3.10)			8 (3.92)	

\*Unpaired *t*-tests were used for this outcome.

**Table 4.** Comparison of post periods for the primary and pre-COVID cohorts.

	South West London			Lincolnshire			Birmingham		
	Primary cohort n=277	Pre-COVID cohort n=218	Gross cohort comparison	Primary cohort n=308	Pre-COVID cohort n=289	Gross cohort comparison	Primary cohort n=387	Pre-COVID cohort n=427	Gross cohort comparison
<b>Primary outcomes</b>									
Mental health trust: number experiencing an informal admission	54	57	OR=1.46, p=.08	51	75	OR=1.49, p<.05	64	81	OR=1.18, p=.37
Acute trust: ED presentations (mean, SD)	0.67 (1.86)	0.80 (1.55)	t=0.81, p=.42	0.45 (1.00)	0.43 (0.93)	t=0.24, p=.81	1.73 (2.20)	1.83 (2.45)	t=0.57, p=.57
<b>Secondary outcomes</b>									
Inpatient admissions	64	63	OR=1.35, p=.14	78	97	OR=1.77, p<.01	95	110	OR=1.07, p=.69
0-5 days inpatient admissions	26	14	OR=0.66, p=.23	11	17	OR=1.69, p=.19	11	8	OR=0.65, p=.36
Compulsory admissions	21	12	OR=0.71, p=.36	31	31	OR=1.08, p=.78	37	36	OR=0.87, p=.57
Length of non-nil inpatient stays (days; mean, SD)	23.35 (26.71), 64	30.49 (42.35), 63	t=1.14, p=.26	35.77 (41.31), 78	46.44 (61.92), 97	t=1.31, p=.19	41.75 (60.88), 95	49.82 (79.73), 110	t=0.85, p=.40
Community contacts with services (mean, SD)	23.31 (27.42)	26.98 (30.38)	t=1.41, p=.16	27.38 (31.37)	21.94 (24.33)	t=2.36, p<.05	5.52 (11.82)	4.98 (8.38)	t=0.76, p=.40
Contact type: face to face (mean, SD)	13.62 (16.61)	17.60 (18.67)	t=2.51, p=.01	12.07 (12.45)	13.98 (15.68)	t=1.65, p=.10	2.45 (6.91)	4.28 (7.43)	t=3.62, p<.001
Contact type: telephone/video (mean, SD)	9.69 (13.68)	9.38 (14.37)	t=0.24, p=.81	15.31 (22.49)	7.96 (11.59)	t=4.97, p<.001	3.06 (7.91)	0.69 (1.94)	t=6.00, p<.001
Community mental health team Crisis resolution or home treatment team; CRHT	7.50 (13.30)	6.28 (10.27)	t=1.11, p=.27	9.75 (22.91)	7.01 (13.79)	t=1.76, p=.08	5.76 (12.29)	5.30 (8.84)	t=1.78, p=.08
Mental health telephone support line	9.18 (12.44)	14.83 (17.74)	t=4.15, p<.001	13.84 (17.64)	12.42 (14.83)	t=1.06, p=.29	8.64 (18.38)	12.31 (23.46)	t=50.03, p<.001
	0.72 (5.90)	0.71 (5.11)	t=0.01, p=.99	-	-	-	-	-	-

**Table 5.** Equalities impact.

	South West London			Lincolnshire			Birmingham			Sheffield		
	Primary cohort	Trust population	Comparison	Primary cohort	Trust population	Comparison	Primary cohort	Trust population	Comparison	Primary cohort	Trust population	Comparison
Sample or population size	277	17,982	–	308	59,824	–	387	69,513	–	204	31,190	–
N female (%)	162 (58.48)	10,362 (57.62)	z=0.22, p=.84	167 (54.25)	36,150 (60.49)	z=1.65, p=.10	171 (44.19)	37,669 (54.19)	z=2.62, p<.01	109 (54.43)	15,978 (51.23)	z=0.67, p=.50
Age mean (SD)	36.74 (13.77)	48.12 (21.53)	t=13.75, p<.001	37.43 (15.22)	43.75 (24.65)	t=7.29, p<.001	34.64 (13.53)	47.66 (21.71)	t=19.17, p<.001	37.63 (13.46)	48.29	t=11.31, p<.001
Asian /British	19 (6.86)	1,452 (8.07)	z=0.19, p=.85	0 (0.0)	270 (0.45)	–	36 (9.30)	9,572 (13.77)	z=0.78, p=.44	10 (4.90)	913 (2.93)	z=0.37, p=.71
Black/Black British n (%)	24 (8.66)	1,895 (10.54)	z=0.23, p=.76	3 (0.97)	171 (0.29)	z=0.219, p=.83	27 (6.98)	4,284 (6.16)	z=0.18, p=.86	11 (5.39)	828 (2.65)	z=2.06, p=.04
White British n (%)	163 (58.84)	9,603 (53.40)	z=1.39, p=.16	265 (86.04)	51,338 (85.82)	z=0.10, p=.92	257 (66.41)	41,019 (59.01)	z=2.41, p=.02	136 (66.67)	18,574 (59.55)	z=1.67, p=.10
White Irish/White other n (%)	40 (14.44)	2,306 (12.82)	z=0.31, p=.76	9 (2.92)	1,853 (3.10)	z=0.03, p=.98	15 (3.88)	2,551 (3.67)	z=0.045, p=.96	5 (2.45)	554 (1.78)	z=0.11, p=.91
Other n (%)	26 (9.39)	1,458 (8.11)	z=0.24, p=.81	0	147 (0.25)	–	8 (2.07)	1,876 (2.70)	z=0.11, p=.91	7 (3.43)	519 (1.66)	z=0.37, p=.71
Mixed n (%)	1 (0.36)	755 (4.31)	–	1 (0.32)	612 (1.02)	–	18 (4.65)	2,347 (3.38)	z=0.30, p=.77	2 (0.98)	582 (1.87)	z=0.09, p=.93
Missing n (%)	4 (1.44)	493 (2.74)	z=0.16, p=.87	30 (9.74)	5,433 (9.08)	z=0.13, p=.90	26 (6.72)	7,864 (11.31)	z=0.74, p=.46	33 (16.18)	9,220 (29.56)	z=1.68, p=.09

aimed to assess the effects of the COVID-19 pandemic on service use post-discharge from a PDU. The population accessing PDUs was consistently younger across all sites than a recent sample of all people using mental health services at each site. In one site, there were significantly fewer female and White British first-time visitors to the unit than one would expect compared to the wider Trust population. In another site, more service users using the unit for the first time were Black compared to the Trust population, although this result was based on small numbers.

Inpatient admissions increased in the period following first visit to a PDU (compared to a similar period pre-first visit) at all sites, including increases in voluntary and compulsory admissions. At three of four sites the number of individuals experiencing a short (0–5 days) admission increased. At all four sites there was an increase in length of inpatient stay. There was a substantial increase in use of community mental health services at all sites, including community-based crisis teams. A wider range of specialist mental health services were accessed by service users in the ‘post’ period compared to the ‘pre’ period. A reduction in ED presentations in the period post-first visit to PDU was observed at two sites, but was unchanged at the other sites.

Overall, the effect of the pandemic on service use post-first visit to a PDU was minimal, especially in terms of inpatient admissions. There was some effect on types of contacts, with fewer face-to-face community appointments and fewer appointments with crisis resolution and home treatment teams in two sites during the pandemic, while in two sites there were increases in remote contacts (face-to-face contacts also increased in two sites).

### Comparison with the literature

People accessing crisis services in mental health Trusts in England are typically younger than the mean age of all people using mental health services we observed in our study (Lloyd-Evans et al., 2020), but our cohort of first time PDU visitors was younger still. We note that a relatively high proportion of our participants had only recently accessed any mental health care. As such it is possible that PDUs offer a first access point to mental health care for some people, and as such units may benefit from being oriented towards working with a younger demographic. An established literature is indicative of inequalities in care pathways in mental health services for people from some racialised communities (Bhui et al., 2003), including among young people accessing mental health care (Chui et al., 2021). There were indications at some sites that people visiting a PDU for the first time were either less likely to be White British, or more likely to be Black, but this was not consistent across sites and based on a small amount of data. Further research is needed to establish whether

PDUs might improve access to mental health care for people who otherwise experience inequalities of access.

We note that many short-stay crisis units internationally are established with the intention of reducing psychiatric inpatient admissions, with evaluations indicating an overall drop in admissions at a service level following opening of the units (Gillig et al., 1989; Lester et al., 2018; Stamy et al., 2021). However, at the individual patient level, our study indicated a general increase in admissions post-first visit to a PDU. Again, we note that for many of our participants this was an early contact with mental health services and so units might function for some as an entry to appropriate care. Some of the units we evaluated also had an explicit assessment and signposting function (Goldsmith et al., 2021; Trethewey et al., 2019), and so onward referral to either community or inpatient care might be appropriate in some cases. The fact that inpatient stays were longer post-first visit to PDU may be indicative of a higher proportion of planned admissions, rather than very brief admissions than can be typical of crisis and for which evidence of beneficial effect is limited (Clibbens et al., 2018). Increased access to community mental health services post-first visit to a PDU was also a potentially positive outcome as lack of engagement with community care has been shown to be a risk factor for admission (Karasch et al., 2020; Walker et al., 2019), while crisis and home treatment teams have had some success in reducing admissions (Stulz et al., 2020), especially where quality of care is high (Lloyd-Evans et al., 2020). Furthermore, in England as elsewhere, some people experience multiple, repeat mental health presentations to ED (Care Quality Commission, 2014; Okorie et al., 2011). More research is needed to establish whether the increase in mental health service use post-first visit to ED we observed is evidence that that group of people are successfully being signposted, from ED, via a short stay crisis unit to appropriate mental health care. We did see some evidence of a reduction in ED presentations in our cohort after the first PDU visit. This appears promising, although this result is not replicated in findings elsewhere (Anderson et al., 2022), and it is worth noting that a large proportion of our participants were referred to the PDU from ED and so this might be an artefact of the research.

### *Short stay crisis units and the COVID-19 pandemic*

The psychological challenges of the COVID-19 pandemic and social distancing measures were widely considered to present huge psychological challenges (Gruber et al., 2021) and population-level mental health in the UK deteriorated during the first wave of the pandemic (Jia et al., 2020; Pierce et al., 2020). In the early phase of the pandemic, immediate service challenges related to controlling infection in inpatient settings and establishing remote working in the community (Johnson et al., 2021). Pooled

data pertaining to the European COVID-19 first wave peak (April 2020) found psychiatrists treated half as many patients in outpatient settings compared to usual (Rojnic Kuzman et al., 2021). As such our finding that there was little overall change in both inpatient and community service use between our primary and pre-pandemic cohorts is interesting (the switch from face-to-face to remote contacts in some sites notwithstanding), suggesting perhaps that crisis care was prioritised and suffered less from the pandemic than routine care (Johnson et al., 2021). Other research has indicated that some people with pre-existing mental health conditions were able to adapt to challenges posed by the pandemic in part by seeking alternative sources of help (Shah et al., 2022). It is possible that PDUs were able to signpost people to a range of helpful community-based support, as suggested by our data.

### *Strengths and limitations*

We note this is the only historically controlled study examining the impact of visiting a short-stay crisis unit on individual service use. Other evaluations typically compare organisational-level outcomes between those who access a crisis unit and those who access other forms of crisis and acute care (Gillig et al., 1989; Kealy-Bateman et al., 2019; Mok & Walker, 1995; Schneider & Ross, 1996; Trethewey et al., 2019). As such, the present paper makes a valuable contribution to understanding the potential impact on individual crisis care pathways, although it may be that studies which take this historical approach compare a period of crisis with a period of relative stability, and are therefore not identifying a true casual effect of the PDU. A strength of the study is that all eligible service users are included in the sample, reducing any potential selection bias and enhancing generalisability and applicability. Use of routinely collected data (e.g. number of appointments and hospital admissions) is another strength where these are comprehensively recorded, although substantial missing data for some important demographic variables including ethnicity and sexual orientation was a limitation. Our analysis of the effect of the COVID-19 pandemic using a pre-post design controls for differences between cohorts as a result of seasonality. However, we cannot be certain that any differences in outcomes between the primary and pre-COVID cohorts are solely related to the pandemic.

### **Conclusions**

Our study suggests that short-stay crisis units might help people in mental health crisis, visiting a crisis unit for the first time, to access a wide range of inpatient and community mental health care. This may especially be the case for people who are new to mental health services and for younger people using mental health services. As such, short-stay crisis units may make a useful contribution to mental health crisis care pathways.

## Acknowledgements

We would like to thank the St George's, University of London Peer Expertise in Education & Research (PEER) group for their contribution to the design of this project and the DECISION lived experience group for coproducing with us.

## Conflicts of interest

None.

## Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was funded by the National Institute of Health Research (NIHR) Health Services and Delivery Research (Grant Number 17/49/70); <https://fundingawards.nihr.ac.uk/award/17/49/70>. The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

## Ethics approval and consent to participate

The project was given favourable opinion from the East Midlands Leicester South Research Ethics Committee (19/EM/0226) on 12th August 2019; subsequent amendments were approved and the study was run in accordance with Good Clinical Practice guidelines.

## ORCID iD

Goldsmith LP  <https://orcid.org/0000-0002-6934-1925>

## Data availability statement

The data that support the findings of this study are available from the corresponding author, upon reasonable request.

## Supplemental material

Supplemental material for this article is available online.

## References

- Anderson, K., Goldsmith, L., Lomani, J., Ali, Z., Clarke, G., Crowe, C., Jarman, H., Johnson, S., & McDaid, D. (2022). Short-stay crisis units for mental health service users on crisis care pathways: Systematic review and meta-analysis. *BJPsych Open*, 8(4), E144. <https://doi.org/doi:10.1192/bjo.2022.534>
- Bhui, K., Stansfeld, S., Hull, S., Priebe, S., Mole, F., & Feder, G. (2003). Ethnic variations in pathways to and use of specialist mental health services in the UK: Systematic review. *British Journal of Psychiatry*, 182(2), 105–116. <https://doi.org/10.1192/bjp.182.2.105>
- Braitberg, G., Gerdts, M., Harding, S., Pincus, S., Thompson, M., & Knott, J. (2018). Behavioural assessment unit improves outcomes for patients with complex psychosocial needs. *Emergency Medicine Australasia*, 30(3), 353–358.
- Browne, V., Knott, J., Dakis, J., Fielding, J., Lyle, D., Daniel, C., Bruce, M., & Virtue, E. (2011). Improving the care of mentally ill patients in a tertiary emergency department: Development of a psychiatric assessment and planning unit. *Australasian Psychiatry*, 19(4), 350–353.
- Care Quality Commission. (2014). *Key findings for the national accident and emergency patient survey*. Author.
- Chui, Z., Gazard, B., MacCrimmon, S., Harwood, H., Downs, J., Bakolis, I., Polling, C., Rhead, R., & Hatch, S. L. (2021). Inequalities in referral pathways for young people accessing secondary mental health services in south east London. *European Child & Adolescent Psychiatry*, 30(7), 1113–1128. <https://doi.org/10.1007/s00787-020-01603-7>
- Clibbens, N., Harrop, D., & Blackett, S. (2018). Early discharge in acute mental health: A rapid literature review. *International Journal of Mental Health Nursing*, 27(5), 1305–1325.
- Crisp, N., Smith, G., & Nicholson, K. (2016). Old problems, new solutions—improving acute psychiatric care for adults in England. *The Commission on Acute Adult Psychiatric Care*. [https://www.rcpsych.ac.uk/docs/default-source/improving-care/better-mh-policy/policy/policy-old-problems-new-solutions-caapc-report-england.pdf?sfvrsn=7563102e\\_2](https://www.rcpsych.ac.uk/docs/default-source/improving-care/better-mh-policy/policy/policy-old-problems-new-solutions-caapc-report-england.pdf?sfvrsn=7563102e_2)
- Department of Health and Concordat Signatories. (2014). *Mental health crisis care Concordat – Improving outcomes for people experiencing mental health crisis*. Department of Health. <https://www.bl.uk/collection-items/mental-health-crisis-care-concordat-improving-outcomes-for-people-experiencing-mental-health-crisis>
- Efron, B., & Tibshirani, R. J. (1994). *An introduction to the bootstrap*. CRC Press.
- Fagerland, M. W., Lydersen, S., & Laake, P. (2013). The McNemar test for binary matched-pairs data: Mid-p and asymptotic are better than exact conditional. *BMC Medical Research Methodology*, 13(1), 91. <https://doi.org/10.1186/1471-2288-13-91>
- Fleury, M. J., Grenier, G., Farand, L., & Ferland, F. (2019). Use of emergency rooms for mental health reasons in Quebec: Barriers and facilitators. *Administration and Policy in Mental Health and Mental Health Services Research*, 46, 18–33.
- Gillig, P., Hillard, J. R., Bell, J., Combs, H. E., Martin, C., & Deddens, J. A. (1989). The psychiatric emergency service holding area: Effect on utilization of inpatient resources. *The American Journal of Psychiatry*, 146(3), 369–372. <https://doi.org/10.1176/ajp.146.3.369>
- Goldsmith, L. P. (2020). *Evaluating mental health decision units in acute care pathways (DECISION): A quasi-experimental and health economic evaluation*. ISCTRN Registry. <https://doi.org/10.1186/ISRCTN53431343>
- Goldsmith, L. P., Anderson, K., Clarke, G., Crowe, C., Jarman, H., Johnson, S., Lloyd-Evans, B., Lomani, J., McDaid, D., Park, A.-L., Smith, J. A., Turner, K., & Gillard, S. (2021). The psychiatric decision unit as an emerging model in mental health crisis care: A national survey in England. *International Journal of Mental Health Nursing*, 30(4), 955–962. <https://doi.org/https://doi.org/10.1111/inm.12849>
- Gruber, J., Prinstein, M. J., Clark, L. A., Rottenberg, J., Abramowitz, J. S., Albano, A. M., Aldao, A., Borelli, J. L., Chung, T., & Davila, J. (2021). Mental health and clinical psychological science in the time of COVID-19: Challenges, opportunities, and a call to action. *American Psychologist*, 76(3), 409.

- Hayes, L., & Berry, G. (2006). Comparing the part with the whole: Should overlap be ignored in public health measures? *Journal of Public Health*, 28(3), 278–282. <https://doi.org/doi.org/10.1093/pubmed/fdl038>
- Jia, R., Ayling, K., Chalder, T., Massey, A., Broadbent, E., Coupland, C., & Vedhara, K. (2020). Mental health in the UK during the COVID-19 pandemic: Cross-sectional analyses from a community cohort study. *BMJ Open*, 10(9), e040620. <https://doi.org/10.1136/bmjopen-2020-040620>
- Johnson, S., Dalton-Locke, C., Vera San Juan, N., Foye, U., Oram, S., Papamichail, A., Landau, S., Rowan Olive, R., Jeynes, T., Shah, P., Sheridan Rains, L., Lloyd-Evans, B., Carr, S., Killaspy, H., Gillard, S., Simpson, A., Bell, A., Bentivegna, F., Botham, J., . . . Tzouvara, V. (2021). Impact on mental health care and on mental health service users of the COVID-19 pandemic: A mixed methods survey of UK mental health care staff. *Social Psychiatry and Psychiatric Epidemiology*, 56(1), 25–37. <https://doi.org/10.1007/s00127-020-01927-4>
- Karasch, O., Schmitz-Buhl, M., Mennicken, R., Zielasek, J., & Gouzoulis-Mayfrank, E. (2020). Identification of risk factors for involuntary psychiatric hospitalization: Using environmental socioeconomic data and methods of machine learning to improve prediction. *BMC Psychiatry*, 20(1), 401. <https://doi.org/10.1186/s12888-020-02803-w>
- Kealy-Bateman, W., McDonald, A., Haber, P., Green, T., White, B., Sundakov, V., O’Cionnaith, C., & Glozier, N. (2019). Development of a joint mental-health and drug health assessment unit and short-stay unit. *Australasian Psychiatry*, 27(4), 374–377.
- Lester, N. A., Thompson, L. R., Herget, K., Stephens, J. A., Campo, J. V., Adkins, E. J., Terndrup, T. E., & Moffatt-Bruce, S. (2018). CALM interventions: Behavioral health crisis assessment, linkage, and management improve patient care. *American Journal of Medical Quality*, 33(1), 65–71.
- Lloyd-Evans, B., Osborn, D., Marston, L., Lamb, D., Ambler, G., Hunter, R., Mason, O., Sullivan, S., Henderson, C., Onyett, S., Johnston, E., Morant, N., Nolan, F., Kelly, K., Christoforou, M., Fullarton, K., Forsyth, R., Davidson, M., Piotrowski, J., . . . Johnson, S. (2020). The CORE service improvement programme for mental health crisis resolution teams: Results from a cluster-randomised trial. *British Journal of Psychiatry*, 216(6), 314–322. <https://doi.org/10.1192/bjp.2019.21>
- McCrone, P., Johnson, S., Nolan, F., Pilling, S., Sandor, A., Hault, J., McKenzie, N., Thompson, M., & Bebbington, P. (2009). Economic evaluation of a crisis resolution service: A randomised controlled trial. *Epidemiology and Psychiatric Sciences*, 18(1), 54–58. <https://doi.org/10.1017/s1121189x00001469>
- McLennan, D., Noble, S., Noble, M., Plunkett, E., Wright, G., & Gutacker, N. (2019). *The English indices of deprivation 2019 technical report*. Ministry of Housing, Communities & Local Government.
- Mok, H., & Walker, C. (1995). Brief psychiatric hospitalization: Preliminary experience with an urban short-stay unit. *Canadian Journal of Psychiatry*, 40, 45–47.
- National Institute for Health Research Funding and Awards Award ID 17/49/70. (2019). *Evaluating Mental Health Decision Units in acute care pathways (DECISION): A quasi-experimental and health economic evaluation*. <https://fundingawards.nihr.ac.uk/award/17/49/70>
- NHS England, Public Health England, Health Education England, Monitor, Care Quality Commission, NHS Trust Development Authority. (2014). *Five year forward view*. <https://www.england.nhs.uk/wp-content/uploads/2014/10/5yfv-web.pdf>
- Nicks, B. A., & Manthey, D. M. (2012). The impact of psychiatric patient boarding in emergency departments. *Emergency Medicine International*, 2012, 360308. <https://doi.org/10.1155/2012/360308>
- Okorie, E. F., McDonald, C., & Dineen, B. (2011). Patients repeatedly attending accident and emergency departments seeking psychiatric care. *The Psychiatrist*, 35(2), 60–62. <https://doi.org/10.1192/pb.bp.108.024455>
- Parwani, V., Tinloy, B., Ulrich, A., D’Onofrio, G., Goldenberg, M., Rothenberg, C., Patel, A., & Venkatesh, A. K. (2018). Opening of psychiatric observation unit eases boarding crisis. *Academic Emergency Medicine*, 25, 456–460. <https://doi.org/10.1111/acem.13369>
- Pierce, M., Hope, H., Ford, T., Hatch, S., Hotopf, M., & John, A. (2020). Mental health before and during the COVID-19 pandemic: A longitudinal probability sample survey of the UK population. *The Lancet Psychiatry*, 7(10), 883–892.
- Rojnic Kuzman, M., Vahip, S., Fiorillo, A., Beezhold, J., Pinto da Costa, M., Skugarevsky, O., Dom, G., Pajevic, I., Peles, A. M., Mohr, P., Kleinberg, A., Chkonja, E., Balazs, J., Flannery, W., Mazaliauskiene, R., Chihai, J., Samochowiec, J., Cozman, D., Mihajlovic, G., . . . Goorwod, P. (2021). Mental health services during the first wave of the COVID-19 pandemic in Europe: Results from the EPA Ambassadors Survey and implications for clinical practice. *European Psychiatry*, 64(1), e41. <https://doi.org/10.1192/j.eurpsy.2021.2215>
- Schneider, S., & Ross, I. (1996). Ultra-short hospitalization for severely mentally ill patients. *Psychiatric Services*, 47(2), 137–138.
- Shah, P., Hardy, J., Birken, M., Foye, U., Rowan Olive, R., Nyikavaranda, P., Dare, C., Stefanidou, T., Schlieff, M., Pearce, E., Lyons, N., Machin, K., Jeynes, T., Chipp, B., Chhappia, A., Barber, N., Gillard, S., Pitman, A., Simpson, A., . . . Lloyd-Evans, B. (2022). What has changed in the experiences of people with mental health problems during the COVID-19 pandemic: A coproduced, qualitative interview study. *Social Psychiatry and Psychiatric Epidemiology*, 57(6), 1291–1303. <https://doi.org/10.1007/s00127-022-02254-6>
- Spooren, D., van Heeringen, K., & Jannes, C. (1997). Short-term outcome following referral to a psychiatric emergency service. *Crisis*, 18(2), 80–85.
- Srebnik, D., & Russo, J. (2008). Use of psychiatric advance directives during psychiatric crisis events. *Administration and Policy in Mental Health*, 35(4), 272–282. <https://doi.org/10.1007/s10488-008-0172-0>
- Stamy, C., Shane, D., Kennedy, L., Van Heukelom, P., Mohr, N., Tate, J., Montross, K., & Lee, S. (2021). Economic evaluation of the emergency department after implementation of an Emergency Psychiatric Assessment, Treatment, and Healing Unit. *Academic Emergency Medicine*, 28, 82–91.
- Stulz, N., Nevely, A., Hilpert, M., Bielinski, D., Spisla, C., Maeck, L., & Hepp, U. (2015). Referral to inpatient treatment does not necessarily imply a need for inpatient treatment. *Administration and Policy in Mental Health and Mental Health Services Research*, 42, 474–483.

- The Strategy Unit. (2019). *Exploring mental health inpatient capacity across sustainability and transformation partnerships in England*. Royal College of Psychiatrists Retrieved from <https://www.strategyunitwm.nhs.uk/publications/exploring-mental-health-inpatient-capacity>
- Stulz, N., Wyder, L., Maeck, L., Hilpert, M., Lerzer, H., Zander, E., Kawohl, W., Grosse Holtforth, M., Schnyder, U., & Hepp, U. (2020). Home treatment for acute mental healthcare: Randomised controlled trial. *British Journal of Psychiatry*, 216(6), 323–330. <https://doi.org/10.1192/bjp.2019.31>
- Thibaut, B., Dewa, L. H., Ramtale, S. C., D’Lima, D., Adam, S., Ashrafian, H., Darzi, A., & Archer, S. (2019). Patient safety in inpatient mental health settings: A systematic review. *BMJ Open*, 9(12), e030230. <https://doi.org/10.1136/bmjopen-2019-030230>
- Thomas, K. C., Owino, H., Ansari, S., Adams, L., Cyr, J. M., Gaynes, B. N., & Glickman, S. W. (2018). Patient-Centered values and experiences with emergency department and mental health crisis care. *Administration and Policy in Mental Health and Mental Health Services Research*, 45(4), 611–622. <https://doi.org/10.1007/s10488-018-0849-y>
- Trethewey, S. P., Deepak, S., Saad, S., Hughes, E., & Tadros, G. (2019). Evaluation of the Psychiatric Decisions Unit (PDU): Effect on emergency department presentations and psychiatric inpatient admissions. *Postgraduate Medical Journal*, 95, 6–11. <https://doi.org/10.1136/postgradmedj-2018-135788>
- Van Der Sande, R., Van Rooijen, L., Buskens, E., & Allart, E. (1997). Intensive in-patient and community intervention versus routine care after attempted suicide. *The British Journal of Psychiatry*, 171, 35–41.
- Walker, S., Mackay, E., Barnett, P., Sheridan Rains, L., Leverton, M., Dalton-Locke, C., Trevillion, K., Lloyd-Evans, B., & Johnson, S. (2019). Clinical and social factors associated with increased risk for involuntary psychiatric hospitalisation: A systematic review, meta-analysis, and narrative synthesis. *Lancet Psychiatry*, 6(12), 1039–1053. [https://doi.org/10.1016/s2215-0366\(19\)30406-7](https://doi.org/10.1016/s2215-0366(19)30406-7)