



City Research Online

City, University of London Institutional Repository

Citation: White, S., Bhattacharya, R., Bremner, S., Faulkner, A., Foster, R., Gibson, S., Goldsmith, L., Harnett, D., Patel, A., Priebe, S., et al (2023). Predictors of engagement with peer support: analysis of data from a randomised controlled trial of one-to-one peer support for discharge from inpatient psychiatric care. *International Journal of Social Psychiatry*, 69(4), pp. 994-1003. doi: 10.1177/00207640221148090

This is the published version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/29729/>

Link to published version: <https://doi.org/10.1177/00207640221148090>

Copyright: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

Reuse: Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Predictors of engagement with peer support: analysis of data from a randomised controlled trial of one-to-one peer support for discharge from inpatient psychiatric care



International Journal of
Social Psychiatry
1–10

© The Author(s) 2023



Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/00207640221148090
journals.sagepub.com/home/isp



Sarah White¹ , Rahul Bhattacharya², Stephen Bremner³,
Alison Faulkner⁴, Rhiannon Foster⁵, Sarah Gibson⁴,
Lucy Goldsmith¹ , Daniella Harnett², Akshaykumar Patel⁶,
Stefan Priebe⁶, Julie Repper⁷, Miles Rinaldi⁸, Anthony Salla⁵,
Alan Simpson⁹, Michael Ussher^{1,10} and Steve Gillard⁵

Abstract

Background: A range of evidence for the effectiveness of one-to-one peer support in mental health services is emerging. Levels of engagement with peer support vary with limited studies showing few individual participant characteristics predicting engagement. Implementation factors that might predict engagement have not been considered.

Methods: Data were analysed from the intervention arm of the ENRICH trial of one-to-one peer support for discharge from acute psychiatric inpatient care. Two outcomes were considered: (1) a measure of 'engaged with peer worker'; (2) number of face-to-face contacts with peer worker post-discharge. Two sets of independent variables were analysed against each outcome: (1) pre-randomisation participant characteristics; (2) implementation factors measured pre-discharge. Analyses used logistic and zero-inflated negative binomial regression models according to outcome structure.

Results: Data were analysed for 265 participants randomised to peer support who had a known peer worker. Non-heterosexual participants had increased odds of engaging with peer support compared to heterosexual participants, $OR=4.38$ (95% CI: 1.13, 16.9, $p=.032$). Longer duration of first contact with peer worker ($OR=1.03$, 95% CI: 1.00, 1.04, $p<.001$) and more relationship building activities in the first contact ($OR=1.4$, 95% CI: 1.13, 1.85, $p=.004$) were associated with greater odds of engaging with peer support. Analysis of number of contacts post-discharge showed consistent findings.

Conclusions: Implementation of peer support should include a focus on relationship building in the first session of peer support. The potential for peer support to break down barriers to accessing mental health services experienced by people from marginalised communities warrants further investigation.

Keywords

Peer support, engagement, secondary analysis, mental health services

¹St George's, University of London, UK

²East London NHS Foundation Trust, UK

³Brighton and Sussex Medical School, University of Sussex, UK

⁴Independent Survivor Researcher, London, UK

⁵City, University of London, UK

⁶Queen Mary, University of London, UK

⁷Implementing Recovery through Organisational Change, Nottingham, UK

⁸South West London & St George's Mental Health NHS Trust, UK

⁹Kings' College London, UK

¹⁰University of Stirling, UK

*Miles Rinaldi is also affiliated to 9 Centre for Work and Mental Health, Nordland Hospital trust, Bodø, Norway; Centre for Research and Education in Forensic Psychiatry, Haukeland University Hospital, Bergen, Norway

Corresponding author:

Sarah White, St George's, University of London, Cranmer Terrace, London SW17 0RE, UK.
Email: swhite@sgul.ac.uk

Introduction

Peer support in state funded or regulated mental health services is often delivered one-to-one by peer workers; people with personal experiences of using mental health services, employed and trained to offer support to others receiving care for similar mental health difficulties. Qualitative research has consistently reported positive experiences of the impacts of peer support (Gidugu et al., 2015; Walker & Bryant, 2013). Evidence is mounting for the effectiveness of one-to-one peer support in mental health services, with a recent systematic review indicating that peer support can lead to modest but significant improvements in personal recovery (Standardised Mean Difference (SMD) of 0.22, 95% Confidence Interval (CI) (0.01, 0.42), based on 593 participants from three trials) and empowerment (SMD=0.23, 95% CI 0.04, 0.42, 519 participants from four trials) (White, Foster et al., 2020). However, no significant effects were observed for hospitalisation, psychiatric symptoms, quality of life, satisfaction with services, social functioning, social network support or working alliance, and numbers of studies remain small and of variable quality (White, Foster et al., 2020). This mixed picture lacks clarity for policy makers and as a guide to implementing peer support into mental health services.

Data from recent trials have indicated a range of rates of engagement with peer support. In two trials of peer support for people at discharge from psychiatric hospital, 45.8% (22 of 48) (O'Connell et al., 2018) and 63% (40/63) (Rogers et al., 2016) of participants respectively took up a minimum of one session of peer support. In studies of community-based peer support, 72.4% (160/221) of participants took up at least three of the peer support sessions on offer in one trial (Johnson et al., 2018), with 75.4% (86/114) of participants taking up a least four sessions in another (Mahlke et al., 2017). Average number of sessions attended per participant as a proportion of maximum or target number of sessions on offer also varies, from 15.4% (mean of 6 out of a maximum of 39 on offer) in one of the hospital discharge trials (Rogers et al., 2016), to 46.9% (mean of 12.2/26) (Mahlke et al., 2017) and 70% (median of 7/10) (Johnson et al., 2018) in the community-based trials. With studies not reporting further analysis of these data, this variation alone warrants investigation.

There is limited literature exploring levels of engagement with peer support and outcome. In an analysis of service use data from people attending a peer-based recovery service for people with a diagnosis of substance use disorder, a positive association was observed between number of peer-supported follow-up sessions attended and recovery capital score (Ashford et al., 2021). A per-protocol analysis from a trial of a peer specialist intervention for people with mental illness and substance use disorders indicated that participants with high engagement with peer

support (12 contacts or more) were more likely than control group participants to show positive change in psychiatric symptoms compared to participants with low engagement (Chinman et al., 2018). A small amount of research to date has explored factors that might predict levels of engagement with peer support. A study of peer support among homeless veterans with mental health conditions and substance use histories found that neither participant socio-demographic characteristics nor mental health and substance use severity were predictive of engagement (Ellison et al., 2016). In a randomised controlled trial of peer specialist services, also for homeless veterans, higher hope for the future, shorter period spent homeless prior to being offered peer support, higher severity of self-reported substance use symptoms and higher level of mental health service use were all associated with engagement with peer support (Chinman et al., 2019). These studies did not explore any association between how peer support was delivered and engagement.

Literature on the values and principles underpinning peer support is clear that choice and control over engaging with peer support is core to the distinctive nature of the peer support relationship (Gillard et al., 2017). If people choose not to engage or continue with peer support once they have been offered it, there remains a need to better understand how peer support might be offered in a way which is more engaging and therefore potentially more effective. This echoes a wider mental health literature indicating how strength of therapeutic relationship is predictive of the effectiveness of many mental health therapies and interventions (Goldsmith et al., 2015; Priebe & McCabe, 2008). In addition, it has been argued that the peer-to-peer relationship should also be grounded in shared experiences of marginalisation related to community (including gender, ethnicity, sexual orientation, age and religion), especially where people are excluded from mental health services (Faulkner & Kalathil, 2012; Hope & Ali, 2019). Questions about potential association between outcomes and this 'matching' of peer workers and participants (Pfeiffer et al., 2011) remain unanswered.

Our study aims to address these questions in order to optimise engagement and, potentially, the outcomes of one-to-one peer support in mental health services.

Methods

We conducted a multisite, individually randomised controlled trial (ENRICH) of peer support for discharge from inpatient to community mental health care, with the full methods described elsewhere (Gillard et al., 2020). The trial recruited 590 participants who were currently psychiatric inpatients with a previous recent admission, randomising half ($n=294$) to one-to-one peer support for discharge in addition to care as usual (CAU; follow-up from community mental health services post-discharge),

and half ($n=296$) to CAU only. Participants allocated to peer support were offered at least one face-to-face contact with a designated peer worker while still in hospital, and then 10 weekly contacts followed by three fortnightly contacts post-discharge. Peer workers had all completed manualised training and provided flexible support focused on building individual strengths and connection to community (Gillard et al., 2020; Marks et al., 2021).

Participants offered peer support were no less likely to be readmitted to inpatient care in the year post-discharge – the primary outcome – compared to those in the CAU group (relative risk (RR) 0.97 95% CI 0.82, 1.14; $p=.68$) (Gillard et al., 2022). A Complier Average Causal Effect (CACE) analysis (Dunn et al., 2015) compared participants who had had at least two contacts with their peer worker, at least one of which was in the community post-discharge, with a subgroup of similar participants in the CAU group (identified using all baseline participant data). These criteria were developed with the study's Lived Experience Advisory Panel – people with direct experience of peer support – and reflected the principle of choice in peer support (i.e. choosing to meet away from the restrictive environment of the ward) (Goldsmith et al., 2019). We found that participants who met these criteria were significantly less likely to be readmitted in the year post-discharge than similar participants not offered peer support (RR 0.88 95% CI 0.76, 0.99) (Gillard et al., 2022).

This association between amount of contact with peer support and outcome emphasises the importance of better understanding the process of engagement with peer support. We also found that Black participants who were offered peer support had lower odds of being admitted to hospital compared to control (odds ratio (OR) 0.40 (95% CI 0.17, 0.94)) than were participants of any other ethnicity (OR 1.12 (95% CI 0.77, 1.63)); interaction $p=.031$) (Gillard et al., 2022). While we did not match participants to peer workers by ethnicity, it is of interest to explore whether incidental matching was predictive of engagement. This paper reports two analyses which aim to identify pre-randomisation and pre-discharge predictors of engagement with peer support.

The ENRICH study was approved by the UK National Research Ethics Service, Research Ethics Committee London - London Bridge on 10 May 2016, reference number 16/LO/0470.

Participants

Our sample were all participants in the ENRICH trial randomised to peer support who had a known allocated peer worker.

Data

Two dependent variables were examined: (1) 'engaged with peer worker', defined in the CACE analysis (Gillard

et al., 2022) as having had at least two contacts with the allocated peer worker, one of which was in the community following discharge; (2) number of face-to-face contacts with peer worker post-discharge. Data were attained from structured questionnaires and standardised measures collected at a baseline interview prior to randomisation, electronic patient record extracted by site information management personnel for the 12 months pre-index admission, and a contact log completed by peer workers using an online survey after each attempted contact.

At baseline socio-demographic data for each participant were collected, and standardised assessments made of social inclusion using the Social Outcomes Index (SIX) (Priebe et al., 2008), severity of symptoms using the Brief Psychiatric Rating Scale (BPRS) (Overall & Gorham, 1962), hope for the future using the Herth Hope Index (Herth, 1992), treatment stigma using the stigma subscale of the Barriers to Care Evaluation (BACE-3) (Clement et al., 2012), and internalised stigma using the Questionnaire on Anticipated Discrimination (QUAD) (Gabbidon et al., 2013). The QUAD scale produces two measures of anticipated stigma, severity (mean of all items) and count of life areas where stigma has been anticipated. Number of psychiatric admissions in the year prior to index admission, and type of index admission were extracted from patient records. Using the contact log, peer workers indicated which of a list of activities were covered in each contact. Based on our theoretical model (Gillard et al., 2015), we calculated a variable 'relationship building' as a count of the activities 'using your own lived experience', 'learning from the lived experience of the person you are supporting', 'negotiating (or renegotiating) the boundaries in your relationship', 'accompanying or being alongside the person you are supporting' and 'using active listening skills' recorded as having taken place in the first peer support contact (range of 0–5).

Pre-randomisation predictor variables were; index admission type (compulsory/voluntary), sexual orientation (heterosexual/not heterosexual/declined to answer), gender (male/female), diagnostic group (psychosis/personality disorder/other non-psychotic), ethnicity (Black/Other), age (years), social inclusion (SIX), severity of symptoms (BPRS), hope (HHI), number of admissions in year prior to index admission, treatment stigma (BACE-3), severity of anticipated stigma and count of life areas of stigma (QUAD). Ethnicity data were dichotomised as any Black ethnicity and any other ethnicity because being of Black ethnicity has been shown to be a significant predictor of psychiatric readmission (Halvorsrud et al., 2018) (our primary outcome) and because, as noted above, Black participants offered peer support were significantly less likely to be readmitted compared to control than participants of any other ethnicity (Gillard et al., 2022).

Pre-discharge predictor variables were: same gender, same age (within 18–35, 36+ age groups), same ethnicity (within Asian, Black, White, other ethnic groups) and

same diagnosis (within diagnostic groups as above) as allocated peer worker; relationship building at first contact with peer worker (equal to 0 if no contact with peer worker); length of first contact with peer worker (minutes, equal to 0 if no contact with peer worker); days between recruitment and discharge; whether the participant had two or more peer worker contacts pre-discharge.

Statistical analysis

Logistic regression was used to model the relationship between the two groups of predictor variables and the 'engaged with peer support' dependent variable. Results are reported with odds ratios (OR) and 95% confidence interval. Zero-inflated negative binomial (ZINB) regression was used to model the relationship between the two groups of predictor variables and the number of face-to-face contacts post-discharge. The dependent variable had a high proportion of zero values (zero-inflated) – participants who, for a range of reasons, might have either chosen or been unable to continue to meet their peer worker post-discharge – and a skewed distribution with a few very high values (over-dispersed). ZINB regression models the two parts of the distribution using two linked models; the first estimates the odds of a zero count (zero face-to-face contacts post-discharge) using logistic regression reported with ORs (odds of zero contacts), the second estimates the non-zero counts (number of face-to-face contacts post-discharge) using negative binomial regression reported by incidence rate ratios (IRR), both with 95% bootstrapped percentile confidence limits (BPCL).

A sensitivity analysis was conducted for the pre-randomisation predictors models due to missing data and a high correlation between two of the predictors. Six was removed as this variable had the most missing data, and the count of life areas QUAD variable was removed as the two versions of the scale, mean and count of life areas were highly correlated, $r=.73$. Analysis was conducted using R v4.10.2 (SAMHSA, 2021).

Results

Five hundred ninety participants were randomised into the ENRICH trial of whom 294 were allocated to peer support. Three of these participants withdrew their data. Of the remainder, 265 had a known peer worker. The 26 participants allocated to peer support for whom peer worker was not known were largely at one study site where reporting was incomplete.

Descriptive data

Summary statistics of the sample are reported in Table 1. In brief, the sample had a mean age of 39.7 years ($SD=14.0$)

ranging from 18 to 74. There were similar numbers of male (125, 48.3%) and female (133, 51.4%) participants, one identifying as transgender. Most of the sample were White, 144/257 (56%), 43/257 (16.7%) were Black. Two hundred seventeen participants (81.9%) were heterosexual. The most common diagnosis was psychosis, with 118 participants (44.5%). Approximately half of participants (126, 47.5%) had been admitted compulsorily under the Mental Health Act. In the year prior to index admission participants had a median of one psychiatric admission and 18 community mental health team contacts

The median duration between recruitment and discharge was 17 days (IQR 6–40.5). One hundred one participants (38.1%) had at least two peer worker contacts on the ward. The mean length of the first contact was 42 minutes ranging from 0 to 200 minutes. Variation in the number of relationship-building activities conducted in the first contact is evident. The extent to which participants and peer workers had the same demographic characteristics ranged from 31.7% to 52.8% across characteristics.

Engagement with peer support

Sixty-two percent, 163/265 of participants received peer support as defined as having had at least two peer worker meetings, at least one of which was in the community following discharge. The mean number of face-to-face contacts was 6.2 (median=5.0, $SD=5.3$, IQR=1.0–10.5) and ranged from 0 to 22 contacts. For the 163 who engaged with peer support their mean number of face-to-face contacts was 9.1 (median=9; $SD=4.7$) ranging from 2 to 22. Post-discharge they had a mean of 7.1 face-to-face contacts (median=7; $SD=4.0$; IQR=1–14) ranging from 1 to 15. Ninety-one participants (34.3% of the sample) had zero face-to-face contacts post-discharge. The mean number of face-to-face contacts post-discharge for those with at least one was 6.7 (median=7.0; $SD=4.1$; IQR=3–10).

Pre-randomisation predictors of engagement

The only demographic characteristic significantly predictive of engaging with peer working was sexual orientation (see Table 2), non-heterosexual participants having 4.4 times the odds of engaging with peer support than heterosexual participants, $OR=4.38$ (95% CI: 1.13, 16.9, $p=.032$). There was weaker evidence that sexual orientation was also associated with not having any contacts post-discharge, $OR=0.72$ (95% CI: 0.00, 25.00, $p=.062$), non-heterosexual participants having lower odds of having zero contacts post-discharge than heterosexual participants. Sexual orientation was not associated with the number of contacts post-discharge assuming participants had at least one contact. There was a non-significant finding that Black participants had lower odds of engaging with their

Table 1. Sample descriptive statistics.

| | | n = 265 | n (%) |
|--|---------------------------------------|---------|-------------------|
| Socio-demographic characteristics | | | |
| Age (years) mean (SD) min–max | | 255 | 39.7 (14.0) 18–74 |
| Gender | Female | 259 | 133 (51.4) |
| | Male | | 125 (48.3) |
| | Transgender | | 1 (0.4) |
| Ethnicity | Asian | 257 | 34 (13.2) |
| | Black | | 43 (16.7) |
| | Mixed | | 29 (11.3) |
| | Other | | 7 (2.7) |
| | White | | 144 (56.0) |
| Sexual orientation | Bisexual | 260 | 18 (6.9) |
| | Gay | | 10 (3.8) |
| | Heterosexual | | 217 (83.5) |
| | Lesbian | | 5 (1.9) |
| | Declined to answer | | 10 (3.8) |
| Admission, service use and clinical characteristics | | | |
| Index admission | Voluntary | 248 | 122 (49.2) |
| | Compulsory | | 126 (50.8) |
| | N | | Median (IQR)* |
| Number of psychiatric admissions in year pre-index admission | 265 | | 1 (0–1) |
| Number of CMHT contacts in year pre-index admission | 265 | | 18 (5–35) |
| DNA rate in year pre-index admission | 265 | | 7.4 (0–21.1) |
| Social inclusion, mean (SD) min–max | 219 | | 3.0 (1.4) 0–6 |
| Severity of symptoms, mean (SD) min–max | 248 | | 33.8 (9.5) 9–62 |
| Diagnosis | F20–F29 (psychosis) | 265 | 118 (44.5) |
| | F60 (Specific personality disorder) | | 51 (19.2) |
| | Other eligible non-psychotic disorder | | 96 (36.2) |
| Hope, mean (SD) min–max | 249 | | 33.2 (8.2) 14–48 |
| Treatment stigma, mean (SD) min–max | 257 | | 1.2 (0.8) 0–3 |
| Severity of anticipated stigma, mean (SD) min–max | 260 | | 1.4 (0.6) 0–3 |
| Count of life areas of stigma, mean (SD) min–max | 260 | | 6.0 (3.7) 0–14 |
| Pre-discharge peer worker contact characteristics | | | |
| Two or more contacts on ward pre-discharge | 265 | | 101 (38.1) |
| Peer worker and peer are same gender | 265 | | 140 (52.8) |
| Peer worker and peer in same age group | 265 | | 119 (44.9) |
| Peer worker and peer have same diagnosis group | 265 | | 84 (31.7) |
| Peer worker and peer are same ethnicity | 265 | | 127 (47.9) |
| Relationship building activity in first contact (number of activities) mean (SD) min–max | 265 | | 2.8 (1.6) 0–5 |
| Length of first contact (minutes) mean (SD) min–max | 265 | | 41.9 (30.8) 0–200 |
| Time between enrolment and discharge (days) | 265 | | 17 (6–40.5) |

*Unless otherwise specified.

peer worker, $OR=0.5$ (95% CI: 0.23, 1.22, $p=.134$). There was weak evidence regarding age, $p=.074$, that for those participants who had at least one contact post-discharge that older age was associated with more contacts, $IRR=1.01$ (95% CI: 1.00, 1.02).

The models in Table 2 were refitted without SIX and count of life areas of stigma (QUAD) in line with pre-specified sensitivity analyses. This increased the sample size to 211 participants. Results were consistent with the reported models in Table 2. Little's test was non-significant, $\chi^2=91.9$, $p=.981$, indicating that there was no evidence

that missing data were not missing completely at random and therefore complete-case analysis is appropriate.

Pre-discharge predictors of engagement

Table 3 reports the results of the regression models exploring predictors relevant to the beginning of the peer support relationship. A shorter time between recruitment and discharge ($OR=0.99$, 95% CI: 0.99, 1.00, $p=.002$), a longer duration of first contact with peer worker ($OR=1.03$, 95% CI: 1.00, 1.04, $p<.001$) and more relationship building

Table 2. Pre-randomisation predictors of engagement.

| (n = 181) | | Number of face-to-face contacts post-discharge | | | | | |
|---|---------------------------------------|--|---------|----------------------|---------|-------------------|---------|
| | | Engaged with peer support | | Zero inflation model | | Count model | |
| | | OR (95% CI) | p-Value | OR (95% CI) | p-Value | IRR (95% CI) | p-Value |
| Index admission | Voluntary | 1 | .279 | 1 | .304 | 1 | .353 |
| | Compulsory | 0.68 (0.34, 1.37) | | 1.50 (0.30, 11.43) | | 0.86 (0.58, 1.24) | |
| Sexual orientation | Heterosexual | 1 | | 1 | | 1 | |
| | Not heterosexual | 4.38 (1.13, 16.9) | .032 | 0.72 (0.00, 25.00) | .062 | 0.94 (0.13, 2.25) | .245 |
| | Declined to answer | 1.91 (0.19, 19.61) | .588 | 1.48 (0.40, 12.0) | .794 | 0.79 (0.55, 1.10) | .862 |
| Gender | Female | 1 | .868 | 1 | .306 | 1 | .099 |
| | Male | 0.87 (0.44, 1.72) | | 0.24 (0.00, 1.03) | | 1.25 (0.80, 1.96) | |
| Diagnosis | F20-F29 | 1 | | 1 | | 1 | |
| | F60 (Specific personality disorder) | 0.60 (0.21, 1.70) | .333 | 1.85 (0.20, 106.28) | .294 | 1.19 (0.64, 1.84) | .496 |
| | Other eligible non-psychotic disorder | 0.65 (0.31, 1.38) | .263 | 1.75 (0.70, 48.59) | .182 | 1.13 (0.80, 1.62) | .433 |
| Ethnicity | Other | 1 | .134 | 1 | .309 | 1 | .218 |
| | Black | 0.53 (0.23, 1.22) | | 1.62 (0.16, 10.28) | | 0.76 (0.36, 1.33) | |
| Age (years) | | 1.01 (0.99, 1.03) | .468 | 0.99 (0.92, 1.04) | .655 | 1.01 (1.00, 1.02) | .074 |
| Social inclusion | | 1.160.89, 1.51) | .277 | 0.76 (0.07, 1.11) | .091 | 0.95 (0.78, 1.09) | .403 |
| Severity of symptoms | | 0.98 (0.95, 1.03) | .449 | 1.02 (0.95, 1.15) | .386 | 1.00 (0.98, 1.02) | .593 |
| Hope | | 1.02 (0.97, 1.07) | .551 | 0.98 (0.87, 1.08) | .444 | 1.00 (0.98, 1.03) | .786 |
| Admissions in year prior to index admission | | 0.97 (0.74, 1.26) | .801 | 1.06 (0.69, 3.00) | .669 | 1.02 (0.92, 1.21) | .677 |
| Treatment stigma | | 1.30 (0.77, 2.20) | .326 | 0.68 (0.02, 1.38) | .212 | 0.99 (0.73, 1.25) | .917 |
| Severity of anticipated stigma | | 1.39 (0.40, 4.91) | .607 | 0.67 (0.00, 13.46) | .587 | 0.99 (0.43, 2.22) | .983 |
| Count of life areas of stigma | | 0.94 (0.77, 1.14) | .511 | 1.08 (0.65, 3.26) | .496 | 0.98 (0.84, 1.13) | .629 |

activities in the first contact ($OR=1.4$, 95% CI: 1.13, 1.85, $p=.004$) were all associated with greater odds of ‘engaging with peer support’.

The two parts of the ZINB regression model indicated complementary results. More activities based on relationship building during the first contact and a longer length of first contact were associated with lower odds of having zero contacts post-discharge, $OR=0.59$ (95% BPCI: 0.07, 1.09, $p=.001$) and $OR=0.96$ (95% BPCI: 0.01, 0.98, $p=.001$) respectively. For those participants with at least one contact, having had more days between recruitment and discharge was associated with a lower number of face-to-face contact post-discharge, $IRR=0.99$ (95% BPCI: 0.99, 1.00, $p=.003$).

Discussion

This study aimed to identify whether pre-randomisation (participant characteristic) or pre-discharge variables (peer support characteristics) were predictive of either ongoing engagement with peer support or number of peer support sessions post-discharge. We observed little association

between participant characteristics and engagement with peer support. Participants who identified as non-heterosexual had significantly higher odds of engaging with peer support post-discharge than heterosexual participants. Notwithstanding low numbers of participants in this group, we note a lack of research explicitly exploring mental health peer support among LGBTQ+ people (Faulkner & Kalathil, 2012). Other research has indicated that people who identify as LGBTQ+ are reluctant to disclose their sexual orientation to healthcare providers (including primary care and mental healthcare providers), with evidence suggesting that LGBTQ+ people can continue to feel pathologised because of their sexuality (Bachmann & Gooch, 2018), and as a result are less likely than heterosexual people to access care that meets their needs (Durso & Meyer, 2013; Salway et al., 2018). We also note that Black participants might have been less likely to engage with peer support, while still benefitting more from the intervention than people of other ethnicity (Gillard et al., 2022). This could reflect heterogeneity resulting from grouping together Black ethnicities, but suggests a need to investigate further the experiences and understandings of

Table 3. Pre-discharge predictors of engagement.

| (N = 265) | | Number of face-to-face contacts post-discharge ^a | | | | | |
|---|-----|---|---------|----------------------|---------|-------------------|---------|
| | | Engaged with peer support | | Zero inflation model | | Count model | |
| | | OR (95% CI) | p-Value | OR (95% BPCI) | p-Value | IRR (95% BPCI) | p-Value |
| Two or more contacts on ward pre-discharge | No | 1 | .100 | 1 | .878 | 1 | .074 |
| | Yes | 1.82 (0.89, 3.69) | | 1.08 (0.02, 6.76) | | 1.30 (0.96, 1.72) | |
| Peer worker and peer has same gender | No | 1 | .526 | 1 | .137 | 1 | .948 |
| | Yes | 1.22 (0.66, 2.27) | | 0.54 (0.08, 1.51) | | 0.99 (0.80, 1.25) | |
| Peer worker and peer in same age group | No | 1 | .443 | 1 | .302 | 1 | .764 |
| | Yes | 0.79 (0.43, 1.45) | | 1.51 (0.44, 7.20) | | 0.97 (0.77, 1.21) | |
| Peer worker and peer has same diagnosis group | No | 1 | .287 | 1 | .889 | 1 | .750 |
| | Yes | 1.44 (0.74, 2.79) | | 1.06 (0.29, 4.86) | | 1.04 (0.82, 1.30) | |
| Peer worker and peer has same ethnicity | No | 1 | .611 | 1 | .645 | 1 | .758 |
| | Yes | 0.85 (0.46, 1.58) | | 1.21 (0.45, 43.33) | | 1.03 (0.84, 1.29) | |
| Relationship building activity in first contact (number of activities, 0–5) | | 1.42 (1.13, 1.85) | .004 | 0.60 (0.22, 1.00) | .002 | 0.97 (0.88, 1.09) | .598 |
| Length of first contact (minutes) | | 1.03 (1.02, 1.05) | <.001 | 0.96 (0.63, 0.98) | .001 | 1.00 (1.00, 1.01) | .272 |
| Time between enrolment and discharge (days) | | 0.99 (0.99, 1.00) | .002 | 1.01 (0.98, 1.05) | .163 | 0.99 (0.99, 1.00) | .003 |

^aZINB regression estimates the odds ratio (OR) of a zero count (zero inflation model) using logistic regression, and the incidence rate ratios (IRR) of non-zero counts (count model) using negative binomial regression.

peer support in different cultural contexts. Other research has indicated that peer support offers a different sort of relationship to a typical patient-clinician relationship, enabling disclosure more generally (Gillard et al., 2015). As such our findings raise the possibility that people who experience marginalisation in mental health care might find it easier to engage with, and/or open up to peer workers than to other mental healthcare providers.

Findings from elsewhere that higher baseline hope scores were associated with a greater level of subsequent engagement in peer support (Chinman et al., 2019) were not replicated in our study. Research has indicated a positive association between hope and intention to seek both informal and formal psychological help (McDermott et al., 2017). We note that mean values for hope were lower in the inpatient population in our study than they were for participants in Chinman et al. (2019) community-based study, that relative lack of optimism in our sample perhaps accounting for some lack of enthusiasm for peer support, possibly indicating that the timing of the offer of peer support was challenging for this group of people.

We observed more associations between pre-discharge characteristics of peer support, and engagement with peer support post-discharge. Longer first contact with peer worker was associated with engagement with peer support and, correspondingly, with less likelihood of having no contacts post-discharge. A higher number of relationship-building activities in that first contact was similarly associated with ongoing engagement with peer support, suggesting that a first session of peer support with a clear focus on relationship building may lead to improved

engagement and potentially better outcomes. Our earlier research was indicative of the importance of relationship building as a peer support mechanism (Gillard et al., 2015, 2017). This informed an emphasis on relationship work in peer worker training and in the supervision of peer workers by an experienced peer, as specified in the intervention handbook for the trial (Marks et al., 2021). The importance of role specific training for peer workers has been widely noted (Simpson et al., 2014; Tse et al., 2014).

Interestingly, we found no association between having multiple contacts with the peer worker pre-discharge and ongoing engagement, while a longer duration between recruitment into the study and discharge was associated with lower odds of remaining engaged with peer support and fewer contacts in total post-discharge. This felt counter-intuitive as more opportunities to meet with the peer worker before discharge might be assumed to enhance relationship building. It is possible that this longer length of stay on the ward reflected a frustrating period of delayed discharge, or a prolonged period of acute mental distress for those individuals, with associated challenges to building a relationship with the peer worker.

We also found no association between matching of participant to peer worker (incidental rather than planned in our study) – in terms of either age, gender, ethnicity or diagnostic group – and engagement with peer support. This is despite research suggesting that peer identification in mental health encapsulates wider aspects of shared identity beyond experiences of mental distress (Faulkner & Kalathil, 2012; Hope & Ali, 2019), and studies reporting peer support targeting specific clinical groups (Chien

et al., 2019; Pfeiffer et al., 2011). As such we saw people engaging in peer support ‘across difference’, again raising the possibility that there is something about the quality of the peer-to-peer relationship – perhaps in contrast to the patient-clinician relationship – that encourages ongoing engagement independent of the identities of the individuals involved.

Strengths and limitations

Strengths of this study were our a priori identification of potential predictor variables grounded in empirical and theoretical research in the field, and our ‘engaged in peer support’ criteria we developed with our lived experience advisory group. Confidence in our findings was enhanced by the complementarity of our analyses; pre-discharge predictors of being engaged with peer support were all supported by either the zero-inflation or count model of number of peer support contacts post-discharge.

Missing data in some baseline variables was a potential weakness of the study, although sensitivity analysis indicated that findings remained robust when measures with missing and highly correlated variables were removed. Low numbers of participants in some groups, particularly, for example, participants identifying as lesbian, gay, or bisexual, suggest that some of our estimates should be treated with caution and seen as avenues for further study, rather than definitive findings. We also note that the socio-demographic categorisation we used for sexual orientation and ethnicity were incomplete and limited our analyses. Incomplete contact logs by peer workers at one site also potentially undermined reliability of findings. Finally, this was a quantitative study; qualitative research exploring people’s experiences and views on engaging with peer support would help interpret these findings.

Implications for practice

Given our findings around an association between length and quality of first peer support contact and ongoing engagement with peer support, a focus on relationship building in peer worker training, supervision and guidance for the crucial first session should inform peer support practice in mental health services going forward. However, our findings also suggest that leaving inpatient care might be a difficult time for people to build and sustain a new relationship. The approach to peer support we developed was very flexible, encouraging peer worker and participant to agree how they worked together, choosing from a range of tools (Gillard et al., 2020; Marks et al., 2021). It is possible that people might have benefited from a more structured and explicitly supportive approach, including a more structured first session to support relationship building, introducing flexibility as the relationship developed.

In addition, our findings with respect to sexual orientation, ethnicity and the lack of association between incidental matching and engagement suggest that peer support has the potential to work ‘across difference’ (limitations in our data notwithstanding). These findings raise the possibility that peer support might offer additional benefits to people who may feel marginalised, unheard or discriminated against by mental health services. Mental health services struggle to address inequalities in access, experience and outcomes of care for people from marginalised communities (Halvorsrud et al., 2018), and as such, support and training for peer workers to work with diversity and break down barriers to communication and relationship building in mental health care would seem to offer an opportunity to address, at least in part, historic inequities in care.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was funded by the UK National Institute for Health Research (NIHR), Programme Grants for Applied Research funding programme (grant number RP-PG-1212-20019). This paper presents independent research funded by NIHR. The views expressed are those of the authors and not necessarily those of the UK National Health Service (NHS), the NIHR or the Department of Health and Social Care.

ORCID iD

Sarah White  <https://orcid.org/0000-0003-2468-6193>

Lucy Goldsmith  <https://orcid.org/0000-0002-6934-1925>

References

- Ashford, R. D., Brown, A., Canode, B., Sledd, A., Potter, J. S., & Bergman, B. G. (2021). Peer-based recovery support services delivered at recovery community organizations: Predictors of improvements in individual recovery capital. *Addictive Behaviors*, 119, 106945.
- Bachmann, C. L., & Gooch, B. (2018). *LGBT in Britain: Trans report*. Stonewall.
- Chien, W. T., Clifton, A. V., Zhao, S., & Lui, S. (2019). Peer support for people with schizophrenia or other serious mental illness. *Cochrane Database of Systematic Reviews*, 4(4), CD010880.
- Chinman, M., McCarthy, S., Bachrach, R. L., Mitchell-Miland, C., Schutt, R. K., & Ellison, M. (2018). Investigating the degree of reliable change among persons assigned to receive mental health peer specialist services. *Psychiatric Services*, 69(12), 1238–1244.
- Chinman, M., McCarthy, S., Mitchell-Miland, C., Bachrach, R. L., Schutt, R. K., & Ellison, M. (2019). Predicting engagement with mental health peer specialist services. *Psychiatric Services*, 70(4), 333–336.
- Clement, S., Brohan, E., Jeffery, D., Henderson, C., Hatch, S. L., & Thornicroft, G. (2012). Development and psychometric properties the barriers to access to care evaluation scale (BACE) related to people with mental ill health. *BMC Psychiatry*, 12(1), 1–11.

- Dunn, G., Emsley, R., Liu, H., Landau, S., Green, J., White, I., & Pickles, A. (2015). *Evaluation and validation of social and psychological markers in randomised trials of complex interventions in mental health: A methodological research programme*. Health Technology Assessment.
- Durso, L. E., & Meyer, I. H. (2013). Patterns and predictors of disclosure of sexual orientation to healthcare providers among lesbians, gay men, and bisexuals. *Sexuality Research and Social Policy*, 10(1), 35–42.
- Ellison, M. L., Schutt, R. K., Glickman, M. E., Schultz, M. R., Chinman, M., Jensen, K., Mitchell-Miland, C., Smelson, D., & Eisen, S. (2016). Patterns and predictors of engagement in peer support among homeless veterans with mental health conditions and substance use histories. *Psychiatric Rehabilitation Journal*, 39(3), 266–273.
- Faulkner, A., & Kalathil, J. (2012). *The freedom to be, the chance to dream: preserving user-led peer support*. Together for Mental Wellbeing.
- Gabbidon, J., Brohan, E., Clement, S., Henderson, R. C., & Thornicroft, G. (2013). The development and validation of the questionnaire on anticipated discrimination (QUAD). *BMC Psychiatry*, 13(1), 1–14.
- Gidugu, V., Rogers, E. S., Harrington, S., Maru, M., Johnson, G., Cohee, J., & Hinkel, J. (2015). Individual peer support: A qualitative study of mechanisms of its effectiveness. *Community Mental Health Journal*, 51(4), 445–452.
- Gillard, S., Bremner, S., Foster, R., Gibson, S. L., Goldsmith, L., Healey, A., Lucock, M., Marks, J., Morshead, R., Patel, A., Priebe, S., Repper, J., Rinaldi, M., Roberts, S., Simpson, A., & White, S. (2020). Peer support for discharge from inpatient to community mental health services: Study protocol clinical trial (SPIRIT compliant). *Medicine*, 99(10), e19192.
- Gillard, S., Bremner, S., Patel, A., Goldsmith, L., Marks, J., Foster, R., Morshead, R., White, S., Gibson, S. L., Healey, A., Lucock, M., Patel, S., Repper, J., Rinaldi, M., Simpson, A., Ussher, M., Worner, J., & Priebe, S.; ENRICH Trial Study Group. (2022). Peer support for discharge from inpatient mental health care versus care as usual in England (ENRICH): A parallel, two-group, individually randomised controlled trial. *The Lancet Psychiatry*, 9(2), 125–136.
- Gillard, S., Foster, R., Gibson, S., Goldsmith, L., Marks, J., & White, S. (2017). Describing a principles-based approach to developing and evaluating peer worker roles as peer support moves into mainstream mental health services. *Mental Health and Social Inclusion*, 21, 133–143.
- Gillard, S., Gibson, S. L., Holley, J., & Lucock, M. (2015). Developing a change model for peer worker interventions in mental health services: A qualitative research study. *Epidemiology and Psychiatric Sciences*, 24(5), 435–445.
- Goldsmith, L. P., Lewis, S. W., Dunn, G., & Bentall, R. P. (2015). Psychological treatments for early psychosis can be beneficial or harmful, depending on the therapeutic alliance: An instrumental variable analysis. *Psychological Medicine*, 45(11), 2365–2373.
- Goldsmith, L. P., Morshead, R., McWilliam, C., Forbes, G., Ussher, M., Simpson, A., Lucock, M., & Gillard, S. (2019). Co-producing randomized controlled trials: How do we work together? *Frontiers in Sociology*, 4, 21.
- Halvorsrud, K., Nazroo, J., Otis, M., Brown Hajdukova, E., & Bhui, K. (2018). Ethnic inequalities and pathways to care in psychosis in England: A systematic review and meta-analysis. *BMC Medicine*, 16(1), 223–317.
- Herth, K. (1992). Abbreviated instrument to measure hope: Development and psychometric evaluation. *Journal of Advanced Nursing*, 17(10), 1251–1259.
- Hope, E., & Ali, S. (2019). Peer support, intersectionality and socially excluded communities. In E. Watson & S. Meddings (Eds.), *Peer support in mental health* (pp. 69–82). Red Globe Press.
- Johnson, S., Lamb, D., Marston, L., Osborn, D., Mason, O., Henderson, C., Ambler, G., Milton, A., Davidson, M., Christoforou, M., Sullivan, S., Hunter, R., Hindle, D., Paterson, B., Leverton, M., Piotrowski, J., Forsyth, R., Mosse, L., Goater, N. . . , Lloyd-Evans, B. (2018). Peer-supported self-management for people discharged from a mental health crisis team: A randomised controlled trial. *Lancet*, 392(10145), 409–418.
- Mahlke, C. I., Priebe, S., Heumann, K., Daubmann, A., Wegscheider, K., & Bock, T. (2017). Effectiveness of one-to-one peer support for patients with severe mental illness - a randomised controlled trial. *European Psychiatry*, 42, 103–110.
- Marks, J., Foster, R., Gibson, S. L., Simpson, A., Rinaldi, M., Repper, J., Worner, J., Patel, S., Lucock, M., Ussher, M., White, S., Goldsmith, L., Barlow, S., & Gillard, S. (2021). Development of a peer support intervention to improve the experience and outcomes of discharge from inpatient mental health care: The role of experiential knowledge in a coproduced approach. *BMC Research Notes*, 14(1), 1–10.
- McDermott, R. C., Cheng, H. L., Wong, J., Booth, N., Jones, Z., & Sevig, T. (2017). Hope for help-seeking: A positive psychology perspective of psychological help-seeking intentions. *The Counseling Psychologist*, 45(2), 237–265.
- O'Connell, M. J., Sledge, W. H., Staeheli, M., Sells, D., Costa, M., Wieland, M., & Davidson, L. (2018). Outcomes of a peer mentor intervention for persons with recurrent psychiatric hospitalization. *Psychiatric Services*, 69(7), 760–767.
- Overall, J. E., & Gorham, D. R. (1962). The brief psychiatric rating scale. *Psychological Reports*, 10(3), 799–812.
- Pfeiffer, P. N., Heisler, M., Piette, J. D., Rogers, M. A., & Valenstein, M. (2011). Efficacy of peer support interventions for depression: A meta-analysis. *General Hospital Psychiatry*, 33(1), 29–36.
- Priebe, S., & McCabe, R. (2008). Therapeutic relationships in psychiatry: The basis of therapy or therapy in itself? *International Review of Psychiatry*, 20(6), 521–526.
- Priebe, S., Watzke, S., Hansson, L., & Burns, T. (2008). Objective social outcomes index (SIX): A method to summarise objective indicators of social outcomes in mental health care. *Acta Psychiatrica Scandinavica*, 118(1), 57–63.
- R Core Team. (2021). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Rogers, E. S., Maru, M., Johnson, G., Cohee, J., Hinkel, J., & Hashemi, L. (2016). A randomized trial of individual peer support for adults with psychiatric disabilities undergoing civil commitment. *Psychiatric Rehabilitation Journal*, 39(3), 248–255.
- Salway, T., Ferlatte, O., Purdie, A., Shoveller, J., Trussler, T., & Gilbert, M. (2018). Healthcare engagement among gay

- and bisexual men with recent suicide ideation or attempts. *American Journal of Orthopsychiatry*, 88(6), 713–722.
- Simpson, A., Quigley, J., Henry, S. J., & Hall, C. (2014). Evaluating the selection, training, and support of peer support workers in the United Kingdom. *Journal of Psychosocial Nursing and Mental Health Services*, 52(1), 31–40.
- Tse, S., Tsoi, E. W. S., Wong, S., Kan, A., & Kwok, C. F. Y. (2014). Training of mental health peer support workers in a non-western high-income city: Preliminary evaluation and experience. *International Journal of Social Psychiatry*, 60(3), 211–218.
- Walker, G., & Bryant, W. (2013). Peer support in adult mental health services: A meta-synthesis of qualitative findings. *Psychiatric Rehabilitation Journal*, 36(1), 28–34.
- White, S., Foster, R., Marks, J., Morshead, R., Goldsmith, L., Barlow, S., Sin, J., & Gillard, S. (2020). The effectiveness of one-to-one peer support in mental health services: A systematic review and meta-analysis. *BMC Psychiatry*, 20(1), 534–620.